


MIT OpenCourseWare
<http://ocw.mit.edu>

18.01 Single Variable Calculus
Fall 2006

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

18.01 Practice Questions for Exam 2 Solutions. Fall 2006

1) $f(x) = 3x^5 - 5x^3 + 1$ $f'(x) = 0$ $x = 0, \pm 1$
 $f'(x) = 15x^4 - 15x^2$ $f''(x) = 0$ $x = 0, \pm \sqrt{2}/2$
 $f''(x) = 60x^3 - 30x$ $f(x) \rightarrow -\infty$ $x \rightarrow -\infty$
 $f(x) \rightarrow +\infty$ $x \rightarrow +\infty$





x	f(x)	f'(x)	f''(x)	
-2	-55	< 0	< 0	
-1	3	0	-30	la. max.
$-\sqrt{2}/2$	$1 + \frac{7\sqrt{2}}{8}$	> 0	0	inflection
0	1	0	0	inflection
$\sqrt{2}/2$	$1 - \frac{7\sqrt{2}}{8}$	> 0	0	inflection
1	-1	0	30	la. min.
2	57	> 0	> 0	

There is an x , $-2 < x < -1$, since $f(-2) < 0$ and $f(-1) > 0$.
 There is an x , $1 < x < 2$, since $f(1) < 0$, $f(2) > 0$.
 There is an x , $0 < x < 1$, since $f(0) > 0$, $f(1) < 0$.

2) $f(x) = 4x^2 - \frac{1}{x}$ $f(x) = 0$, $x = \sqrt[3]{4}$
 $f'(x) = 8x + \frac{1}{x^2}$ $f'(x) = 0$, $x = -1/2$
 $f''(x) = 8 - \frac{2}{x^3}$ $f''(x) = 0$, $x = \sqrt[3]{4}$
 as $x \rightarrow \infty$, $f(x) \rightarrow \infty$ asymptote at
 as $x \rightarrow -\infty$, $f(x) \rightarrow \infty$ $x = 0$

$f''(-1/2) > 0$




$A = \frac{1}{2} x_{int} y_{int} = 2 - \frac{2}{m} - \frac{m}{2}$
 $-\infty < m < 0$

$\frac{dA}{dm} = -\frac{1}{2} + \frac{2}{m^2}$

$\frac{dA}{dm} = 0$ at $m = -2$
 So $m = -2$, $A = 4$ is a local min.

$\left. \frac{d^2A}{dm^2} \right|_{m=-2} = \frac{4}{m^3} \Big|_{m=-2} = \frac{1}{2} > 0$

$A \rightarrow \infty$ as $m \rightarrow 0$ or as $m \rightarrow -\infty$. So $m = -2$, $A = 4$ is the global min.



$L = 2L + 4 - x = 2\sqrt{4+x^2} + 4 - x$


$L(\sqrt{3}) \approx 9.2 < 10$

$0 \leq x \leq 4$ $\frac{dL}{dx} = \frac{2x}{\sqrt{4+x^2}} - 1$

$L(0) = 10$
 $L(4) = 10$
 $\frac{dL}{dx} = 0$ at $x = \sqrt{3}$

$L(\sqrt{3}) = 3\sqrt{3} + 4 \approx 9.2$

Since L at the endpoints is larger than at the unique interior crit. pt., this unique crit. pt. is a min.

5)  $\cos \theta = \frac{x}{200}$
 $-\sin \theta \frac{d\theta}{dt} = \frac{1}{200} \frac{dx}{dt}$
 $\frac{dx}{dt} \Big|_{\theta=\pi/6} = 50$ $\frac{d\theta}{dt} \Big|_{\theta=\pi/6} = \frac{1}{200} \cdot 50 \cdot (-2) = -\frac{1}{2} \frac{\text{rad}}{\text{sec}}$

7) $f(x) = e^{-2x} (1+2\sin x)^{-1}$
 $\approx (1-2x)(1+2x)^{-1}$
 $\approx (1-2x)(1-2x)$

$= 1 - (2+2)x + 2\lambda x^2$
 So f is const. to first order if

$\lambda = -2$

To estimate

$f(x) = e^{2x} (1+2\sin x)^{-1}$

we use 2nd order approx.


$f(x) = e^{2x} (1+2\sin x)^{-1}$
 $\approx (1+2x+\frac{(2x)^2}{2})(1+2x)^{-1}$

$\approx (1+2x+2x^2)(1-2x+2x^2)$

Think

$= 1+2x^2 + \dots$

So $f(.1) \approx 1+2(.1)^2 = 1.02$

6)  $r=h$ since this is a right circular cone.
 $V = \frac{1}{3} \pi r^2 h = \frac{1}{3} \pi h^3$
 $3 = \frac{dV}{dt} = \pi h^2 \frac{dh}{dt}$ $\frac{dh}{dt} \Big|_{h=2} = \frac{3}{4\pi}$

8) Assume the rate of evaporation is proportional to the surface area, i.e. $\frac{dV}{dt} = C \pi r^2 = C \pi h^2$ (C is some negative constant).

$C \pi h^2 = \frac{dV}{dt} = \pi h^2 \frac{dh}{dt}$ So $\frac{dh}{dt} = \text{const}$

9) From M.V.T.: for any $a < b$ there is a c $a < c < b$ s.t.

a) $f(b) - f(a) = f'(c)(b-a) > 0$

since $f'(c) > 0$, $b-a > 0$.

So $f(b) > f(a)$, i.e. f is increasing.

b) For $x > 0$ $\frac{e^x - e^0}{x - 0} = \frac{d}{dx}(e^x) \Big|_{x=c} = e^c$

$0 < c < x$

or $e^x = 1 + e^c x$. For $0 < c, e^c > 1$,

so $e^x = 1 + e^c x > 1 + x$.

10) $\frac{dy}{dx} = x(y^2+1)$ $\frac{dy}{y^2+1} = x dx$ $\tan^{-1} y = \frac{x^2}{2} + C$

$y = \tan(\frac{x^2}{2} + C)$ $1 = \tan C, C = \pi/4$

$y = \tan(\frac{x^2}{2} + \pi/4)$.

9) a) $\int \frac{dx}{(3x+2)^2} = \frac{1}{3} \int \frac{du}{u^2} = -\frac{1}{3} \frac{1}{u} + C$
 $u = 3x+2$ $du = 3dx$ $= -\frac{1}{3} \frac{1}{3x+2} + C$

b) $\int \sin(2x) \sin x dx = \int 2 \sin^2 x \cos x dx$ $u = \sin x$ $du = \cos x dx$
 $= 2 \int u^2 du = \frac{2}{3} u^3 + C = \frac{2}{3} \sin^3 x + C$

c) $\int \frac{(\ln x)^2}{x} dx = \int u^2 du = \frac{1}{3} u^3 + C$
 $u = \ln x$ $du = \frac{dx}{x}$ $= \frac{1}{3} (\ln x)^3 + C$

11) $\frac{dT}{dt} = k(T-T_e)$ $\frac{dT}{T-T_e} = k dt$ $\ln(T-T_e) = kt + C$

$T = T_e + A e^{kt}$ (let $A = e^C$)
 $20 = T(0) = 100 + A$ so $A = -80$
 $30 = T(5) = 100 - 80 e^{5k}$ $k = \frac{1}{5} \ln \frac{7}{6}$

$60 = T(t_{\text{done}})$
 $= 100 - 80 e^{kt_{\text{done}}}$
 $t_{\text{done}} = \frac{5 \ln \frac{2}{3}}{\ln \frac{7}{6}} \approx 26$