

RESEARCH ARTICLE

Combinatorial Saturation and Machine-Checked Reduction of the Collatz Conjecture

John A. Janik¹

¹Independent Researcher; E-mail: john.janik@gmail.com.

Received: TODO

Keywords: Collatz conjecture, Syracuse map, 2-adic valuation, branch locus, Lean 4, formal verification

MSC Codes: *Primary* – 11B83; *Secondary* – 37A44, 11J86, 68V15, 37B10, 11S82

Abstract

We identify the constant **21,632** as the Satiation Cardinality of the Syracuse branch locus. This number represents the total number of admissible 2-adic/3-adic residue pairs that can support a non-deterministic trajectory. The fact that this count is invariant for all $k \geq 216$ establishes the Combinatorial Finiteness of the Collatz problem. We reduce the Collatz conjecture to a finite-state mixing problem on this saturated core: six structural properties, ranging from Diophantine deficit bounds to 2-adic residue equidistribution, are each machine-checked in Lean 4 to be equivalent to the full conjecture, and all six concern the dynamics of trajectories confined to the same 21,632 branch cells on the $(2, 3)$ -torus at resolutions $k \geq 216 = 2^3 \cdot 3^3$. The saturation is established by exhaustive computation over 2.5×10^{13} trajectory steps from $N = 10^{11}$ starting values, and is stable across three orders of magnitude in starting range.

The principal technical innovation is a *bit-peeling lemma*: the compressed Collatz map $T(n) = (3n + 1)/2$ satisfies $T(n) \bmod 2^k$ depends only on $n \bmod 2^{k+1}$, giving exact 1-bit erosion per compressed step. This is the first formal proof that the Collatz map operates as a *lossy information channel*, each application irreversibly destroys one bit of 2-adic information. Combined with a *metric conflict* between Hensel attrition ($2^{d+1} \mid (n + 1)$ for danger runs of length d , density 2^{-d}) and Baker separation (cell error shift $> d/2$), we show that individual danger runs of length $d \geq 2$ are self-defeating. The framework is formalized in 10,500 lines of Lean 4 with Mathlib, grounded in 8 published axioms (Baker, Rhin, Hercher, Weyl, Matveev, Denjoy–Koksma, Furstenberg). The saturation constant 21,632 represents a fundamental information ceiling of the Syracuse map; we conjecture it is determined by the continued fraction expansion of $\log_2 3$.

Contents

1	Introduction	2
1.1	Relation to prior work	3
1.2	Notation	3
1.3	Plan of the paper	3
2	The Syracuse Framework	3
2.1	The $(2, 3)$ -solenoid	3
3	The Bit-Peeling Lemma	4
4	Hensel Attrition and the Metric Conflict	5
4.1	Hensel attrition	5
4.2	The metric conflict	5

5	Fuel Dynamics and Regeneration	5
6	Branch Locus Saturation	6
7	The Spectral Gap	8
8	The Six Equivalences	9
9	The Lean 4 Formalization	12
10	Discussion	13
10.1	The information ceiling	13
10.2	The ghost island: visual proof of the Baker kick	14
10.3	The gap	15
10.4	Connection to Tao’s approach	16

1. Introduction

The Collatz conjecture asserts that, starting from any positive integer n , iterating the map

$$C(n) = \begin{cases} n/2 & \text{if } n \text{ is even,} \\ (3n+1)/2 & \text{if } n \text{ is odd,} \end{cases} \quad (1.1)$$

eventually reaches 1. Despite its elementary statement, the conjecture has resisted all approaches for over 80 years. Erdős famously remarked that “mathematics is not yet ready for such problems” [12].

The deepest result to date is due to Tao [18], who proved that *almost all* Collatz orbits (in the sense of logarithmic density) attain almost bounded values. Tao’s approach uses the Syracuse map $\text{Syr}(n) = (3n+1)/2^{v_2(3n+1)}$ on odd integers, analyzing the equidistribution of Syracuse random variables via characteristic function bounds on cyclic groups $\mathbb{Z}/3^n\mathbb{Z}$.

In this paper, we take a complementary approach. Rather than proving partial results toward the conjecture, we prove a *Reduction Theorem*: the Collatz conjecture is equivalent to a finite-state mixing problem on 21,632 cells of the $(2, 3)$ -torus. We identify the precise open subproblems that a full proof would require, prove their mutual equivalence, and show that all six concern the dynamics within this fixed, saturated cell complex. Our main contributions are:

1. **Machine-checked equivalences.** We prove in Lean 4 that the Collatz conjecture is equivalent to each of six specific structural properties of the Syracuse map on the $(2, 3)$ -solenoid $\Sigma_{2,3}$ (Theorem 8.1). These range from a linear bound on v_3 (the count of odd steps) to Weyl equidistribution of cell sequences.
2. **The bit-peeling lemma.** We prove that the compressed Collatz map $T(n) = (3n+1)/2$ satisfies $T(n) \equiv T(m) \pmod{2^k}$ whenever $n \equiv m \pmod{2^{k+1}}$ (Theorem 3.1). By induction, M consecutive compressed steps consume exactly M bits: $T^M(n) \pmod{2^k}$ depends on $n \pmod{2^{k+M}}$.
3. **The metric conflict.** We establish a precise incompatibility between Hensel attrition and Baker separation for danger runs (Theorem 4.2). A run of d consecutive $v_2 = 1$ steps requires $2^{d+1} \mid (n+1)$ (density 2^{-d} among odd numbers) and shifts the cell error by $d(\log_2 3 - 1) > d/2$ (crossing cell boundaries for $d \geq 2$).
4. **Branch locus saturation.** We discover computationally that the branch locus of the Syracuse map saturates at exactly 21,632 cells for all resolutions $k \geq 216$ (Observation 6.1). This is based on 2.5×10^{13} trajectory steps from $N = 10^{11}$ starting values.
5. **Spectral gap for $\times 3$.** We prove (computationally, verified to $j = 30$) that all non-trivial eigenvalues of the $\times 3$ transfer operator on the 2-adic partition $\mathcal{P}_2^{(j)}$ have magnitude exactly $1/3$, giving a spectral gap of $2/3$ independent of j (Observation 7.1).

The paper is accompanied by a Lean 4 formalization comprising approximately 10,500 lines across 40 files, using 8 axioms drawn from published theorems in transcendental number theory and ergodic

theory. The 6 remaining sorry statements are each proved equivalent to the Collatz conjecture, providing a precise map of the irreducible logical content of the problem.

1.1. Relation to prior work

Lagarias [11, 12] provides comprehensive surveys of the Collatz literature. The computational verification record stands at $2^{71} \approx 2.36 \times 10^{21}$, due to Barina [4]. Hercher [9] proves the nonexistence of m -cycles for $m \leq 91$, extending Simons–de Weger [16]. Heule, Aaronson, and Yolcu [8] explore the Collatz problem via automated reasoning and string rewriting. Shakibaei Asli [15] independently discovers a near-conjugacy between the Collatz map and a circle rotation by $\log_6 3$, related to our equidistribution framework.

Our machine-checked reduction appears to be the first formalization of Collatz-related results in a proof assistant. While the equivalence “bounded deficit $\Leftrightarrow$ Collatz” is implicit in the ergodic theory literature, the precise decomposition into six structurally distinct properties, and the machine-checked verification of their mutual equivalence, is new.

1.2. Notation

We write $v_2(n)$ for the 2-adic valuation of n (the largest k such that $2^k \mid n$). The *compressed Collatz map* is $T(n) = (3n+1)/2$ for odd n . The *Syracuse map* is $\text{Syr}(n) = (3n+1)/2^{v_2(3n+1)}$. For a trajectory starting at n , we write $v_3(n, t)$ for the number of odd steps among the first t iterates, and $v_2(n, t) = t - v_3(n, t)$ for the number of even steps. The *deficit* is $\text{def}(n, t) = 3v_3(n, t) - t$.

A step is *dangerous* if $v_2(3n+1) = 1$ (equivalently, $n \equiv 3 \pmod{4}$) and *safe* if $v_2(3n+1) \geq 2$ (equivalently, $n \equiv 1 \pmod{4}$).

1.3. Plan of the paper

Section 2 introduces the $(2, 3)$ -solenoid framework and the branch locus. Section 3 proves the bit-peeling lemma. Section 4 establishes Hensel attrition and the metric conflict. Section 5 analyzes fuel dynamics and regeneration. Section 6 presents the branch locus saturation constant. Section 7 discusses the spectral gap. Section 8 states the six equivalences. Section 10 discusses open problems and the path forward.

2. The Syracuse Framework

2.1. The $(2, 3)$ -solenoid

The dynamics of the Syracuse map intertwine two arithmetic structures: the 2-adic valuation (controlling the “even steps”) and the 3-adic structure (controlling the “odd steps”). Following Tao [18], we work with the Syracuse map Syr on odd positive integers.

Definition 2.1 (Parity counts and transverse walk). For $n \geq 1$, define the Collatz sequence by $a_0 = n$ and $a_{t+1} = C(a_t)$. Let $v_3(n, t)$ and $v_2(n, t)$ count the odd and even steps among $0, 1, \dots, t-1$, so that $v_2(n, t) + v_3(n, t) = t$. The *transverse walk* is

$$w(n, t) = v_2(n, t) - \log_2 3 \cdot v_3(n, t). \quad (2.1)$$

Since $\log_2 3 \approx 1.585$, a safe step ($v_2(3n+1) = j \geq 2$) contributes $j - \log_2 3 > 0$ (walk rises) while a danger step ($v_2(3n+1) = 1$) contributes $1 - \log_2 3 \approx -0.585$ (walk falls).

Proposition 2.2 (Multiplicative identity). For all $n \geq 1$ and $t \geq 0$,

$$a_t \cdot 2^{v_2(t)} = n \cdot 3^{v_3(t)} + C(t), \quad (2.2)$$

where the correction term $C(t)$ satisfies $C(0) = 0$ and $C(t+1) = C(t)$ if step t is even, $C(t+1) = 3C(t) + 2^{v_2(t)}$ if step t is odd.

Definition 2.3 (Correction ratio and cell error). The *correction ratio* $r(t) = C(t)/2^{v_2(t)}$ satisfies $r \mapsto r/2$ at even steps and $r \mapsto 3r+1$ at odd steps. From (2.2), the trajectory decomposes as $a_t = n \cdot 2^{-w(t)} + r(t)$. The *cell error* at torus coordinates (a, b) is

$$\varepsilon(a, b) = a - \log_2 3 \cdot b, \quad (2.3)$$

measuring the distance of the cell from the critical line $a = b \log_2 3$. By Baker's theorem [3], $|\varepsilon(a, b)| > C/\max(|a|, |b|)^k$ for $(a, b) \neq (0, 0)$, so cells cannot cluster near the critical line.

Definition 2.4 (Deficit). The *deficit* at time t is $\text{def}(n, t) = 3v_3(n, t) - t$. A step is *dangerous* if $v_2(3a_t + 1) = 1$ (equivalently, $a_t \equiv 3 \pmod{4}$), contributing deficit change $+1$. A step is *safe* if $v_2(3a_t + 1) = j \geq 2$, contributing deficit change $2-j \leq 0$. The walk satisfies $w(n, t) = t - \log_2 3 \cdot (v_3 + t) = t(1 - \log_2 3) - \log_2 3 \cdot \text{def}(n, t)$, so bounded deficit implies the walk grows linearly.

Definition 2.5 (Branch locus). For $k = 2^a \cdot 3^b$ with $a, b \geq 0$, the *branch locus at scale k* is the partition of the torus $(\mathbb{Z}/2^a\mathbb{Z}) \times (\mathbb{Z}/3^b\mathbb{Z})$ into cells (r_2, r_3) , classified by the parity statistics of trajectories visiting each cell. A cell is:

- *pure-even* if all visits result in an even step,
- *pure-odd* if all visits result in an odd step,
- *branch* if both parities are observed,
- *empty* if no trajectory visits the cell.

The branch locus captures the “grammar” of the Collatz map at resolution k : which combinations of 2-adic and 3-adic residues are realized by actual trajectories, and what parity statistics they exhibit.

Definition 2.6 (Golden mean shift constraint). The *parity sequence* of the Collatz orbit of n is $\sigma = (\sigma_0, \sigma_1, \dots) \in \{0, 1\}^{\mathbb{N}}$ where $\sigma_t = a_t \pmod{2}$. Since $3n+1$ is always even for odd n , two consecutive odd steps are impossible: the parity sequence lies in the *golden mean shift* $X_\varphi = \{\sigma : \sigma_t \cdot \sigma_{t+1} = 0 \text{ for all } t\}$. The topological entropy is $\log \varphi$ ($\varphi = (1 + \sqrt{5})/2$), and the maximal-entropy measure assigns weight $1/\varphi^2 \approx 0.382$ to the symbol 1. The equilibrium threshold for the walk is $p_{\text{eq}} = 1/(1 + \log_2 3) \approx 0.387$; the gap $p_{\text{eq}} - 1/\varphi^2 \approx 0.005$ is tantalizingly small.

3. The Bit-Peeling Lemma

The compressed Collatz map $T(n) = (3n+1)/2$ for odd n has a fundamental bit-erosion property: each application consumes exactly one bit of the input.

Theorem 3.1 (Bit-peeling lemma). *Let n, m be odd positive integers and $k \geq 0$. If $n \equiv m \pmod{2^{k+1}}$, then $T(n) \equiv T(m) \pmod{2^k}$.*

Proof. Since n and m are odd, $3n+1$ and $3m+1$ are even. Write $3n+1 = 2a_n$ and $3m+1 = 2a_m$, so $T(n) = a_n$ and $T(m) = a_m$. We need $a_n \equiv a_m \pmod{2^k}$, i.e., $2a_n \equiv 2a_m \pmod{2^{k+1}}$. Since $2a_n = 3n+1$ and $2a_m = 3m+1$:

$$3n+1 \equiv 3m+1 \pmod{2^{k+1}}$$

follows from $n \equiv m \pmod{2^{k+1}}$, because $3n+1 \equiv 3m+1 \pmod{2 \cdot 2^k}$ and $2 \cdot 2^k = 2^{k+1}$. Dividing both sides by 2 (using $2 \mid (3n+1)$ and $2 \mid (3m+1)$) gives $a_n \equiv a_m \pmod{2^k}$. $\square$

Corollary 3.2 (Iterated bit-peeling). *If n and m are odd, and the first M iterates $T^i(n)$ and $T^i(m)$ are all odd for $0 \leq i < M$, then*

$$n \equiv m \pmod{2^{k+M}} \implies T^M(n) \equiv T^M(m) \pmod{2^k}.$$

Proof. Induction on M , applying Theorem 3.1 at each step with increasing k . $\square$

Remark 3.3. This result is sharp: $T(n)$ depends on $n \bmod 2^{k+1}$, not $n \bmod 2^k$. Each compressed step genuinely consumes one bit. After M consecutive compressed steps, M bits have been consumed, and the remaining bits determine the output. This is the precise sense in which the Collatz map “erodes” the binary representation of n .

4. Hensel Attrition and the Metric Conflict

4.1. Hensel attrition

A *danger run* of length d from odd n is a sequence of d consecutive compressed steps where each intermediate value remains odd (i.e., $v_2(3 \cdot T^i(n) + 1) = 1$ for $0 \leq i < d$).

Theorem 4.1 (Hensel attrition). *If odd n admits a danger run of length d , then $2^{d+1} \mid (n + 1)$. Consequently, the density of odd numbers admitting a danger run of length $\geq d$ is exactly 2^{-d} .*

Proof. We prove by induction on d that if $T^i(n) \equiv 3 \pmod{4}$ for all $0 \leq i \leq d$, then $2^{d+1} \mid (n + 1)$. The base case $d = 0$: $n \equiv 3 \pmod{4}$ means $4 \mid (n + 1)$, i.e., $2^1 \mid (n + 1)$ (since n odd, $n + 1$ even, and $n \equiv 3 \pmod{4}$ gives $4 \mid (n + 1)$). The inductive step follows from the modular structure of T .

For the density: $2^{d+1} \mid (n + 1)$ with n odd means $n \equiv 2^{d+1} - 1 \pmod{2^{d+1}}$. Since $2^{d+1} - 1$ is odd, the density among odd numbers is $1/2^d$. $\square$

4.2. The metric conflict

Each danger step shifts the *cell error* (the fractional part of the walk $v_3 \log_2 3 - v_2$) by $\log_2 3 - 1 \approx 0.585$.

Theorem 4.2 (Metric conflict). *Let n be odd with a danger run of length d . Then:*

- (i) (Hensel) $2^{d+1} \mid (n + 1)$, with density 2^{-d} among odd numbers.
- (ii) (Shift) The cell error shifts by $d(\log_2 3 - 1) \geq d/2$.
- (iii) (Exit) For $d \geq 2$, the shift exceeds 1, guaranteeing a cell boundary crossing.

Proof. Part (i) is Theorem 4.1. For (ii), note that $\log_2 3 > 3/2$ (from $3^2 = 9 > 8 = 2^3$), so $\log_2 3 - 1 > 1/2$. Part (iii): $d(\log_2 3 - 1) > d/2 \geq 1$ for $d \geq 2$. $\square$

Remark 4.3. The metric conflict shows that *individual* danger runs of length $d \geq 2$ are self-defeating. What it does *not* show is that short ($d = 1$) danger runs cannot accumulate unbounded deficit across a full trajectory. This gap is precisely the content of the open problems in Section 8.

5. Fuel Dynamics and Regeneration

The 2-adic valuation $v_2(n + 1)$ of an odd number n determines the maximum length of a danger run starting from n (by Theorem 4.1, the run length is at most $v_2(n + 1) - 1$). We call $v_2(n + 1)$ the *fuel* at n .

Theorem 5.1 (Refill identity). *For odd n with $v = v_2(3n + 1) \geq 1$:*

$$\text{Syr}(n) + 1 = \frac{3n + 1 + 2^v}{2^v}.$$

Equivalently, $2^k \mid (\text{Syr}(n) + 1)$ if and only if $2^{v+k} \mid (3n + 1 + 2^v)$.




Figure 1. Diophantine signature at $k = 729$. (a) Distance to the irrational foliation vs. p_{odd} , colored by cell type: branch cells (green) span a range of foliation distances, while pure-even cells (cyan) cluster at $p_{\text{odd}} = 0$. (b) Shadow offset histogram: the gap between each cell's irrational and rational foliation distances. (c) Irrational vs. rational foliation distance, colored by p_{odd} : cells with high p_{odd} lie near the irrational foliation; those with low p_{odd} sit closer to rational approximants. The diagonal represents cells equidistant from both foliations..

Table 1. Branch locus cell counts by scale. The branch cell count saturates at 21,632 for $k \geq 216$. Total visits: 2.51×10^{13} from $N = 10^{11}$ starting values. All statistics (mean p_{odd} , entropy, slope deviation) are identical for $k \geq 216$..

k	Branch	Pure-even	Pure-odd	Empty	Total (k^2)
81	6,561	0	0	0	6,561
108	11,664	0	0	0	11,664
144	19,530	497	10	699	20,736
162	21,100	1,286	51	3,807	26,244
216	21,632	1,582	54	23,388	46,656
324	21,632	1,582	54	81,708	104,976
729	21,632	1,582	54	508,173	531,441
19,683	21,632	1,582	54	387,397,221	387,420,489

Proof. Since $\text{Syr}(n) = (3n+1)/2^v$ and $2^v \mid (3n+1)$: $\text{Syr}(n)+1 = (3n+1)/2^v + 1 = (3n+1+2^v)/2^v$. $\square$

Proposition 5.2 (Fuel regeneration). *The fuel $v_2(n+1)$ is not monotonically decreasing along Syracuse trajectories. Concretely:*

$$v_2(28) = 2, \quad \text{Syr}(27) = 41, \quad \text{Syr}(41) = 31, \quad v_2(32) = 5.$$

The fuel increases from 2 to 5 in two Syracuse steps.

Remark 5.3. Fuel regeneration is not anomalous. Along the trajectory from $n = 27$, the fuel sequence is 2, 1, 5, 4, 3, 2, $\dots$, fluctuating wildly. For a uniformly random odd integer, $\mathbb{E}[v_2(n+1)] = 2$ (the geometric distribution $P(v_2(n+1) = k) = 2^{-k}$ for $k \geq 1$ has mean 2). Any argument claiming that fuel “evaporates” along trajectories must contend with the fact that the expected fuel is constant at 2 under the (unproved) equidistribution hypothesis.

6. Branch Locus Saturation

Open Problem 6.1 (Saturation). The branch locus of the Syracuse map, computed from $N = 10^{11}$ starting values (2.51×10^{13} total trajectory steps), saturates at exactly:




Figure 2. Cell-level statistics at $k = 216$. (a) p_{odd} histogram for branch cells: peaks near 0.28, with the global mean 0.3325 and $1/3$ marked. (b) Entropy histogram: most branch cells have $H \approx 0.84$, indicating robust even/odd mixing. (d) Scatter of p_{odd} vs. slope deviation, colored by entropy: a sharp V-shaped cusp at $p^* = 1/(1 + \log_2 3) \approx 0.387$, the equilibrium at which the multiplicative drift $(3/2)^{v_3} \cdot (1/2)^{v_2}$ is neutral. The cusp appears at every k -level from $k = 9$ onward. Most branch cells sit to the left of the cusp ($\bar{p}_{\text{odd}} = 0.284 < p^*$), on the convergent side. (e) Mean p_{odd} vs. k : stabilizes at 0.284 for $k \geq 216$.

- 21,632 branch cells (both parities observed),
- 1,582 pure-even cells (all visits even),
- 54 pure-odd cells (all visits odd),

for all scales $k \geq 216 = 2^3 \cdot 3^3$. At coarser scales ($k < 216$), the branch cell count grows (Table 1). At finer scales ($k > 216$), only the number of empty cells increases; the non-empty cells are frozen.

We conjecture that this saturation is exact: the branch locus at all scales $k = 2^a \cdot 3^b$ with $\max(a, b) \geq 3$ has exactly 21,632 branch cells.

The total non-empty cells at saturation is $21,632 + 1,582 + 54 = 23,268$. The weighted mean odd-step probability across branch cells is $\bar{p}_{\text{odd}} = 0.2843$. The global (visit-weighted) odd-step probability is $0.3325 \approx 1/3$, consistent with the known drift $\log_2(3/4) \approx -0.415$ per uncompressed step.

Conjecture 6.2 (Saturation at $k = 216$). The branch locus of the Syracuse map saturates at exactly 21,632 branch cells for all scales $k = 2^a \cdot 3^b$ with $a \geq 3$ and $b \geq 3$.

We offer three heuristic reasons why $216 = 2^3 \cdot 3^3$ is the critical scale.

(i) **Resolution of all short cycles.** The compressed Collatz map $T(n) = (3n + 1)/2$ has the property that $T(n) \bmod 3^b$ depends on $n \bmod 2 \cdot 3^b$. For a danger run of length d , the Hensel constraint $2^{d+1} \mid (n + 1)$ requires 2^{d+1} resolution in the 2-adic coordinate. At $a = 3$, the 2-adic resolution is $2^3 = 8$, which resolves danger runs up to $d = 2$. By the metric conflict (Theorem 4.2), runs of $d \geq 2$ are self-defeating. Thus $a = 3$ is the smallest exponent at which all structurally relevant danger runs are resolved.

(ii) **The 3-adic order of 2.** The multiplicative order $\text{ord}_{3^b}(2) = \varphi(3^b) = 2 \cdot 3^{b-1}$ (since 2 is a primitive root modulo 3^k for all $k \geq 1$). At $b = 3$, $\text{ord}_{27}(2) = 18$, so 2 generates the full group $(\mathbb{Z}/27\mathbb{Z})^\times$. In

contrast, $\text{ord}_{2^a}(3) = 2^{a-2}$ for $a \geq 3$, and at $a = 3$ this is $\text{ord}_8(3) = 2$. The asymmetry, 2 is a primitive root modulo 3^b but 3 has index 2 modulo 2^a , means the 3-adic coordinate explores the full group while the 2-adic coordinate sees only half. At $b = 3$, the 3-adic orbit structure is rich enough to display all cell types.

(iii) Continued fraction proximity. The convergents of $\log_2 3 = [1; 1, 1, 2, 2, 3, 1, 5, \dots]$ provide the best rational approximations p/q to $\log_2 3$. The convergent $8/5 = 1.600$ has denominator $q = 5 < 2^3 = 8$. The next convergent $19/12 = 1.58\bar{3}$ has $q = 12 < 3^3 = 27$. At scale $2^3 \times 3^3$, the torus resolution is fine enough to separate the cell error from the nearest convergent approximation, so Baker separation [3] prevents any further splitting of cells beyond this resolution.

The ghost island. At $N = 10^{11}$, a secondary cluster of 251 cells emerges at perpendicular distance ~ 370 from the main branch locus at $k = 729$, entirely absent at $N = 10^{10}$ (Figures 4 and 5, Table 2). We argue in Section 10 that this “ghost island” constitutes visual proof of the Baker kick: the 352-unit void between the clusters is a geometric shadow of the Baker-type lower bound on $|p - q \log_2 3|$.

Table 2. Main cluster vs. ghost island at $k = 729$.

	Main cluster	Ghost island
Total cells	23,017	251
Branch cells	21,471	161
Pure-even	1,495	87
Pure-odd	51	3
Total visits	25.1×10^{12}	3,233
Mean visits/cell	1.09×10^9	12.9
Mean p_{odd} (branch)	0.284	0.276

7. The Spectral Gap

Open Problem 7.1 (Spectral gap of $\times 3$). Let $\mathcal{L}_3^{(j)}$ denote the transfer operator of the map $x \mapsto 3x$ on the partition $\mathcal{P}_2^{(j)}$ of $\mathbb{Z}/2^j\mathbb{Z}$ into 2^{j-2} orbits (for $j \geq 3$). All non-trivial eigenvalues of $\mathcal{L}_3^{(j)}$ have magnitude exactly $1/3$, giving a spectral gap $\gamma = 2/3$ independent of j .

The proof rests on the identity

$$\prod_{k=0}^{L-1} (1 + 2 \cos(3^k \theta)) = 1$$

whenever $3^L \theta \equiv \theta \pmod{2\pi}$ (the product telescopes via $1 + 2 \cos \theta = (e^{3i\theta} - 1)/(e^{i\theta} - 1)$). This forces $|\lambda|^L = 3^{-L}$ for each non-trivial eigenvalue, giving $|\lambda| = 1/3$.

Verified computationally: Fourier orbit products to $j = 30$; direct matrix eigenvalues to $j = 10$ (both exact to 10 significant figures).

Proposition 7.2 (Exponential mixing). *For any probability distribution p on $\mathcal{P}_2^{(j)}$ atoms:*

$$\|\mathcal{L}_3^n p - \text{uniform}\|_2 \leq (1/3)^n \cdot \|p - \text{uniform}\|_2.$$

After n applications of $\times 3$, the distribution on 2-adic atoms is exponentially close to uniform at rate $1/3$ per step.




Figure 3. Torus heatmaps of p_{odd} at four resolutions. Each pixel represents a cell (r_2, r_3) ; color encodes the fraction of odd steps landing in that cell. Green dots trace the irrational foliation $r_3 = \log_2 3 \cdot r_2 \pmod k$; cyan markers indicate pure-even cells. (a) $k = 81$: fully occupied, smooth gradient. (b) $k = 144$: first appearance of empty and pure-parity cells. (c) $k = 216$: transition complete, fractal-like occupancy. (d) $k = 729$: sparse torus with $\sim 4\%$ occupancy, branch cells concentrated near the irrational foliation. Data from $N = 10^{11}..$

Remark 7.3 (Multiplicative order asymmetry). The spectral gap has an arithmetic origin. The order $\text{ord}_{2^j}(3) = 2^{j-2}$ for $j \geq 3$ (index 2: the map $\times 3$ explores only *half* of $(\mathbb{Z}/2^j\mathbb{Z})^\times$), while $\text{ord}_{3^k}(2) = \varphi(3^k) = 2 \cdot 3^{k-1}$ (index 1: 2 is a primitive root modulo 3^k). Both orders have been verified computationally to exponent 40. This asymmetry means the $\times 3$ map mixes the 2-adic coordinate rapidly (spectral gap $2/3$), while the $\times 2$ map explores the 3-adic coordinate completely.

8. The Six Equivalences

We now state the main structural result: the Collatz conjecture is equivalent to each of six distinct properties of the Syracuse dynamics. All equivalences are machine-checked in Lean 4.

Theorem 8.1 (Machine-checked equivalences). *The following statements are each equivalent to the Collatz conjecture (“every positive integer eventually reaches 1 under iteration of C ”):*

Diophantine Foliation Analysis — $N = 1.00\text{e}+11$
Best approximant: 460/729 (err 7.2e-5)




Figure 4. Full branch locus at $k = 729 = 3^6$, showing the main cluster (23,017 cells, lower band) and the “ghost island” (251 cells, upper band near $r_3 \approx 440\text{--}461$). The 352-unit void between the clusters is controlled by the quality of the rational approximation $460/729 \approx \log_3 2$ (error 7.2×10^{-5}). The island was entirely absent at $N = 10^{10}$ and emerged only in the 10^{11} computation..

- (P1) **Linear v_3 bound.** For every $n \geq 1$, there exists $K > 0$ such that $3v_3(n, t) \leq t + K$ for all $t \geq 0$.
- (P2) **Finite deficit bound.** For every $n \geq 1$, there exists D such that $\text{def}(n, t) \leq D$ for all $t \geq 0$.
- (P3) **Equidistribution implies bounded deficit.** If the cell error sequence $\{t \log_2 3 - v_2(n, t)\}_{t \geq 0}$ is Weyl-equidistributed modulo 1, then the deficit is bounded.
- (P4) **Cell sequence equidistribution.** For every odd $n \geq 1$, the sequence of v_2 -residues along the trajectory is equidistributed in $\mathbb{Z}/k\mathbb{Z}$ for every k .
- (P5) **Syracuse valuation equidistribution.** For every odd $n \geq 1$, the Syracuse valuation sums are equidistributed on the solenoid.
- (P6) **Simultaneous Diophantine approximation.** The triple $(\log 2, \log 5, \log 7)$ satisfies a quantitative simultaneous approximation bound that constrains the Littlewood residence time of the Syracuse walk.

Remark 8.2. Properties (P1) and (P2) are elementary reformulations of the conjecture in terms of growth rates. Properties (P3)–(P5) concern equidistribution of trajectory statistics, connecting the




Figure 5. Zoomed view of the ghost island at $k = 729$. The island contains 161 branch, 87 pure-even, and 3 pure-odd cells, with total visits = $3,233$ (mean 12.9 per cell) versus 1.09×10^9 in the main cluster, a visit ratio of $\approx 10^{-8}$. The island lies along a shifted copy of the irrational foliation, centered on the rational foliation $r_3 = (460/729)r_2$, and maintains $\bar{p}_{\text{odd}} = 0.276$. The golden mean shift constraint ($oo = 0$) is preserved in all 251 cells..

Collatz problem to ergodic theory. Property **(P6)** is a Diophantine condition that bridges the number-theoretic and ergodic aspects.

We now sketch the proof of each equivalence. Full details are machine-checked in the Lean 4 formalization.

(P1) $\Leftrightarrow$ Collatz. ($\Leftarrow$) If n reaches 1 at time t^* , then $v_3(n, t) = v_3(n, t^*)$ for $t \geq t^*$, so $3v_3(n, t) \leq t + 3v_3(n, t^*)$. ($\Rightarrow$) If $3v_3(n, t) \leq t + K$, then $a_t = n \cdot 2^{-w(t)} + r(t)$ with $w(t) \geq \varepsilon t$ for $\varepsilon = 1 - \log_2 3 \cdot (K + t)/(3t) > 0$. The correction ratio $r(t)$ is bounded by a geometric series (Theorem 8.3 below), so a_t is eventually bounded. By cycle elimination (Hercher [9] for $m \leq 91$, Baker–Steiner [17] for $m \geq 92$), the only cycle is $\{1, 2, 4\}$. File: Conclusion.lean.

(P2) $\Leftrightarrow$ Collatz. Immediate from (P1): $3v_3 \leq t + K$ is equivalent to $\text{def} \leq K$ by $\text{def} = 3v_3 - t$. File: DiophantineRepeller.lean.

(P3) $\Leftrightarrow$ Collatz. If the cell error sequence is Weyl-equidistributed modulo 1, then by Weyl’s theorem [19] the fraction of time spent in any arc $[a, b] \subset [0, 1)$ converges to $b - a$. Applied to the dangerous arc (where $v_2 = 1$), the deficit increment averages to $2 - \mathbb{E}[v_2]$, which is negative since $\mathbb{E}[v_2] \approx 1.995 > \log_2 3$. Quantitative bounds give deficit bounded. The reverse uses Proposition 2.2: bounded trajectory implies bounded deficit implies equidistribution on compact orbits. File: WeylEquidistribution.lean.

(P4) $\Leftrightarrow$ Collatz. The cell sequence $v_2(n, t) \bmod k$ records which column of the $(2, 3)$ -torus the trajectory visits. Equidistribution modulo every k implies that the time spent in dangerous cells has the correct density, yielding (P3). Conversely, bounded deficit forces the walk to stay near the linear trajectory $w(t) \approx \varepsilon t$, and the irrationality of $\log_2 3$ (Baker) forces the cell visits to equidistribute. File: WeylEquidistribution.lean.

(P5) $\Leftrightarrow$ Collatz. The Syracuse valuation sum $\sum_{i=0}^{t-1} v_2(3a_{s_i} + 1)$ (summed over odd steps s_i) records the total even-step allocation. Equidistribution of this sum on the solenoid $\Sigma_{2,3}$ is equivalent to equidistribution of the cell sequence (P4) by the Chinese Remainder Theorem decomposition of cells into 2-adic and 3-adic components. The proof lifts Denjoy–Koksma bounds [10] on Birkhoff sums to the Syracuse cocycle on the skew product. File: SolenoidMixing.lean.

(P6) $\Leftrightarrow$ Collatz. The Littlewood residence time bounds how long a trajectory can linger in a narrow strip of the torus. A simultaneous approximation bound on $(\log 2, \log 5, \log 7)$ constrains the return time to

Table 3. Axioms used in the formalization. Each corresponds to a published theorem in transcendental number theory or ergodic theory..

#	Name	Reference	Used for
A1	Baker’s theorem	Baker 1975	Cycle elimination
A2	Rhin’s bound	Rhin 1987	Irrationality measure
A3	Hercher’s bound	Hercher 2023	m -cycle elimination ($m \leq 91$)
A4	Weyl equidistribution	Weyl 1916	Cell error equidistribution
A5	Baker–Steiner	Steiner 1977	Cycle length lower bound
A6	Matveev three-log	Matveev 2000	Linear forms in three logarithms
A7	Denjoy–Koksma	Herman 1979	Birkhoff sum bounds
A8	Furstenberg unique erg.	Furstenberg 1961	Skew product ergodicity

Table 4. Open problems (sorry statements) in the formalization. Each is machine-checked to be equivalent to the Collatz conjecture..

#	Name	File	Property
S1	nu3_linear_bound	Drift.lean	(P1)
S2	finite_deficit_bound	DiophantineRepeller.lean	(P2)
S3	equidist_implies_deficit	WeylEquidistribution.lean	(P3)
S4	cellSeqNu2_equidist	WeylEquidistribution.lean	(P4)
S5	syracuseValSum_equidist	SolenoidMixing.lean	(P5)
S6	simult_approx_log2_5_7	LittlewoodInduction.lean	(P6)

dangerous cells, via Matveev’s theorem on linear forms in three logarithms [13]. The reduction proceeds through the continued fraction expansion of $\log_2 3$ and the residence time analysis in 2-dimensional torus cells. File: `LittlewoodInduction.lean`.

Theorem 8.3 (Correction ratio bound). *If the walk has eventually linear drift with rate $\varepsilon > 0$ (i.e., $w(n, t) \geq \varepsilon t$ for $t \geq T_0$), then $r(t)$ is eventually bounded. Specifically, letting $\alpha > \log_2 3$ denote the minimum safe-step valuation sum per odd step, we have $\rho = 3 \cdot 2^{-\alpha} < 1$, and for all k sufficiently large,*

$$r(\tau_k + 1) \leq \frac{1}{1 - \rho} + 1, \quad (8.1)$$

where τ_k is the time of the k -th odd step.

Proof sketch. After the k -th odd step, $C(\tau_k + 1) = \sum_{j=1}^k 3^{k-j} \cdot 2^{U_j}$ where U_j is the number of even steps before the j -th odd step. Dividing by $2^{\nu_2(\tau_k + 1)}$ and using $w(t) \geq \varepsilon t$ gives $r(\tau_k + 1) \leq \sum_{j=0}^{\infty} \rho^j = 1/(1 - \rho)$ plus a finite correction. $\square$

9. The Lean 4 Formalization

The formalization comprises approximately 10,500 lines across 40 files, built on Lean 4 (v4.27.0) with Mathlib. Table 3 lists the 8 axioms (published theorems used without proof) and Table 4 lists the 6 open problems (each proved equivalent to the Collatz conjecture).

The primary technical innovation in the formalization is the **bit-peeling lemma** (`CarryBitScrambling.lean`), the first machine-checked proof that the Collatz map operates as a *lossy information channel*. Specifically: $T(n) \bmod 2^k$ depends only on $n \bmod 2^{k+1}$, so each compressed step irreversibly destroys exactly one bit of 2-adic information. By induction, M consecutive compressed steps consume M bits: the output modulo 2^k is determined by the input modulo 2^{k+M} , and the erased

bits are lost forever. This gives the Collatz map a precise channel capacity: 1 bit consumed per step, with no possibility of reconstruction. The bit-peeling lemma is the engine behind both the metric conflict (the erased bit shifts the cell error by $\log_2 3 - 1$) and the finite saturation of the branch locus (cells at resolution 2^a can only refine a times before the bit budget is exhausted).

The remaining 0-sorry infrastructure (fully proved theorems) includes:

- Cycle elimination for 1- and 2-cycles (`CycleElimination.lean`),
- Hensel attrition and the backward induction theorem (`HenselAttrition.lean`),
- The metric conflict: Hensel $\times$ Baker incompatibility (`MetricConflict.lean`),
- The fuel refill identity and divisibility transfer (`FuelDynamics.lean`),
- The spectral gap of $\times 3$: orbit cosine product identity (`SpectralGap.lean`),
- The non-coboundary theorem for the Collatz cocycle (`UniqueErgodicity.lean`),
- Sublinear drift from deficit boundedness (`SublinearDrift.lean`),
- The E_8 root system: 240 roots, norm, inner product (`E8Lattice.lean`).

10. Discussion

10.1. The information ceiling

The saturation of the branch locus at $k = 216$ suggests that the Syracuse map admits a finite *generating partition* in the sense of Adler and Weiss [1, 2]. The constant complexity of 21,632 cells represents the *state complexity* of the underlying arithmetic automaton—a limit on base-reconciliation first theorized by Cobham [5], whose theorem implies that sets defined simultaneously in base 2 and base 3 can only overlap in eventually periodic ways. Furthermore, the spectral gap of $2/3$ (Observation 7.1) ensures that the system is *uniquely ergodic* at the partition level, a property that allows lifting from statistical mixing of the $(2, 3)$ -solenoid to individual trajectories, following the foundational work of Furstenberg [6, 7]. The 1967 paper provides the “disjointness” framework (the 2-adic and 3-adic metrics do not interact), while the 1961 paper provides the unique ergodicity that converts a spectral gap into a pointwise guarantee.

The saturation of the branch locus at 21,632 cells thus suggests that the Collatz map has *finite combinatorial complexity*: beyond a certain resolution ($k = 216$), no new qualitative behavior emerges. Every trajectory, regardless of its starting value, visits only the same 23,268 non-empty cells on the $(2, 3)$ -torus. This is a strong form of “topological finiteness” that, to our knowledge, has not been previously observed.

Table 5. Computational evidence. Statistics are stable across three orders of magnitude in starting range..

Quantity	$N = 10^{10}$	$N = 10^{11}$	Note
Total trajectory steps	2.27×10^{12}	25.1×10^{12}	
Mean p_{odd}	0.3324	0.3325	Stable; well below $p_{\text{eq}} \approx 0.387$
Branch cells ($k \geq 216$)	21,632	21,632	Unchanged
Pure-even cells	1,582	1,582	Unchanged
Pure-odd cells	54	54	Unchanged
Mean $\bar{p}_{\text{odd}}$ (branch)	0.2843	0.2843	All 4 decimal places stable
Mean entropy $\bar{H}$	0.8380	0.8380	All 4 decimal places stable
“11” SFT violations	0	0	Over 25.1×10^{12} transitions

We emphasize that saturation of the branch locus does *not* imply convergence. A finite graph can support non-terminating paths, and the transition structure within the 21,632 branch cells is not Markovian (transitions depend on the full value of n , not just its cell). The gap between “finite static complexity” and “universal convergence” is precisely the content of properties (P1)–(P6).



Figure 6. Convergence of cell counts with increasing N . **(a)** Branch cell count at $k = 216$ vs. checkpoint N : rapid initial growth, saturating at 21,632 by $N \approx 10^8$. **(b)** Multiple k -levels on log scale: small- k levels saturate immediately at k^2 ; larger levels ($k = 162, 216$) take longer but all converge. **(c)** Empty cell count at $k = 729$ vs. N : slowly decreasing, but the branch cell count has already frozen. **(d)** Pure-even (1,582) and pure-odd (54) counts at $k = 216$: both stabilize early, confirming robust classification...

10.2. The ghost island: visual proof of the Baker kick

At $N = 10^{11}$, a secondary cluster of 251 cells emerges at perpendicular distance ~ 370 from the main branch locus at $k = 729$, entirely absent at $N = 10^{10}$ (Figures 4 and 5). This *ghost island* is a direct, visible consequence of Baker’s theorem in action.

The island lies along the rational approximant $460/729 \approx \log_3 2$ (error 7.2×10^{-5}), and the 352-unit void separating it from the main cluster is a geometric shadow of the Baker-type lower bound $|p - q \log_2 3| > C/q^k$. No cells can exist in the void: Baker separation forbids it. The ghost island is populated only by trajectories that reach the second-nearest rational approximant of $\log_2 3$, an event so rare that it requires $> 10^{10}$ starting values to observe even a single visit.

The ghost island is significant because it provides *visual proof* that the “Baker kick” is real and operates at exactly the scale predicted by the theory. Each danger step shifts the cell error by $\log_2 3 - 1 \approx 0.585$, and this shift accumulates until the trajectory is ejected from the main cluster entirely. The 352-unit void is the minimum ejection distance, set by the quality of the best rational approximation at scale $k = 729$. The ghost island’s statistics ($\bar{p}_{\text{odd}} = 0.276$, identical SFT constraint, cell type distribution




Figure 7. Transition and shadow analysis at $k = 729$, $N = 10^{11}$. **Top left:** Transition valence map, each non-empty cell is colored by the number of distinct transition types (ee, eo, oe) it exhibits; the ghost island appears as a secondary band near $r_3 \approx 440-460$. **Top center:** "11" successor analysis verifying the golden mean shift at each cell. **Top right:** Shadow offset spatial map, red cells are closer to the rational foliation 460/729, blue cells closer to the irrational foliation $\log_2 3$. **Bottom right:** Shadow offset distribution with gap near zero and asymmetric shoulder at +0.007 from the ghost island..

proportional to the main cluster) confirm that it is governed by the same dynamics, merely displaced by one Baker kick.

10.3. The gap

The honest assessment: each of the six open problems (P1)–(P6) is equivalent to the Collatz conjecture. No reformulation, however elegant, changes this. The contribution of this paper is not a proof of the conjecture, but a *complete structural map* of its irreducible content.

The structure decomposes into “microscopic” and “macroscopic” layers:

Table 6. Three structural forces constraining Syracuse trajectories..

Force	Metric	Effect	Status
Hensel attrition	2-adic	Danger runs of length d require $2^{d+1} \mid (n + 1)$; density 2^{-d} among odd numbers	Proved
Baker separation	Archimedean	$ \varepsilon(a, b) > C/\max^k$; cells cannot cluster near critical line	Proved (mod A1)
Weyl equidistribution	Ergodic	Cell visits uniform mod k ; no systematic avoidance of safe cells	Open ($\equiv$ Collatz)

Table 7. Comparison of approaches to the Collatz conjecture..

	Tao [18]	This paper
<i>Result type</i>	Density (almost all orbits)	Equivalence (all orbits)
<i>Strength</i>	Almost bounded values	Full conjecture $\Leftrightarrow$ 6 properties
<i>Map</i>	Syracuse Syr on $\mathbb{Z}/3^n\mathbb{Z}$	Syracuse on $(2, 3)$ -solenoid
<i>Key tool</i>	Characteristic functions	Baker separation + Hensel attrition
<i>Equidistribution</i>	Syracuse r.v.'s on $\mathbb{Z}/3^n\mathbb{Z}$	Cell sequences on torus
<i>Mixing source</i>	Entropy of Syracuse r.v.'s	Spectral gap $2/3$ of $\times 3$
<i>Gap</i>	Logarithmic density $\rightarrow$ natural	Partition mixing $\rightarrow$ trajectory
<i>Computation</i>	None	2.5×10^{13} steps, GPU
<i>Formalization</i>	None	10,500 lines Lean 4
<i>Cycle elimination</i>	Not needed (density result)	Baker + Hercher ($m \leq 91$)

The first two forces (“microscopic”) are fully proved. Hensel attrition shows that individual danger runs are exponentially rare, and Baker separation shows that they are spatially isolated on the torus. Carry-bit scrambling gives exact 1-bit erosion per compressed step, so the Collatz map genuinely “forgets” information.

The third force (“macroscopic”) is the open problem: proving that the microscopic mixing, which is entirely deterministic, not random, implies trajectory-level equidistribution of cell visits. This is the irreducible content of the Collatz conjecture, common to all six formulations (P1)–(P6).

10.4. Connection to Tao’s approach

Tao’s method [18] controls the *typical* behavior of Syracuse random variables via characteristic function bounds on $\mathbb{Z}/3^n\mathbb{Z}$. His result, that almost all orbits (in the sense of logarithmic density) attain almost bounded values, is the deepest partial result toward the conjecture. Our framework complements this in three ways:

1. **Individual vs. typical.** Tao’s approach proves that most starting values behave well; ours identifies what would be needed for *every* starting value. The six equivalences (P1)–(P6) constitute a precise “wishlist” for an individual-trajectory proof.
2. **Mixing mechanism.** Tao uses the entropy of Syracuse random variables on $\mathbb{Z}/3^n\mathbb{Z}$; we identify the spectral gap $\gamma = 2/3$ of the $\times 3$ transfer operator on 2-adic partitions (Observation 7.1) as the structural source of mixing. Both approaches ultimately rely on the multiplicative independence of 2 and 3.
3. **Machine-checked reduction.** The equivalence chain is verified in Lean 4, providing a formal “map” of the irreducible content. Any future proof of the conjecture must (implicitly or explicitly) establish one of (P1)–(P6).

The spectral gap $\gamma = 2/3$ for $\times 3$ on 2-adic partitions provides the mixing rate at the partition level. Lifting this to trajectory-level equidistribution remains the central challenge: the gap between Tao’s “almost all” and our “every trajectory” is precisely the gap between statistical and pointwise ergodic theorems for the Syracuse dynamics.

Acknowledgments. The Lean 4 formalization uses the Mathlib library. Computations were performed on an Intel Core Ultra 9 275HX (24 cores) with 128 GB DDR5 RAM and an NVIDIA RTX 5090 (24 GB GDDR7) for GPU-accelerated branch locus computation.

Competing interests. None.

Data availability statement. The Lean 4 formalization, C source code for all computations, and computed branch locus data are available at <https://github.com/johnjanik/syracuse-confinement>.

References

- [1] R. L. Adler and B. Weiss, Entropy, a complete metric invariant for automorphisms of the torus, *Proc. Nat. Acad. Sci. U.S.A.* **57** (1967), 1573–1576.
- [2] R. L. Adler and B. Weiss, *Similarity of Automorphisms of the Torus*, Memoirs of the American Mathematical Society, No. 98, AMS, 1970.
- [3] A. Baker, *Transcendental Number Theory*, Cambridge University Press, 1975.
- [4] D. Barina, Improved verification limit for the convergence of the Collatz conjecture, *J. Supercomput.* **81** (2025), 810.
- [5] A. Cobham, On the base-dependence of sets of numbers recognizable by finite automata, *Math. Systems Theory* **3** (1969), 186–192.
- [6] H. Furstenberg, Strict ergodicity and transformation of the torus, *Amer. J. Math.* **83** (1961), 573–601.
- [7] H. Furstenberg, Disjointness in ergodic theory, minimal sets, and a problem in Diophantine approximation, *Math. Systems Theory* **1** (1967), 1–49.
- [8] M. Heule, S. Aaronson, and E. Yolcu, An automated approach to the Collatz conjecture, *J. Automat. Reason.* (2023).
- [9] J. Hercher, There are no Collatz m -cycles with $m \leq 91$, *J. Integer Seq.* **26** (2023).
- [10] M. Herman, Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations, *Publ. Math. IHÉS* **49** (1979), 5–233.
- [11] J. C. Lagarias, The $3x + 1$ problem: An annotated bibliography (1963–1999), [arXiv:math/0309224](https://arxiv.org/abs/math/0309224), 2003.
- [12] J. C. Lagarias (ed.), *The Ultimate Challenge: The $3x + 1$ Problem*, AMS, 2010.
- [13] E. M. Matveev, An explicit lower bound for a homogeneous rational linear form in the logarithms of algebraic numbers. II, *Izv. Ross. Akad. Nauk Ser. Mat.* **64** (2000), 125–180.
- [14] G. Rhin, Approximants de Padé et mesures effectives d’irrationalité, *Séminaire de Théorie des Nombres, Paris 1985–86*, Progress in Math. **71** (1987), 155–164.
- [15] B. Shakibaei Asli, An explicit near-conjugacy between the Collatz map and a circle rotation, [arXiv:2601.04289](https://arxiv.org/abs/2601.04289), 2026.
- [16] J. Simons and B. M. M. de Weger, Theoretical and computational bounds for m -cycles of the $3n + 1$ problem, *Acta Arith.* **117** (2005), 51–70.
- [17] R. P. Steiner, A theorem on the Syracuse problem, in *Proceedings of the 7th Manitoba Conference on Numerical Mathematics*, 1977, pp. 553–559.
- [18] T. Tao, Almost all orbits of the Collatz map attain almost bounded values, *Forum Math. Pi* **10** (2022), e12.
- [19] H. Weyl, Über die Gleichverteilung von Zahlen mod. Eins, *Math. Ann.* **77** (1916), 313–352.