

Bachelor's Thesis  
Academic Year 2022

Modeling Head-Bobbing in Pigeon Locomotion  
using Reinforcement Learning

Faculty of Environment and Information Sciences,  
Keio University

Mioto Takahashi

A Bachelor's Thesis  
submitted to Faculty of Environment and Information Sciences, Keio University  
in partial fulfillment of the requirements for the degree of  
BACHELOR of Environment and Information Sciences

Mioto Takahashi

Thesis Committee:

Professor Tatsuya Hagino (Supervisor)

Professor Takashi Hattori (Co-Supervisor)

Abstract of Bachelor's Thesis of Academic Year 2022

# Modeling Head-Bobbing in Pigeon Locomotion using Reinforcement Learning

Category: Science / Engineering

## Summary

Lorem ipsum dolor sit amet, consectetur adipiscing elit. In efficitur porta augue, at interdum nunc lobortis at. Morbi feugiat facilisis justo, vitae maximus dolor. Cras convallis at elit in porta. Fusce lobortis tortor nibh, quis imperdiet arcu luctus quis. Mauris imperdiet urna eu mauris aliquet, vitae tincidunt orci dapibus. Vestibulum convallis elit ut velit accumsan cursus. Pellentesque lacus lacus, blandit eu felis vitae, pellentesque dignissim est.

## Keywords:

Reinforcement Learning, Biomimetic, Pigeon, Locomotion

Faculty of Environment and Information Sciences, Keio University

Mioto Takahashi

# Contents

<b>1</b>	<b>Background</b>	<b>1</b>
1.1.	Reinforcement Learning . . . . .	1
<b>2</b>	<b>Preliminary Research</b>	<b>3</b>
2.1.	Modeling Biological Phenomena using Robotics . . . . .	3
2.2.	Head-Bobbing in Pigeons . . . . .	3
2.2.1	Hold Phase . . . . .	4
2.2.2	Thrust Phase . . . . .	4
2.2.3	Regarding Kinematic Functionalities . . . . .	4
<b>3</b>	<b>Approach</b>	<b>6</b>
	<b>Acknowledgements</b>	<b>7</b>
	<b>References</b>	<b>8</b>
	<b>Appendix</b>	<b>9</b>
A.	equa porro quisquam est qu . . . . .	9

# List of Figures

# List of Tables

# Chapter 1

## Background

### 1.1. Reinforcement Learning

Reinforcement learning is a type of control system that attempts to execute tasks described by manually set cost functions by minimizing them using optimization algorithms or learning algorithms. In the case of deep reinforcement learning, the controller is modeled using deep neural networks and the cost function is minimized using gradient descent algorithms.

Reinforcement learning divides control systems into an agent and an environment. The agent acts the controller of the system that inspects its environment's state and sends output signals, or actions, that affect the environment. This mutual interactions creates a feedback loop between the two modules.

In the context of a pigeon tasked to move forward, the pigeon's brain and its nervous system connected to each limb act as the agent, and its surroundings, such as the ground and arbitrary objects on it, act as the environment. The environment outputs a state, such as the global position of the pigeon, which is used as the input for the agent. Using the provided state, the agent calculates and outputs an action, such as the torque of each joint in the pigeon's body. The action alters the state of the environment, and the environment outputs a new state and a reward. The reward describes how well the pigeon was able to execute its task, such as the current position of the pigeon relative to its previous timestep. The agent is updated to output sequences of actions that maximizes the cumulative reward, or return. The return can be interpreted as the inverted

or negated cost function.



# Chapter 2

## Preliminary Research

### 2.1. Modeling Biological Phenomena using Robotics

As Webb discusses on the topic of utilizing robotics to aid research in biology [3], robotic modeling can be used for proposing or testing hypotheses. This can be accomplished by depicting the hypotheses in the form of algorithms or hardware configurations in robots and observing their emergent behaviors. The generated behaviors can then be compared with their biological counterparts. In particular, robotic models implemented based on preliminary hypotheses can be used as null hypotheses, which can be validated after observing behaviors exhibited by them.

While Webb argues about the potential of using robotics for research in biology, he also highlights the limitations of such methods. Since robotic models naturally contain information on all of the assumptions made on the hypotheses of the biological phenomena they reference, discrepancies between the model and the phenomena must be carefully taken into consideration during experiments. For example, if a bipedal robot were to be used to model a pigeon's forward locomotion, it should account for noises and disturbances in the pigeon's sensory input signals and neurological output signals that stimulate their muscles. Additionally, environmental factors such as rough terrain that pigeons walk on should be simulated in the experiment.

From such perspective algorithmic models are inferior to models that utilize real robots since, as Webb states, they cannot confront problems seen in physical environments, such as noises or disturbances, that are not seen in virtual environ-

ments, such as physics engines. As such, algorithms in biology are often used for modeling neurological functionalities. In the context of modeling pigeons' locomotion, algorithmic models can represent the neurological controllers for the pigeons. In this case, the pigeons' interaction between their brains, their morphology, and their surrounding environments can be seen as a control system.

## 2.2. Head-Bobbing in Pigeons

The "head-bob" behavior in pigeons consists of stabilizing the global location and orientation of the head and altering them periodically. Such are dubbed as the hold phase and the thrust phase, respectively [2].

Frost and Davies' have proposed hypotheses regarding the functionalities of such behavior have been proposed [2] [1]. Both proposals highlight the use of the hold phase as a means to stabilize vision and the use of the thrust phase as a means to detect motion parallax and determine the distance between objects.

### 2.2.1 Hold Phase

Frost's hypothesis links the functionality of the hold phase to the detection of backward motion within the eye [2]. Pigeons' heads, while flying or moving forward, would be detecting objects whose movements can be distinguished from the surrounding stationary objects. Since stationary objects would be moving backwards relative to the pigeons' eyes, desensitizing backward motion would be necessary for such distinctions to be detected. However, this desensitization would be detrimental to the pigeons' object recognition while the pigeons' heads are stationary. The hypothesis highlights the existence of "backward notch" cells which counteract the aforementioned desensitization. Such cells would be activated when the pigeons' vision is stabilized and allowing them to distinguish objects moving backward relative to stationary objects, hence the necessity of the hold phase during locomotion.

Davies' hypothesis challenges this notion and highlights the lack of necessary conditions to induce a hold phase by stating that "they would fail to detect objects moving backwards through the visual field at velocities similar to that of the bird, as their responses could not be discriminated from those caused by self-induced

motion” [1]. Davies proposes the existence of cells that detect objects’ movement relative to stationary backgrounds regardless of their directions.

In the context of our model, combining the two hypotheses leads to a mechanism that stabilizes the head of the pigeon relative to arbitrary stationary objects and activate cells that detect arbitrary motion during the hold phase.

### **2.2.2 Thrust Phase**

### **2.2.3 Regarding Kinematic Functionalities**

When building our pigeon model, in addition to the hypotheses proposed regarding the hold phase and the thrust phase, we must take the effect of the torques of the neck joints and the movement of the head generated by them. Intuitively such motion would alter the balance of the entire pigeon, leading to mutual adjustments between the bipedal walk cycle and the neck control for head positioning. Additionally, the head-bobbing motion could be hypothesized to function as a means to balance the pigeon’s forward locomotion. However as Davies argues in his paper [1], since head-bobbing is not exhibited during fast forward locomotion, such as flying, it is unlikely that such behavior has kinematic purposes. Frost’s findings [2] further support this idea by demonstrating that pigeons stabilize their head in one global coordinate regardless of the body’s global velocity, it is likely that head-bobbing’s functionalities are solely based on vision.

# Chapter 3

## Approach

We define a simplified 2-dimensional model of pigeons based on incremental modeling. The pigeon model consists of 3 joints connecting one body representing the head, 2 bodies representing the neck, and one body representing the torso. The model's physics, mainly the collision and gravity, is simulated in a 2 dimensional physics engine. The torso's orientation and y-position is fixed while its x-position is incremented by a constant value. This represents forward locomotion at a constant speed.

Additionally, we build control systems for the model using reinforcement learning. By using reinforcement learning, we can train the controller to maximize reward functions that represent hypotheses or manually-defined trajectories for the bodies in the model to follow.

As a baseline for the model's control system, we attempt to recreate the head-bob movement by setting a target position for the head's position to match every timestep. The target position is first defined at a set location in front of the pigeon model  $T$  relative to the position of its torso. The target then acts as a static position in the global coordinate for the head to follow. If the distance between the target position and the torso's position goes below a set threshold value, the target is repositioned at the same location  $T$  relative to the torso's position.

We compare the trajectories of the bodies in the baseline control system to those generated by the control system that represents preliminary hypotheses.

# Acknowledgements

Lorem ipsum dolor sit amet, consectetur adipiscing elit. In efficitur porta augue, at interdum nunc lobortis at. Morbi feugiat facilisis justo, vitae maximus dolor. Cras convallis at elit in porta. Fusce lobortis tortor nibh, quis imperdiet arcu luctus quis. Mauris imperdiet urna eu mauris aliquet, vitae tincidunt orci dapibus. Vestibulum convallis elit ut velit accumsan cursus. Pellentesque lacus lacus, blandit eu felis vitae, pellentesque dignissim est.

# References

- [1] DAVIES, M. N., and GREEN, P. R. Head-bobbing during walking, running and flying: relative motion perception in the pigeon. *Journal of Experimental Biology* 138, 1 (1988), 71–91.
- [2] Frost, B. The optokinetic basis of head-bobbing in the pigeon. *Journal of Experimental Biology* 74, 1 (1978), 187–195.
- [3] Webb, B. What does robotics offer animal behaviour? *Animal behaviour* 60, 5 (2000), 545–558.

# Appendix

## A. eque porro quisquam est qu

Morbi tincidunt nec libero vel sagittis. Mauris laoreet mattis felis vitae feugiat. Aliquam erat volutpat. Interdum et malesuada fames ac ante ipsum primis in faucibus. Cras nec tellus justo. Nullam malesuada felis ut nunc dictum dapibus. Morbi dictum libero ac elementum sodales. Integer augue dui, auctor quis velit sit amet, aliquet maximus arcu. Vivamus id consequat nisi, et feugiat ex. Suspendisse justo sapien, condimentum mattis aliquet quis, elementum a diam. Duis metus dui, commodo sit amet ultrices vel, laoreet sed sem. Pellentesque rhoncus sodales lorem, vitae placerat est luctus quis. Mauris mi sem, sodales mattis velit sit amet, sodales aliquet magna. Phasellus sit amet rhoncus justo. In vehicula elit placerat porttitor au