

Color Images

A color image $f(x,y)$ has 3 values at each pixel

$$f(x,y) = [R(x,y), G(x,y), B(x,y)]$$

where $R(x,y), G(x,y), B(x,y)$ specifies the color at (x,y) .

Color Images

A color image $f(x,y)$ has 3 values at each pixel

$$f(x,y) = [R(x,y), G(x,y), B(x,y)]$$

where $R(x,y), G(x,y), B(x,y)$ specifies the color at (x,y) .

If each of the red, green, and blue bands has 8 bits/pixel,
then $f(x,y)$ has 24 bits/pixel (> 16 million colors)

Color Images

A color image $f(x,y)$ has 3 values at each pixel

$$f(x,y) = [R(x,y), G(x,y), B(x,y)]$$

where $R(x,y), G(x,y), B(x,y)$ specifies the color at (x,y) .

If each of the red, green, and blue bands has 8 bits/pixel,
then $f(x,y)$ has 24 bits/pixel (> 16 million colors)

A color is a point in a 3-D space.

Color Images

A color image $f(x,y)$ has 3 values at each pixel

$$f(x,y) = [R(x,y), G(x,y), B(x,y)]$$

where $R(x,y), G(x,y), B(x,y)$ specifies the color at (x,y) .

If each of the red, green, and blue bands has 8 bits/pixel,
then $f(x,y)$ has 24 bits/pixel (> 16 million colors)

A color is a point in a 3-D space.




FIGURE 6.7

Schematic of the RGB color cube. Points along the main diagonal have gray values, from black at the origin to white at point $(1, 1, 1)$.

Color Images

A color image $f(x,y)$ has 3 values at each pixel

$$f(x,y) = [R(x,y), G(x,y), B(x,y)]$$

where $R(x,y), G(x,y), B(x,y)$ specifies the color at (x,y) .

If each of the red, green, and blue bands has 8 bits/pixel,
then $f(x,y)$ has 24 bits/pixel (> 16 million colors)

A color is a point in a 3-D space.




FIGURE 6.7

Schematic of the RGB color cube. Points along the main diagonal have gray values, from black at the origin to white at point $(1, 1, 1)$.







FIGURE 6.8 RGB 24-bit color cube.

FIGURE 6.7

Schematic of the RGB color cube. Points along the main diagonal have gray values, from black at the origin to white at point $(1, 1, 1)$.


**FIGURE 6.8** RGB 24-bit color cube.

Each individual band of a color image, e.g. $R(x,y)$, can be displayed as a gray level image.

a
b

FIGURE 6.9

- (a) Generating the RGB image of the cross-sectional color plane $(127, G, B)$.
(b) The three hidden surface planes in the color cube of Fig. 6.8.



Ex) Limit R,G,B to the 6 values 0,51,102,153,204,255




FIGURE 6.11

Ex) Limit R,G,B to the 6 values 0,51,102,153,204,255

blue
255 204 153 102 51 0 R=255 R=204 R=153
255
204
153
102
51
0

green




FIGURE 6.10

Ex) Limit R,G,B to the 6 values 0,51,102,153,204,255

blue						R=255	R=204	R=153
255	204	153	102	51	0	R=102	R=51	R=0
255	204	153	102	51	0			
green								
153								
102								
51								
0								

Where are the
gray colors?





FIGURE 6.10

Human Color Vision

The human retina has 3 kinds of color receptors called cones. Each cone type has a certain photopigment with a characteristic spectral response.


Human Color Vision

The human retina has 3 kinds of color receptors called cones. Each cone type has a certain photopigment with a characteristic spectral response.



Human Color Vision


The human retina has 3 kinds of color receptors called cones. Each cone type has a certain photopigment with a characteristic spectral response.



At a retina location (x,y) , the light stimulus $I(x,y,\lambda)$ is encoded as 3 numbers R_L, R_M, R_S

Human Color Vision

The human retina has 3 kinds of color receptors called cones. Each cone type has a certain photopigment with a characteristic spectral response.



At a retina location (x, y) , the light stimulus $I(x, y, \lambda)$ is encoded as 3 numbers R_L, R_M, R_S

$$R_L = \int I(\lambda) L(\lambda) d\lambda \quad R_M = \int I(\lambda) M(\lambda) d\lambda \quad R_S = \int I(\lambda) S(\lambda) d\lambda$$

$L(\lambda)$ = response of long wavelength cones

$M(\lambda)$ = response of middle wavelength cones

$S(\lambda)$ = response of short wavelength cones




FIGURE 6.3 Absorption of light by the red, green, and blue cones in the human eye as a function of wavelength.

Stimuli $I_1(\lambda)$ and $I_2(\lambda)$ will appear identical if they give the same values R_L, R_M, R_S .

Stimuli $I_1(\lambda)$ and $I_2(\lambda)$ will appear identical if they give the same values R_L, R_M, R_S .

If $I_1(\lambda) = I_2(\lambda)$, then $I_1(\lambda)$ and $I_2(\lambda)$ are isomers.

Stimuli $I_1(\lambda)$ and $I_2(\lambda)$ will appear identical if they give the same values R_L, R_M, R_S .


If $I_1(\lambda) = I_2(\lambda)$, then $I_1(\lambda)$ and $I_2(\lambda)$ are isomers.

If $I_1(\lambda) \neq I_2(\lambda)$ but $I_1(\lambda)$ and $I_2(\lambda)$ give the same R_L, R_M, R_S then $I_1(\lambda)$ and $I_2(\lambda)$ are metamers.

Stimuli $I_1(\lambda)$ and $I_2(\lambda)$ will appear identical if they give the same values R_L, R_M, R_S .

If $I_1(\lambda) = I_2(\lambda)$, then $I_1(\lambda)$ and $I_2(\lambda)$ are isomers.

If $I_1(\lambda) \neq I_2(\lambda)$ but $I_1(\lambda)$ and $I_2(\lambda)$ give the same R_L, R_M, R_S then $I_1(\lambda)$ and $I_2(\lambda)$ are metamers.



If $I_1(\lambda)$ and $I_2(\lambda)$ appear identical (isomeric or metameric match), we write $I_1(\lambda) \stackrel{M}{=} I_2(\lambda)$

If $I_1(\lambda)$ and $I_2(\lambda)$ appear identical (isomeric or metameric match), we write $I_1(\lambda) \stackrel{M}{=} I_2(\lambda)$

Human vision is assumed to be trichromatic meaning

- 1) Any color can be matched by one additive combination of 3 fixed independent primaries
- 2) Color processing is linear

Color Matching Experiments




Consider the monochromatic primaries $R'(\lambda_1), G'(\lambda_2), B'(\lambda_3)$

where $\lambda_1 = 700\text{nm}$, $\lambda_2 = 546.1\text{nm}$, $\lambda_3 = 435.8\text{nm}$

Color Matching Experiments

Consider the monochromatic primaries $R'(\lambda_1), G'(\lambda_2), B'(\lambda_3)$




where $\lambda_1 = 700\text{nm}$, $\lambda_2 = 546.1\text{nm}$, $\lambda_3 = 435.8\text{nm}$



Color Matching Experiments

Consider the monochromatic primaries $R'(\lambda_1), G'(\lambda_2), B'(\lambda_3)$

where $\lambda_1 = 700\text{nm}$, $\lambda_2 = 546.1\text{nm}$, $\lambda_3 = 435.8\text{nm}$






Any unit energy monochromatic test stimulus $Q(\lambda)$ can be matched using

$$Q(\lambda) \stackrel{M}{=} x'R'(\lambda_1) + y'G'(\lambda_2) + z'B'(\lambda_3)$$

Color Matching Experiments

Consider the monochromatic primaries $R'(\lambda_1), G'(\lambda_2), B'(\lambda_3)$


where $\lambda_1 = 700\text{nm}$, $\lambda_2 = 546.1\text{nm}$, $\lambda_3 = 435.8\text{nm}$



Any unit energy monochromatic test stimulus $Q(\lambda)$ can be matched using

$$Q(\lambda) \stackrel{M}{=} x'R'(\lambda_1) + y'G'(\lambda_2) + z'B'(\lambda_3)$$

As λ varies, we get functions $x'(\lambda), y'(\lambda), z'(\lambda)$
called color matching functions for these primaries.



Example: Match $Q(\lambda)$ if $\lambda = 460 \text{ nm} \rightarrow x' = -0.026 \quad y' = 0.015 \quad z' = 0.298$


$$Q(460) \stackrel{M}{=} -0.026R'(\lambda_1) + 0.015G'(\lambda_2) + 0.298B'(\lambda_3)$$

FIGURE 6.5


Chromaticity diagram.

(Courtesy of the General Electric Co., Lamp Business Division.)

(C.I.E. CHROMATICITY DIAGRAM)




$$Q(460) \stackrel{M}{=} -0.026 R'(\lambda_1) + 0.015 G'(\lambda_2) + 0.298 B'(\lambda_3)$$



To avoid negative values of color matching functions,
CIE (Commission Internationale de l'Eclairage)
in 1931 defined all positive color matching functions
 $x(\lambda), y(\lambda), z(\lambda)$ for hypothetical primaries $R(\lambda_1), G(\lambda_2), B(\lambda_3)$

To avoid negative values of color matching functions,
CIE (Commission Internationale de l'Eclairage)
in 1931 defined all positive color matching functions
 $x(\lambda), y(\lambda), z(\lambda)$ for hypothetical primaries $R(\lambda_1), G(\lambda_2), B(\lambda_3)$



Let $I(\lambda)$ be an arbitrary (not monochromatic) light

$$I(\lambda) \stackrel{M}{=} X_I R(\lambda_1) + Y_I G(\lambda_2) + Z_I B(\lambda_3)$$

Let $I(\lambda)$ be an arbitrary (not monochromatic) light

$$I(\lambda) \stackrel{M}{=} X_I R(\lambda_1) + Y_I G(\lambda_2) + Z_I B(\lambda_3)$$

where $X_I = \int x(\lambda) I(\lambda) d\lambda$ $Y_I = \int y(\lambda) I(\lambda) d\lambda$

$$Z_I = \int z(\lambda) I(\lambda) d\lambda$$

Let $I(\lambda)$ be an arbitrary (not monochromatic) light

$$I(\lambda) \stackrel{M}{=} X_I R(\lambda_1) + Y_I G(\lambda_2) + Z_I B(\lambda_3)$$

where $X_I = \int x(\lambda) I(\lambda) d\lambda$ $Y_I = \int y(\lambda) I(\lambda) d\lambda$


$$Z_I = \int z(\lambda) I(\lambda) d\lambda$$

Problem: Given $I(\lambda)$, find r, g, b so that

$$I(\lambda) \stackrel{M}{=} r D_R(\lambda) + g D_G(\lambda) + b D_B(\lambda)$$

where $D_R(\lambda), D_G(\lambda), D_B(\lambda)$ are the known emitted energy distributions for the display guns for a color monitor.

$D_R(\lambda)$, $D_G(\lambda)$, $D_B(\lambda)$ for Plasma Display



Represent $D_R(\lambda)$, $D_G(\lambda)$, $D_B(\lambda)$ using X, Y, Z system

$$D_R(\lambda) \stackrel{M}{=} X_R R(\lambda_1) + Y_R G(\lambda_2) + Z_R B(\lambda_3)$$

$$D_G(\lambda) \stackrel{M}{=} X_G R(\lambda_1) + Y_G G(\lambda_2) + Z_G B(\lambda_3)$$

$$D_B(\lambda) \stackrel{M}{=} X_B R(\lambda_1) + Y_B G(\lambda_2) + Z_B B(\lambda_3)$$

Represent $D_R(\lambda)$, $D_G(\lambda)$, $D_B(\lambda)$ using X, Y, Z system

$$D_R(\lambda) \stackrel{M}{=} X_R R(\lambda_1) + Y_R G(\lambda_2) + Z_R B(\lambda_3)$$

$$D_G(\lambda) \stackrel{M}{=} X_G R(\lambda_1) + Y_G G(\lambda_2) + Z_G B(\lambda_3)$$

$$D_B(\lambda) \stackrel{M}{=} X_B R(\lambda_1) + Y_B G(\lambda_2) + Z_B B(\lambda_3)$$

$$X_R = \int x(\lambda) D_R(\lambda) d\lambda \quad Y_R = \int y(\lambda) D_R(\lambda) d\lambda \quad Z_R = \int z(\lambda) D_R(\lambda) d\lambda$$

$$X_G = \int x(\lambda) D_G(\lambda) d\lambda \quad Y_G = \int y(\lambda) D_G(\lambda) d\lambda \quad Z_G = \int z(\lambda) D_G(\lambda) d\lambda$$

$$X_B = \int x(\lambda) D_B(\lambda) d\lambda \quad Y_B = \int y(\lambda) D_B(\lambda) d\lambda \quad Z_B = \int z(\lambda) D_B(\lambda) d\lambda$$

Represent $D_R(\lambda)$, $D_G(\lambda)$, $D_B(\lambda)$ using X, Y, Z system

$$D_R(\lambda) \stackrel{M}{=} X_R R(\lambda_1) + Y_R G(\lambda_2) + Z_R B(\lambda_3)$$

$$D_G(\lambda) \stackrel{M}{=} X_G R(\lambda_1) + Y_G G(\lambda_2) + Z_G B(\lambda_3)$$

$$D_B(\lambda) \stackrel{M}{=} X_B R(\lambda_1) + Y_B G(\lambda_2) + Z_B B(\lambda_3)$$

$$X_R = \int x(\lambda) D_R(\lambda) d\lambda \quad Y_R = \int y(\lambda) D_R(\lambda) d\lambda \quad Z_R = \int z(\lambda) D_R(\lambda) d\lambda$$

$$X_G = \int x(\lambda) D_G(\lambda) d\lambda \quad Y_G = \int y(\lambda) D_G(\lambda) d\lambda \quad Z_G = \int z(\lambda) D_G(\lambda) d\lambda$$

$$X_B = \int x(\lambda) D_B(\lambda) d\lambda \quad Y_B = \int y(\lambda) D_B(\lambda) d\lambda \quad Z_B = \int z(\lambda) D_B(\lambda) d\lambda$$

$$\begin{bmatrix} D_R(\lambda) \\ D_G(\lambda) \\ D_B(\lambda) \end{bmatrix} \stackrel{M}{=} \underbrace{\begin{bmatrix} X_R & Y_R & Z_R \\ X_G & Y_G & Z_G \\ X_B & Y_B & Z_B \end{bmatrix}}_T \begin{bmatrix} R(\lambda_1) \\ G(\lambda_2) \\ B(\lambda_3) \end{bmatrix}$$

Represent $D_R(\lambda)$, $D_G(\lambda)$, $D_B(\lambda)$ using X, Y, Z system

$$D_R(\lambda) \stackrel{M}{=} X_R R(\lambda_1) + Y_R G(\lambda_2) + Z_R B(\lambda_3)$$

$$D_G(\lambda) \stackrel{M}{=} X_G R(\lambda_1) + Y_G G(\lambda_2) + Z_G B(\lambda_3)$$

$$D_B(\lambda) \stackrel{M}{=} X_B R(\lambda_1) + Y_B G(\lambda_2) + Z_B B(\lambda_3)$$

$$X_R = \int x(\lambda) D_R(\lambda) d\lambda \quad Y_R = \int y(\lambda) D_R(\lambda) d\lambda \quad Z_R = \int z(\lambda) D_R(\lambda) d\lambda$$

$$X_G = \int x(\lambda) D_G(\lambda) d\lambda \quad Y_G = \int y(\lambda) D_G(\lambda) d\lambda \quad Z_G = \int z(\lambda) D_G(\lambda) d\lambda$$

$$X_B = \int x(\lambda) D_B(\lambda) d\lambda \quad Y_B = \int y(\lambda) D_B(\lambda) d\lambda \quad Z_B = \int z(\lambda) D_B(\lambda) d\lambda$$

$$\begin{bmatrix} D_R(\lambda) \\ D_G(\lambda) \\ D_B(\lambda) \end{bmatrix} \stackrel{M}{=} \underbrace{\begin{bmatrix} X_R & Y_R & Z_R \\ X_G & Y_G & Z_G \\ X_B & Y_B & Z_B \end{bmatrix}}_T \begin{bmatrix} R(\lambda_1) \\ G(\lambda_2) \\ B(\lambda_3) \end{bmatrix}$$

T matrix depends on monitor and properties of the human vision system $(x(\lambda), y(\lambda), z(\lambda))$

$$\begin{bmatrix} R(\lambda_1) \\ G(\lambda_2) \\ B(\lambda_3) \end{bmatrix} \stackrel{M}{=} T^{-1} \begin{bmatrix} D_R(\lambda) \\ D_G(\lambda) \\ D_B(\lambda) \end{bmatrix}$$

$$\begin{bmatrix} R(\lambda_1) \\ G(\lambda_2) \\ B(\lambda_3) \end{bmatrix} \stackrel{M}{=} T^{-1} \begin{bmatrix} D_R(\lambda) \\ D_G(\lambda) \\ D_B(\lambda) \end{bmatrix}$$

$$I(\lambda) \stackrel{M}{=} [X_I \ Y_I \ Z_I] \begin{bmatrix} R(\lambda_1) \\ G(\lambda_2) \\ B(\lambda_3) \end{bmatrix}$$

$$\begin{bmatrix} R(\lambda_1) \\ G(\lambda_2) \\ B(\lambda_3) \end{bmatrix} \stackrel{M}{=} T^{-1} \begin{bmatrix} D_R(\lambda) \\ D_G(\lambda) \\ D_B(\lambda) \end{bmatrix}$$

$$I(\lambda) \stackrel{M}{=} [X_I \ Y_I \ Z_I] \begin{bmatrix} R(\lambda_1) \\ G(\lambda_2) \\ B(\lambda_3) \end{bmatrix}$$

$$\stackrel{M}{=} [X_I \ Y_I \ Z_I] T^{-1} \begin{bmatrix} D_R(\lambda) \\ D_G(\lambda) \\ D_B(\lambda) \end{bmatrix}$$

$$\begin{bmatrix} R(\lambda_1) \\ G(\lambda_2) \\ B(\lambda_3) \end{bmatrix} \stackrel{M}{=} T^{-1} \begin{bmatrix} D_R(\lambda) \\ D_G(\lambda) \\ D_B(\lambda) \end{bmatrix}$$

$$I(\lambda) \stackrel{M}{=} [X_I \ Y_I \ Z_I] \begin{bmatrix} R(\lambda_1) \\ G(\lambda_2) \\ B(\lambda_3) \end{bmatrix}$$

$$\stackrel{M}{=} [X_I \ Y_I \ Z_I] T^{-1} \begin{bmatrix} D_R(\lambda) \\ D_G(\lambda) \\ D_B(\lambda) \end{bmatrix}$$

$$[r \ g \ b] = [X_I \ Y_I \ Z_I] T^{-1}$$

Relationship between $x(\lambda), y(\lambda), z(\lambda)$ and $s(\lambda), M(\lambda), L(\lambda)$

In the X, Y, Z system we represent an input $I(\lambda)$ using the primaries $R(\lambda_1), G(\lambda_2), B(\lambda_3)$ as

$$I(\lambda) \stackrel{M}{=} X_I R(\lambda_1) + Y_I G(\lambda_2) + Z_I B(\lambda_3)$$

Relationship between $x(\lambda), y(\lambda), z(\lambda)$ and $S(\lambda), M(\lambda), L(\lambda)$

In the X, Y, Z system we represent an input $I(\lambda)$ using the primaries $R(\lambda_1), G(\lambda_2), B(\lambda_3)$ as

$$I(\lambda) \stackrel{M}{=} X_I R(\lambda_1) + Y_I G(\lambda_2) + Z_I B(\lambda_3)$$

$$\text{where } X_I = \int x(\lambda) I(\lambda) d\lambda \quad Y_I = \int y(\lambda) I(\lambda) d\lambda$$

$$Z_I = \int z(\lambda) I(\lambda) d\lambda$$

Relationship between $x(\lambda), y(\lambda), z(\lambda)$ and $S(\lambda), M(\lambda), L(\lambda)$

In the X, Y, Z system we represent an input $I(\lambda)$ using the primaries $R(\lambda_1), G(\lambda_2), B(\lambda_3)$ as

$$I(\lambda) \stackrel{M}{=} X_I R(\lambda_1) + Y_I G(\lambda_2) + Z_I B(\lambda_3)$$

$$\text{where } X_I = \int x(\lambda) I(\lambda) d\lambda \quad Y_I = \int y(\lambda) I(\lambda) d\lambda$$

$$Z_I = \int z(\lambda) I(\lambda) d\lambda$$

Two stimuli $I_1(\lambda)$ and $I_2(\lambda)$ will appear identical if and only if they have the same X, Y, Z .

Relationship between $x(\lambda), y(\lambda), z(\lambda)$ and $S(\lambda), M(\lambda), L(\lambda)$

In the X, Y, Z system we represent an input $I(\lambda)$ using the primaries $R(\lambda_1), G(\lambda_2), B(\lambda_3)$ as

$$I(\lambda) \stackrel{M}{=} X_I R(\lambda_1) + Y_I G(\lambda_2) + Z_I B(\lambda_3)$$

$$\text{where } X_I = \int x(\lambda) I(\lambda) d\lambda \quad Y_I = \int y(\lambda) I(\lambda) d\lambda$$

$$Z_I = \int z(\lambda) I(\lambda) d\lambda$$

Two stimuli $I_1(\lambda)$ and $I_2(\lambda)$ will appear identical if and only if they have the same X, Y, Z .

Stimuli $I_1(\lambda)$ and $I_2(\lambda)$ will appear identical if they give the same values R_L, R_M, R_S .

Relationship between $x(\lambda), y(\lambda), z(\lambda)$ and $s(\lambda), m(\lambda), l(\lambda)$

In the X, Y, Z system we represent an input $I(\lambda)$ using the primaries $R(\lambda_1), G(\lambda_2), B(\lambda_3)$ as

$$I(\lambda) \stackrel{M}{=} X_I R(\lambda_1) + Y_I G(\lambda_2) + Z_I B(\lambda_3)$$

$$\text{where } X_I = \int x(\lambda) I(\lambda) d\lambda \quad Y_I = \int y(\lambda) I(\lambda) d\lambda$$

$$Z_I = \int z(\lambda) I(\lambda) d\lambda$$

Two stimuli $I_1(\lambda)$ and $I_2(\lambda)$ will appear identical if and only if they have the same X, Y, Z .

Stimuli $I_1(\lambda)$ and $I_2(\lambda)$ will appear identical if they give the same values R_L, R_M, R_S .

$$R_L = \int I(\lambda) L(\lambda) d\lambda \quad R_M = \int I(\lambda) M(\lambda) d\lambda \quad R_S = \int I(\lambda) S(\lambda) d\lambda$$

$x(\lambda), y(\lambda), z(\lambda)$ play a similar role to $L(\lambda), M(\lambda), S(\lambda)$

$x(\lambda), y(\lambda), z(\lambda)$ play a similar role to $L(\lambda), M(\lambda), S(\lambda)$

When we change primaries, the new representation is a linear transformation of X_I, Y_I, Z_I

$x(\lambda), y(\lambda), z(\lambda)$ play a similar role to $L(\lambda), M(\lambda), S(\lambda)$

When we change primaries, the new representation is a linear transformation of X_I, Y_I, Z_I

Ex $[r \ g \ b] = [X_I \ Y_I \ Z_I] T^{-1}$

$x(\lambda), y(\lambda), z(\lambda)$ play a similar role to $L(\lambda), M(\lambda), S(\lambda)$

When we change primaries, the new representation is a linear transformation of X_I, Y_I, Z_I

Ex $[r \ g \ b] = [X_I \ Y_I \ Z_I] T^{-1}$

$$I(\lambda) \stackrel{M}{=} X_I R(\lambda_1) + Y_I G(\lambda_2) + Z_I B(\lambda_3)$$

$$I(\lambda) \stackrel{M}{=} \hat{x}' \hat{R}'(\lambda) + \hat{y}' \hat{G}'(\lambda) + \hat{z}' \hat{B}'(\lambda)$$

$x(\lambda), y(\lambda), z(\lambda)$ play a similar role to $L(\lambda), M(\lambda), S(\lambda)$

When we change primaries, the new representation is a linear transformation of X_I, Y_I, Z_I

Ex $[r \ g \ b] = [X_I \ Y_I \ Z_I] T^{-1}$

$$I(\lambda) \stackrel{M}{=} X_I R(\lambda_1) + Y_I G(\lambda_2) + Z_I B(\lambda_3)$$

$$I(\lambda) \stackrel{M}{=} \hat{x}' \hat{R}'(\lambda) + \hat{y}' \hat{G}'(\lambda) + \hat{z}' \hat{B}'(\lambda)$$

$$\hat{x}' = a_{11} X_I + a_{12} Y_I + a_{13} Z_I$$

$$\hat{y}' = a_{21} X_I + a_{22} Y_I + a_{23} Z_I$$

$$\hat{z}' = a_{31} X_I + a_{32} Y_I + a_{33} Z_I$$

$$\begin{aligned}\hat{x}' &= a_{11} \int x(\lambda) I(\lambda) d\lambda + a_{12} \int y(\lambda) I(\lambda) d\lambda + a_{13} \int z(\lambda) I(\lambda) d\lambda \\ &= \underbrace{\int_L (a_{11}x(\lambda) + a_{12}y(\lambda) + a_{13}z(\lambda)) I(\lambda) d\lambda}_{\text{new color matching function}}\end{aligned}$$

$$\hat{x}' = a_{11} \int x(\lambda) I(\lambda) d\lambda + a_{12} \int y(\lambda) I(\lambda) d\lambda + a_{13} \int z(\lambda) I(\lambda) d\lambda$$
$$= \underbrace{\int_L (a_{11}x(\lambda) + a_{12}y(\lambda) + a_{13}z(\lambda)) I(\lambda) d\lambda}_{\text{new color matching function}}$$

$$\hat{y}' = a_{21} \int x(\lambda) I(\lambda) d\lambda + a_{22} \int y(\lambda) I(\lambda) d\lambda + a_{23} \int z(\lambda) I(\lambda) d\lambda$$
$$= \underbrace{\int_L (a_{21}x(\lambda) + a_{22}y(\lambda) + a_{23}z(\lambda)) I(\lambda) d\lambda}_{\text{new color matching function}}$$

$$\begin{aligned}\hat{x}' &= a_{11} \int x(\lambda) I(\lambda) d\lambda + a_{12} \int y(\lambda) I(\lambda) d\lambda + a_{13} \int z(\lambda) I(\lambda) d\lambda \\ &= \int_L \underbrace{(a_{11}x(\lambda) + a_{12}y(\lambda) + a_{13}z(\lambda))}_\text{new color matching function} I(\lambda) d\lambda\end{aligned}$$

$$\begin{aligned}\hat{y}' &= a_{21} \int x(\lambda) I(\lambda) d\lambda + a_{22} \int y(\lambda) I(\lambda) d\lambda + a_{23} \int z(\lambda) I(\lambda) d\lambda \\ &= \int_L \underbrace{(a_{21}x(\lambda) + a_{22}y(\lambda) + a_{23}z(\lambda))}_\text{new color matching function} I(\lambda) d\lambda\end{aligned}$$

$$\begin{aligned}\hat{z}' &= a_{31} \int x(\lambda) I(\lambda) d\lambda + a_{32} \int y(\lambda) I(\lambda) d\lambda + a_{33} \int z(\lambda) I(\lambda) d\lambda \\ &= \int_L \underbrace{(a_{31}x(\lambda) + a_{32}y(\lambda) + a_{33}z(\lambda))}_\text{new color matching function} I(\lambda) d\lambda\end{aligned}$$

$$\begin{aligned}\hat{x}' &= a_{11} \int x(\lambda) I(\lambda) d\lambda + a_{12} \int y(\lambda) I(\lambda) d\lambda + a_{13} \int z(\lambda) I(\lambda) d\lambda \\ &= \int_L \underbrace{(a_{11}x(\lambda) + a_{12}y(\lambda) + a_{13}z(\lambda))}_\text{new color matching function} I(\lambda) d\lambda\end{aligned}$$

$$\begin{aligned}\hat{y}' &= a_{21} \int x(\lambda) I(\lambda) d\lambda + a_{22} \int y(\lambda) I(\lambda) d\lambda + a_{23} \int z(\lambda) I(\lambda) d\lambda \\ &= \int_L \underbrace{(a_{21}x(\lambda) + a_{22}y(\lambda) + a_{23}z(\lambda))}_\text{new color matching function} I(\lambda) d\lambda\end{aligned}$$

$$\begin{aligned}\hat{z}' &= a_{31} \int x(\lambda) I(\lambda) d\lambda + a_{32} \int y(\lambda) I(\lambda) d\lambda + a_{33} \int z(\lambda) I(\lambda) d\lambda \\ &= \int_L \underbrace{(a_{31}x(\lambda) + a_{32}y(\lambda) + a_{33}z(\lambda))}_\text{new color matching function} I(\lambda) d\lambda\end{aligned}$$

For different primaries, new color matching functions are a linear transformation of $x(\lambda), y(\lambda), z(\lambda)$

$$\begin{aligned}\hat{x}' &= a_{11} \int x(\lambda) I(\lambda) d\lambda + a_{12} \int y(\lambda) I(\lambda) d\lambda + a_{13} \int z(\lambda) I(\lambda) d\lambda \\ &= \underbrace{\int_L (a_{11} x(\lambda) + a_{12} y(\lambda) + a_{13} z(\lambda)) I(\lambda) d\lambda}_{\text{new color matching function}}\end{aligned}$$




$$\begin{aligned}\hat{y}' &= a_{21} \int x(\lambda) I(\lambda) d\lambda + a_{22} \int y(\lambda) I(\lambda) d\lambda + a_{23} \int z(\lambda) I(\lambda) d\lambda \\ &= \underbrace{\int_L (a_{21} x(\lambda) + a_{22} y(\lambda) + a_{23} z(\lambda)) I(\lambda) d\lambda}_{\text{new color matching function}}\end{aligned}$$

$$\begin{aligned}\hat{z}' &= a_{31} \int x(\lambda) I(\lambda) d\lambda + a_{32} \int y(\lambda) I(\lambda) d\lambda + a_{33} \int z(\lambda) I(\lambda) d\lambda \\ &= \underbrace{\int_L (a_{31} x(\lambda) + a_{32} y(\lambda) + a_{33} z(\lambda)) I(\lambda) d\lambda}_{\text{new color matching function}}\end{aligned}$$




For different primaries, new color matching functions are a linear transformation of $x(\lambda), y(\lambda), z(\lambda)$

$x(\lambda), y(\lambda), z(\lambda)$ are a linear transformation of $L(\lambda), M(\lambda), S(\lambda)$
so that the color matching experiments and physiological data agree.

(Ex) Suppose the human vision system has the idealized response functions



(Ex) Suppose the human vision system has the idealized response functions



so that any stimuli $I(\lambda)$ having the same R_S, R_M, R_L

$$R_S = \int I(\lambda)S(\lambda)d\lambda \quad R_M = \int I(\lambda)M(\lambda)d\lambda \quad R_L = \int I(\lambda)L(\lambda)d\lambda$$


will appear the same.

Consider 3 primaries

$P_1(\lambda)$ = monochromatic unit energy light at 450 nm

$P_2(\lambda)$ = monochromatic unit energy light at 550 nm

$P_3(\lambda)$ = monochromatic unit energy light at 650 nm



a) Find the color matching functions $x'(\lambda), y'(\lambda), z'(\lambda)$ corresponding to $P_1(\lambda), P_2(\lambda), P_3(\lambda)$ respectively that will be measured for these primaries.

a) Find the color matching functions $x'(\lambda), y'(\lambda), z'(\lambda)$ corresponding to $P_1(\lambda), P_2(\lambda), P_3(\lambda)$ respectively that will be measured for these primaries.

Let $Q(\lambda_o)$ = monochromatic unit energy light at λ_o nm

$$Q(\lambda_o) \stackrel{M}{=} x'(\lambda_o)P_1(\lambda) + y'(\lambda_o)P_2(\lambda) + z'(\lambda_o)P_3(\lambda)$$

a) Find the color matching functions $x'(\lambda), y'(\lambda), z'(\lambda)$ corresponding to $P_1(\lambda), P_2(\lambda), P_3(\lambda)$ respectively that will be measured for these primaries.

Let $Q(\lambda_o)$ = monochromatic unit energy light at λ_o nm

$$Q(\lambda_o) \stackrel{M}{=} x'(\lambda_o)P_1(\lambda) + y'(\lambda_o)P_2(\lambda) + z'(\lambda_o)P_3(\lambda)$$

For a match

$$(1) \int Q(\lambda_o)S(\lambda)d\lambda = \int [x'(\lambda_o)P_1(\lambda) + y'(\lambda_o)P_2(\lambda) + z'(\lambda_o)P_3(\lambda)]S(\lambda)d\lambda$$

$$(2) \int Q(\lambda_o)M(\lambda)d\lambda = \int [x'(\lambda_o)P_1(\lambda) + y'(\lambda_o)P_2(\lambda) + z'(\lambda_o)P_3(\lambda)]M(\lambda)d\lambda$$

$$(3) \int Q(\lambda_o)L(\lambda)d\lambda = \int [x'(\lambda_o)P_1(\lambda) + y'(\lambda_o)P_2(\lambda) + z'(\lambda_o)P_3(\lambda)]L(\lambda)d\lambda$$

$$(1) \int Q(\lambda_0) S(\lambda) d\lambda = \int x'(\lambda_0) P_1(\lambda) S(\lambda) d\lambda$$

$$(2) \int Q(\lambda_0) M(\lambda) d\lambda = \int y'(\lambda_0) P_2(\lambda) M(\lambda) d\lambda$$

$$(3) \int Q(\lambda_0) L(\lambda) d\lambda = \int z'(\lambda_0) P_3(\lambda) L(\lambda) d\lambda$$

$$(1) \int Q(\lambda_o) S(\lambda) d\lambda = \int x'(\lambda_o) P_1(\lambda) S(\lambda) d\lambda$$

$$(2) \int Q(\lambda_o) M(\lambda) d\lambda = \int y'(\lambda_o) P_2(\lambda) M(\lambda) d\lambda$$

$$(3) \int Q(\lambda_o) L(\lambda) d\lambda = \int z'(\lambda_o) P_3(\lambda) L(\lambda) d\lambda$$

If $400 \text{ nm} < \lambda_o < 500 \text{ nm}$ then

$$(1) \quad I = \int x'(\lambda_o) P_1(\lambda) S(\lambda) d\lambda \rightarrow x'(\lambda_o) = I$$

$$(2) \quad 0 = \int y'(\lambda_o) P_2(\lambda) M(\lambda) d\lambda \rightarrow y'(\lambda_o) = 0$$

$$(3) \quad 0 = \int z'(\lambda_o) P_3(\lambda) L(\lambda) d\lambda \rightarrow z'(\lambda_o) = 0$$

If $500 \text{ nm} < \lambda_0 < 600 \text{ nm}$ then

$$(1) \quad 0 = \int x'(\lambda_0) P_1(\lambda) S(\lambda) d\lambda \rightarrow x'(\lambda_0) = 0$$

$$(2) \quad 1 = \int y'(\lambda_0) P_2(\lambda) M(\lambda) d\lambda \rightarrow y'(\lambda_0) = 1$$

$$(3) \quad 0 = \int z'(\lambda_0) P_3(\lambda) L(\lambda) d\lambda \rightarrow z'(\lambda_0) = 0$$

If $500 \text{ nm} < \lambda_o < 600 \text{ nm}$ then

$$(1) \quad 0 = \int x'(\lambda_o) P_1(\lambda) S(\lambda) d\lambda \rightarrow x'(\lambda_o) = 0$$

$$(2) \quad 1 = \int y'(\lambda_o) P_2(\lambda) M(\lambda) d\lambda \rightarrow y'(\lambda_o) = 1$$

$$(3) \quad 0 = \int z'(\lambda_o) P_3(\lambda) L(\lambda) d\lambda \rightarrow z'(\lambda_o) = 0$$

If $600 \text{ nm} < \lambda_o < 700 \text{ nm}$ then

$$(1) \quad 0 = \int x'(\lambda_o) P_1(\lambda) S(\lambda) d\lambda \rightarrow x'(\lambda_o) = 0$$

$$(2) \quad 0 = \int y'(\lambda_o) P_2(\lambda) M(\lambda) d\lambda \rightarrow y'(\lambda_o) = 0$$

$$(3) \quad 1 = \int z'(\lambda_o) P_3(\lambda) L(\lambda) d\lambda \rightarrow z'(\lambda_o) = 1$$

If $500 \text{ nm} < \lambda_o < 600 \text{ nm}$ then

$$(1) \quad 0 = \int x'(\lambda_o) P_1(\lambda) S(\lambda) d\lambda \rightarrow x'(\lambda_o) = 0$$

$$(2) \quad 1 = \int y'(\lambda_o) P_2(\lambda) M(\lambda) d\lambda \rightarrow y'(\lambda_o) = 1$$




$$(3) \quad 0 = \int z'(\lambda_o) P_3(\lambda) L(\lambda) d\lambda \rightarrow z'(\lambda_o) = 0$$

If $600 \text{ nm} < \lambda_o < 700 \text{ nm}$ then




$$(1) \quad 0 = \int x'(\lambda_o) P_1(\lambda) S(\lambda) d\lambda \rightarrow x'(\lambda_o) = 0$$

$$(2) \quad 0 = \int y'(\lambda_o) P_2(\lambda) M(\lambda) d\lambda \rightarrow y'(\lambda_o) = 0$$




$$(3) \quad 1 = \int z'(\lambda_o) P_3(\lambda) L(\lambda) d\lambda \rightarrow z'(\lambda_o) = 1$$



b) Consider display guns $D_1(\lambda)$, $D_2(\lambda)$, $D_3(\lambda)$ with spectral distributions






b) Consider display guns $D_1(\lambda)$, $D_2(\lambda)$, $D_3(\lambda)$ with spectral distributions



Find a 3×3 matrix C such that

$$[D_1(\lambda) \ D_2(\lambda) \ D_3(\lambda)] \stackrel{M}{=} [P_1(\lambda) \ P_2(\lambda) \ P_3(\lambda)] C$$

b) Consider display guns $D_1(\lambda)$, $D_2(\lambda)$, $D_3(\lambda)$ with spectral distributions



Find a 3×3 matrix C such that

$$[D_1(\lambda) \ D_2(\lambda) \ D_3(\lambda)] \stackrel{M}{=} [P_1(\lambda) \ P_2(\lambda) \ P_3(\lambda)] C$$

$$C = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$$

$$D_1(\lambda) \stackrel{M}{=} c_{11} P_1(\lambda) + c_{21} P_2(\lambda) + c_{31} P_3(\lambda)$$

$$D_2(\lambda) \stackrel{M}{=} c_{12} P_1(\lambda) + c_{22} P_2(\lambda) + c_{32} P_3(\lambda)$$

$$D_3(\lambda) \stackrel{M}{=} c_{13} P_1(\lambda) + c_{23} P_2(\lambda) + c_{33} P_3(\lambda)$$

$$D_1(\lambda) \stackrel{M}{=} c_{11} P_1(\lambda) + c_{21} P_2(\lambda) + c_{31} P_3(\lambda)$$

$$D_2(\lambda) \stackrel{M}{=} c_{12} P_1(\lambda) + c_{22} P_2(\lambda) + c_{32} P_3(\lambda)$$

$$D_3(\lambda) \stackrel{M}{=} c_{13} P_1(\lambda) + c_{23} P_2(\lambda) + c_{33} P_3(\lambda)$$

$$c_{11} = \int D_1(\lambda) x'(\lambda) d\lambda = 50 \quad c_{21} = \int D_1(\lambda) y'(\lambda) d\lambda = 50 \quad c_{31} = \int D_1(\lambda) z'(\lambda) d\lambda = 0$$

$$D_1(\lambda) \stackrel{M}{=} c_{11} P_1(\lambda) + c_{21} P_2(\lambda) + c_{31} P_3(\lambda)$$

$$D_2(\lambda) \stackrel{M}{=} c_{12} P_1(\lambda) + c_{22} P_2(\lambda) + c_{32} P_3(\lambda)$$

$$D_3(\lambda) \stackrel{M}{=} c_{13} P_1(\lambda) + c_{23} P_2(\lambda) + c_{33} P_3(\lambda)$$

$$c_{11} = \int D_1(\lambda) x'(\lambda) d\lambda = 50 \quad c_{21} = \int D_1(\lambda) y'(\lambda) d\lambda = 50 \quad c_{31} = \int D_1(\lambda) z'(\lambda) d\lambda = 0$$

$$c_{12} = \int D_2(\lambda) x'(\lambda) d\lambda = 0 \quad c_{22} = \int D_2(\lambda) y'(\lambda) d\lambda = 50 \quad c_{32} = \int D_2(\lambda) z'(\lambda) d\lambda = 50$$

$$D_1(\lambda) \stackrel{M}{=} c_{11} P_1(\lambda) + c_{21} P_2(\lambda) + c_{31} P_3(\lambda)$$

$$D_2(\lambda) \stackrel{M}{=} c_{12} P_1(\lambda) + c_{22} P_2(\lambda) + c_{32} P_3(\lambda)$$

$$D_3(\lambda) \stackrel{M}{=} c_{13} P_1(\lambda) + c_{23} P_2(\lambda) + c_{33} P_3(\lambda)$$

$$c_{11} = \int D_1(\lambda) x'(\lambda) d\lambda = 50 \quad c_{21} = \int D_1(\lambda) y'(\lambda) d\lambda = 50 \quad c_{31} = \int D_1(\lambda) z'(\lambda) d\lambda = 0$$

$$c_{12} = \int D_2(\lambda) x'(\lambda) d\lambda = 0 \quad c_{22} = \int D_2(\lambda) y'(\lambda) d\lambda = 50 \quad c_{32} = \int D_2(\lambda) z'(\lambda) d\lambda = 50$$

$$c_{13} = \int D_3(\lambda) x'(\lambda) d\lambda = 100 \quad c_{23} = \int D_3(\lambda) y'(\lambda) d\lambda = 100 \quad c_{33} = \int D_3(\lambda) z'(\lambda) d\lambda = 100$$

$$D_1(\lambda) \stackrel{M}{=} c_{11} P_1(\lambda) + c_{21} P_2(\lambda) + c_{31} P_3(\lambda)$$

$$D_2(\lambda) \stackrel{M}{=} c_{12} P_1(\lambda) + c_{22} P_2(\lambda) + c_{32} P_3(\lambda)$$

$$D_3(\lambda) \stackrel{M}{=} c_{13} P_1(\lambda) + c_{23} P_2(\lambda) + c_{33} P_3(\lambda)$$

$$c_{11} = \int D_1(\lambda) x'(\lambda) d\lambda = 50 \quad c_{21} = \int D_1(\lambda) y'(\lambda) d\lambda = 50 \quad c_{31} = \int D_1(\lambda) z'(\lambda) d\lambda = 0$$

$$c_{12} = \int D_2(\lambda) x'(\lambda) d\lambda = 0 \quad c_{22} = \int D_2(\lambda) y'(\lambda) d\lambda = 50 \quad c_{32} = \int D_2(\lambda) z'(\lambda) d\lambda = 50$$

$$c_{13} = \int D_3(\lambda) x'(\lambda) d\lambda = 100 \quad c_{23} = \int D_3(\lambda) y'(\lambda) d\lambda = 100 \quad c_{33} = \int D_3(\lambda) z'(\lambda) d\lambda = 100$$

$$C = \begin{bmatrix} 50 & 0 & 100 \\ 50 & 50 & 100 \\ 0 & 50 & 100 \end{bmatrix}$$

c) For an arbitrary $I(\lambda)$, find a_1, a_2, a_3 so that

$$I(\lambda) \stackrel{M}{=} a_1 D_1(\lambda) + a_2 D_2(\lambda) + a_3 D_3(\lambda)$$

c) For an arbitrary $I(\lambda)$, find a_1, a_2, a_3 so that

$$I(\lambda) \stackrel{M}{=} a_1 D_1(\lambda) + a_2 D_2(\lambda) + a_3 D_3(\lambda)$$

$$I(\lambda) \stackrel{M}{=} [P_1(\lambda) \ P_2(\lambda) \ P_3(\lambda)] [X_I \ Y_I \ Z_I]^T$$

$$X_I = \int I(\lambda) x'(\lambda) d\lambda \quad Y_I = \int I(\lambda) y'(\lambda) d\lambda \quad Z_I = \int I(\lambda) z'(\lambda) d\lambda$$

c) For an arbitrary $I(\lambda)$, find a_1, a_2, a_3 so that

$$I(\lambda) \stackrel{M}{=} a_1 D_1(\lambda) + a_2 D_2(\lambda) + a_3 D_3(\lambda)$$

$$I(\lambda) \stackrel{M}{=} [P_1(\lambda) \ P_2(\lambda) \ P_3(\lambda)] [X_I \ Y_I \ Z_I]^T$$

$$X_I = \int I(\lambda) x'(\lambda) d\lambda \quad Y_I = \int I(\lambda) y'(\lambda) d\lambda \quad Z_I = \int I(\lambda) z'(\lambda) d\lambda$$

From b), $[P_1(\lambda) \ P_2(\lambda) \ P_3(\lambda)] \stackrel{M}{=} [D_1(\lambda) \ D_2(\lambda) \ D_3(\lambda)] C^{-1}$

c) For an arbitrary $I(\lambda)$, find a_1, a_2, a_3 so that

$$I(\lambda) \stackrel{M}{=} a_1 D_1(\lambda) + a_2 D_2(\lambda) + a_3 D_3(\lambda)$$

$$I(\lambda) \stackrel{M}{=} [P_1(\lambda) \ P_2(\lambda) \ P_3(\lambda)] [X_I \ Y_I \ Z_I]^T$$

$$X_I = \int I(\lambda) x'(\lambda) d\lambda \quad Y_I = \int I(\lambda) y'(\lambda) d\lambda \quad Z_I = \int I(\lambda) z'(\lambda) d\lambda$$

From b), $[P_1(\lambda) \ P_2(\lambda) \ P_3(\lambda)] \stackrel{M}{=} [D_1(\lambda) \ D_2(\lambda) \ D_3(\lambda)] C^{-1}$

$$I(\lambda) \stackrel{M}{=} [D_1(\lambda) \ D_2(\lambda) \ D_3(\lambda)] C^{-1} [X_I \ Y_I \ Z_I]^T$$

c) For an arbitrary $I(\lambda)$, find a_1, a_2, a_3 so that

$$I(\lambda) \stackrel{M}{=} a_1 D_1(\lambda) + a_2 D_2(\lambda) + a_3 D_3(\lambda)$$

$$I(\lambda) \stackrel{M}{=} [P_1(\lambda) \ P_2(\lambda) \ P_3(\lambda)] [X_I \ Y_I \ Z_I]^T$$

$$X_I = \int I(\lambda) x'(\lambda) d\lambda \quad Y_I = \int I(\lambda) y'(\lambda) d\lambda \quad Z_I = \int I(\lambda) z'(\lambda) d\lambda$$

From b), $[P_1(\lambda) \ P_2(\lambda) \ P_3(\lambda)] \stackrel{M}{=} [D_1(\lambda) \ D_2(\lambda) \ D_3(\lambda)] C^{-1}$

$$I(\lambda) \stackrel{M}{=} [D_1(\lambda) \ D_2(\lambda) \ D_3(\lambda)] C^{-1} [X_I \ Y_I \ Z_I]^T$$

$$[a_1 \ a_2 \ a_3]^T = C^{-1} [X_I \ Y_I \ Z_I]^T$$

c) For an arbitrary $I(\lambda)$, find a_1, a_2, a_3 so that

$$I(\lambda) \stackrel{M}{=} a_1 D_1(\lambda) + a_2 D_2(\lambda) + a_3 D_3(\lambda)$$

$$I(\lambda) \stackrel{M}{=} [P_1(\lambda) \ P_2(\lambda) \ P_3(\lambda)] [X_I \ Y_I \ Z_I]^T$$

$$X_I = \int I(\lambda) x'(\lambda) d\lambda \quad Y_I = \int I(\lambda) y'(\lambda) d\lambda \quad Z_I = \int I(\lambda) z'(\lambda) d\lambda$$

From b), $[P_1(\lambda) \ P_2(\lambda) \ P_3(\lambda)] \stackrel{M}{=} [D_1(\lambda) \ D_2(\lambda) \ D_3(\lambda)] C^{-1}$

$$I(\lambda) \stackrel{M}{=} [D_1(\lambda) \ D_2(\lambda) \ D_3(\lambda)] C^{-1} [X_I \ Y_I \ Z_I]^T$$

$$[a_1 \ a_2 \ a_3]^T = C^{-1} [X_I \ Y_I \ Z_I]^T$$

$$C^{-1} = \frac{1}{100} \begin{bmatrix} 0 & 2 & -2 \\ -2 & 2 & 0 \\ 1 & -1 & 1 \end{bmatrix}$$