MATH 152 **EXAM II** NAME _____

Mrs. Bonny Tighe 50 points

7.7-11.7 TA_____ Wed 11/3/2021

PART II

1. (10 points) Complete the following tests

GEOMETRIC SERIES $\sum_{n=0}^{\infty} a_n (r)^n$ is _____ if $|r| \geq 1$

THE COMPARISON TEST Suppose $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are series with positive terms.

i) If $\sum_{n=1}^{\infty} b_n$ is convergent and a_n ____ b_n for all n, then $\sum_{n=1}^{\infty} a_n$ is also convergent

ii) If $\sum_{n=1}^{\infty} b_n$ is divergent and a_n ____ b_n for all n, then $\sum_{n=1}^{\infty} a_n$ is also divergent

THE LIMIT COMPARISON TEST

Suppose that $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are series with positive terms . If $\lim_{n\to\infty} \frac{a_n}{b_n} = c$ where c is _____, then either both series converge or both diverge.

Name the test:_____ If $\lim_{n\to\infty} a_n \neq 0$, then $\sum_{n=1}^{\infty} a_n$ is divergent

THE INTEGRAL TEST Suppose f is a _____, _____ and _____function on [1, ∞) and let $a_n = f(n)$ then the series $\sum_{n=1}^{\infty} a_n$ is convergent if and only if _____is convergent.

P-SERIES $\sum_{n=1}^{\infty} \frac{1}{n^p}$ is _____ if $p > 1$

RATIO TEST

If $\lim_{n\to\infty} \left| \frac{a_{n+1}}{a_n} \right| = L$ $\begin{cases} L < 1 \text{ then } \sum a_n \text{ is } _____ \\ L > 1 \text{ then } \sum a_n \text{ is } _____ \\ L = 1 \text{ then the test is } _____ \end{cases}$

ROOT TEST

If $\lim_{n\to\infty} \sqrt[n]{|a_n|} = L$ and $\begin{cases} L < 1 \text{ then } \sum a_n \text{ is } _____ \\ L > 1 \text{ then } \sum a_n \text{ is } _____ \\ L = 1 \text{ then the root test is } _____ \end{cases}$

THE ALTERNATING SERIES TEST

If i) if $b_{n+1} \leq b_n$ for all n and ii) $\lim_{n\to\infty} b_n \neq 0$ Then $\sum_{n=1}^{\infty} (-1)^{n-1} b_n$ is _____

REMAINDER ESTIMATE FOR THE INTEGRAL TEST

Suppose $a_n = f(n)$ where f is a continuous, positive, decreasing function for $x \geq n$, and $\sum a_n$ is convergent . If $R_n = S - S_n$, then _____ $\leq R_n \leq$ _____

2. (5 points). Use the **Integral test** to determine if the series is divergent or convergent.

$$\sum_{n=1}^{\infty} \frac{e^n}{e^{2n}+1}$$

3. (10 points) Find the length of the arc of the curve $x = \frac{y^4}{8} + \frac{1}{4y^2}$ $1 \le y \le 2$

4. (5 points) Determine if the following improper integral is convergent or divergent. If convergent, evaluate.

$$\int_2^3 \frac{5x}{\sqrt{x^2 - 4}}\,dx$$

5. (10 points) Determine whether the series is convergent or divergent and if it is convergent **find the sum**.

a) $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n^2 + 2n}$

b) $\displaystyle\sum_{n=1}^{\infty} \frac{5^{n+1}}{4^{2n}}$

6. (10 points) Test the following series for absolute convergence, conditional convergence or divergence. State the test you are using and show all your work.

a) $\sum_{n=1}^{\infty} \frac{1}{n^2 + 3^n}$

b) $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n^2 + 3n + 1}}$

c) $\sum_{n=1}^{\infty} \frac{(3 \cdot 7 \cdot 11 \cdot 15 \cdot (4n - 1))}{5^n \, n!}$

7. (5 points) The given curve is rotated about the y-axis. Find the area of the resulting surface. $S = 2\pi \int radius \ (arclength)$ $\quad y^{2/3} + x^{2/3} = 1, \ \ 0 \le y \le 1$

MATH 152

Mrs. Bonny Tighe

EXAM II

50 points

7.7-11.7

NAME _____

TA_____ Wed 11/3/2021

PART I

1. (10 points) Use (a) the Midpoint Rule (b) the Trapezoidal Rule and (c) Simpson's Rule to approximate the given integral with the specified value of n. Find the error bounds for the Midpoint rule. Suppose $|f''(x)| \le K \quad for \quad a \le x \le b$, then

$\frac{6-2}{4} = 1$

$$|E_M| \le \frac{K(b-a)^3}{24n^2} \qquad \int_2^6 \ln x\, dx, \quad n = 4$$



$$M_4 = 1\left[\ln\left(\frac{5}{2}\right) + \ln\left(\frac{7}{2}\right) + \ln\left(\frac{9}{2}\right) + \ln\left(\frac{11}{2}\right)\right]$$

$$T_4 = \frac{1}{2}\left[\ln(2) + 2\ln(3) + 2\ln(4) + 2\ln(5) + 6\right]$$

$$S_4 = \frac{1}{3}\left[\ln(2) + 4\ln(3) + 2\ln(4) + 4\ln(5) + 6\right]$$

$f(x) = \ln x$

$f'(x) = \frac{1}{x}$

$f''(x) = -\frac{1}{x^2}$

$\left|-\frac{1}{x^2}\right|$ largest when $x = 2$

$K = \frac{1}{4}$

$$|E_M| \le \frac{\frac{1}{4}(4)^3}{24(4)^2} \qquad \frac{\cancel{\frac{1}{4}(4)^3}}{24\cancel{(4)^2}} = \boxed{\frac{1}{24}}$$

How large should we take n in order to guarantee that the Midpoint approximation for $\int_2^6 \ln x\, dx$ is accurate to within 0.001?

$$\frac{\frac{1}{4}(4)^3}{24n^2} \le \frac{1}{10^3} \qquad \frac{16}{24n^2} \le \frac{1}{10^3} \Rightarrow \frac{2}{3n^2} \le \frac{1}{10^3}$$

$$\boxed{\sqrt{\frac{2000}{3}} \le n}$$

next largest integer

2. (15 points) Multiple ~~Choice: Circle the correct letter~~

a) Find a sequence for $1, \frac{1\cdot 5}{1}, \frac{1\cdot 5\cdot 9}{1\cdot 2}, \frac{1\cdot 5\cdot 9\cdot 13}{1\cdot 2\cdot 3}, \ldots$ is

a) $a_n = \frac{4n-3}{n!}$

b) $a_n = \frac{(1 \cdot 5 \cdot 9 \cdots (4n-3))}{(n-1)!}$

c) $a_n = \frac{(1 \cdot 5 \cdot 9 \cdots (4n-3))}{n!}$

d) $a_n = \frac{4n-1}{(n-1)!}$

e) None of the above

b) $\sum_{n=1}^{\infty} \frac{(-3)^{n+1}}{2^{2n}}$ equals

a) $\frac{-12}{7}$

b) $\frac{9}{7}$

c) 12

d) $\frac{-9}{7}$

e) None of the above

c) Which of the following sequences is not convergent?

a) $a_n = \frac{5^n + 2}{3^n}$

b) $a_n = \frac{4n-1}{2n+1}$

c) $a_n = \frac{3}{n!}$

d) $a_n = \ln(n+2) - \ln(n+3)$

e) They are all convergent

d) Which test would you use to determine the absolute convergence of the series

$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\sqrt{n^3 + 4n + 2}}$

a) Alternating Series Test

b) Integral Test

c) Limit Comparison Test

d) Ratio Test

e) None of the above tests

e) Which test would you use to determine the convergence of the series $\sum_{n=0}^{\infty} \frac{(-1)^n}{e^n}$

a) Limit Comparison Test
b) Root Test
c) Integral Test
d) Ratio Test
e) All of the above

3. (5 points) Circle True or False

a) **True** **False** The first test you try when determining the convergence of a series is the limit comparison test.

b) **True** **False** Given $a_n = n^2 + 2n + 1$, then $a_{n+1} = n^2 + 4n + 4$.

c) **True** **False** To determine the convergence of $\sum_{n=0}^{\infty} \frac{n-1}{n^2 + n^3 + 1}$ I could use the Integral Test.

d) **True** **False** If $\lim_{n \to \infty} a_n = 0$, then the series $\sum_{n=0}^{\infty} a_n$ is convergent.

e) **True** **False** If $\lim_{n \to \infty} a_n = 0$, then the sequence $\{a_n\}$ is convergent.

4. (10 points) Circle True or False

True **False** a) The formula for the Trapezoid rule for approximation of a definite integral with four subdivisions is $\Delta x \left[f(x_0) + 2f(x_1) + 2f(x_2) + 2f(x_3) + f(x_4) \right]$.

True **False** b) We would compare $\int_1^{\infty} \frac{x}{\sqrt{x^6 + 2x - 3}} dx$ to $\int_1^{\infty} \frac{1}{x^2} dx$ to determine convergence.

True **False** c) The arc length of the function $y = \sin x$ on the interval $0 \le y \le 1$ is $\int_0^1 \sqrt{1 + \left(\frac{1}{1 - y^2}\right)} dy$.

$x = \sin^{-1} y$

True **False** d) $\sum_{n=2}^{\infty} \frac{(-1)^n n}{\ln(n)}$ is conditionally convergent. divergent

True **False** e) The Surface Area for the function $y = e^{2x}$ on the interval $0 \le x \le 1$ is $2\pi \int_0^1 e^{2x} \sqrt{1 + 4e^{2x^2}} dx$

5. (15 points) Multiple Choice – (3 points each) Circle the correct letter

a) When using the comparison test for the improper integral $\int_2^{\infty} \frac{x}{\sqrt{x^5 - 2x^3 - 3}} dx$ the inequality will be

a) $\frac{1}{\sqrt{x^5}} \le \frac{x}{\sqrt{x^5 - 2x^3 - 3}}$

b) $\frac{1}{\sqrt{x^5}} \ge \frac{x}{\sqrt{x^5 - 2x^3 - 3}}$

c) $\frac{1}{\sqrt{x^3}} \le \frac{x}{\sqrt{x^5 - 2x^3 - 3}}$

d) $\frac{1}{\sqrt{x^3}} \ge \frac{x}{\sqrt{x^5 - 2x^3 - 3}}$

e) None of the above

$r = x$

b) The surface area for the function $x = e^y$ when rotated about the y-axis on the interval $2 \le y \le 3$ is:

a) $2\pi \int_2^3 y\sqrt{1 + e^{2y}} dy$

b) $2\pi \int_2^3 e^y \sqrt{1 + e^{2y}} dy$

c) $2\pi \int_2^3 x\sqrt{1+\left(\frac{1}{x^2}\right)}\,dx$

d) $2\pi \int_2^3 \ln x\sqrt{1+\left(\frac{1}{x^2}\right)}\,dx$

e) None of the above

c) The K for the error bound for the midpoint formula for approximating $\int_1^4 \frac{1}{\sqrt{x}}\,dx$ with 4 subdivisions is:

a) $K = -\frac{3}{4}$

b) $K = \frac{3}{128}$

c) $K = \frac{3}{4}$

d) $K = \frac{1}{2}$

e) None of the above

d) Which of the following is **not** an improper integral?

a) $\int_{-2}^{5} \frac{1}{\sqrt{x+2}}\,dx$

b) $\int_1^3 \frac{x}{x^2+3x-4}\,dx$

c) $\int_1^3 \frac{x}{x^2-3x+4}\,dx$

d) $\int_0^1 \frac{1}{\ln x}\,dx$

e) They are all improper integrals

e) Represent $0.\overline{746}$ as a series

a) $\sum_{n=0}^{\infty} \left(\frac{746}{1000}\right)^n$

b) $\sum_{n=0}^{\infty} \frac{746}{1000}\left(\frac{1}{10}\right)^n$

c) $\sum_{n=0}^{\infty} 746\left(\frac{1}{1000}\right)^n$

d) $\sum_{n=0}^{\infty} \frac{746}{1000}\left(\frac{1}{1000}\right)^n$

e) None of the above

MATH 152 **EXAM IIA** NAME _____

Mrs. Bonny Tighe 50 points

7.7-11.7 TA_____ Wed 11/3/2021

PART II

1. (10 points) Complete the following tests

GEOMETRIC SERIES $\sum_{n=0}^{\infty} a_n (r)^n$ is _____ if $|r| < 1$

THE COMPARISON TEST Suppose $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are series with positive terms.

i) If $\sum_{n=1}^{\infty} b_n$ is convergent and a_n ____ b_n for all n, then $\sum_{n=1}^{\infty} a_n$ is also convergent

ii) If $\sum_{n=1}^{\infty} b_n$ is divergent and a_n ____ b_n for all n, then $\sum_{n=1}^{\infty} a_n$ is also divergent

THE LIMIT COMPARISON TEST

Suppose that $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are series with positive terms . If $\lim_{n\to\infty} \frac{a_n}{b_n} = c$ where c is _____, then either both series converge or both diverge.

Name the test:_____ **If** $\lim_{n\to\infty} a_n \neq 0$, then $\sum_{n=1}^{\infty} a_n$ is divergent

THE INTEGRAL TEST Suppose f is a _____, _____ and _____function on [1, ∞) and let $a_n = f(n)$ then the series $\sum_{n=1}^{\infty} a_n$ is convergent if and only if _____is convergent.

P-SERIES $\sum_{n=1}^{\infty} \frac{1}{n^p}$ is _____ if $p \leq 1$

RATIO TEST

If $\lim_{n\to\infty} \left| \frac{a_{n+1}}{a_n} \right| = L$ $\begin{cases} L < 1 \text{ then } \sum a_n \text{ is } _____ \\ L > 1 \text{ then } \sum a_n \text{ is } _____ \\ L = 1 \text{ then the test is } _____ \end{cases}$

ROOT TEST

If $\lim_{n\to\infty} \sqrt[n]{|a_n|} = L$ and $\begin{cases} L < 1 \text{ then } \sum a_n \text{ is } _____ \\ L > 1 \text{ then } \sum a_n \text{ is } _____ \\ L = 1 \text{ then the root test is } _____ \end{cases}$

THE ALTERNATING SERIES TEST

If i) if $b_{n+1} \leq b_n$ for all n and ii) $\lim_{n\to\infty} b_n = 0$ Then $\sum_{n=1}^{\infty} (-1)^{n-1} b_n$ is _____

REMAINDER ESTIMATE FOR THE INTEGRAL TEST

Suppose $a_n = f(n)$ where *f* is a continuous, positive, decreasing function for $x \geq n$, and $\sum a_n$ is convergent . If $R_n = S - S_n$, then _____ $\leq R_n \leq$ _____

2. (5 points). Use the **Integral test** to determine if the series is divergent or convergent.

$$\sum_{n=1}^{\infty} \frac{e^{1/n}}{n^2}$$

3. (10 points) Find the length of the arc of the curve $y = \frac{x^3}{6} + \frac{1}{2x}, \ \frac{1}{2} \le x \le 1$

4. (5 points) Determine if the following improper integral is convergent or divergent. If convergent, evaluate.

$$\int_1^3 \frac{x}{\sqrt{x^2 - 1}}\,dx$$

5. (10 points) Determine whether the series is convergent or divergent and if it is convergent **find the sum**.

a) $\displaystyle\sum_{n=1}^{\infty} \frac{1}{n^2 + 3n}$

b) $\displaystyle\sum_{n=1}^{\infty} \frac{(-4)^n}{e^{2n+2}}$

6. (10 points) Test the following series for absolute convergence, conditional convergence or divergence. State the test you are using and show all your work.

a) $\sum_{n=1}^{\infty} \frac{n+1}{\sqrt{n^5+3n^2+1}}$

b) $\sum_{n=0}^{\infty} \frac{(-1)^n}{2n+3}$

c) $\sum_{n=1}^{\infty} \frac{4 \cdot 7 \cdot 10 \cdots (3n+1)}{4^n n!}$

7. (5 points) The given curve is rotated about the x-axis. Find the area of the resulting surface. $S = 2\pi \int_a^b radius\, arclength$ $\qquad y^{2/3} + x^{2/3} = 1,\ 0 \le x \le 1$

MATH 152 **EXAM IIA** NAME _____

Mrs. Bonny Tighe 50 points

7.7-11.7 TA_____ Wed 11/3/2021

PART I

1. (10 points) Use (a) the Midpoint Rule (b) the Trapezoidal Rule and (c) Simpson's Rule to approximate the given integral with the specified value of n. Find the error bounds for the Midpoint rule. Suppose $|f''(x)| \le K \quad for \quad a \le x \le b$, then

$|E_M| \le \dfrac{K(b-a)^3}{24n^2}$ $\qquad \displaystyle\int_3^5 \ln x \, dx, \quad n = 4$

How large should we take *n* in order to guarantee that the Midpoint approximation for $\displaystyle\int_3^5 \ln x \, dx, \quad n = 4$ is accurate to within 0.001?

2. (15 points) Multiple Choice: Circle the correct letter

a) Find a sequence for $1, \frac{1 \cdot 5}{1}, \frac{1 \cdot 5 \cdot 9}{1 \cdot 2}, \frac{1 \cdot 5 \cdot 9 \cdot 13}{1 \cdot 2 \cdot 3}, \ldots$ is

a) $a_n = \frac{4n - 3}{n!}$

b) $a_n = \frac{(1 \cdot 5 \cdot 9 \cdots (4n - 3))}{(n - 1)!}$

c) $a_n = \frac{(1 \cdot 5 \cdot 9 \cdots (4n - 3))}{n!}$

d) $a_n = \frac{4n - 1}{(n - 1)!}$

e) None of the above

b) Which of the following sequences is not convergent?

a) $a_n = \frac{5^n + 2}{3^n}$

b) $a_n = \frac{4n - 1}{2n + 1}$

c) $a_n = \frac{3}{n!}$

d) $a_n = \ln(n + 2) - \ln(n + 3)$

e) They are all convergent

c) $\sum_{n=1}^{\infty} \frac{(-3)^{n+1}}{2^{2n}}$ equals

a) $\frac{-12}{7}$

b) $\frac{9}{7}$

c) 12

d) $\frac{-9}{7}$

e) None of the above

d) Which test would you use to determine the absolute convergence of the series $\sum_{n=0}^{\infty} \frac{(-1)^n}{e^n}$

a) Limit Comparison Test

b) Root Test

c) Integral Test
d) Ratio Test
e) None of the above

e) Which test would you use to determine the convergence of the series $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\sqrt{n^3+4n+2}}$

a) Alternating Series Test
b) Integral Test
c) Limit Comparison Test
d) Ratio Test
e) All of the above tests

3. (5 points) Circle True or False

a) **True False** The first test you try when determining the convergence of a series is the ratio test.
b) **True False** Given $a_n = n^2 + 3n + 1$, then $a_{n+1} = n^2 + 5n + 5$.
c) **True False** To determine the convergence of $\sum_{n=0}^{\infty} \frac{n-2}{n^2+n^3+1}$ I could use the Integral Test.
d) **True False** If $\lim_{n\to\infty} a_n = 0$, then the sequence $\{a_n\}$ is convergent.
e) **True False** If $\lim_{n\to\infty} a_n = 0$, then the series $\sum_{n=0}^{\infty} a_n$ is convergent.

4. (10 points) Circle True or False

True False a) The formula for the Trapezoid rule for approximation of a definite integral with four subdivisions is $\Delta x\left[f(x_0)+2f(x_1)+2f(x_2)+2f(x_3)+f(x_4)\right]$.

True **False** b) We would compare $\int_1^{\infty} \frac{x}{\sqrt{x^6 + 2x - 3}} dx$ to $\int_1^{\infty} \frac{1}{x^2} dx$ to determine convergence.

True **False** c) The Surface Area for the function $y = e^{2x}$ on the interval $0 \le x \le 1$ is $2\pi \int_0^1 e^{2x} \sqrt{1 + 4e^{2x^2}}\, dx$

True **False** d) $\sum_{n=2}^{\infty} \frac{(-1)^n n}{\ln(n)}$ is conditionally convergent .

True **False** e) The arc length for the function $y = \tan x$ on the interval $0 \le y \le 1$ is $\int_0^1 \sqrt{1 + (\sec^4 x)} dx$

5. (15 points) Multiple Choice – (3 points each) Circle the correct letter

a) When using the comparison test for the improper integral $\int_2^{\infty} \frac{x}{\sqrt{x^5 - 2x^3 - 3}} dx$ the inequality will be

a) $\frac{1}{\sqrt{x^5}} \le \frac{x}{\sqrt{x^5 - 2x^3 - 3}}$

b) $\frac{1}{\sqrt{x^5}} \ge \frac{x}{\sqrt{x^5 - 2x^3 - 3}}$

c) $\frac{1}{\sqrt{x^3}} \le \frac{x}{\sqrt{x^5 - 2x^3 - 3}}$

d) $\frac{1}{\sqrt{x^3}} \ge \frac{x}{\sqrt{x^5 - 2x^3 - 3}}$

e) None of the above

b) Represent $0.\overline{746}$ as a series

a) $\sum_{n=0}^{\infty} \left(\frac{746}{1000}\right)^n$

b) $\sum_{n=0}^{\infty} \frac{746}{1000}\left(\frac{1}{10}\right)^n$ c) $\sum_{n=0}^{\infty} 746\left(\frac{1}{1000}\right)^n$

d) $\sum_{n=0}^{\infty} \frac{746}{1000}\left(\frac{1}{1000}\right)^n$

e) None of the above

c) The surface area for the function $x = e^y$ when rotated about the y-axis on the interval $2 \le y \le 3$ is:

a) $2\pi \int_2^3 y\sqrt{1+e^{2y}}\,dy$

b) $2\pi \int_2^3 e^y \sqrt{1+e^{2y}}\,dy$

c) $2\pi \int_2^3 x \sqrt{1+\left(\frac{1}{x^2}\right)}\,dx$

d) $2\pi \int_2^3 \ln x \sqrt{1+\left(\frac{1}{x^2}\right)}\,dx$

e) None of the above

d) The K for the error bound for the midpoint formula for approximating $\int_1^4 \frac{1}{\sqrt{x}}\,dx$ with 4 subdivisions is:

a) $K = -\frac{3}{4}$

b) $K = \frac{3}{128}$

c) $K = \frac{3}{4}$

d) $K = \frac{1}{2}$

e) None of the above

e) Which of the following is **not** an improper integral?

a) $\int_{-2}^{5} \frac{1}{\sqrt{x+2}}\,dx$

b) $\int_1^3 \frac{x}{x^2+3x-4}\,dx$

c) $\int_1^3 \frac{x}{x^2-3x+4}\,dx$

d) $\int_0^1 \frac{1}{\ln x} dx$

e) They are all improper integrals