


## Key Facts: Maternal Mental Health (MMH) Conditions



### 1 in 5 Mothers are Impacted by Mental Health Conditions

Maternal mental health (MMH) conditions are the **MOST COMMON** complication of pregnancy and birth, affecting 800,000 families each year in the U.S.<sup>1,2</sup>




### Most Women are Untreated, Increasing Risk of Negative Impacts

75% of women impacted by MMH conditions **REMAIN UNTREATED**, increasing the risk of long-term negative impacts on mothers, babies, and families.<sup>4</sup>



### Timing and Onset of Anxiety and Depression<sup>6</sup>



If untreated, symptoms of MMH conditions can last up to 3 years.<sup>7</sup>



### Mental Health Conditions are a Leading Cause of Maternal Deaths

Mental health conditions are a **LEADING CAUSE** of maternal mortality, accounting for 22% of pregnancy-related deaths.<sup>3</sup>



### \$14 Billion: The Cost of Untreated MMH Conditions

The cost of not treating MMH conditions is \$32,000 per mother-infant pair, or **\$14 BILLION** each year in the U.S.<sup>5</sup>



### High-Risk Groups

Specific groups have a higher risk for maternal mental health conditions as determined by race, ethnicity, family history of mental illness, pregnancy and birth experience, and other risk factors. [Learn More](#)

Terms	Definitions
<b>Perinatal</b>	From conception through full year postpartum.
<b>Antenatal / prenatal</b>	During pregnancy.
<b>Postpartum / postnatal</b>	First year following pregnancy.
<b>Postpartum Depression / PPD / Postpartum</b>	An umbrella term describing mood changes following pregnancy.
<b>Perinatal mood disorders (PMDs) or perinatal mood and anxiety disorders (PMADs)</b> <b>Maternal mental health (MMH) or perinatal mental health (PMH) challenges / complications / conditions / disorders / illnesses</b>	Various terms used to describe mental health conditions during the perinatal timeframe.

# Range of MMH Conditions, Prevalence, and Symptoms

## Baby Blues<sup>20</sup>

- **Up to 85% of pregnant or postpartum women.**
- Normal period of transition.
- Typically include emotional sensitivity, weepiness, and / or feeling overwhelmed.
- Likely associated with the significant changes in hormones in the immediate postpartum period.
- Resolves without treatment within 2-3 weeks following childbirth.

## Anxiety Disorders<sup>20, 21</sup>

- **6-8% of pregnant or postpartum women.**
- Feeling easily stressed, worried, overwhelmed, tense.
- Panic attacks, including shortness of breath, rapid pulse, dizziness, chest or stomach pain.
- Fear of going crazy or dying.
- Intrusive or scary thoughts; thoughts of harming self or baby.
- Fear of going outside.
- Sleep disturbances; difficulty falling or staying asleep, even if baby is sleeping.

## Obsessive-Compulsive Disorder<sup>20</sup>

- **4% of pregnant or postpartum women.**
- Disturbing, repetitive, intrusive thoughts which may include thoughts of harming self or baby; these thoughts cause the individual great distress (i.e. thoughts are ego-dystonic).
- Compulsive behaviors, such as checking, in response to intrusive thoughts or in an attempt to make the thoughts stop or go away.

## Substance Use Disorder (SUD)<sup>22</sup>

- **Often co-morbid.**
- Most-frequently used substances: tobacco, alcohol, marijuana, cocaine, opioids.
- Women are at the highest risk for SUD during reproductive years, especially if access to mental health services is limited.
- Most women who use substances often decrease their use during pregnancy. Those who can quit on their own usually do so, which is the distinguishing factor between substance use and SUD.

## Depression<sup>20, 21</sup>

- **14% of pregnant or postpartum women.**
- Change in appetite, sleep, energy, motivation, concentration.
- Negative thinking including guilt, helplessness, hopelessness, worthlessness.
- Irritable, angry, rageful.
- Lack of interest in the baby.
- Low self-care.
- Intrusive or scary thoughts; thoughts of harming self or baby.

## Post-Traumatic Stress Disorders<sup>20</sup>

- **9% of pregnant or postpartum women.**
- Change in cognition, mood, arousal associated with traumatic events, typically around childbirth.
- Avoidance of stimuli associated with the traumatic event.
- Feeling constantly keyed up or on guard.
- Learn more about birth trauma and PTSD with MMHLA's [Birth Trauma and Maternal Mental Health Fact Sheet](#).

## Bipolar Disorder<sup>20, 21</sup>

- **3% of pregnant or postpartum women.**
- Manic or hypomanic episodes alternate with depressive episodes.
- Unusual shifts in mood, energy, activity levels, and ability to carry out day-to-day tasks.
- NOTE: Women with bipolar disorder are extremely vulnerable to recurrence during pregnancy and have an increased risk for postpartum depression and psychosis.

## Psychosis — MEDICAL EMERGENCY<sup>20, 21</sup>

- **1-2 women per 1,000 births.**
- Most significant and least frequent mental health condition occurring during the perinatal period.
- Increases the risk of infanticide and/or suicide.
- Symptoms include delusions, hallucinations, paranoia, rapid mood swings, cognitive impairment, focus on death, reckless behavior.
- Thoughts are ego-syntonic, meaning they do not cause the individual distress.
- Onset is sudden, usually within 1-2 weeks following childbirth.
- The mother should be under the care of a medical provider or taken to the emergency room for assessment and care.
- Learn more with MMHLA's [Pregnancy and Postpartum Psychosis Fact Sheet](#).

# Causes of MMH Conditions


MMH conditions are caused by a combination of bio-psycho-social factors.

**Biological:** The dramatic change in hormones during pregnancy and in the immediate postpartum period can have a significant impact on mood.<sup>23</sup>

**Psychological:** Some women struggle with changes in roles, relationships, and responsibilities that come with the transition to motherhood.<sup>24</sup>

**Social:** The childbearing years often include changes in jobs, homes, and finances that can add stress. External factors, such as isolation during the COVID-19 pandemic, can add to or increase feelings of anxiety or depression.<sup>25</sup>

## The Biopsychosocial Model



# Consequences of Untreated MMH Conditions

## On Mothers

Women with untreated MMH conditions during pregnancy are more likely to:<sup>26, 27</sup>

- Have poor prenatal care.
- Use substances such as alcohol, tobacco, or drugs.
- Experience physical, emotional, or sexual abuse.

Women with untreated MMH conditions postpartum are more likely to:<sup>28</sup>

- Be less responsive to their baby's cues.
- Have fewer positive interactions with their baby.
- Experience breastfeeding challenges.
- Question their competences as mothers.

## On Children

Infants born to mothers with untreated MMH conditions are at higher risk for:

- Preterm birth, small for gestational size, low birth weight.<sup>27, 29</sup>
- Stillbirth.<sup>27</sup>
- Longer stay in the neonatal intensive care unit.<sup>30</sup>
- Excessive crying.<sup>31</sup>

Untreated MMH conditions in the parent can increase the risk for:

- Impaired parent-child interactions.<sup>31</sup>
- Behavioral, cognitive, emotional delays in the child.<sup>32</sup>
- Adverse childhood experiences.<sup>33</sup>

## On Parents

Parents who are depressed or anxious are more likely to:<sup>34, 35</sup>

- Make more trips to the emergency department or doctor's office.
- Find it challenging to manage their child's chronic health conditions.
- Not adhere to guidance for safe infant sleep and car seat usage.

# Women experiencing MMH conditions might say...

“Having a baby was a mistake.”

“I'm such a bad mother, my baby and family would be better off without me.”

“I'm exhausted but can't sleep, even when the baby sleeps.”

“I feel like I'm drowning.”

“I'm afraid to be alone with my baby.”

“I want to run away.”

“I'm not bonding with my baby.”

“I was so embarrassed to say that I have postpartum depression out loud. It felt dirty, like it was a contagious disease.”

# Women at Increased Risk for MMH Conditions



The number one predictor for experiencing a maternal mental health condition is a personal or family history of mental health disorders.<sup>9</sup>

- Women with a personal or family history of mental illness.<sup>8</sup>
- Women of color.<sup>9-11</sup>
- Women who live in low-income neighborhoods.<sup>9-11</sup>
- Military servicemembers and their spouses.<sup>12</sup>
- Women veterans.<sup>44</sup>
- Immigrant women.<sup>13</sup>
- Women with a baby in the neonatal intensive care unit.<sup>14</sup>
- Women who lack social support, especially from their partner.<sup>8</sup>
- Women who have experienced birth trauma or previous sexual trauma in their lifetime.<sup>15</sup>

## Racial and Cultural Considerations

**Increased Risk:** Women of color are 3-4 times more likely to experience complications during pregnancy and childbirth and die from these complications than white women.<sup>36</sup>

**Intergenerational Trauma:** Black women enter pregnancy and childbirth suffering the impacts of intergenerational trauma, including the knowledge that many obstetric and gynecologic procedures were tested on Black women without their consent and without pain medication.<sup>37</sup>

**Institutional Racism:** Institutional racism in health care settings contributes to Black women receiving lower quality of care – such as giving birth in lower-quality hospitals – as well as being subject to dangerous, demeaning, or humiliating treatment.<sup>36, 37</sup>

## Impact on Non-Birthing Parents

**Fathers, Partners, Adoptive Parents At-Risk:** Non-birthing parents – including fathers, partners, adoptive parents – are also at risk for experiencing mental health conditions related to pregnancy and parenting.<sup>38, 39</sup>

**1 in 10 Fathers:** As many as 1 in 10 fathers experience postpartum depression, with maternal depression as the #1 predictor of paternal depression.<sup>38</sup>

**Grief and Loss:** Parents involved in adoption – both the birthing parents and the adopting parents – can also experience strong emotions, including grief and loss.<sup>39</sup>

## Barriers to Accessing Care

- Feelings of shame, stigma, guilt.<sup>16</sup>
- Expense and/or lack of access to healthcare.<sup>16</sup>
- Social biases in the healthcare system.<sup>16, 17</sup>
- Logistical challenges, such as lack of transportation or childcare.<sup>17</sup>
- Distrust of the healthcare system.<sup>16</sup>
- Fear that child protective services or immigration agencies will become involved.<sup>18, 19</sup>
- Fear of being considered a “bad mom.”<sup>16</sup>
- Racial, cultural, and religious beliefs.<sup>16</sup>



Women of color and women of low income are **MORE LIKELY** to experience maternal mental health conditions and **LESS LIKELY** to be able to access care.<sup>16, 17</sup>

# Treatment for Maternal Mental Health (MMH) Conditions

Most MMH conditions are temporary and treatable. Almost all women who experience MMH conditions can recover from a combination of self-care, social support, therapy / counseling, and medication.

Self-Care	Peer / Social Support <sup>20, 28</sup>
<p>Basic self-care – such as regular and adequate sleep, nutrition and exercise – may be challenging during the first few days and weeks with an infant, but are necessary to recover from the physical and emotional demands of pregnancy and childbirth.<sup>40</sup></p> <ul style="list-style-type: none"> <li>• <b>SLEEP.</b> Getting 4-5 hours of uninterrupted sleep is one of the most effective, least expensive things a new mother can do to start feeling better. <sup>20, 40</sup></li> <li>• <b>NUTRITION.</b> Breastfeeding mothers should eat / drink every time the baby eats to maintain calorie intake and hydration.<sup>40</sup></li> <li>• <b>MOVEMENT.</b> Light exercise (stretching, walking) and getting outdoors every day can have a significant positive impact on mood. <sup>20, 40</sup></li> <li>• <b>LIGHT.</b> Going outdoors for 20-60 minutes or using bright light therapy can help with perinatal depression.<sup>41</sup></li> <li>• <b>TIME FOR ONESELF.</b> Taking even a few minutes to recharge and rejuvenate – such as taking an interrupted shower – can increase feelings of well-being.<sup>20, 40</sup></li> </ul>	<p>New parents can feel isolated and alone during the intense period of caring for a newborn. Social support is vital during this time, and can include emotional support, companionship, information and resources, and tangible support such as preparing meals or running errands.</p>
	Mindfulness & Mindful Breathing <sup>20, 42</sup>
	<p>Mindfulness-based interventions have shown to be helpful with stress, anxiety, and depression in the perinatal population.</p>
	Therapy / Counseling <sup>21</sup>
	<p>Counseling during the perinatal period is often short-term, pragmatic, and focused on symptom relief and coping skills. Cognitive behavioral therapy and interpersonal therapy are evidence-based therapeutic techniques proven supportive during the perinatal timeframe.</p>
	Medication <sup>21, 43</sup>
	<p>Sometimes medication is required to treat MMH conditions; fortunately, there are safe and effective medications to manage mood during pregnancy and lactation. Decisions about medication are best made in consultation with obstetric and psychiatric providers.</p>

## Maternal Mental Health Resources



### National Maternal Mental Health Hotline

For individuals who are not in crisis but need real-time support and assistance for maternal mental health conditions.

- 1-833-TLC-MAMA (1-833-852-6262)
- 24 / 7 / 365 response within 5 minutes
- Voice and text
- English and Spanish
- Other languages available via translator



For more resources go to [mmhla.org/resource-hub](https://mmhla.org/resource-hub).



### Postpartum Support International Helpline

For individuals who are not in crisis but need resources and referrals for maternal mental health conditions.

- 1-800-944-4773
- Online support groups
- Peer mentor program
- Volunteer coordinators in all states
- Provider directory



# Special Thanks to Our Funders

This fact sheet was funded by grants from the [California Health Care Foundation](#) and the [ZOMA Foundation](#).

## Citations

1. Fawcett, E. J., Fairbrother, N., Cox, M. L., White, I. R., & Fawcett, J. M. (2019). The Prevalence of Anxiety Disorders During Pregnancy and the Postpartum Period: A Multivariate Bayesian Meta-Analysis. *The Journal of Clinical Psychiatry*, 80(4). <https://doi.org/10.4088/JCP.18r12527>
2. Gavin, N. I., Gaynes, B. N., Lohr, K. N., Meltzer-Brody, S., Gartlehner, G., & Swinson, T. (2005). Perinatal Depression: A Systematic Review of Prevalence and Incidence. *Obstetrics and Gynecology*, 106(5 Part 1), 1071–1083. <https://doi.org/10.1097/01.AOG.0000183597.31630.db>
3. Trost SL, Busacker A, Leonard M, et al. (2024). Pregnancy-Related Deaths: Data From Maternal Mortality Review Committees in 38 U.S. States, 2020. *U.S. Centers for Disease Control and Prevention*. <https://www.cdc.gov/maternal-mortality/php/data-research/index.html>
4. Byatt, N., Levin, L. L., Ziedonis, D., Moore Simas, T. A., & Allison, J. (2015). Enhancing Participation in Depression Care in Outpatient Perinatal Care Settings: A Systematic Review. *Obstetrics and Gynecology*, 126(5), 1048–1058. <https://doi.org/10.1097/AOG.0000000000001067>
5. Luca, D. L., Margiotta, C., Staatz, C., Garlow, E., Christensen, A., & Zivin, K. (2020). Financial Toll of Untreated Perinatal Mood and Anxiety Disorders Among 2017 Births in the United States. *American Journal of Public Health*, 110(6), 888–896. <https://doi.org/10.2105/AJPH.2020.305619>
6. Wisner, K. L., Sit, D. K. Y., McShea, M. C., Rizzo, D. M., Zoretich, R. A., Hughes, C. L., Eng, H. F., Luther, J. F., Wisniewski, S. R., Costantino, M. L., Confer, A. L., Moses-Kolko, E. L., Famy, C. S., & Hanusa, B. H. (2013). Onset Timing, Thoughts of Self-harm, and Diagnoses in Postpartum Women With Screen-Positive Depression Findings. *JAMA Psychiatry*, 70(5), 490–498. <https://doi.org/10.1001/jamapsychiatry.2013.87>
7. Putnick, D. L., Sundaram, R., Bell, E. M., Ghassabian, A., Goldstein, R. B., Robinson, S. L., Vafai, Y., Gilman, S. E., & Yeung, E. (2020). Trajectories of Maternal Postpartum Depressive Symptoms. *Pediatrics*, 146(5), e20200857. <https://doi.org/10.1542/peds.2020-0857>
8. Agrawal, I., Mehendale, A. M., & Malhotra, R. (2022). Risk Factors of Postpartum Depression. *Cureus*, 14(10), e30898. <https://doi.org/10.7759/cureus.30898>
9. Hutchens, B. F., & Kearney, J. (2020). Risk Factors for Postpartum Depression: An Umbrella Review. *Journal of Midwifery & Women's Health*, 65(1), 96–108. <https://doi.org/10.1111/jmwh.13067>
10. Howell, E. A., Mora, P. A., Horowitz, C. R., & Leventhal, H. (2005). Racial and Ethnic Disparities in Factors Associated With Early Postpartum Depressive Symptoms. *Obstetrics and Gynecology*, 105(6), 1442–1450. <https://doi.org/10.1097/01.AOG.0000164050.34126.37>
11. Kozhimannil, K. B., Trinacty, C. M., Busch, A. B., Huskamp, H. A., & Adams, A. S. (2011). Racial and Ethnic Disparities in Postpartum Depression Care Among Low-income Women. *Psychiatric Services*, 62(6), 619–25. <https://doi.org/10.1176/appi.ps.62.6.619>
12. United States Government Accountability Office (2022). *Defense Health Care, Prevalence of and Efforts to Screen and Treat Mental Health Conditions in Prenatal and Postpartum TRICARE Beneficiaries*. <https://www.gao.gov/assets/gao-22-105136.pdf>
13. Falah-Hassani, K., Shiri, R., Vigod, S., & Dennis, C.-L. (2015). Prevalence of Postpartum Depression Among Immigrant Women: A Systematic Review and Meta-Analysis. *Journal of Psychiatric Research*, 70, 67–82. <https://doi.org/10.1016/j.jpsychires.2015.08.010>
14. Grunberg, V. A., Geller, P. A., Hoffman, C., Njoroge, W., Ahmed, A., & Patterson, C. A. (2022). Parental Mental Health Screening in the NICU: A Psychosocial Team Initiative. *Journal of Perinatology*, 42(3), 401–409. <https://doi.org/10.1038/s41372-021-01217-0>
15. Wosu, A. C., Gelaye, B., & Williams, M. A. (2015). History of childhood sexual abuse and risk of prenatal and postpartum depression or depressive symptoms: an epidemiologic review. *Archives of Women's Mental Health*, 18(5), 659–671. <https://doi.org/10.1007/s00737-015-0533-0>
16. Hansotte, E., Payne, S. I., & Babich, S. M. (2017). Positive postpartum depression screening practices and subsequent mental health treatment for low-income women in Western countries: a systematic literature review. *Public Health Reviews*, 38(1), 3. <https://doi.org/10.1186/s40985-017-0050-y>
17. Taylor, J., Novoa, C., Hamm, K., & Phadke, S. (2019). Eliminating Racial Disparities in Maternal and Infant Mortality A Comprehensive Policy Blueprint. *Center for American Progress*. [www.americanprogress.org/article/eliminating-racial-disparities-maternal-infant-mortality/](https://www.americanprogress.org/article/eliminating-racial-disparities-maternal-infant-mortality/)
18. Dwarakanath, M., Hossain, F., Balascio, P., Moore, M. C., Hill, A. V., & De Genna, N. M. (2023). Experiences of postpartum mental health sequelae among black and biracial women during the COVID-19 pandemic. *BMC Pregnancy and Childbirth*, 23(1), 636. <https://doi.org/10.1186/s12884-023-05929-3>
19. Heslehurst, N., Brown, H., Pemu, A., Coleman, H., & Rankin, J. (2018). Perinatal health outcomes and care among asylum seekers and refugees: a systematic review of systematic reviews. *BMC Medicine*, 16(1), 89. <https://doi.org/10.1186/s12916-018-1064-0>
20. Byatt, N., Mittal, L., Brenckle, L., Logan, D., Masters, G., Bergman, A., & Moore Simas, T. (2019). Lifeline for Moms Perinatal Mental Health Toolkit. *Psychiatry Information in Brief*, 16(7). <https://doi.org/10.7191/pib.1140>
21. American College of Obstetricians and Gynecologists. (2023). Treatment and Management of Mental Health Conditions During Pregnancy and Postpartum. Clinical Practice Guideline Number 5. *Obstetrics & Gynecology*, 141(6), 1262–1288. <https://doi.org/10.1097/AOG.0000000000005202>
22. Prince, M. K., Daley, S. F., & Ayers, D. (2023). *Substance Use in Pregnancy*. StatPearls Publishing. <https://www.ncbi.nlm.nih.gov/books/NBK542330/>
23. Trifu, S., Vladuti, A., & Popescu, A. (2019). The Neuroendocrinological Aspects of Pregnancy and Postpartum Depression. *Acta Endocrinol (Buchar)*, 15(3), 410–415. <https://doi.org/10.4183/aeb.2019.410>
24. Epifanio, M. S., Genna, V., De Luca, C., Roccella, M., & La Grutta, S. (2015). Paternal and Maternal Transition to Parenthood: The Risk of Postpartum Depression and Parenting Stress. *Pediatric Reports*, 7(2), 5872. <https://doi.org/10.4081/pr.2015.5872>
25. Firestein, M. R., Dumitriu, D., Marsh, R., & Monk, C. (2022). Maternal Mental Health and Infant Development During the COVID-19 Pandemic. *JAMA Psychiatry*, 79(10), 1040–1045. <https://doi.org/10.1001/jamapsychiatry.2022.2591>
26. Zhou, J., Ko, J. Y., Haight, S. C., & Tong, V. T. (2019). Treatment of Substance Use Disorders Among Women of Reproductive Age by Depression and Anxiety Disorder Status, 2008–2014. *Journal of Women's Health*, 28(8), 1068–1076. <https://doi.org/10.1089/jwh.2018.7597>
27. Jahan, N., Went, T. R., Sultan, W., Sapkota, A., Khurshid, H., Qureshi, I. A., & Alfonso, M. (2021). Untreated Depression During Pregnancy and Its Effect on Pregnancy Outcomes: A Systematic Review. *Cureus*, 13(8), e17251. <https://doi.org/10.7759/cureus.17251>
28. Fitelson, E., Kim, S., Scott Baker, A., & Leight, K. (2010). Treatment of Postpartum Depression: Clinical, Psychological, and Pharmacological Options. *International Journal of Women's Health*, 2011(3), 1–14. <https://doi.org/10.2147/IJWH.S6938>
29. Langham, J., Gurol-Urganci, I., Muller, P., Webster, K., Tassie, E., Heslin, M., Byford, S., Khalil, A., Harris, T., Sharp, H., Pasupathy, D., van der Meulen, J., Howard, L. M., & O'Mahen, H. A. (2023). Obstetric and neonatal outcomes in pregnant women with and without a history of specialist mental health care: a national population-based cohort study using linked routinely collected data in England. *The Lancet Psychiatry*, 10(10), P748–759. [https://doi.org/10.1016/S2215-0366\(23\)00200-6](https://doi.org/10.1016/S2215-0366(23)00200-6)
30. Cherry, A., Mignogna, M., Roddenberry Vaz, A., Hetherington, C., McCaffree, M. A., Anderson, M., & Gillaspay, S. (2016). The Contribution of Maternal Psychological Functioning to Infant Length of Stay in the Neonatal Intensive Care Unit. *International Journal of Women's Health*, 8, 233–242. <https://doi.org/10.2147/IJWH.S91632>
31. Olmestig, T. K., Siersma, V., Birkmoose, A. R., Kragstrup, J., & Ertmann, R. K. (2021). Infant Crying Problems Related to Maternal Depressive and Anxiety Symptoms During Pregnancy: A Prospective Cohort Study. *BMC Pregnancy and Childbirth*, 21(1), 777. <https://doi.org/10.1186/s12884-021-04252-z>
32. Mughal, M. K., Giallo, R., Arnold, P. D., Kehler, H., Bright, K., Benzie, K., Wajid, A., & Kingston, D. (2019). Trajectories of Maternal Distress and Risk of Child Developmental Delays: Findings From the All Our Families (AOF) Pregnancy Cohort. *Journal of Affective Disorders*, 248, 1–12. <https://doi.org/10.1016/j.jad.2018.12.132>
33. Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V., Koss, M. P., & Marks, J. S. (1998). Relationship of Childhood Abuse and Household Dysfunction to Many of the Leading Causes of Death in Adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventative Medicine*, 14(4), 245–58. [https://doi.org/10.1016/s0749-3797\(98\)00017-8](https://doi.org/10.1016/s0749-3797(98)00017-8)
34. Field, T. (2010). Postpartum Depression Effects on Early Interactions, Parenting, and Safety Practices: A Review. *Infant and Behavior Development*, 33(1), 1–6. <https://doi.org/10.1016/j.infbeh.2009.10.005>
35. Sriraman, N. K., Pham, D.-Q., & Kumar, R. (2017). Postpartum Depression: What Do Pediatricians Need to Know? *Pediatrics in Review*, 38(12), 541–551. <https://doi.org/10.1542/pir.2015-0133>
36. MacDorman, M. F., Thoma, M., Declercq, E., & Howell, E. A. (2021). Racial and Ethnic Disparities in Maternal Mortality in the United States Using Enhanced Vital Records, 2016–2017. *American Journal of Public Health*, 111(9), 1673–1681. <https://doi.org/10.2105/AJPH.2021.306375>
37. Chambers, B. D., Taylor, B., Nelson, T., Harrison, J., Bell, A., O'Leary, A., Arega, H. A., Hashemi, S., McKenzie-Sampson, S., Scott, K. A., Raine-Bennett, T., Jackson, A. V., Kuppermann, M., & McLemore, M. R. (2022). Clinicians' Perspectives on Racism and Black Women's Maternal Health. *Women's Health Reports*, 3(1), 476–482. <https://doi.org/10.1089/whr.2021.0148>
38. Da Costa, D., Danieli, C., Abrahamowicz, M., Dasgupta, K., Sewitch, M., Lowenstein, I., & Zekowitz, P. (2019). A Prospective Study of Postnatal Depressive Symptoms and Associated Risk Factors in First-time Fathers. *Journal of Affective Disorders*, 249, 371–377. <https://doi.org/10.1016/j.jad.2019.02.033>
39. Foli, K. J. (2021). Understanding Parental Postadoption Depression. *Adoption Advocate*, 158, 1–12. <https://adoptioncouncil.org/wp-content/uploads/2021/08/Adoption-Advocate-No-158.pdf>
40. Barkin, J. L., & Wisner, K. L. (2013). The role of maternal self-care in new motherhood. *Midwifery*, 29(9), 1050–1055. <https://doi.org/10.1016/j.midw.2012.10.001>
41. Nonacs, R. (2020, September). *Can Bright Light Therapy Be Used for the Treatment of Depression During Pregnancy?* MGH Center for Women's Mental Health. <https://womensmentalhealth.org/posts/bright-light-therapy/>
42. Shi, Z., & MacBeth, A. (2017). The Effectiveness of Mindfulness-Based Interventions on Maternal Perinatal Mental Health Outcomes: A Systematic Review. *Mindfulness*, 8(4), 823–847. <https://doi.org/10.1007/s12671-016-0673-y>
43. Payne, J. (2017). Psychopharmacology During Pregnancy and Breastfeeding. *Psychiatric Clinics of North America*, 40(2), 217–238. <https://doi.org/10.1016/j.psc.2017.01.001>
44. Miller, L. J., Rowlands, S., Esposito, L. et al. (2022). The Veterans Health Administration Reproductive Mental Health Consultation Program: an Innovation to Improve Access to Specialty Care. *Journal of General Internal Medicine*, 37 (Suppl 3), 833–836. <https://doi.org/10.1007/s11606-022-07583-5>