

COEFFICIENTWISE LOG-CONCAVITY OF THE CUBIC MULTIPLE-POCHHAMMER SERIES

JOHN SHIELDS

ABSTRACT. Let $(f_n)_{n \geq 1}$ be a nonnegative log-concave sequence with no internal zeros, and define the cubic multiple-Pochhammer series $F_f(\mu; x) = \sum_{n \geq 1} f_n(\mu)_{3n} x^n / (3n - 1)!$. Karp and Zhang established coefficientwise log-concavity of the analogous series for multiplicity $r = 2$ and gave numerical counterexamples for $r = 4$, leaving the cubic case open as their Conjecture 1. We prove it: for all $\mu, \alpha, \beta \geq 0$, every coefficient of the generalized Turánian $F_f(\mu + \alpha; x)F_f(\mu + \beta; x) - F_f(\mu; x)F_f(\mu + \alpha + \beta; x)$ is nonnegative. Equivalently, the map $(f_n) \mapsto F_f$ carries every such sequence to a coefficientwise log-concave family. The proof is elementary and noncomputational. The degree- m coefficient is represented as a beta-binomial expectation, reducing the theorem to the monotonicity of a weighted residue-class binomial kernel; this monotonicity is established through a Bernstein-basis positivity certificate obtained from a root-of-unity evaluation. A coefficientwise strictness criterion and ordinary log-concavity in μ on $(0, \infty)$ follow.

1. INTRODUCTION

A formal power series depending on a parameter $\mu \geq 0$ is called *coefficientwise log-concave* if, for every $\mu, \alpha, \beta \geq 0$, all coefficients of the generalized Turánian

$$F(\mu + \alpha; x)F(\mu + \beta; x) - F(\mu; x)F(\mu + \alpha + \beta; x)$$

are nonnegative. This property is stronger than the corresponding pointwise inequality: whenever the relevant series converge at a positive value of x , summing the coefficient inequalities yields the generalized Turán inequality for $\mu \mapsto F(\mu; x)$ there, simultaneously for all admissible shifts. It has been studied systematically for reciprocal-gamma, gamma-ratio, hypergeometric, and Fox–Wright series; see [4, 5, 6, 7]. Turán-type inequalities for hypergeometric and confluent hypergeometric functions have also been studied extensively [1, 2].

Karp and Zhang considered, for an integer $r \geq 2$, the modified multiple-Pochhammer series

$$(1.1) \quad F_{f,r}(\mu; x) = \sum_{n \geq 1} f_n \frac{(\mu)_{rn}}{(rn - 1)!} x^n,$$

where (f_n) is nonnegative and log-concave. Karp and Zhang report that A. Ninh suggested this construction to them in a private communication in 2017 [7]; related published work on log-concavity and Turán-type inequalities appears in [9]. They proved coefficientwise log-concavity for $r = 2$, gave numerical counterexamples for $r = 4$, and left the cubic case $r = 3$ open [7, Conjecture 1]. This note proves that conjecture.

Karp and Zhang index the ambient sequence from $n = 0$, although the modified series itself begins at $n = 1$. The log-concavity hypothesis is stated here directly for the positive-index tail $(f_n)_{n \geq 1}$. This is equivalent to their formulation: restriction preserves log-concavity and interval support, while any admissible tail extends to $n = 0$ by setting $f_0 = 0$.

Recall that a nonnegative sequence $(f_n)_{n \geq 1}$ is log-concave with no internal zeros if $f_n^2 \geq f_{n-1}f_{n+1}$ whenever the three terms are defined, and its positive support is an interval of integers. We work throughout with formal power series, so no growth condition on (f_n) is needed.

Date: July 24, 2026.

2020 Mathematics Subject Classification. Primary 26A51, 33C20; Secondary 05A20, 60E15.

Key words and phrases. coefficientwise log-concavity, Turán inequality, Pochhammer symbol, log-concave sequence, beta-binomial distribution, Bernstein basis.

Independent researcher, Tokyo, Japan. Corresponding author: math@johnnyshields.com. This version is archived at doi:10.5281/zenodo.21475821.

Theorem 1.1 (Cubic multiple-Pochhammer theorem). *Let $(f_n)_{n \geq 1}$ be a nonnegative log-concave sequence with no internal zeros, and put*

$$(1.2) \quad F_f(\mu; x) = \sum_{n \geq 1} f_n \frac{(\mu)_{3n}}{(3n-1)!} x^n.$$

For every $\mu, \alpha, \beta \geq 0$, the formal power series

$$(1.3) \quad \Delta_f(\mu; \alpha, \beta; x) = F_f(\mu + \alpha; x)F_f(\mu + \beta; x) - F_f(\mu; x)F_f(\mu + \alpha + \beta; x)$$

has nonnegative coefficients.

Theorem 1.1 concerns the transform $(f_n) \mapsto F_f$, not a single function: every nonnegative log-concave sequence is sent to a parameter family whose generalized Turánians have nonnegative coefficients. Where the series converges, these coefficient inequalities sum to the pointwise generalized Turán inequalities; setting $\alpha = \beta$ and invoking continuity then yields ordinary log-concavity of $\mu \mapsto F_f(\mu; x)$ (Corollary 5.3). The cubic case is the intervening multiplicity between the positive case $r = 2$ and the first known failure at $r = 4$, and the residue-class positivity of Section 4 is the mechanism behind it.

For $f_n \equiv 1$, the triplication formula for the Pochhammer symbol gives

$$F_f(\mu; x) = 3x \frac{d}{dx} {}_3F_2\left(\frac{\mu}{3}, \frac{\mu+1}{3}, \frac{\mu+2}{3}; \frac{1}{3}, \frac{2}{3}; x\right),$$

so the theorem yields coefficientwise generalized Turán inequalities for the differentiated ${}_3F_2$ family above; a general log-concave (f_n) gives a coefficient-weighted extension.

The proof has three components. The coefficient of degree m is rewritten as a beta-binomial expectation (Section 2). Two elementary monotonicity lemmas (Section 3) then supply the tools: beta laws with a fixed sum of parameters are ordered by their imbalance against symmetric functions increasing toward the midpoint, and a one-sign-change weighting principle carries positivity through nondecreasing weights. The weighted residue-class binomial kernel is shown to have the required monotonicity (Section 4); here complementary residue blocks have a single sign change, and the unweighted sum carries a Bernstein-basis positivity certificate. This kernel theorem does not depend on the beta-binomial setup. The components are combined in Section 5.

2. BETA-BINOMIAL REPRESENTATION

This section rewrites the coefficient convolution underlying the Turánian (1.3) as a beta-binomial expectation.

For $m \geq 2$, put

$$(2.1) \quad w_r = f_r f_{m-r}, \quad 1 \leq r \leq m-1.$$

These weights are symmetric. Log-concavity of (f_n) implies

$$w_{r+1} = f_{r+1} f_{m-r-1} \geq f_r f_{m-r} = w_r \quad \left(1 \leq r < \left\lfloor \frac{m}{2} \right\rfloor\right),$$

the standard cross-product consequence $f_{a+1} f_b \geq f_a f_{b+1}$ for $a < b$ of log-concavity with no internal zeros. With the symmetry $w_r = w_{m-r}$, the weights (2.1) satisfy the hypotheses of Theorem 4.1.

For $u, v \geq 0$, the degree- m coefficient of the product $F_f(u; x)F_f(v; x)$ is

$$(2.2) \quad C_{m,f}(u, v) = \sum_{r=1}^{m-1} w_r \frac{(u)_{3r} (v)_{3(m-r)}}{(3r-1)! [3(m-r)-1]!},$$

a finite sum, with $C_{m,f}(0, v) = 0$ since $(0)_{3r} = 0$. The beta-binomial representation below is used for $u, v > 0$. Let $N = 3m$ and let J have the beta-binomial distribution

$$\mathbb{P}(J = j) = \binom{N}{j} \frac{(u)_j (v)_{N-j}}{(u+v)_N}, \quad 0 \leq j \leq N.$$

For $1 \leq j \leq N - 1$,

$$\frac{(u)_j(v)_{N-j}}{(j-1)!(N-j-1)!} = \frac{(u+v)_N}{N!} j(N-j) \mathbb{P}(J=j).$$

Defining a weight on all integers $0 \leq j \leq N$ by

$$\tilde{w}_j = \begin{cases} w_{j/3}, & 0 < j < N \text{ and } 3 \mid j, \\ 0, & \text{otherwise,} \end{cases}$$

we obtain

$$(2.3) \quad C_{m,f}(u, v) = \frac{(u+v)_N}{N!} \mathbb{E}[J(N-J)\tilde{w}_J].$$

The factor $J(N-J)$ vanishes at $j=0$ and $j=N$, so the endpoint terms do not contribute.

Use the standard mixture representation $P \sim \text{Beta}(u, v)$ and $J \mid P \sim \text{Bin}(N, P)$. The identity

$$j(N-j) \binom{N}{j} = N(N-1) \binom{N-2}{j-1}$$

then transforms (2.3) into

$$(2.4) \quad C_{m,f}(u, v) = \frac{(u+v)_N}{N!} N(N-1) \mathbb{E} G_{m,w}(P),$$

where the kernel

$$(2.5) \quad G_{m,w}(p) = \sum_{r=1}^{m-1} w_r \binom{3m-2}{3r-1} p^{3r} (1-p)^{3(m-r)}, \quad 0 \leq p \leq 1,$$

is defined for arbitrary symmetric nonnegative weights $w_r = w_{m-r}$ and is symmetric under $p \mapsto 1-p$. The congruence $j \equiv 0 \pmod{3}$ becomes $j-1 \equiv 2 \pmod{3}$, which explains the binomial coefficient $\binom{3m-2}{3r-1}$.

By (2.4), Theorem 1.1 reduces to the monotonicity of $G_{m,w}$ on $[0, 1/2]$ (Section 4), via the imbalance ordering of Section 3.

3. TWO MONOTONICITY LEMMAS

This section records two elementary monotonicity facts: a weighting principle for the paired residue blocks of Section 4, due to [5, Lem. 6] and applied to multiplicity two in [7], and an ordering of beta laws by imbalance used in Section 5, a fixed-sum likelihood-ratio comparison for folded beta laws [3].

Lemma 3.1 (One-sign-change weighting). *Let $0 \leq w_1 \leq w_2 \leq \dots \leq w_L$ and suppose the real sequence $A_1, \dots, A_L$ has no more than one sign change, necessarily from nonpositive to nonnegative. If*

$$\sum_{k=1}^L A_k \geq 0,$$

then

$$\sum_{k=1}^L w_k A_k \geq 0.$$

Proof. The claim is immediate if all A_k are nonnegative. Otherwise, let q be the largest index for which $A_q < 0$. Put $c = w_q$. For $k \leq q$ we have $w_k \leq c$ and $A_k \leq 0$, while for $k > q$ we have $w_k \geq c$ and $A_k \geq 0$. Therefore

$$\sum_{k=1}^L w_k A_k \geq c \sum_{k=1}^L A_k \geq 0.$$

□

Lemma 3.2 (Symmetric beta imbalance ordering). *Fix $s > 0$. For $0 \leq d < s/2$, let $P_d \sim \text{Beta}(s/2 + d, s/2 - d)$. If $H : [0, 1] \rightarrow \mathbb{R}$ is symmetric, $H(p) = H(1 - p)$, and nondecreasing on $[0, 1/2]$, then*

$$(3.1) \quad d_1 \leq d_2 \implies \mathbb{E}H(P_{d_1}) \geq \mathbb{E}H(P_{d_2}).$$

If H is strictly increasing on $(0, 1/2)$ and $d_1 < d_2$, then the inequality is strict.

Proof. The case $d_1 = d_2$ is immediate, so assume $d_1 < d_2$. Put $Q_d = P_d(1 - P_d) \in (0, 1/4)$. For

$$p_{\pm}(q) = \frac{1 \pm \sqrt{1 - 4q}}{2}, \quad \ell(q) = \log \frac{p_+(q)}{p_-(q)},$$

the density of Q_d is

$$g_d(q) = \frac{2}{B(s/2 + d, s/2 - d)} q^{s/2-1} (1 - 4q)^{-1/2} \cosh(d\ell(q)).$$

The densities g_{d_1} and g_{d_2} share the factor $q^{s/2-1} (1 - 4q)^{-1/2}$, so

$$\frac{g_{d_2}(q)}{g_{d_1}(q)} = \frac{B(s/2 + d_1, s/2 - d_1)}{B(s/2 + d_2, s/2 - d_2)} \frac{\cosh(d_2\ell(q))}{\cosh(d_1\ell(q))}.$$

The function ℓ is strictly decreasing, since $p_+(q)/p_-(q)$ decreases to 1 as $q \uparrow 1/4$. For $d_2 > d_1$, $\cosh(d_2\ell)/\cosh(d_1\ell)$ is strictly increasing in $\ell > 0$, because $d \mapsto d \tanh(d\ell)$ is strictly increasing; hence g_{d_2}/g_{d_1} is strictly decreasing in q . Thus Q_{d_2} is smaller than Q_{d_1} in the likelihood-ratio order, and therefore in the usual stochastic order [11, Thm. 1.C.1]. Since both densities integrate to one and their ratio is strictly monotone, it crosses 1 exactly once; hence $g_{d_1} - g_{d_2}$ has a single sign change, negative near 0 and positive near $1/4$, so for every nondecreasing ϕ ,

$$\mathbb{E}\phi(Q_{d_1}) - \mathbb{E}\phi(Q_{d_2}) = \int_0^{1/4} (\phi(q) - \phi(q^*)) (g_{d_1}(q) - g_{d_2}(q)) dq \geq 0,$$

where q^* is the crossing point and the integrand is nonnegative throughout. Finally, symmetry of H gives $H(p) = \tilde{H}(p(1-p))$, with $\tilde{H}$ nondecreasing on $[0, 1/4]$ precisely when H is nondecreasing on $[0, 1/2]$; taking $\phi = \tilde{H}$ proves (3.1). If H is strictly increasing and $d_1 < d_2$, the likelihood ratio is nonconstant, the crossing is nondegenerate, and the inequality is strict. $\square$

4. THE CUBIC RESIDUE KERNEL

For symmetric nonnegative weights $w_r = w_{m-r}$, the kernel of (2.5) is

$$G_{m,w}(p) = \sum_{r=1}^{m-1} w_r \binom{3m-2}{3r-1} p^{3r} (1-p)^{3(m-r)}, \quad G_{m,w}(p) = G_{m,w}(1-p).$$

This section proves that $G_{m,w}$ is monotone on $[0, 1/2]$ whenever the weights increase toward the center.

Theorem 4.1 (Weighted cubic residue monotonicity). *Suppose*

$$(4.1) \quad 0 \leq w_1 \leq w_2 \leq \cdots \leq w_{\lfloor m/2 \rfloor}, \quad w_r = w_{m-r}.$$

Then $G_{m,w}$ is nondecreasing on $[0, 1/2]$. If the weights are not all zero, then $G_{m,w}$ is strictly increasing on $[0, 1/2]$.

4.1. The constant-weight kernel. Let $w_r \equiv 1$, and write $G_m = G_{m,1}$. Set $n = 3m - 2$ and $t = p/(1-p)$. For $0 \leq p \leq 1/2$ one has $0 \leq t \leq 1$, and

$$G_m\left(\frac{t}{1+t}\right) = \frac{1}{(1+t)^{3m}} \sum_{r=1}^{m-1} \binom{n}{3r-1} t^{3r}.$$

Differentiation gives

$$(4.2) \quad \frac{d}{dt} G_m\left(\frac{t}{1+t}\right) = \frac{3J_m(t)}{(1+t)^{3m+1}},$$

where

$$(4.3) \quad J_m(t) = \sum_{r=1}^{m-1} \binom{n}{3r-1} t^{3r-1} (r - (m-r)t).$$

Lemma 4.2 (Bernstein certificate). *For every $m \geq 2$,*

$$J_m(t) > 0 \quad (0 < t < 1).$$

Consequently, G_m is strictly increasing on $[0, 1/2]$.

Proof. Writing $k = 3r - 1$ in (4.3) gives

$$J_m(t) = \frac{1}{3} \sum_{\substack{0 \leq k \leq n \\ k \equiv 2(3)}} \binom{n}{k} t^k ((k+1) - (n-k+1)t).$$

Introduce the projective Bernstein transform

$$(4.4) \quad \mathcal{P}_m(x) = (1+x)^{n+1} J_m\left(\frac{x}{1+x}\right).$$

Then

$$(4.5) \quad \mathcal{P}_m(x) = \frac{1}{3} \sum_{k \equiv 2(3)} \binom{n}{k} x^k (1+x)^{n-k} ((k+1) + (2k-n)x).$$

For $0 \leq j \leq n+1$, coefficient extraction from (4.5) gives

$$(4.6) \quad \begin{aligned} [x^j] \mathcal{P}_m(x) &= \frac{1}{3} \sum_{\substack{0 \leq k \leq j \\ k \equiv 2(3)}} \binom{n}{k} \left[(k+1) \binom{n-k}{j-k} + (2k-n) \binom{n-k}{j-k-1} \right] \\ &= \frac{\binom{n+1}{j}}{3(n+1)} S_{n,j}, \end{aligned}$$

where

$$(4.7) \quad S_{n,j} = \sum_{\substack{0 \leq k \leq j \\ k \equiv 2(3)}} \binom{j}{k} (n-k+1)(2k+1-j).$$

The second equality in (4.6) follows from the elementary identities

$$\binom{n}{k} \binom{n-k}{j-k} = \binom{n}{j} \binom{j}{k}, \quad \binom{n}{k} \binom{n-k}{j-k-1} = \binom{n+1}{j} \binom{j}{k} \frac{j-k}{n+1}.$$

The transform is Bernstein-theoretic: writing $J_m(t) = \sum_j b_j \binom{n+1}{j} t^j (1-t)^{n+1-j}$ in the degree- $(n+1)$ Bernstein basis, one has $[x^j] \mathcal{P}_m(x) = b_j \binom{n+1}{j}$, so nonnegativity of the Bernstein coefficients b_j certifies $J_m(t) \geq 0$ on $[0, 1]$ [10]. We claim that every $S_{n,j}$ is nonnegative.

Put $d = n+1-j$. A third-root-of-unity filter evaluates the sum over the class $k \equiv 2 \pmod{3}$; its trigonometric terms are periodic modulo 6 in j , so the closed form splits into six residue classes,

$$(4.8) \quad 3S_{n,j} = \begin{cases} d(2^j - 3j - 1) - 3j(j-1), & j \equiv 0 \pmod{6}, \\ d(2^j - 2) - 3j(j-1), & j \equiv 1 \pmod{6}, \\ d(2^j + 3j - 1), & j \equiv 2 \pmod{6}, \\ d(2^j + 3j + 1) + 3j(j-1), & j \equiv 3 \pmod{6}, \\ d(2^j + 2) + 3j(j-1), & j \equiv 4 \pmod{6}, \\ d(2^j - 3j + 1), & j \equiv 5 \pmod{6}. \end{cases}$$

In detail, for

$$R_a(j) = \sum_{k \equiv a(3)} \binom{j}{k}$$

and $\omega = e^{2\pi i/3}$,

$$R_a(j) = \frac{1}{3} \sum_{\ell=0}^2 \omega^{-a\ell} (1 + \omega^\ell)^j = \frac{1}{3} \left(2^j + 2 \cos \frac{(j-2a)\pi}{3} \right),$$

where $2 \cos(x\pi/3)$ runs through $2, 1, -1, -2, -1, 1$ as $x \equiv 0, 1, 2, 3, 4, 5 \pmod{6}$. These values also follow with no appeal to $\mathbb{C}$: Pascal's rule gives $R_a(j+1) = R_a(j) + R_{a-1}(j)$ (subscripts modulo 3), and this recurrence with $R_0(0) = 1$, $R_1(0) = R_2(0) = 0$ fixes them by induction. The first two moments over the residue class are

$$\sum_{k \equiv 2(3)} k \binom{j}{k} = j R_1(j-1), \quad \sum_{k \equiv 2(3)} k^2 \binom{j}{k} = j(j-1) R_0(j-2) + j R_1(j-1).$$

Writing $d = n+1-j$ and expanding the summand in (4.7) gives, for $j \geq 2$,

$$(4.9) \quad \begin{aligned} S_{n,j} &= d(2j R_1(j-1) + (1-j) R_2(j)) \\ &\quad + j(j-1)(-2R_0(j-2) + 3R_1(j-1) - R_2(j)), \end{aligned}$$

with $j = 0, 1$ following directly from (4.7). Substituting the values of $R_2(j)$, $R_1(j-1)$, and $R_0(j-2)$ by residue $j \pmod{6}$ yields (4.8).

It remains only to inspect the six cases. The cases $j \equiv 2, 3, 4 \pmod{6}$ are immediate. If $j \equiv 5 \pmod{6}$, then either $d = 0$, or $j \geq 5$ and $2^j - 3j + 1 > 0$. If $j \equiv 1 \pmod{6}$, then $d \geq 1$; the case $j = 1$ gives zero, while for $j \geq 7$,

$$(4.10) \quad 2^j - 2 \geq 3j(j-1).$$

Finally, $n+1 = 3m-1$ is congruent to 2 or 5 modulo 6. Thus, when $j \equiv 0 \pmod{6}$ and $j > 0$, one has $d \geq 2$, and for $j \geq 6$,

$$(4.11) \quad 2(2^j - 3j - 1) \geq 3j(j-1).$$

The inequalities (4.10) and (4.11) hold with equality at $j = 7$ and $j = 6$, respectively. Advancing j by one increases the left-minus-right difference by $2^j - 6j$ in (4.10) and by $2^{j+1} - 6j - 6$ in (4.11), both positive for $j \geq 6$; the inequalities therefore persist for all larger j . Likewise $2^j - 3j + 1 > 0$ at $j = 5$, with increment $2^j - 3 > 0$ thereafter. Hence $S_{n,j} \geq 0$ in every case.

By (4.6), all coefficients of $\mathcal{P}_m$ are nonnegative. The coefficient at x^2 is positive, since $d = n-1 > 0$ and $3S_{n,2} = 9d$. Therefore $\mathcal{P}_m(x) > 0$ for $x > 0$. The substitution $t = x/(1+x)$ in (4.4) proves $J_m(t) > 0$ for $0 < t < 1$, and (4.2) completes the proof. $\square$

4.2. Insertion of log-concave weights. Arbitrary weights satisfying (4.1) preserve positivity of the derivative.

For $0 < t < 1$, differentiation of (2.5) gives

$$(4.12) \quad \frac{d}{dt} G_{m,w} \left(\frac{t}{1+t} \right) = \frac{3J_{m,w}(t)}{(1+t)^{3m+1}},$$

where

$$(4.13) \quad J_{m,w}(t) = \sum_{r=1}^{m-1} w_r \binom{3m-2}{3r-1} t^{3r-1} (r - (m-r)t).$$

Pair the terms indexed by r and $m-r$. For $1 \leq r < m/2$, let

$$(4.14) \quad \begin{aligned} B_{m,r}(t) &= \binom{3m-2}{3r-1} \left[t^{3r-1} (r - (m-r)t) \right. \\ &\quad \left. + t^{3(m-r)-1} ((m-r) - rt) \right]. \end{aligned}$$

If m is even, the central block is

$$(4.15) \quad B_{m,m/2}(t) = \binom{3m-2}{3m/2-1} \frac{m}{2} t^{3m/2-1} (1-t) > 0.$$

Then

$$J_{m,w}(t) = \sum_{r=1}^{\lfloor m/2 \rfloor} w_r B_{m,r}(t),$$

with the convention (4.15) at the center.

Lemma 4.3 (Single sign change of the paired blocks). *For each fixed $m \geq 2$ and $0 < t < 1$, the sequence*

$$B_{m,1}(t), B_{m,2}(t), \dots, B_{m,\lfloor m/2 \rfloor}(t)$$

has no more than one sign change, from negative to positive.

Proof. For $r < m/2$, the sign of $B_{m,r}(t)$ is the sign of

$$H_{m,r}(t) = r - (m-r)t + t^{3(m-2r)}((m-r) - rt).$$

Put $d = m - 2r > 0$ and $t = e^{-x}$, $x > 0$. Direct rearrangement gives

$$(4.16) \quad \begin{aligned} H_{m,r}(t) &= \frac{1}{2}(1+t)(1+t^{3d}) \left[m \frac{1-t}{1+t} - d \frac{1-t^{3d}}{1+t^{3d}} \right] \\ &= \frac{1}{2}(1+t)(1+t^{3d}) \left[m \tanh \frac{x}{2} - d \tanh \frac{3dx}{2} \right]. \end{aligned}$$

The function $d \mapsto d \tanh(3dx/2)$ is strictly increasing for $d > 0$. Therefore, for $2 \leq r < m/2$, if $B_{m,r}(t) \leq 0$, then the same bracket in (4.16) is strictly negative after replacing d by $d+2$, which corresponds to replacing r by $r-1$. Thus

$$B_{m,r}(t) \leq 0 \implies B_{m,r-1}(t) < 0 \quad (2 \leq r < m/2).$$

The final block is always positive. For even m this is (4.15). For odd m , the final block has $d = 1$, and its bracket in (4.16) is $m \tanh(x/2) - \tanh(3x/2) > 0$, because $m \geq 3$ and $\tanh(3y) < 3 \tanh y$ for $y > 0$. Hence the blocks have the asserted sign pattern. $\square$

Proof of Theorem 4.1. By Lemma 4.2,

$$\sum_{r=1}^{\lfloor m/2 \rfloor} B_{m,r}(t) = J_m(t) > 0 \quad (0 < t < 1).$$

The block sequence has one sign change by Lemma 4.3. Applying Lemma 3.1 with the nondecreasing weights in (4.1) gives

$$J_{m,w}(t) = \sum_{r=1}^{\lfloor m/2 \rfloor} w_r B_{m,r}(t) \geq 0.$$

Equation (4.12) proves that $G_{m,w}$ is nondecreasing on $[0, 1/2]$.

Suppose now that the weights are not all zero, and put $L = \lfloor m/2 \rfloor$. Then $w_L > 0$, and $B_{m,L}(t) > 0$ by Lemma 4.3. If every block is nonnegative, then $J_{m,w}(t) \geq w_L B_{m,L}(t) > 0$. Otherwise let q be the last index with $B_{m,q}(t) < 0$, and put $c = w_q$. The decomposition

$$\sum_{r=1}^L w_r B_{m,r} = c \sum_{r=1}^L B_{m,r} + \sum_{r \leq q} (w_r - c) B_{m,r} + \sum_{r > q} (w_r - c) B_{m,r}$$

displays the weighted sum as three nonnegative terms: the first because $\sum_r B_{m,r} = J_m(t) > 0$ and $c \geq 0$; the second because $w_r \leq c$ and $B_{m,r} \leq 0$ for $r \leq q$; the third because $w_r \geq c$ and $B_{m,r} \geq 0$ for $r > q$. If $c > 0$, the first term is strictly positive; if $c = 0$, the third term is strictly positive, because $q < L$ and it contains $(w_L - c) B_{m,L}(t) = w_L B_{m,L}(t) > 0$. Hence $J_{m,w}(t) > 0$ for $0 < t < 1$, and $G_{m,w}$ is strictly increasing on $[0, 1/2]$. $\square$

5. PROOF OF THE THEOREM

The representation (2.4) and the two monotonicity facts of Sections 3 and 4 combine to show that the coefficient convolution is Schur-concave in its two parameters [8, Def. 3.A.1 and Thm. 3.A.4] — equivalently, nonincreasing under increasing imbalance at fixed parameter sum — which gives Theorem 1.1.

Proposition 5.1 (Schur-concavity of the coefficient convolution). *For each $m \geq 2$ and each log-concave sequence (f_n) as in Theorem 1.1, the symmetric function $(u, v) \mapsto C_{m,f}(u, v)$ is Schur-concave on $(0, \infty)^2$. Equivalently, at fixed $s = u + v$,*

$$d \mapsto C_{m,f}\left(\frac{s}{2} + d, \frac{s}{2} - d\right)$$

is nonincreasing for $0 \leq d < s/2$. If at least one product $f_r f_{m-r}$ is positive, then this function is strictly decreasing.

Proof. The prefactor in (2.4) depends only on $s = u + v$. By Theorem 4.1, $G_{m,w}$ is symmetric and nondecreasing on $[0, 1/2]$. The conclusion follows immediately from Lemma 3.2. If at least one weight $w_r = f_r f_{m-r}$ is positive, then Theorem 4.1 makes $G_{m,w}$ strictly increasing, and the strict part of Lemma 3.2 gives strict decrease in d . $\square$

Proof of Theorem 1.1. The coefficients of degrees 0 and 1 in (1.3) vanish. For $m \geq 2$, direct multiplication gives

$$(5.1) \quad \begin{aligned} [x^m] \Delta_f(\mu; \alpha, \beta; x) &= C_{m,f}(\mu + \alpha, \mu + \beta) \\ &\quad - C_{m,f}(\mu, \mu + \alpha + \beta). \end{aligned}$$

Assume first that $\mu > 0$ and $\alpha, \beta > 0$. The two parameter pairs in (5.1) have the same sum

$$s = 2\mu + \alpha + \beta.$$

Their imbalances are

$$d_1 = \frac{|\alpha - \beta|}{2}, \quad d_2 = \frac{\alpha + \beta}{2},$$

and $d_1 \leq d_2$. Proposition 5.1 therefore gives

$$C_{m,f}(\mu + \alpha, \mu + \beta) \geq C_{m,f}(\mu, \mu + \alpha + \beta).$$

This proves nonnegativity of every coefficient. If $\alpha = 0$ or $\beta = 0$, the Turánian is identically zero. The boundary case $\mu = 0$ follows either by continuity in μ or directly from $F_f(0; x) = 0$ and the nonnegative coefficients of $F_f(\alpha; x)F_f(\beta; x)$. $\square$

Corollary 5.2 (Strict coefficient classification). *Under the hypotheses of Theorem 1.1, let $\mu \geq 0$, $\alpha, \beta > 0$, and $m \geq 2$. Then*

$$[x^m] \Delta_f(\mu; \alpha, \beta; x) > 0$$

if and only if there exists $1 \leq r \leq m - 1$ with $f_r f_{m-r} > 0$.

Proof. If every product $f_r f_{m-r}$ vanishes, then both terms of (5.1) vanish. Conversely, suppose one such product is positive. If $\mu > 0$, the two imbalances in the proof of Theorem 1.1 satisfy

$$\frac{|\alpha - \beta|}{2} < \frac{\alpha + \beta}{2},$$

so the strict part of Proposition 5.1 gives strict inequality. If $\mu = 0$, then $F_f(0; x) = 0$, so the coefficient equals $C_{m,f}(\alpha, \beta)$, a sum of nonnegative terms with at least one strictly positive. $\square$

Corollary 5.3 (Ordinary log-concavity). *Suppose (f_n) is not identically zero, and let R be the radius of convergence of $\sum_{n \geq 1} f_n x^n$. For every $0 < x < R$, the function $\mu \mapsto F_f(\mu; x)$ is finite, continuous, and positive on $(0, \infty)$, and $\mu \mapsto \log F_f(\mu; x)$ is concave there.*

Proof. For each fixed $\mu > 0$, Stirling's formula gives

$$\frac{(\mu)_{3n}}{(3n-1)!} = \frac{\Gamma(3n+\mu)}{\Gamma(\mu)\Gamma(3n)} \sim \frac{(3n)^\mu}{\Gamma(\mu)} \quad (n \rightarrow \infty).$$

Hence $F_f(\mu; \cdot)$ has the same radius of convergence R as $\sum_{n \geq 1} f_n x^n$. On every compact set of positive μ and for $0 < x < R$, the series converges locally uniformly in μ , since its extra factor grows at most polynomially in n ; thus $\mu \mapsto F_f(\mu; x)$ is continuous. Since (f_n) is not identically zero, its nonnegative coefficients give $F_f(\mu; x) > 0$ for $\mu, x > 0$. By Theorem 1.1,

$$F_f(\mu + \alpha; x)F_f(\mu + \beta; x) \geq F_f(\mu; x)F_f(\mu + \alpha + \beta; x)$$

for all nonnegative shifts. Taking $\alpha = \beta = h > 0$ gives

$$F_f(\mu + h; x)^2 \geq F_f(\mu; x)F_f(\mu + 2h; x),$$

so $\mu \mapsto \log F_f(\mu; x)$ is midpoint concave, and by continuity concave, on $(0, \infty)$. $\square$

6. CONCLUDING REMARKS

Remark 6.1 (Why the cubic case closes). The proof hinges on the special feature of the residue class modulo 3. After the beta-binomial reduction, the entire conjecture is controlled by the monotonicity of

$$p(1-p)\mathbb{P}\{\text{Bin}(3m-2, p) \equiv 2 \pmod{3}\}.$$

The projective transform (4.4) converts its derivative into a polynomial whose Bernstein coefficients are nonnegative. This positivity is not termwise obvious in the monomial basis; it appears only after the degree lift and the mod-6 root-of-unity evaluation.

Remark 6.2 (The role of log-concavity). The log-concavity assumption is used exactly once: it makes the symmetric weights $w_r = f_r f_{m-r}$ nondecreasing as r approaches $m/2$. The paired derivative blocks have a single sign change, so Lemma 3.1 transfers the constant-weight positivity certificate to the full weighted kernel.

Remark 6.3 (Higher multiplicities). For a general integer r , the same beta-binomial reduction produces, in the constant-weight case,

$$p(1-p)\mathbb{P}\{\text{Bin}(rm-2, p) \equiv r-1 \pmod{r}\},$$

and in general its analogue weighted by the products $f_k f_{m-k}$. The cubic proof suggests testing whether a suitable projective degree lift has a positive coefficient array. The known failure at $r = 4$ shows that such positivity cannot persist without an additional restriction; locating the first coefficient obstruction in the lifted array may explain the failure at $r = 4$.

ACKNOWLEDGEMENTS

During the preparation of this work, the author used OpenAI's ChatGPT (GPT-5.6 Sol) and Anthropic's Claude (Claude Opus 4.8) to assist with manuscript drafting and the development of auxiliary verification code.

STATEMENTS AND DECLARATIONS

Funding. This research received no external funding.

Competing interests. The author has no relevant interests to disclose.

Data availability. No datasets were generated or analysed.

Computational verification. The proofs do not depend on numerical computation. Verification scripts are available at <https://github.com/johnnyshields/papers>.

REFERENCES

- [1] Á. Baricz, *Turán type inequalities for hypergeometric functions*, Proc. Amer. Math. Soc. **136** (2008), no. 9, 3223–3229, DOI: 10.1090/S0002-9939-08-09353-2.
- [2] R. W. Barnard, M. B. Gordy, and K. C. Richards, *A note on Turán type and mean inequalities for the Kummer function*, J. Math. Anal. Appl. **349** (2009), no. 1, 259–263, DOI: 10.1016/j.jmaa.2008.08.024.
- [3] K. S. Berenhaut and L. D. Bergen, *Stochastic orderings, folded beta distributions and fairness in coin flips*, Statist. Probab. Lett. **81** (2011), no. 6, 632–638, DOI: 10.1016/j.spl.2011.01.002.
- [4] S. I. Kalmykov and D. B. Karp, *Log-concavity for series in reciprocal gamma functions and applications*, Integral Transforms Spec. Funct. **24** (2013), no. 11, 859–872, DOI: 10.1080/10652469.2013.764874.
- [5] S. I. Kalmykov and D. B. Karp, *Log-convexity and log-concavity for series in gamma ratios and applications*, J. Math. Anal. Appl. **406** (2013), no. 2, 400–418, DOI: 10.1016/j.jmaa.2013.04.061.
- [6] D. Karp and S. M. Sitnik, *Log-convexity and log-concavity of hypergeometric-like functions*, J. Math. Anal. Appl. **364** (2010), no. 2, 384–394, DOI: 10.1016/j.jmaa.2009.10.057.
- [7] D. Karp and Y. Zhang, *Log-concavity and log-convexity of series containing multiple Pochhammer symbols*, Fract. Calc. Appl. Anal. **27** (2024), no. 1, 458–486, DOI: 10.1007/s13540-023-00238-0.
- [8] A. W. Marshall, I. Olkin, and B. C. Arnold, *Inequalities: Theory of Majorization and Its Applications*, 2nd ed., Springer Series in Statistics, Springer, New York, 2011, DOI: 10.1007/978-0-387-68276-1.
- [9] A. Ninh and M. Pham, *Logconcavity, twice-logconcavity and Turán-type inequalities*, Ann. Oper. Res. **354** (2025), no. 3, 1271–1283, DOI: 10.1007/s10479-018-2923-y.
- [10] V. Powers and B. Reznick, *Polynomials that are positive on an interval*, Trans. Amer. Math. Soc. **352** (2000), no. 10, 4677–4692, DOI: 10.1090/S0002-9947-00-02595-2.
- [11] M. Shaked and J. G. Shanthikumar, *Stochastic Orders*, Springer Series in Statistics, Springer, New York, 2007, DOI: 10.1007/978-0-387-34675-5.