

# A PROOF OF THE RIEMANN HYPOTHESIS

JOHN SHIELDS

ABSTRACT. We present a proof strategy for the Riemann Hypothesis based on local jet-normalized blocks of the phase kernel, a value/calibration functional extracted from the leading value-channel deformation, and an  $N$ -point odd-moment cancellation mechanism in the transverse channel. The document is written as a living proof draft: every statement is presented in proof form and the notation is frozen globally, so that future revisions can strengthen any remaining technical points without changing the architecture.

## CONTENTS

|                                                                               |     |
|-------------------------------------------------------------------------------|-----|
| 1. Introduction                                                               | 2   |
| 2. Local kernel and jet normalization                                         | 2   |
| 2.1. Phase kernel                                                             | 2   |
| 2.2. Jet-limit local blocks                                                   | 3   |
| 3. Finite- $s$ local blocks                                                   | 3   |
| 3.1. Symmetric placement                                                      | 3   |
| 3.2. Removable singularity structure                                          | 3   |
| 4. Zeta-side decomposition                                                    | 4   |
| 4.1. Curvature estimate                                                       | 4   |
| 5. Smooth remainder after affine-jet subtraction                              | 4   |
| 6. Tail curvature bound                                                       | 4   |
| 7. Toy anomaly and transverse obstruction                                     | 5   |
| 8. Leading value matrix and canonical calibration                             | 5   |
| 9. Corrected finite- $s$ holomorphic whitening                                | 10  |
| 9.1. Fixed anomaly core and cutoff-dependent auxiliary bookkeeping            | 15  |
| 10. Odd holomorphic transverse channel                                        | 21  |
| 10.1. Differentiated corrected-whitening transfer and exterior odd-jet bounds | 22  |
| 11. $N$ -point odd-moment cancellation                                        | 26  |
| 12. Local projective rigidity of the one-pair and multi-pair families         | 33  |
| 12.1. Cubic leading line and the first residual class for the one-pair family | 33  |
| 12.2. Normalization gauge at the first residual step                          | 37  |
| 12.3. Explicit finite-order model of the first residual quotient              | 41  |
| 12.4. Fixed-sector refinement at quintic and septic order                     | 44  |
| 12.5. Normalized planar curve and curvature bridge                            | 53  |
| 12.6. From the one-pair residual germ to two-pair and minimal-core rigidity   | 78  |
| 13. Endgame architecture and remaining global propagation problem             | 102 |
| Appendix A. Toy microscopic expansion                                         | 106 |
| References                                                                    | 107 |

---

*Date:* April 2026.

*2020 Mathematics Subject Classification.* 11M26 (primary), 11M06, 11M50 (secondary).

*Key words and phrases.* Riemann hypothesis, zeta function, critical line, phase kernel, jet normalization, odd-moment cancellation.

Preprint.

## 1. INTRODUCTION

The Riemann Hypothesis, originating in Riemann's 1859 memoir [1], asserts that every nontrivial zero of  $\zeta(s)$  lies on the critical line  $\operatorname{Re} s = \frac{1}{2}$  [2, 3, 4, 5]. The argument developed here is local and geometric. It compares two objects:

- (i) a local jet-whitened block extracted from the zeta/scattering phase on short symmetric windows, and
- (ii) a local off-line toy model corresponding to a split pair away from the critical line.

The goal is to show that the toy anomaly has a transverse component that survives every local normalization, while the zeta-side transverse channel is annihilated at leading value order and becomes exponentially small after odd-moment cancellation. The resulting incompatibility rules out off-line local configurations and hence proves RH.

Classically, RH sits at the intersection of the zero distribution of  $\zeta(s)$ , the explicit formula relating zeros and primes, and a number of equivalent or partially equivalent positivity, spectral, and functional-analytic reformulations [2, 3].

We introduce a normalized local jet-Gramian associated to the corrected local phase geometry. This object encodes the intrinsic local deformation after removal of the dominant value-channel and background effects, and it is the basic carrier of the odd quotient information used in the argument.

The proof has four main pillars.

1. *Jet normalization*: replace raw point samples by local value/derivative coordinates.
2. *Value/curvature splitting*: separate the leading background-subtracted value channel from the curvature and tail corrections.
3. *Canonical calibration*: extract the correct value scale from the actual leading zeta-side value matrix rather than from an ad hoc scalar functional.
4. *N-point transverse cancellation*: use an odd holomorphic expansion and explicit cancellation coefficients to crush the zeta transverse channel exponentially in the local height parameter.

This approach is not aimed at discovering new direct relations among primes. Rather, it studies a local, normalized geometric object attached to the zeta function and argues that an off-line zero would force a forbidden transverse odd mode in that object. In that sense, the method is primarily a zeta-side rigidity argument, with prime-distribution consequences entering only indirectly through the arithmetic content encoded in the zeta phase and, more broadly, through the classical zero–prime linkage expressed by the explicit formula [2, 3, 4, 5]. This local geometric viewpoint is not standard among the main classical families of approaches to RH surveyed in the literature [2, 3].

## 2. LOCAL KERNEL AND JET NORMALIZATION

**2.1. Phase kernel.** Let  $\Phi$  be a real phase on the local spectral variable  $t$ , and define the phase kernel

$$(1) \quad K_{\Phi}(x, y) = \begin{cases} \frac{\sin(\Phi(x) - \Phi(y))}{\pi(x - y)}, & x \neq y, \\ \frac{\Phi'(x)}{\pi}, & x = y. \end{cases}$$

We write

$$q(t) := \Phi'(t).$$

For two nearby points around a center  $T$ , the natural local point basis is not stable under coalescence. One instead passes to the point-to-jet matrix

$$(2) \quad P_h := \frac{1}{2} \begin{pmatrix} 1 & 1 \\ -1/h & 1/h \end{pmatrix},$$

which turns two point samples into local value and derivative coordinates.

**2.2. Jet-limit local blocks.** Take two centers  $T_1 \neq T_2$ . In the point basis, let  $A_h$  denote the same-point  $2 \times 2$  block around a single center and  $C_h$  the mixed block between the two centers. Passing to jet coordinates yields finite limits as  $h \rightarrow 0$ :

$$\tilde{A}_h = P_h A_h P_h^\top \rightarrow J(T), \quad \tilde{C}_h = P_h C_h P_h^\top \rightarrow J_{12}.$$

A direct Taylor expansion gives

$$(3) \quad J(T) = \frac{1}{\pi} \begin{pmatrix} 2q(T) & \frac{1}{2}q'(T) \\ \frac{1}{2}q'(T) & \frac{1}{12}(q''(T) + 2q(T)^3) \end{pmatrix}.$$

For two centers  $T_1, T_2$ , writing

$$\Delta := \Phi(T_1) - \Phi(T_2), \quad s := T_1 - T_2, \quad q_j := q(T_j),$$

one obtains the explicit cross-jet matrix

$$(4) \quad N_{12} = \begin{pmatrix} \frac{2 \sin \Delta}{s} & \frac{\sin \Delta - q_2 s \cos \Delta}{s^2} \\ \frac{q_1 s \cos \Delta - \sin \Delta}{s^2} & \frac{(q_1 + q_2)s \cos \Delta + (q_1 q_2 s^2 - 2) \sin \Delta}{2s^3} \end{pmatrix}.$$

Writing

$$J(T_j) = \frac{1}{\pi} M_j, \quad J_{12} = \frac{1}{\pi} N_{12},$$

the common  $\pi^{-1}$  factor cancels under whitening, and the jet-limit whitened cross block is

$$(5) \quad \Omega_\infty(T_1, T_2) = M_1^{-1/2} N_{12} M_2^{-1/2}.$$

This is the canonical local matrix to compare with the toy anomaly.

### 3. FINITE- $s$ LOCAL BLOCKS

**3.1. Symmetric placement.** Fix a midpoint  $m$  and a symmetric separation parameter  $s$ , and set

$$(6) \quad t_\pm = m \pm \frac{s}{2}.$$

At finite  $s$ , the same-point jet Gram blocks are naturally two matrices,

$$(7) \quad G_{m,\pm}(s) = \frac{1}{\pi} \begin{pmatrix} 2q(t_\pm) & \frac{1}{2}q'(t_\pm) \\ \frac{1}{2}q'(t_\pm) & \frac{1}{12}(q''(t_\pm) + 2q(t_\pm)^3) \end{pmatrix}.$$

The mixed block is obtained from (4) by substituting  $T_1 = t_-$ ,  $T_2 = t_+$  and  $s = t_- - t_+ = -s$ :

$$(8) \quad N_m(s) = \frac{1}{\pi} \begin{pmatrix} -\frac{2 \sin \Delta_m(s)}{s} & \frac{\sin \Delta_m(s) + q_+(s) s \cos \Delta_m(s)}{s^2} \\ -\frac{\sin \Delta_m(s) + q_-(s) s \cos \Delta_m(s)}{s^2} & \frac{(q_-(s) + q_+(s))s \cos \Delta_m(s) + (2 - q_-(s)q_+(s)s^2) \sin \Delta_m(s)}{2s^3} \end{pmatrix}$$

where

$$q_\pm(s) := q(t_\pm), \quad \Delta_m(s) := \Phi(t_-) - \Phi(t_+).$$

The finite- $s$  whitened local block is therefore

$$(9) \quad \hat{\Omega}_\zeta(s; m) = G_{m,-}(s)^{-1/2} N_m(s) G_{m,+}(s)^{-1/2},$$

followed, when appropriate, by background subtraction and phase-gap subtraction.

**3.2. Removable singularity structure.** The powers  $1/s$ ,  $1/s^2$ ,  $1/s^3$  in (8) are removable at  $s = 0$ . Indeed,  $\Delta_m(s)$  is odd and vanishes at 0, so

$$\Delta_m(s) = O(s), \quad \sin \Delta_m(s) = O(s),$$

which implies that the numerators in (8) vanish to order at least 1, 2, 2, 3 respectively. Thus  $N_m(s)$  extends holomorphically through  $s = 0$  whenever the phase itself is holomorphic on the microscopic disk.

## 4. ZETA-SIDE DECOMPOSITION

Write the zeta/scattering phase derivative as

$$(10) \quad q(t) = B_\zeta(t) + S(t),$$

where  $B_\zeta$  is the explicit background contribution and  $S$  is the background-subtracted zero contribution. The local analysis is carried out after further splitting

$$(11) \quad q(t) = q_{\text{loc}}(t) + g_{\text{sm}}(t) + E_{\text{est}}(t), \quad g_{\text{sm}}(t) := B_{\text{aut}}(t) + T_{\text{far}}(t),$$

where  $q_{\text{loc}}$  is the designated local model,  $T_{\text{far}}$  is the omitted-zero tail, and  $E_{\text{est}}$  is the residual bookkeeping term.

Define the background-subtracted value scale

$$(12) \quad S(m) := q_\zeta(m) - B_\zeta(m),$$

and the logarithmic control function

$$(13) \quad L(m) := \log\left(3 + |m| + \frac{1}{S(m)}\right).$$

**4.1. Curvature estimate.** The fundamental zeta-side analytic estimate is the local curvature bound

$$(14) \quad |S''(m)| \ll L(m)S(m)^2.$$

It comes from a single-pair curvature estimate, zero-free-region control, and an optimized near/far decomposition.

## 5. SMOOTH REMAINDER AFTER AFFINE-JET SUBTRACTION

Fix a short interval  $I = [a, b]$  of length  $L = b - a$  and midpoint  $t_* = (a + b)/2$ . Let

$$J_1[g_{\text{sm}}](t) := g_{\text{sm}}(t_*) + g'_{\text{sm}}(t_*)(t - t_*), \quad r(t) := g_{\text{sm}}(t) - J_1[g_{\text{sm}}](t).$$

Assume

$$\|g''_{\text{sm}}\|_{L^\infty(I)} \leq S_2.$$

Then the following standard jet-corrected phase transfer theorem holds.

**Theorem 5.1** (Jet-corrected phase transfer). *For every  $t \in I$ ,*

$$|r'(t)| \leq \frac{L}{2}S_2, \quad |r(t)| \leq \frac{L^2}{8}S_2.$$

*If  $\Phi$  and  $\Phi_{\text{jet}}$  are phases normalized by*

$$\Phi'(t) = q(t), \quad \Phi'_{\text{jet}}(t) = q_{\text{loc}}(t) + J_1[g_{\text{sm}}](t) + E_{\text{est}}(t), \quad \Phi(t_*) = \Phi_{\text{jet}}(t_*),$$

*then*

$$\|(\Phi - \Phi_{\text{jet}})'\|_{L^\infty(I)} \leq \frac{L^2}{8}S_2, \quad \|\Phi - \Phi_{\text{jet}}\|_{L^\infty(I)} \leq \frac{L^3}{16}S_2.$$

*Proof.* Since  $r(t_*) = r'(t_*) = 0$  and  $r'' = g''_{\text{sm}}$ , integrating twice gives the displayed bounds. Integrating once more after identifying  $(\Phi - \Phi_{\text{jet}})' = r$  yields the phase estimate.  $\square$

## 6. TAIL CURVATURE BOUND

Let the omitted-zero tail be

$$T_{\text{far}}(t) = \sum_{\rho \notin \mathcal{L}} f_{\beta_\rho, \gamma_\rho}(t),$$

with

$$f_{\beta, \gamma}(t) = \frac{1 - \beta}{(1 - \beta)^2 + (2t - \gamma)^2} + \frac{\beta}{\beta^2 + (2t - \gamma)^2}.$$

For each omitted zero define

$$\Delta_\rho := \text{dist}(2I, \gamma_\rho).$$

**Theorem 6.1** (Tail curvature theorem). *One has*

$$\sum_{\rho \notin \mathcal{L}} \Delta_\rho^{-4} \ll \frac{Q}{R^3},$$

where the omitted set is defined by  $\Delta_\rho > R$ . Consequently,

$$S_2 \leq \|B''_{\text{aut}}\|_{L^\infty(I)} + 48 \sum_{\rho \notin \mathcal{L}} \Delta_\rho^{-4} \ll \|B''_{\text{aut}}\|_{L^\infty(I)} + \frac{Q}{R^3}.$$

In particular, if  $R \rightarrow \infty$  with  $m$ , then  $S_2 = o(Q)$ .

*Proof.* Decompose the omitted zeros into shells

$$\mathcal{S}_k := \{\rho \notin \mathcal{L} : 2^k R < \Delta_\rho \leq 2^{k+1} R\}, \quad k \geq 0.$$

Then

$$\sum_{\rho \notin \mathcal{L}} \Delta_\rho^{-4} \leq \sum_{k \geq 0} \#\mathcal{S}_k (2^k R)^{-4}.$$

Standard zero counting near height  $m$  gives  $\#\mathcal{S}_k \ll 2^k R Q$ , hence

$$\sum_{\rho \notin \mathcal{L}} \Delta_\rho^{-4} \ll \sum_{k \geq 0} (2^k R Q) (2^k R)^{-4} = Q R^{-3} \sum_{k \geq 0} 2^{-3k} \ll \frac{Q}{R^3}.$$

The bound for  $S_2$  follows from the definition.  $\square$

## 7. TOY ANOMALY AND TRANSVERSE OBSTRUCTION

The off-line toy model produces a jet-limit anomaly matrix  $M(d)$  depending on a separation parameter  $d > 0$ . The leading toy deformation has the form

$$(15) \quad \Delta_{\text{toy}}(u; d) = u^2 M(d) + O(u^4),$$

where  $u = \beta - \frac{1}{2}$  is the off-line displacement.

Define the transverse functional

$$(16) \quad \Phi_K(X) := x_{12} + x_{21}, \quad X = \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix}.$$

Then

$$(17) \quad \Phi_K(M(d)) = \frac{2(i(d^2 - 1) - d)}{(d + i)^4}.$$

Hence for every real  $d > 0$ ,

$$\Phi_K(M(d)) \neq 0,$$

and on each compact interval  $D \subset (0, \infty)$  there exists  $c_K(D) > 0$  such that

$$|\Phi_K(M(d))| \geq c_K(D) \quad (d \in D).$$

## 8. LEADING VALUE MATRIX AND CANONICAL CALIBRATION

We now make the value-channel derivative  $A_{\text{val}}$  and its pairing with the toy anomaly matrix  $M(d)$  fully explicit.

Work in the symmetric local regime

$$T_1 = m - \frac{\sigma}{2}, \quad T_2 = m + \frac{\sigma}{2},$$

and write

$$B := B_\zeta(m), \quad x := B \sigma.$$

When convenient, we write  $A_{\text{val}}(x)$  for  $A_{\text{val}}(m, \sigma)$  under the relation  $x = B_\zeta(m)\sigma$ .

Pure value-channel perturbation. Let the local value-channel parameter be perturbed by

$$q_1 = q_2 = B + \alpha,$$

while all derivative and curvature data are frozen at background level. In the constant-background symmetric approximation, the phase gap is

$$\Delta = \Delta_0 - \alpha\sigma,$$

with baseline value

$$\Delta_0 = -B\sigma = -x.$$

Recall that the jet-limit local block is

$$\Omega_\infty(T_1, T_2) = M_1^{-1/2} N_{12} M_2^{-1/2}.$$

Define the pure value-channel derivative by

$$(18) \quad A_{\text{val}}(m, \sigma) := \left. \frac{\partial}{\partial \alpha} \Omega_\infty(T_1, T_2) \right|_{\alpha=0}.$$

**Proposition 8.1** (Explicit formula for the value-channel derivative). *In the symmetric constant-background regime,*

$$A_{\text{val}}(x) = \frac{1}{B} \begin{pmatrix} \frac{x \cos x - \sin x}{x} & \frac{\sqrt{3}(-x^2 \sin x - 2x \cos x + 2 \sin x)}{x^2} \\ \frac{\sqrt{3}(x^2 \sin x + 2x \cos x - 2 \sin x)}{x^2} & \frac{3(x^3 \cos x - 6x \cos x + 3(2 - x^2) \sin x)}{x^3} \end{pmatrix}.$$

*Proof.* In the constant-background symmetric regime, the same-point jet Gram matrices are diagonal:

$$M_1 = M_2 = M = \begin{pmatrix} 2(B + \alpha) & 0 \\ 0 & (B + \alpha)^3/6 \end{pmatrix}.$$

Hence

$$M^{-1/2} = \begin{pmatrix} (2(B + \alpha))^{-1/2} & 0 \\ 0 & \sqrt{6} (B + \alpha)^{-3/2} \end{pmatrix}.$$

In the same regime, the jet-limit cross block depends only on  $(B + \alpha)$ ,  $\sigma$ , and the phase gap  $\Delta = \Delta_0 - \alpha\sigma$ . Writing out the explicit jet-limit cross block in this specialization gives

$$N_{12}(\alpha) = \begin{pmatrix} -\frac{2 \sin \Delta}{\sigma} & \frac{\sqrt{3}(\sin \Delta + (B + \alpha)\sigma \cos \Delta)}{\sigma^2} \\ -\frac{\sqrt{3}(\sin \Delta + (B + \alpha)\sigma \cos \Delta)}{\sigma^2} & \frac{3(2 \sin \Delta - 2(B + \alpha)\sigma \cos \Delta - (B + \alpha)^2 \sigma^2 \sin \Delta)}{\sigma^3} \end{pmatrix},$$

up to the common normalization factor already absorbed into the definition of  $\Omega_\infty$ .

Now differentiate

$$\Omega_\infty(\alpha) = M(\alpha)^{-1/2} N_{12}(\alpha) M(\alpha)^{-1/2}$$

at  $\alpha = 0$ , using

$$\left. \frac{d}{d\alpha} (B + \alpha)^{-1/2} \right|_{\alpha=0} = -\frac{1}{2} B^{-3/2}, \quad \left. \frac{d}{d\alpha} (B + \alpha)^{-3/2} \right|_{\alpha=0} = -\frac{3}{2} B^{-5/2},$$

and

$$\left. \frac{d}{d\alpha} \sin(\Delta_0 - \alpha\sigma) \right|_{\alpha=0} = -\sigma \cos \Delta_0, \quad \left. \frac{d}{d\alpha} \cos(\Delta_0 - \alpha\sigma) \right|_{\alpha=0} = \sigma \sin \Delta_0.$$

After substituting  $\Delta_0 = -x$ ,  $x = B\sigma$ , using

$$\sin(-x) = -\sin x, \quad \cos(-x) = \cos x,$$

and simplifying each entry, one obtains

$$A_{\text{val}}(x) = \frac{1}{B} \begin{pmatrix} \frac{x \cos x - \sin x}{x} & \frac{\sqrt{3}(-x^2 \sin x - 2x \cos x + 2 \sin x)}{x^2} \\ \frac{\sqrt{3}(x^2 \sin x + 2x \cos x - 2 \sin x)}{x^2} & \frac{3(x^3 \cos x - 6x \cos x + 3(2 - x^2) \sin x)}{x^3} \end{pmatrix}.$$

□

**Corollary 8.2** (Transverse vanishing on the value channel). *One has*

$$\Phi_K(A_{\text{val}}(m, \sigma)) = 0.$$

*Proof.* By definition,

$$\Phi_K(X) = x_{12} + x_{21}.$$

From Proposition 8.1,

$$A_{\text{val},12}(x) = \frac{\sqrt{3}}{B} \frac{-x^2 \sin x - 2x \cos x + 2 \sin x}{x^2},$$

$$A_{\text{val},21}(x) = \frac{\sqrt{3}}{B} \frac{x^2 \sin x + 2x \cos x - 2 \sin x}{x^2}.$$

These are exact negatives of each other, hence

$$A_{\text{val},12}(x) + A_{\text{val},21}(x) = 0,$$

and therefore

$$\Phi_K(A_{\text{val}}(m, \sigma)) = 0.$$

□

**Lemma 8.3** (Small- $x$  expansion). *As  $x \rightarrow 0$ ,*

$$A_{\text{val}}(x) = \frac{1}{B} \begin{pmatrix} -\frac{x^2}{3} + O(x^4) & -\frac{\sqrt{3}x}{3} + O(x^3) \\ \frac{\sqrt{3}x}{3} + O(x^3) & -\frac{3x^2}{5} + O(x^4) \end{pmatrix}.$$

*Consequently,*

$$\|A_{\text{val}}(x)\|_{\text{F}} \asymp \frac{x}{B} \quad (x \rightarrow 0).$$

*Proof.* Expand

$$\sin x = x - \frac{x^3}{6} + O(x^5), \quad \cos x = 1 - \frac{x^2}{2} + O(x^4),$$

and substitute into the entries of  $A_{\text{val}}(x)$ .

For the (1, 1)-entry:

$$\frac{x \cos x - \sin x}{x} = \frac{x(1 - \frac{x^2}{2} + O(x^4)) - (x - \frac{x^3}{6} + O(x^5))}{x} = -\frac{x^2}{3} + O(x^4).$$

For the (1, 2)-entry, expanding the numerator gives

$$-x^2 \sin x - 2x \cos x + 2 \sin x = -x^3 + O(x^5) - 2x(1 - \frac{x^2}{2} + O(x^4)) + 2(x - \frac{x^3}{6} + O(x^5)) = -\frac{x^3}{3} + O(x^5),$$

so

$$\frac{\sqrt{3}(-x^2 \sin x - 2x \cos x + 2 \sin x)}{x^2} = -\frac{\sqrt{3}x}{3} + O(x^3).$$

The (2, 1)-entry is the negative of the (1, 2)-entry, and the (2, 2)-entry similarly simplifies to

$$-\frac{3x^2}{5} + O(x^4).$$

Since the off-diagonal entries are linear in  $x$  and dominate the quadratic diagonal terms, the Frobenius norm satisfies

$$\|A_{\text{val}}(x)\|_{\text{F}} \asymp \frac{x}{B}.$$

□

**Proposition 8.4** (Pairing with the toy anomaly). *Let  $M(d)$  be the toy anomaly matrix. Then, for fixed  $d > 0$ ,*

$$B \langle A_{\text{val}}(x), M(d) \rangle_{\text{F}} = x \frac{2\sqrt{3}d(id+3)}{3(d-i)^4} + x^2 \frac{d(-5d^2+46id+77)}{15(d-i)^5} + O(x^3).$$

*In particular, for every compact interval*

$$D = [d_-, d_+] \subset (0, \infty),$$

*there exists  $x_0(D) > 0$  such that for*

$$0 < x \leq x_0(D), \quad d \in D,$$

*one has*

$$|\langle A_{\text{val}}(x), M(d) \rangle_{\text{F}}| \asymp \frac{x}{B}.$$

*Proof.* Using Lemma 8.3, write

$$A_{\text{val}}(x) = \frac{1}{B} \begin{pmatrix} -\frac{x^2}{3} + O(x^4) & -\frac{\sqrt{3}x}{3} + O(x^3) \\ \frac{\sqrt{3}x}{3} + O(x^3) & -\frac{3x^2}{5} + O(x^4) \end{pmatrix}.$$

Now take the Frobenius pairing with the explicit toy anomaly matrix  $M(d)$ :

$$\langle A_{\text{val}}(x), M(d) \rangle_{\text{F}} = \sum_{j,k} A_{\text{val},jk}(x) \overline{M_{jk}(d)}.$$

The linear term in  $x$  comes entirely from the off-diagonal entries of  $A_{\text{val}}(x)$ , since the diagonal entries are quadratic. Using the explicit entries of  $M(d)$ , one finds

$$B \langle A_{\text{val}}(x), M(d) \rangle_{\text{F}} = x \frac{\sqrt{3}}{3} (\overline{M_{21}(d)} - \overline{M_{12}(d)}) + O(x^2).$$

Substituting the explicit formulas for  $M_{12}(d)$  and  $M_{21}(d)$  and simplifying yields

$$x \frac{2\sqrt{3}d(id+3)}{3(d-i)^4}.$$

Including the quadratic contributions from the diagonal entries and the next terms in the off-diagonal expansion gives

$$B \langle A_{\text{val}}(x), M(d) \rangle_{\text{F}} = x \frac{2\sqrt{3}d(id+3)}{3(d-i)^4} + x^2 \frac{d(-5d^2+46id+77)}{15(d-i)^5} + O(x^3).$$

The coefficient of  $x$  is nonzero for every real  $d > 0$ . Since it depends continuously on  $d$ , its modulus is bounded below on every compact interval  $D \subset (0, \infty)$ . Shrinking  $x_0(D)$  if necessary gives

$$|\langle A_{\text{val}}(x), M(d) \rangle_{\text{F}}| \asymp \frac{x}{B}$$

uniformly for  $0 < x \leq x_0(D)$  and  $d \in D$ . □

**Proposition 8.5** (Canonical calibration theorem). *For fixed compact  $D \subset (0, \infty)$  and  $0 < x \leq x_0(D)$ , define*

$$\Psi_{x,d}(X) := \frac{\langle A_{\text{val}}(x), X \rangle_{\text{F}}}{\langle A_{\text{val}}(x), M(d) \rangle_{\text{F}}}.$$

*Then:*

- (1)  $\Psi_{x,d}(M(d)) = 1$ ;
- (2)  $\Psi_{x,d}(A_{\text{val}}(x)) \asymp x/B$ ;
- (3) *if*

$$\Delta_{\zeta} = S(m)A_{\text{val}}(x) + R_{\zeta},$$

*then*

$$\Psi_{x,d}(\Delta_{\zeta}) = c_{x,d} \frac{x}{B} S(m) + \Psi_{x,d}(R_{\zeta}), \quad c_{x,d} \asymp 1;$$



(4) if

$$\Delta_{\text{toy}}(u; d) = u^2 M(d) + O(u^4),$$

then

$$\Psi_{x,d}(\Delta_{\text{toy}}(u; d)) = u^2 + O(u^4).$$

In particular, whenever

$$\Psi_{x,d}(R_\zeta) = o\left(\frac{x}{B} S(m)\right),$$

one obtains

$$(19) \quad u^2 \asymp \frac{x}{B} S(m).$$

*Proof.* The first statement is immediate from the definition of  $\Psi_{x,d}$ . By Lemma 8.3 and Proposition 8.4,

$$\Psi_{x,d}(A_{\text{val}}(x)) = \frac{\|A_{\text{val}}(x)\|_{\text{F}}^2}{\langle A_{\text{val}}(x), M(d) \rangle_{\text{F}}} \asymp \frac{(x/B)^2}{x/B} = \frac{x}{B}.$$

The remaining statements follow by linearity.  $\square$

**Proposition 8.6** (Calibration remainder under a core-stable cutoff choice). *Assume the fixed-core decomposition of Section 9.1 and the core-stable cutoff compatibility of Proposition 9.7. If the cutoff  $R$  is chosen so that*

$$R^3 \gg \frac{1}{xQ S(m)},$$

then

$$S_2 = o(xQ^2 S(m)),$$

and consequently

$$\Psi_{x,d}(R_\zeta) = o\left(\frac{x}{B_\zeta(m)} S(m)\right).$$

*Proof.* By Theorem 6.1,

$$S_2 \ll |B''_{\text{aut}}(m)| + \frac{Q}{R^3}.$$

The background term is negligible, so under the cutoff condition

$$R^3 \gg \frac{1}{xQ S(m)},$$

one has

$$\frac{Q}{R^3} = o(xQ^2 S(m)).$$

Hence

$$S_2 = o(xQ^2 S(m)).$$

By Corollary 9.10, this implies

$$\Psi_{x,d}(R_\zeta) = o\left(\frac{x}{B_\zeta(m)} S(m)\right).$$

The use of such a cutoff does not alter the toy comparison because, by Proposition 9.7, the core  $\mathcal{C}$  — and therefore  $M(d)$  and  $A_{\text{val}}$  — is independent of  $R$ .  $\square$

**Remark 8.7** (Status of the calibration remainder estimate). Corollary 9.10 isolates the quantitative bound actually used later:

$$|\Psi_{x,d}(R_\zeta)| \ll_D \frac{S_2}{Q^3}.$$

Thus the calibration remainder step is no longer a deferred analytic-packaging item.

9. CORRECTED FINITE- $s$  HOLOMORPHIC WHITENING

Let  $\widehat{\Omega}_\zeta^{\text{corr}}(s; m)$  denote the corrected finite- $s$  local block built from the explicit local model, the affine jet of the smooth remainder, and the jet-corrected phase remainder.

**Proposition 9.1** (Microscopic denominator comparability and omitted-smooth holomorphy). *Fix an admissible midpoint  $m$  and write*

$$t_\pm = m \pm \frac{s}{2}, \quad Q \asymp \log |m|.$$

*For each omitted zero  $\rho = \beta_\rho + i\gamma_\rho$ , set*

$$D_{\rho,\pm}(s) := ((2m - \gamma_\rho) \pm s)^2 + a_\rho^2, \quad a_\rho \in \{\beta_\rho, 1 - \beta_\rho\}.$$

*Assume the zero-free region gives*

$$a_\rho \geq \frac{\sigma_0}{Q}$$

*for some absolute  $\sigma_0 > 0$  on the admissible midpoint class. If*

$$|s| \leq \frac{\rho_0}{Q}, \quad \rho_0 \leq \frac{\sigma_0}{\sqrt{6}},$$

*then*

$$\frac{1}{2} \left( (2m - \gamma_\rho)^2 + a_\rho^2 \right) \leq |D_{\rho,\pm}(s)| \leq \frac{3}{2} \left( (2m - \gamma_\rho)^2 + a_\rho^2 \right).$$

*Consequently, for every derivative order  $r \leq r_*$  needed in the corrected finite- $s$  block entries, the omitted-zero sums defining*

$$T_{\text{far}}(t_\pm), \quad \partial_t T_{\text{far}}(t_\pm), \quad \dots, \quad \partial_t^{r_*} T_{\text{far}}(t_\pm)$$

*converge uniformly on  $|s| < \rho_0/Q$ , provided the corresponding real-line shell sums converge. In particular,*

$$g_{\text{sm}}(t_\pm) = B_{\text{aut}}(t_\pm) + T_{\text{far}}(t_\pm)$$

*and its required  $t$ -derivatives extend holomorphically to*

$$|s| < \rho_0/Q.$$

*Proof.* Set

$$x := 2m - \gamma_\rho, \quad a := a_\rho.$$

Then

$$D_{\rho,\pm}(s) = (x^2 + a^2) \pm 2xs + s^2.$$

Using

$$|2xs + s^2| \leq 2|x||s| + |s|^2 \leq \frac{1}{2}x^2 + 3|s|^2,$$

and the assumptions

$$a \geq \frac{\sigma_0}{Q}, \quad |s| \leq \frac{\rho_0}{Q}, \quad \rho_0 \leq \frac{\sigma_0}{\sqrt{6}},$$

we obtain

$$3|s|^2 \leq \frac{1}{2}a^2.$$

Hence

$$|2xs + s^2| \leq \frac{1}{2}(x^2 + a^2),$$

which gives the stated lower and upper bounds.

Now every differentiated omitted-zero term is a rational combination of powers of  $D_{\rho,\pm}(s)^{-1}$  with coefficients depending only on the derivative order. By the above comparability, on  $|s| < \rho_0/Q$  these are uniformly comparable to their real-line counterparts. Therefore the same shell decomposition used on the real line yields uniform convergence of the differentiated omitted-zero sums on the complex disk, and hence holomorphy of  $T_{\text{far}}(t_\pm)$  and the required  $t$ -derivatives there. Adding the holomorphic background term  $B_{\text{aut}}(t_\pm)$  gives the corresponding statement for  $g_{\text{sm}}(t_\pm)$ .  $\square$

**Proposition 9.2** (Entry-by-entry perturbation estimate for the corrected same-point blocks).  
In the corrected finite- $s$  normalization, write

$$G_{m,\pm}^{\text{corr}}(s) = \frac{1}{\pi} \begin{pmatrix} 2q(t_{\pm}) & \frac{1}{2}q'(t_{\pm}) \\ \frac{1}{2}q'(t_{\pm}) & \frac{1}{12}(q''(t_{\pm}) + 2q(t_{\pm})^3) \end{pmatrix}, \quad t_{\pm} = m \pm \frac{s}{2}.$$

Let

$$q(t) = q_0(t) + \delta q(t),$$

where  $q_0$  is the baseline local-model-plus-affine-jet part and  $\delta q$  is the corrected remainder. Define the baseline same-point blocks

$$G_{m,\pm}^{(0)}(s) = \frac{1}{\pi} \begin{pmatrix} 2q_0(t_{\pm}) & \frac{1}{2}q_0'(t_{\pm}) \\ \frac{1}{2}q_0'(t_{\pm}) & \frac{1}{12}(q_0''(t_{\pm}) + 2q_0(t_{\pm})^3) \end{pmatrix},$$

and set

$$R_{m,\pm}(s) := G_{m,\pm}^{\text{corr}}(s) - G_{m,\pm}^{(0)}(s).$$

Assume that on the microscopic window one has

$$|\delta q(t_{\pm})| \ll L_I^2 S_2, \quad |\delta q'(t_{\pm})| \ll L_I S_2, \quad |\delta q''(t_{\pm})| \ll S_2,$$

where  $L_I$  is the local window length, and that in the fixed-scaled regime

$$L_I \asymp \frac{x_0}{Q}, \quad |q_0(t_{\pm})| \asymp Q.$$

Then

$$\|R_{m,\pm}(s)\| \ll S_2$$

uniformly for  $|s| < \rho_0/Q$ .

*Proof.* We estimate the entries of  $R_{m,\pm}(s)$  one by one.

**(1,1)-entry.** Since

$$R_{11} = \frac{1}{\pi} \cdot 2\delta q(t_{\pm}),$$

we have

$$|R_{11}| \ll |\delta q(t_{\pm})| \ll L_I^2 S_2 \ll \frac{S_2}{Q^2} \ll S_2.$$

**(1,2)- and (2,1)-entries.** Since

$$R_{12} = R_{21} = \frac{1}{\pi} \cdot \frac{1}{2}\delta q'(t_{\pm}),$$

we get

$$|R_{12}| = |R_{21}| \ll |\delta q'(t_{\pm})| \ll L_I S_2 \ll \frac{S_2}{Q} \ll S_2.$$

**(2,2)-entry.** We have

$$R_{22} = \frac{1}{12\pi} \left( \delta q''(t_{\pm}) + 2 \left( (q_0 + \delta q)^3 - q_0^3 \right) (t_{\pm}) \right).$$

Expand the cubic difference:

$$(q_0 + \delta q)^3 - q_0^3 = 3q_0^2\delta q + 3q_0(\delta q)^2 + (\delta q)^3.$$

Hence

$$R_{22} = \frac{1}{12\pi} \left( \delta q'' + 6q_0^2\delta q + 6q_0(\delta q)^2 + 2(\delta q)^3 \right) (t_{\pm}).$$

Now estimate each term.

First,

$$|\delta q''(t_{\pm})| \ll S_2.$$

Second, using  $|q_0(t_{\pm})| \asymp Q$  and  $|\delta q(t_{\pm})| \ll L_I^2 S_2 \ll S_2/Q^2$ ,

$$|q_0(t_{\pm})^2 \delta q(t_{\pm})| \ll Q^2 \cdot \frac{S_2}{Q^2} = S_2.$$

Third,

$$|q_0(t_{\pm})(\delta q(t_{\pm}))^2| \ll Q \left( \frac{S_2}{Q^2} \right)^2 = \frac{S_2^2}{Q^3}.$$

Since from the tail-curvature theorem one already has  $S_2 = o(Q)$ , for  $Q$  sufficiently large this is  $O(S_2/Q^2)$  and hence  $\ll S_2$ .

Finally,

$$|(\delta q(t_{\pm}))^3| \ll \left( \frac{S_2}{Q^2} \right)^3 = \frac{S_2^3}{Q^6} \ll S_2.$$

Therefore

$$|R_{22}| \ll S_2.$$

Collecting the four entrywise bounds, we obtain

$$|R_{11}|, |R_{12}|, |R_{21}|, |R_{22}| \ll S_2.$$

Since all norms on  $2 \times 2$  matrices are equivalent,

$$\|R_{m,\pm}(s)\| \ll S_2,$$

uniformly on the microscopic disk. This proves the proposition.  $\square$

**Proposition 9.3** (Entry-by-entry perturbation estimate for the corrected mixed block). *Let*

$$q(t) = q_0(t) + \delta q(t), \quad \Phi(t) = \Phi_0(t) + \delta \Phi(t),$$

where  $q_0$  is the baseline local-model-plus-affine-jet phase derivative and  $\delta q$  is the corrected remainder. Write

$$t_{\pm} = m \pm \frac{s}{2}, \quad \Delta(s) := \Phi(t_-) - \Phi(t_+), \quad \Delta_0(s) := \Phi_0(t_-) - \Phi_0(t_+),$$

and set

$$\delta \Delta(s) := \Delta(s) - \Delta_0(s).$$

Let  $N_m^{\text{corr}}(s)$  be the corrected finite- $s$  mixed block built from  $q, \Delta$ , and let  $N_m^{(0)}(s)$  be the corresponding block built from  $q_0, \Delta_0$ . Define

$$\tilde{R}_m(s) := N_m^{\text{corr}}(s) - N_m^{(0)}(s).$$

Assume that on the microscopic window one has

$$|\delta q(t_{\pm})| \ll L_I^2 S_2, \quad |\delta q'(t_{\pm})| \ll L_I S_2, \quad |\delta q''(t_{\pm})| \ll S_2,$$

and

$$|\delta \Delta(s)| \ll L_I^3 S_2,$$

with

$$L_I \asymp \frac{x_0}{Q}, \quad |q_0(t_{\pm})| \asymp Q, \quad |s| \asymp L_I.$$

Then

$$\|\tilde{R}_m(s)\| \ll S_2$$

uniformly for  $|s| < \rho_0/Q$ .

*Proof.* Write the corrected mixed block entrywise as

$$N_m^{\text{corr}}(s) = \frac{1}{\pi} \begin{pmatrix} -\frac{2 \sin \Delta}{s} & \frac{\sin \Delta + q(t_+)s \cos \Delta}{s^2} \\ -\frac{\sin \Delta + q(t_-)s \cos \Delta}{s^2} & \frac{(q(t_-) + q(t_+))s \cos \Delta + (2 - q(t_-)q(t_+)s^2) \sin \Delta}{2s^3} \end{pmatrix},$$

and analogously for  $N_m^{(0)}(s)$  with  $(q_0, \Delta_0)$ .

We estimate the perturbation entry by entry. Throughout, use

$$\sin(\Delta_0 + \delta \Delta) - \sin \Delta_0 = O(|\delta \Delta|), \quad \cos(\Delta_0 + \delta \Delta) - \cos \Delta_0 = O(|\delta \Delta|).$$

**(1,1)-entry.**

$$\tilde{R}_{11} = -\frac{2}{\pi s}(\sin \Delta - \sin \Delta_0),$$

hence

$$|\tilde{R}_{11}| \ll \frac{|\delta \Delta|}{|s|} \ll \frac{L_I^3 S_2}{L_I} \ll L_I^2 S_2 \ll S_2.$$

**(1,2)-entry.** The numerator perturbation is

$$(\sin \Delta - \sin \Delta_0) + (q(t_+) - q_0(t_+))s \cos \Delta + q_0(t_+)s(\cos \Delta - \cos \Delta_0).$$

Therefore

$$|\tilde{R}_{12}| \ll \frac{|\delta \Delta| + |\delta q(t_+)| |s| + |q_0(t_+)| |s| |\delta \Delta|}{|s|^2}.$$

Using

$$|\delta \Delta| \ll L_I^3 S_2, \quad |\delta q(t_+)| \ll L_I^2 S_2, \quad |q_0(t_+)| \asymp Q, \quad |s| \asymp L_I \asymp Q^{-1},$$

we get

$$|\tilde{R}_{12}| \ll \frac{L_I^3 S_2}{L_I^2} + \frac{L_I^3 S_2}{L_I^2} + \frac{Q L_I^4 S_2}{L_I^2} \ll \frac{S_2}{Q} \ll S_2.$$

The same argument gives

$$|\tilde{R}_{21}| \ll S_2.$$

**(2,2)-entry.** Write the numerator as

$$F(q_-, q_+, \Delta; s) := (q_- + q_+)s \cos \Delta + (2 - q_- q_+ s^2) \sin \Delta.$$

Then

$$\tilde{R}_{22} = \frac{F(q(t_-), q(t_+), \Delta; s) - F(q_0(t_-), q_0(t_+), \Delta_0; s)}{2\pi s^3}.$$

A first-order expansion of  $F$  in  $(q_-, q_+, \Delta)$  gives

$$|F - F_0| \ll |s| |\delta q(t_-)| + |s| |\delta q(t_+)| + |s|^2 Q |\delta q(t_-)| + |s|^2 Q |\delta q(t_+)| + (|s|Q + 1) |\delta \Delta|.$$

Using

$$|\delta q(t_{\pm})| \ll L_I^2 S_2, \quad |\delta \Delta| \ll L_I^3 S_2, \quad |s| \asymp L_I \asymp Q^{-1},$$

we obtain

$$|F - F_0| \ll L_I^3 S_2.$$

Hence

$$|\tilde{R}_{22}| \ll \frac{L_I^3 S_2}{|s|^3} \ll S_2.$$

Collecting the four entrywise bounds and using equivalence of norms on  $2 \times 2$  matrices gives

$$\|\tilde{R}_m(s)\| \ll S_2,$$

uniformly on the microscopic disk.  $\square$

**Lemma 9.4** (Baseline same-point variation). *Let  $G_{m,\pm}^{(0)}(s)$  be the corrected same-point baseline blocks built from the explicit local model together with the affine jet  $J_1[g_{\text{sm}}]$ . Then, on the microscopic disk  $|s| < \rho_0/Q$ ,*

$$\sup_{|s| < \rho_0/Q} \|G_{m,\pm}^{(0)}(s) - G_{m,\pm}^{(0)}(0)\| \ll \rho_0 Q.$$

*Proof.* Differentiate the explicit same-point block formula entrywise:

$$G_{m,\pm}^{(0)}(s) = \frac{1}{\pi} \begin{pmatrix} 2q_0(t_{\pm}) & \frac{1}{2}q_0'(t_{\pm}) \\ \frac{1}{2}q_0'(t_{\pm}) & \frac{1}{12}(q_0''(t_{\pm}) + 2q_0(t_{\pm})^3) \end{pmatrix}, \quad t_{\pm} = m \pm \frac{s}{2}.$$

By the chain rule,

$$\partial_s G_{m,\pm}^{(0)}(s) = \pm \frac{1}{2\pi} \begin{pmatrix} 2q_0'(t_{\pm}) & \frac{1}{2}q_0''(t_{\pm}) \\ \frac{1}{2}q_0''(t_{\pm}) & \frac{1}{12}(q_0'''(t_{\pm}) + 6q_0(t_{\pm})^2 q_0'(t_{\pm})) \end{pmatrix}.$$

For the explicit local model plus affine jet, one has the baseline local scales

$$|q_0(t_\pm)| \ll Q, \quad |q'_0(t_\pm)| \ll Q^2, \quad |q''_0(t_\pm)| \ll Q^3, \quad |q'''_0(t_\pm)| \ll Q^4,$$

uniformly on the admissible midpoint class. The cubic combination  $q'''_0(t_\pm) + 6q_0(t_\pm)^2 q'_0(t_\pm)$  therefore has size  $O(Q^2)$  in the current normalization, and hence

$$\sup_{|s| < \rho_0/Q} \|\partial_s G_{m,\pm}^{(0)}(s)\| \ll Q^2.$$

Integrating along the segment from 0 to  $s$  gives

$$\|G_{m,\pm}^{(0)}(s) - G_{m,\pm}^{(0)}(0)\| \leq |s| \sup_{|u| < \rho_0/Q} \|\partial_u G_{m,\pm}^{(0)}(u)\| \ll \frac{\rho_0}{Q} \cdot Q^2 = \rho_0 Q.$$

□

**Proposition 9.5** (Corrected finite- $s$  holomorphic whitening). *The corrected same-point and mixed blocks are holomorphic on*

$$|s| < \rho_0/Q.$$

Moreover, after shrinking  $\rho_0$  if necessary, the same-point blocks

$$G_{m,\pm}^{\text{corr}}(s)$$

remain uniformly positive definite on this disk, and the inverse square roots

$$G_{m,\pm}^{\text{corr}}(s)^{-1/2}$$

are holomorphic there. Consequently

$$\widehat{\Omega}_\zeta^{\text{corr}}(s; m) = G_{m,-}^{\text{corr}}(s)^{-1/2} N_m^{\text{corr}}(s) G_{m,+}^{\text{corr}}(s)^{-1/2}$$

is holomorphic on  $|s| < \rho_0/Q$ .

*Proof.* Holomorphy of the omitted smooth part and its required  $t$ -derivatives follows from Proposition 9.1. The only apparent poles at  $s = 0$  in the corrected mixed block come from powers of  $1/s$ ,  $1/s^2$ ,  $1/s^3$ , and these are removable because  $\Delta_m(s)$  is odd in  $s$ ,  $\sin \Delta_m(s) = O(s)$ , and the off-diagonal and  $(2, 2)$  numerators vanish to orders 2 and 3, respectively. Thus both  $G_{m,\pm}^{\text{corr}}(s)$  and  $N_m^{\text{corr}}(s)$  are holomorphic on the disk.

By Proposition 9.2 and the tail-curvature theorem,

$$\sup_{|s| < \rho_0/Q} \|R_{m,\pm}(s)\| = o(Q).$$

By Lemma 9.4,

$$\sup_{|s| < \rho_0/Q} \|G_{m,\pm}^{(0)}(s) - G_{m,\pm}^{(0)}(0)\| \ll \rho_0 Q.$$

At  $s = 0$ , the baseline jet-limit block has spectral gap

$$\lambda_{\min}(G_{m,\pm}^{(0)}(0)) \gg Q$$

on the admissible midpoint class. Therefore

$$\|G_{m,\pm}^{\text{corr}}(s) - G_{m,\pm}^{(0)}(0)\| \leq \|G_{m,\pm}^{(0)}(s) - G_{m,\pm}^{(0)}(0)\| + \|R_{m,\pm}(s)\| \ll \rho_0 Q + o(Q).$$

For  $\rho_0$  sufficiently small and  $m$  sufficiently large, this is smaller than half the baseline spectral gap, hence

$$\lambda_{\min}(G_{m,\pm}^{\text{corr}}(s)) \gg Q$$

uniformly on the disk. Holomorphic functional calculus then gives the holomorphic inverse square roots, and the whitened block is holomorphic as a product of holomorphic factors. □

**9.1. Fixed anomaly core and cutoff-dependent auxiliary bookkeeping.** To separate the anomaly-carrying local model from the cutoff-dependent tail bookkeeping, we fix once and for all a finite local core

$$\mathcal{C}$$

containing the candidate off-line anomaly (for example, the designated off-line pair together with any symmetry-forced companions needed for the local model).

For each cutoff  $R \geq R_0$ , we then decompose the zero contribution into:

- (1) the fixed core contribution  $q_{\text{core}}(t)$ , coming from the zeros in  $\mathcal{C}$ ;
- (2) the auxiliary near contribution  $q_{\text{aux},R}(t)$ , coming from the zeros outside  $\mathcal{C}$  but inside the near shell determined by  $R$ ;
- (3) the far contribution  $T_{\text{far},R}(t)$ , coming from the zeros outside the near shell;
- (4) the background contribution  $B_{\text{bg}}(t)$ ;
- (5) the estimation term  $E_{\text{est},R}(t)$ .

Thus the full phase derivative is decomposed as

$$q(t) = q_{\text{core}}(t) + q_{\text{aux},R}(t) + g_{\text{sm},R}(t) + E_{\text{est},R}(t),$$

where

$$g_{\text{sm},R}(t) := B_{\text{bg}}(t) + T_{\text{far},R}(t).$$

The local toy comparison is attached only to the fixed core  $\mathcal{C}$ . The auxiliary near part  $q_{\text{aux},R}$ , the far part  $T_{\text{far},R}$ , and the estimation term  $E_{\text{est},R}$  are absorbed into the corrected zeta-side scalar defined below.

**Proposition 9.6** (Corrected local deformation decomposition with fixed core). *Fix the anomaly-carrying core  $\mathcal{C}$  as above, and for each cutoff  $R \geq R_0$  write*

$$q(t) = q_{\text{core}}(t) + q_{\text{aux},R}(t) + g_{\text{sm},R}(t) + E_{\text{est},R}(t), \quad g_{\text{sm},R}(t) = B_{\text{bg}}(t) + T_{\text{far},R}(t).$$

*Let  $J_1[g_{\text{sm},R}]$  denote the affine jet of  $g_{\text{sm},R}$  at the midpoint, and let  $\Phi_{\text{corr},R}$  be the corrected phase obtained from this decomposition.*

*Using  $\Phi_{\text{corr},R}$ , define the corrected same-point blocks*

$$G_{m,\pm}^{\text{corr},R}(s),$$

*the corrected mixed block*

$$N_m^{\text{corr},R}(s),$$

*and the corrected whitened block*

$$\hat{\Omega}_\zeta^{\text{corr},R}(s; m) = G_{m,-}^{\text{corr},R}(s)^{-1/2} N_m^{\text{corr},R}(s) G_{m,+}^{\text{corr},R}(s)^{-1/2}.$$

*Let*

$$\hat{\Omega}_{\text{bg}}^{\text{corr},R}(s; m)$$

*denote the corresponding corrected whitened block at background value-channel parameter  $S(m) = 0$ , and define the corrected local deformation*

$$\Delta_\zeta(m, \sigma) := \hat{\Omega}_\zeta^{\text{corr},R}(s; m) - \hat{\Omega}_{\text{bg}}^{\text{corr},R}(s; m), \quad s = \sigma.$$

*Then the corrected local deformation admits the decomposition*

$$\Delta_\zeta(m, \sigma) = S(m) A_{\text{val}}(m, \sigma) + R_\zeta(m, \sigma),$$

*where  $A_{\text{val}}(m, \sigma)$  is the pure value-channel derivative of the jet-limit block associated with the fixed core  $\mathcal{C}$ , and where*

$$R_\zeta(m, \sigma) := \Delta_\zeta(m, \sigma) - S(m) A_{\text{val}}(m, \sigma).$$

*Thus, by definition,  $R_\zeta(m, \sigma)$  is exactly the non-leading part of the corrected local deformation after the explicit value-channel linear term is removed. In particular,  $R_\zeta(m, \sigma)$  is internal to the same full corrected cutoff-dependent object*

$$H_{m,R}(s) := \Phi_K(\hat{\Omega}_\zeta^{\text{corr},R}(s; m)),$$

so once the corrected odd holomorphic package is established for  $H_{m,R}(s)$ , no external tail/error term survives outside the resulting  $N$ -point scalar.

*Proof.* Consider the analytic whitening map

$$\mathcal{W}(G_-, N, G_+) := G_-^{-1/2} N G_+^{-1/2}.$$

By the corrected finite- $s$  holomorphic-whitening theorem, this map is holomorphic on the open set where  $G_{\pm}$  remain uniformly positive definite.

For each cutoff  $R$ , the corrected blocks are built from the decomposition

$$q = q_{\text{core}} + q_{\text{aux},R} + g_{\text{sm},R} + E_{\text{est},R}.$$

Thus every auxiliary near contribution from  $q_{\text{aux},R}$ , every omitted-zero tail contribution from  $T_{\text{far},R}$ , every affine-jet correction from  $J_1[g_{\text{sm},R}]$ , and every explicit estimation term  $E_{\text{est},R}$  is already absorbed into the corrected phase  $\Phi_{\text{corr},R}$ , hence into the corrected same-point blocks, the corrected mixed block, the corrected whitened block, and therefore into the corrected transverse scalar

$$H_{m,R}(s) = \Phi_K(\widehat{\Omega}_{\zeta}^{\text{corr},R}(s; m)).$$

Now expand the corrected whitened block around the background value-channel parameter  $S(m) = 0$ . By analyticity of  $\mathcal{W}$ ,

$$\widehat{\Omega}_{\zeta}^{\text{corr},R}(s; m) = \widehat{\Omega}_{\text{bg}}^{\text{corr},R}(s; m) + S(m) DW_{(\text{bg})}[\dot{G}_-, \dot{N}, \dot{G}_+] + R_{\zeta}(m, \sigma),$$

where the first-order coefficient

$$DW_{(\text{bg})}[\dot{G}_-, \dot{N}, \dot{G}_+]$$

is exactly the pure value-channel derivative  $A_{\text{val}}(m, \sigma)$  associated with the fixed core  $\mathcal{C}$ . We therefore define

$$R_{\zeta}(m, \sigma) := \Delta_{\zeta}(m, \sigma) - S(m) A_{\text{val}}(m, \sigma),$$

so that  $R_{\zeta}(m, \sigma)$  is, by definition, the full non-leading remainder inside the same corrected cutoff-dependent object.

Subtracting the background block gives

$$\Delta_{\zeta}(m, \sigma) = S(m) A_{\text{val}}(m, \sigma) + R_{\zeta}(m, \sigma),$$

as claimed.  $\square$

**Proposition 9.7** (Core-stable cutoff compatibility). *Fix the anomaly-carrying core  $\mathcal{C}$ , and for each cutoff  $R \geq R_0$  define the corrected scalar*

$$H_{m,R}(s) := \Phi_K(\widehat{\Omega}_{\zeta}^{\text{corr},R}(s; m))$$

*from the decomposition*

$$q = q_{\text{core}} + q_{\text{aux},R} + g_{\text{sm},R} + E_{\text{est},R}.$$

*Then enlarging  $R$  changes only the auxiliary near part  $q_{\text{aux},R}$  and the far part  $T_{\text{far},R}$ , while leaving the anomaly-carrying core  $q_{\text{core}}$  unchanged. Consequently:*

- (1) *the toy anomaly matrix  $M(d)$  and the calibration matrix  $A_{\text{val}}(m, \sigma)$  depend only on the fixed core  $\mathcal{C}$ , not on the cutoff  $R$ ;*
- (2) *all  $R$ -dependent auxiliary, tail, jet-correction, and estimation contributions are internal to the full corrected scalar  $H_{m,R}(s)$ ;*
- (3) *if the corrected odd holomorphic package is proved for  $H_{m,R}(s)$ , then the  $N$ -point odd-moment cancellation theorem applies directly to the full corrected scalar for that cutoff, with no external remainder term.*

*In particular, the cutoff  $R$  may be chosen for quantitative convenience in the curvature/tail estimates without changing the fixed toy comparison attached to the core  $\mathcal{C}$ .*



*Proof.* Let  $R_1 < R_2$ . Then the only change in the decomposition

$$q = q_{\text{core}} + q_{\text{aux},R} + g_{\text{sm},R} + E_{\text{est},R}$$

is that some non-core zeros move from the far part into the auxiliary near part:

$$q_{\text{aux},R_2} - q_{\text{aux},R_1} = -(T_{\text{far},R_2} - T_{\text{far},R_1}).$$

The core contribution  $q_{\text{core}}$  is unchanged by construction.

Thus the toy anomaly matrix  $M(d)$ , which is attached only to the fixed core, is unchanged, and the value-channel derivative  $A_{\text{val}}(m, \sigma)$  associated with that core is unchanged as well. All cutoff dependence is confined to the auxiliary near contribution, the far tail, the affine-jet correction built from  $g_{\text{sm},R}$ , and the explicit estimation term  $E_{\text{est},R}$ . These enter only through the corrected phase  $\Phi_{\text{corr},R}$ , hence only through the full corrected scalar  $H_{m,R}(s)$ .

Therefore the corrected odd holomorphic package, once established for  $H_{m,R}(s)$ , applies to the complete cutoff-dependent object. The resulting  $N$ -point scalar already contains all auxiliary/tail/error effects for that cutoff, and no external remainder survives. Since the fixed core is unchanged, the toy comparison is stable under enlarging  $R$ . This proves the claim.  $\square$

**Remark 9.8** (Cutoff choice). Proposition 9.7 is a structural statement. It does not assert any particular optimal choice of  $R$ ; it only says that once the core  $\mathcal{C}$  is fixed, the cutoff may be adjusted in the perturbative bookkeeping without altering the underlying toy comparison.

**Proposition 9.9** (Whitened mixed-block transfer with preserved  $Q^{-3}$  scale). *Let*

$$\mathcal{W}(G_-, N, G_+) := G_-^{-1/2} N G_+^{-1/2}.$$

*Write the corrected finite- $s$  blocks as*

$$G_{\pm}^{\text{corr}}(s) = G_{\pm}^{(0)}(s) + R_{\pm}(s), \quad N^{\text{corr}}(s) = N^{(0)}(s) + \tilde{R}(s),$$

*and set*

$$\widehat{\Omega}_{\zeta}^{\text{corr}}(s; m) := \mathcal{W}(G_-^{\text{corr}}, N^{\text{corr}}, G_+^{\text{corr}}), \quad \widehat{\Omega}_{\zeta}^{(0)}(s; m) := \mathcal{W}(G_-^{(0)}, N^{(0)}, G_+^{(0)}).$$

*Then*

$$\widehat{\Omega}_{\zeta}^{\text{corr}}(s; m) - \widehat{\Omega}_{\zeta}^{(0)}(s; m) = \mathcal{T}_-(s; m) + \mathcal{T}_{\text{mix}}(s; m) + \mathcal{T}_+(s; m) + \mathcal{E}_2(s; m),$$

*where*

$$\begin{aligned} \mathcal{T}_-(s; m) &:= X_-(s) N^{(0)}(s) G_+^{(0)}(s)^{-1/2}, \\ \mathcal{T}_{\text{mix}}(s; m) &:= G_-^{(0)}(s)^{-1/2} \tilde{R}(s) G_+^{(0)}(s)^{-1/2}, \\ \mathcal{T}_+(s; m) &:= G_-^{(0)}(s)^{-1/2} N^{(0)}(s) X_+(s), \end{aligned}$$

*with*

$$X_{\pm}(s) := D(G^{-1/2})_{G_{\pm}^{(0)}(s)}[R_{\pm}(s)],$$

*and  $\mathcal{E}_2$  quadratic in  $(R_-, \tilde{R}, R_+)$ .*

*Moreover,*

$$\Phi_K(\mathcal{T}_-(s; m)) \ll \frac{S_2}{Q^3}, \quad \Phi_K(\mathcal{T}_{\text{mix}}(s; m)) \ll \frac{S_2}{Q^3}, \quad \Phi_K(\mathcal{T}_+(s; m)) \ll \frac{S_2}{Q^3},$$

*and*

$$\Phi_K(\mathcal{E}_2(s; m)) \ll \frac{S_2^2}{Q^3}.$$

*Hence*

$$\boxed{\Phi_K(\widehat{\Omega}_{\zeta}^{\text{corr}}(s; m) - \widehat{\Omega}_{\zeta}^{(0)}(s; m)) \ll \frac{S_2}{Q^3}.}$$

*Proof.* We first collect the scales used below from results established earlier. The same-point perturbation bound  $\|R_\pm(s)\| \ll S_2$  is Proposition 9.2; the mixed-block perturbation bound  $\|\tilde{R}(s)\| \ll S_2$  is Proposition 9.3; and the smallness  $S_2 = o(Q)$  is Theorem 6.1. The baseline entry scales

$$G_\pm^{(0)}(s) = \begin{pmatrix} O(Q) & O(1) \\ O(1) & O(Q^3) \end{pmatrix}, \quad G_\pm^{(0)}(s)^{-1/2} = \begin{pmatrix} O(Q^{-1/2}) & O(Q^{-3/2}) \\ O(Q^{-3/2}) & O(Q^{-3/2}) \end{pmatrix},$$

$$N^{(0)}(s) = \begin{pmatrix} O(Q) & O(Q^2) \\ O(Q^2) & O(Q^3) \end{pmatrix}$$

are read off directly from the explicit same-point formula (7) and the mixed-block formula (8), given the baseline local scales of  $q_0, q'_0, q''_0$  stated in the proof of Lemma 9.4. Uniform positive-definiteness of  $G_\pm^{(0)}(s)$  on the microscopic disk is Proposition 9.5.

The whitening map admits the first-order expansion

$$\hat{\Omega}_\zeta^{\text{corr}} - \hat{\Omega}_\zeta^{(0)} = D\mathcal{W}_{(0)}[R_-, \tilde{R}, R_+] + \mathcal{E}_2,$$

where

$$D\mathcal{W}_{(0)}[R_-, \tilde{R}, R_+] = X_- N^{(0)} G_+^{(0)-1/2} + G_-^{(0)-1/2} \tilde{R} G_+^{(0)-1/2} + G_-^{(0)-1/2} N^{(0)} X_+.$$

We estimate the three first-order pieces separately.

**Step 1: the mixed-block term.** By Proposition 9.3 (entrywise form),

$$\tilde{R}_{11} \ll \frac{S_2}{Q^2}, \quad \tilde{R}_{12}, \tilde{R}_{21} \ll \frac{S_2}{Q}, \quad \tilde{R}_{22} \ll S_2.$$

Combining with the baseline inverse-square-root scales above, the off-diagonal entries of

$$\mathcal{T}_{\text{mix}} := G_-^{(0)-1/2} \tilde{R} G_+^{(0)-1/2}$$

are all  $O(S_2/Q^3)$ . Hence

$$\Phi_K(\mathcal{T}_{\text{mix}}) \ll \frac{S_2}{Q^3}.$$

**Step 2: the same-point Fréchet derivative.** We prove the corresponding estimate for the left same-point term

$$\mathcal{T}_- := X_- N^{(0)} G_+^{(0)-1/2}, \quad X_- := D(G^{-1/2})_{G_-^{(0)}}[R_-].$$

The proof for  $\mathcal{T}_+$  is identical.

Write one baseline same-point block in the form

$$G_0 = \begin{pmatrix} a & b \\ b & c \end{pmatrix}, \quad a \asymp Q, \quad b = O(1), \quad c \asymp Q^3,$$

and one perturbation as

$$R = \begin{pmatrix} r_{11} & r_{12} \\ r_{12} & r_{22} \end{pmatrix}, \quad r_{11} \ll \frac{S_2}{Q^2}, \quad r_{12} \ll \frac{S_2}{Q}, \quad r_{22} \ll S_2.$$

Diagonalize

$$G_0 = U \Lambda U^\top, \quad \Lambda = \text{diag}(\lambda_1, \lambda_2),$$

with  $U$  orthogonal. Since

$$a \asymp Q, \quad c \asymp Q^3, \quad b = O(1),$$

we have

$$\lambda_1 \asymp Q, \quad \lambda_2 \asymp Q^3.$$

Moreover, if  $U$  is a rotation by angle  $\theta$ , then

$$\tan(2\theta) = \frac{2b}{c-a} = O(Q^{-3}),$$

hence

$$\theta = O(Q^{-3}), \quad U = I + O(Q^{-3}).$$

Set

$$H := U^\top R U.$$

A direct computation gives

$$\begin{aligned} H_{11} &= r_{11} + 2\theta r_{12} + O(\theta^2 r_{22}), \\ H_{12} &= r_{12} + \theta(r_{22} - r_{11}) + O(\theta^2 r_{12}), \\ H_{22} &= r_{22} - 2\theta r_{12} + O(\theta^2 r_{11}). \end{aligned}$$

Using

$$\theta = O(Q^{-3}), \quad r_{11} \ll \frac{S_2}{Q^2}, \quad r_{12} \ll \frac{S_2}{Q}, \quad r_{22} \ll S_2,$$

we obtain

$$H_{11} \ll \frac{S_2}{Q^2}, \quad H_{12} \ll \frac{S_2}{Q}, \quad H_{22} \ll S_2.$$

Now let  $f(x) = x^{-1/2}$ . For a diagonal matrix  $\Lambda$ , the Fréchet derivative is given entrywise by the Löwner formula

$$Df_\Lambda(H)_{ij} = f^{[1]}(\lambda_i, \lambda_j) H_{ij},$$

where

$$f^{[1]}(u, v) = \begin{cases} \frac{u^{-1/2} - v^{-1/2}}{u - v}, & u \neq v, \\ -\frac{1}{2u^{3/2}}, & u = v. \end{cases}$$

Since

$$\lambda_1 \asymp Q, \quad \lambda_2 \asymp Q^3,$$

we have

$$f^{[1]}(\lambda_1, \lambda_1) \asymp Q^{-3/2}, \quad f^{[1]}(\lambda_2, \lambda_2) \asymp Q^{-9/2}, \quad f^{[1]}(\lambda_1, \lambda_2) \asymp Q^{-7/2}.$$

Therefore, if

$$Y := Df_\Lambda(H),$$

then

$$\begin{aligned} Y_{11} &\ll Q^{-3/2} \cdot \frac{S_2}{Q^2} = \frac{S_2}{Q^{7/2}}, \\ Y_{12} &\ll Q^{-7/2} \cdot \frac{S_2}{Q} = \frac{S_2}{Q^{9/2}}, \\ Y_{22} &\ll Q^{-9/2} \cdot S_2 = \frac{S_2}{Q^{9/2}}. \end{aligned}$$

Since

$$X = U Y U^\top$$

and  $U = I + O(Q^{-3})$ , these scales persist in the original basis:

$$X_{11} \ll \frac{S_2}{Q^{7/2}}, \quad X_{12}, X_{21} \ll \frac{S_2}{Q^{9/2}}, \quad X_{22} \ll \frac{S_2}{Q^{9/2}}.$$

**Step 3: the left same-point term  $\mathcal{T}_-$ .** The baseline mixed-block scales are

$$N^{(0)} = \begin{pmatrix} O(Q) & O(Q^2) \\ O(Q^2) & O(Q^3) \end{pmatrix}, \quad G_+^{(0)-1/2} = \begin{pmatrix} O(Q^{-1/2}) & O(Q^{-3/2}) \\ O(Q^{-3/2}) & O(Q^{-3/2}) \end{pmatrix}.$$

We estimate only the off-diagonal entries, since

$$\Phi_K(Y) = Y_{12} + Y_{21}.$$

For  $(\mathcal{T}_-)_{12}$ , every summand is  $O(S_2/Q^3)$ . For example,

$$X_{11} N_{12} (G_+^{-1/2})_{22} \ll \frac{S_2}{Q^{7/2}} \cdot Q^2 \cdot Q^{-3/2} = \frac{S_2}{Q^3},$$

and

$$X_{12} N_{22} (G_+^{-1/2})_{22} \ll \frac{S_2}{Q^{9/2}} \cdot Q^3 \cdot Q^{-3/2} = \frac{S_2}{Q^3}.$$

All other terms are smaller. Thus

$$(\mathcal{T}_-)_{12} \ll \frac{S_2}{Q^3}.$$

The same argument gives

$$(\mathcal{T}_-)_{21} \ll \frac{S_2}{Q^3},$$

and hence

$$\Phi_K(\mathcal{T}_-) \ll \frac{S_2}{Q^3}.$$

By symmetry, the same estimate holds for the right same-point term:

$$\Phi_K(\mathcal{T}_+) \ll \frac{S_2}{Q^3}.$$

**Step 4: the quadratic remainder.** Every term in  $\mathcal{E}_2$  contains at least two perturbation factors from  $(R_-, \tilde{R}, R_+)$ , and whitening contributes only polynomial  $Q$ -weights. Therefore

$$\Phi_K(\mathcal{E}_2) \ll \frac{S_2^2}{Q^3}.$$

By Theorem 6.1,  $S_2 = o(Q)$ , so this is lower order than  $S_2/Q^3$ .

Combining the estimates for  $\mathcal{T}_-$ ,  $\mathcal{T}_{\text{mix}}$ ,  $\mathcal{T}_+$ , and  $\mathcal{E}_2$  gives

$$\Phi_K(\hat{\Omega}_\zeta^{\text{corr}}(s; m) - \hat{\Omega}_\zeta^{(0)}(s; m)) \ll \frac{S_2}{Q^3},$$

as claimed. □

**Corollary 9.10** (Sharpened calibration remainder estimate). *Fix a compact interval*

$$D \subset (0, \infty),$$

*and work in the fixed-scaled regime*

$$0 < x \leq x_0(D), \quad d \in D, \quad x = B_\zeta(m)\sigma, \quad |\sigma| \asymp Q^{-1}.$$

*Write the corrected local deformation in the form*

$$\Delta_\zeta(m, \sigma) = S(m)A_{\text{val}}(x) + R_\zeta(m, \sigma).$$

*Then*

$$|\langle A_{\text{val}}(x), R_\zeta(m, \sigma) \rangle_{\text{F}}| \ll_D \frac{x}{B_\zeta(m)} \frac{S_2}{Q^3},$$

*and hence*

$$|\Psi_{x,d}(R_\zeta(m, \sigma))| \ll_D \frac{S_2}{Q^3}.$$

*In particular, if*

$$S_2 = o(xQ^2S(m)),$$

*then*

$$\Psi_{x,d}(R_\zeta(m, \sigma)) = o\left(\frac{x}{B_\zeta(m)} S(m)\right).$$

*Proof.* By Proposition 9.6, after subtraction of the explicit leading value-channel term  $S(m)A_{\text{val}}(x)$ , the remaining contribution  $R_\zeta(m, \sigma)$  is internal to the same corrected transferred package controlled by the whitening perturbation analysis.

We therefore use the entry scales already established in the proof of Proposition 9.9. First, from Proposition 9.3 and the baseline inverse-square-root scales,

$$G_\pm^{(0)}(s)^{-1/2} = \begin{pmatrix} O(Q^{-1/2}) & O(Q^{-3/2}) \\ O(Q^{-3/2}) & O(Q^{-3/2}) \end{pmatrix}, \quad \tilde{R}(s) = \begin{pmatrix} O(S_2/Q^2) & O(S_2/Q) \\ O(S_2/Q) & O(S_2) \end{pmatrix},$$

one gets

$$\mathcal{T}_{\text{mix}}(s; m) = G_-^{(0)}(s)^{-1/2} \tilde{R}(s) G_+^{(0)}(s)^{-1/2} = \begin{pmatrix} O(S_2/Q^3) & O(S_2/Q^3) \\ O(S_2/Q^3) & O(S_2/Q^3) \end{pmatrix}.$$

Next, from the same proof,

$$X_{\pm}(s) = \begin{pmatrix} O(S_2/Q^{7/2}) & O(S_2/Q^{9/2}) \\ O(S_2/Q^{9/2}) & O(S_2/Q^{9/2}) \end{pmatrix}, \quad N^{(0)}(s) = \begin{pmatrix} O(Q) & O(Q^2) \\ O(Q^2) & O(Q^3) \end{pmatrix},$$

and multiplying once more by  $G_{\pm}^{(0)}(s)^{-1/2}$  gives

$$\mathcal{T}_{-}(s; m) = \begin{pmatrix} O(S_2/Q^3) & O(S_2/Q^3) \\ O(S_2/Q^3) & O(S_2/Q^3) \end{pmatrix}, \quad \mathcal{T}_{+}(s; m) = \begin{pmatrix} O(S_2/Q^3) & O(S_2/Q^3) \\ O(S_2/Q^3) & O(S_2/Q^3) \end{pmatrix}.$$

Finally, every entry of the quadratic remainder  $\mathcal{E}_2(s; m)$  is

$$O(S_2^2/Q^3),$$

which is lower order because  $S_2 = o(Q)$  by Theorem 6.1.

Thus the part of the corrected deformation seen after removal of the leading value-channel term satisfies

$$R_{\zeta}(m, \sigma) = \begin{pmatrix} O(S_2/Q^3) & O(S_2/Q^3) \\ O(S_2/Q^3) & O(S_2/Q^3) \end{pmatrix}$$

up to lower-order  $O(S_2^2/Q^3)$  terms.

Now use the explicit small- $x$  structure of the value-channel derivative: by Lemma 8.3,

$$A_{\text{val}}(x) = \frac{1}{B_{\zeta}(m)} \begin{pmatrix} O(x^2) & O(x) \\ O(x) & O(x^2) \end{pmatrix}.$$

Therefore

$$|\langle A_{\text{val}}(x), R_{\zeta}(m, \sigma) \rangle_{\text{F}}| \ll \frac{x}{B_{\zeta}(m)} \frac{S_2}{Q^3}.$$

By Proposition 8.4,

$$|\langle A_{\text{val}}(x), M(d) \rangle_{\text{F}}| \asymp_D \frac{x}{B_{\zeta}(m)},$$

so dividing gives

$$|\Psi_{x,d}(R_{\zeta}(m, \sigma))| \ll_D \frac{S_2}{Q^3}.$$

Finally, since  $B_{\zeta}(m) \asymp Q$ ,

$$\frac{S_2/Q^3}{(x/B_{\zeta}(m))S(m)} \asymp \frac{S_2}{xQ^2S(m)} \rightarrow 0$$

under the hypothesis  $S_2 = o(xQ^2S(m))$ . This proves the final claim.  $\square$

## 10. ODD HOLOMORPHIC TRANSVERSE CHANNEL

Define the corrected transverse scalar

$$(20) \quad H_m(s) := \Phi_K(\widehat{\Omega}_{\zeta}^{\text{corr}}(s; m)).$$

By symmetry under  $s \mapsto -s$ , this is odd. By Corollary 8.2, the leading value channel is annihilated by  $\Phi_K$ , so only the curvature, tail, and corrected finite- $s$  remainder channels contribute.

**Proposition 10.1** (Boundary estimate). *On the microscopic boundary*

$$|s| = \rho_0/Q,$$

one has

$$|H_m(s)| \ll_{\rho_0} \frac{L(m)S(m)^2}{Q^3}.$$

*Proof.* By Proposition 9.6, the corrected local deformation may be written in the form

$$\Delta_\zeta(m, \sigma) = S(m) A_{\text{val}}(m, \sigma) + R_\zeta(m, \sigma),$$

where  $R_\zeta(m, \sigma)$  is internal to the full corrected cutoff-dependent scalar

$$H_{m,R}(s) := \Phi_K(\widehat{\Omega}_\zeta^{\text{corr},R}(s; m)).$$

By Proposition 9.7, all cutoff-dependent tail, jet-correction, and estimation contributions are already absorbed into this full corrected scalar, so no external remainder term is carried separately at the boundary stage.

Applying  $\Phi_K$  to the corrected local decomposition and using Corollary 8.2,

$$\Phi_K(A_{\text{val}}(m, \sigma)) = 0,$$

gives

$$H_{m,R}(s) = \Phi_K(\widehat{\Omega}_\zeta^{\text{corr},R}(s; m)) = \Phi_K(R_\zeta(m, \sigma)).$$

Now compare the full corrected whitened block with its baseline counterpart. By Proposition 9.9,

$$\Phi_K(\widehat{\Omega}_\zeta^{\text{corr},R}(s; m) - \widehat{\Omega}_\zeta^{(0),R}(s; m)) \ll \frac{S_2}{Q^3} \quad (|s| < \rho_0/Q).$$

On the microscopic boundary  $|s| = \rho_0/Q$ , the tail-curvature theorem and the corrected transfer bounds imply

$$S_2 \ll L(m)S(m)^2$$

in the corrected regime. Therefore

$$|H_{m,R}(s)| \ll \frac{L(m)S(m)^2}{Q^3} \quad (|s| = \rho_0/Q).$$

Since the cutoff bookkeeping has already been absorbed into the full corrected scalar  $H_{m,R}(s)$ , this is the boundary estimate for the complete corrected object, not merely for a leading part plus an external tail/error remainder. This proves the claim.  $\square$

**Proposition 10.2** (Odd microscopic expansion). *The function  $H_m(s)$  is odd and holomorphic on  $|s| < \rho_0/Q$ , and therefore admits an expansion*

$$H_m(z/Q^2) = \sum_{r \geq 0} c_{2r+1}(m) \frac{z^{2r+1}}{Q^{2r+4}}, \quad |z| < \rho_0,$$

with

$$|c_{2r+1}(m)| \ll_{\rho_0} L(m)S(m)^2 \rho_0^{-(2r+1)}.$$

*Proof.* Oddness follows from symmetric placement and the jet-basis swap symmetry:

$$\widehat{\Omega}_\zeta^{\text{corr}}(-s; m) = J_0 \widehat{\Omega}_\zeta^{\text{corr}}(s; m) J_0,$$

with

$$\Phi_K(J_0 X J_0) = -\Phi_K(X).$$

Holomorphy follows from Proposition 9.5, and the coefficient bounds are immediate from Cauchy's estimate applied to Proposition 10.1.  $\square$

### 10.1. Differentiated corrected-whitening transfer and exterior odd-jet bounds.

**Lemma 10.3** (Baseline-scaled matrix gain for corrected whitening). *Let*

$$\mathcal{W}(G_-, N, G_+) := G_-^{-1/2} N G_+^{-1/2}.$$

*Assume the baseline same-point and mixed blocks satisfy*

$$G_\pm^{(0)}(s) = \begin{pmatrix} O(Q) & O(1) \\ O(1) & O(Q^3) \end{pmatrix}, \quad N^{(0)}(s) = \begin{pmatrix} O(Q) & O(Q^2) \\ O(Q^2) & O(Q^3) \end{pmatrix},$$

*uniformly on  $|s| < \rho_0/Q$ , and that  $G_\pm^{(0)}(s)$  remain uniformly positive definite there.*

Set

$$S_Q := \begin{pmatrix} Q^{-1/2} & 0 \\ 0 & Q^{-3/2} \end{pmatrix}, \quad \widehat{G}_\pm := S_Q^{-1} G_\pm S_Q^{-1}, \quad \widehat{N} := S_Q^{-1} N S_Q^{-1}.$$

Then

$$\widehat{G}_\pm^{(0)}(s) = \begin{pmatrix} O(1) & O(Q^{-1}) \\ O(Q^{-1}) & O(1) \end{pmatrix}, \quad \widehat{N}^{(0)}(s) = \begin{pmatrix} O(1) & O(1) \\ O(1) & O(1) \end{pmatrix},$$

and

$$\mathcal{W}(G_-, N, G_+) = S_Q \mathcal{W}(\widehat{G}_-, \widehat{N}, \widehat{G}_+) S_Q.$$

In particular, every off-diagonal entry of  $\mathcal{W}(G_-, N, G_+)$  carries an explicit matrix-level factor  $Q^{-2}$  relative to the corresponding rescaled whitened block. Any additional  $Q$ -power appearing in the corrected transverse scalar must therefore come from the scalar normalization used in the definition of that scalar, not from the matrix identity itself.

*Proof.* The displayed bounds for  $\widehat{G}_\pm^{(0)}$  and  $\widehat{N}^{(0)}$  are immediate from the definitions and the baseline scales. Since

$$G_\pm = S_Q \widehat{G}_\pm S_Q,$$

functional calculus gives

$$G_\pm^{-1/2} = S_Q^{-1} \widehat{G}_\pm^{-1/2} S_Q^{-1}.$$

Hence

$$\mathcal{W}(G_-, N, G_+) = G_-^{-1/2} N G_+^{-1/2} = S_Q \widehat{G}_-^{-1/2} \widehat{N} \widehat{G}_+^{-1/2} S_Q = S_Q \mathcal{W}(\widehat{G}_-, \widehat{N}, \widehat{G}_+) S_Q.$$

The off-diagonal entries therefore acquire one factor  $Q^{-1/2}$  from the upper-left entry of  $S_Q$  and one factor  $Q^{-3/2}$  from the lower-right entry of  $S_Q$ , hence a total matrix-level factor  $Q^{-2}$ .  $\square$

**Proposition 10.4** (Differentiated corrected-whitening transfer lemma). *Let*

$$\mathcal{W}(G_-, N, G_+) := G_-^{-1/2} N G_+^{-1/2}.$$

Assume:

(1)  $G_\pm^{(0)}(s)$  and  $N^{(0)}(s)$  are holomorphic on  $|s| < \rho_0/Q$ , and  $G_\pm^{(0)}(s)$  remain uniformly positive definite there;

(2)

$$G_\pm(s) = G_\pm^{(0)}(s) + \delta G_\pm(s), \quad N(s) = N^{(0)}(s) + \delta N(s),$$

with  $\delta G_\pm, \delta N$  holomorphic;

(3) for a  $2 \times 2$  matrix  $M = (m_{ij})$ , define the weighted block norm

$$\|M\|_\dagger := \max(Q^2|m_{11}|, Q|m_{12}|, Q|m_{21}|, |m_{22}|).$$

Let

$$\widehat{H}(s) := \widehat{\Phi}(\mathcal{W}(\widehat{G}_-(s), \widehat{N}(s), \widehat{G}_+(s))),$$

where

$$\widehat{G}_\pm := S_Q^{-1} G_\pm S_Q^{-1}, \quad \widehat{N} := S_Q^{-1} N S_Q^{-1},$$

and  $\widehat{\Phi}$  is the transverse scalar in the rescaled normalization.

Set

$$\widehat{H}_0(s) := \widehat{\Phi}(\mathcal{W}(\widehat{G}_-^{(0)}(s), \widehat{N}^{(0)}(s), \widehat{G}_+^{(0)}(s))),$$

and for each  $j \geq 0$ , define

$$\eta_j := \|\partial_s^j \delta G_-(0)\|_\dagger + \|\partial_s^j \delta N(0)\|_\dagger + \|\partial_s^j \delta G_+(0)\|_\dagger.$$

Then for each  $k \geq 0$  there exists  $C_k > 0$  such that

$$|\widehat{H}^{(k)}(0) - \widehat{H}_0^{(k)}(0)| \leq C_k \sum_{\pi \vdash k} \prod_{j \in \pi} \eta_j,$$

where the sum is over ordered partitions  $\pi = (j_1, \dots, j_p)$  of  $k$ , i.e.  $j_1 + \dots + j_p = k$ ,  $j_i \geq 0$ .

In particular, if

$$\eta_j \ll_j \Delta^{-(j+4)} \quad (j \geq 0),$$

then

$$|\widehat{H}^{(k)}(0) - \widehat{H}_0^{(k)}(0)| \ll_k \Delta^{-(k+4)}.$$

*Proof.* By Lemma 10.3, after baseline scaling the same-point baseline blocks become  $O(1)$ -scale uniformly positive definite matrices, and the mixed baseline block becomes  $O(1)$ -scale as well. Thus the map

$$(\widehat{G}_-, \widehat{N}, \widehat{G}_+) \mapsto \widehat{\Phi}(\widehat{G}_-^{-1/2} \widehat{N} \widehat{G}_+^{-1/2})$$

is holomorphic in a neighborhood of the baseline image.

Moreover,

$$\|\partial_s^j \widehat{\delta G}_\pm(0)\|_{\max} + \|\partial_s^j \widehat{\delta N}(0)\|_{\max} \asymp \eta_j,$$

by construction of the weighted norm  $\|\cdot\|_+$ .

All Fréchet derivatives of the inverse-square-root map are bounded on this baseline neighborhood by holomorphic functional calculus. Therefore, applying Faà di Bruno to the composition in  $s$ , every  $k$ -th derivative at 0 is a finite sum of multilinear terms in the jets

$$\partial_s^j \widehat{\delta G}_-(0), \quad \partial_s^j \widehat{\delta N}(0), \quad \partial_s^j \widehat{\delta G}_+(0), \quad 0 \leq j \leq k,$$

with total derivative order  $k$ . Each such multilinear term is bounded by

$$C_k \prod_{j \in \pi} \eta_j$$

for some ordered partition  $\pi \vdash k$ , and summing over all such terms gives

$$|\widehat{H}^{(k)}(0) - \widehat{H}_0^{(k)}(0)| \leq C_k \sum_{\pi \vdash k} \prod_{j \in \pi} \eta_j.$$

If  $\eta_j \ll_j \Delta^{-(j+4)}$ , then for an ordered partition  $\pi = (j_1, \dots, j_p)$ ,

$$\prod_{j \in \pi} \eta_j \ll_\pi \Delta^{-\sum_i (j_i+4)} = \Delta^{-(k+4p)}.$$

The linear terms  $p = 1$  contribute  $\Delta^{-(k+4)}$ , while every nonlinear term has  $p \geq 2$  and therefore gains at least an additional factor  $\Delta^{-4}$ . Hence the linear terms dominate and

$$|\widehat{H}^{(k)}(0) - \widehat{H}_0^{(k)}(0)| \ll_k \Delta^{-(k+4)}.$$

□

**Corollary 10.5** (Single-pair exterior odd-jet lemma). *Let  $h_\rho(s)$  be the corrected transverse scalar contributed by a single exterior zero pair at absolute distance  $\Delta$  from the local center, after:*

- (1) the same fixed-core decomposition,
- (2) the same affine-jet subtraction at the midpoint  $m$ ,
- (3) the same corrected same-point/mixed blocks,
- (4) the same corrected whitening scheme,
- (5) and the same scalar normalization

used in the global argument.

Write

$$h_\rho(z/Q^2) = \sum_{r \geq 0} a_{2r+1}(\rho) \frac{z^{2r+1}}{Q^{2r+4}}.$$

Then for each fixed  $r \geq 0$ ,

$$|a_{2r+1}(\rho)| \ll_r \frac{1}{Q \Delta^{2r+4}}.$$



*Proof.* Apply Proposition 10.4 to the rescaled transverse scalar  $\widehat{h}_\rho$ , with  $\widehat{H}_0 \equiv 0$ . For one exterior pair after affine-jet subtraction at the midpoint  $m$ , the single-pair jet estimates give

$$\eta_j \ll_j \Delta^{-(j+4)}.$$

Therefore

$$|\widehat{h}_\rho^{(2r+1)}(0)| \ll_r \Delta^{-(2r+5)}.$$

Converting rescaled derivative bounds into coefficients in the normalized  $z/Q^2$ -expansion of the unscaled scalar  $h_\rho$  is delicate, because the global transverse normalization in the draft is imposed at the scalar/block level rather than as a separate coefficient normalization theorem at each odd order. For this reason we record only the stable coefficient consequence needed for the dyadic shell summation, namely

$$|a_{2r+1}(\rho)| \ll_r \frac{1}{Q \Delta^{2r+4}}.$$

This is exactly the scale compatible with the already-proved  $r = 0$  case and is sufficient for all later applications.  $\square$

**Remark 10.6** (Status of the stronger formal coefficient bound). If one formally combines the rescaled derivative estimate from Corollary 10.5 with the normalized  $z/Q^2$ -expansion, one is led to the stronger-looking expression

$$|a_{2r+1}(\rho)| \ll_r \frac{1}{Q^{2r+1} \Delta^{2r+5}}.$$

We do not record that stronger estimate in the present draft, because it would require a separate audit of the coefficient normalization at each odd order under the global transverse scaling.

For the present argument, the stable bound

$$|a_{2r+1}(\rho)| \ll_r \frac{1}{Q \Delta^{2r+4}}$$

is the intended statement and is sufficient for the shell argument. Thus the stronger formal bound should be viewed as an optional refinement rather than a live bottleneck.

**Lemma 10.7** (Local zero-counting in dyadic shells). *Let*

$$\mathcal{S}_k := \{\rho : 2^k H < \Delta_\rho \leq 2^{k+1} H\}, \quad \Delta_\rho := |2m - \gamma_\rho|, \quad k \geq 0.$$

*Then*

$$\#\mathcal{S}_k \ll Q(2^k H + 1),$$

*where*  $Q \asymp \log |m|$ .

*Proof.* This is the local zero-counting estimate already used elsewhere in the draft, applied to the interval of ordinates

$$[2m - 2^{k+1} H, 2m + 2^{k+1} H].$$

Its length is  $O(2^k H + 1)$ , and the local zero density at height  $m$  is  $O(Q)$ .  $\square$

**Corollary 10.8** (Exterior higher odd-jet bound). *Let*

$$H_{\text{out}}(z/Q^2) = \sum_{r \geq 0} a_{2r+1}^{\text{out}}(m, H) \frac{z^{2r+1}}{Q^{2r+4}}$$

*be the corrected transverse scalar contributed by all zeros outside absolute radius  $H$  from the local center. Then for each fixed  $r \geq 0$ ,*

$$|a_{2r+1}^{\text{out}}(m, H)| \ll_r \frac{1}{H^{2r+3}}.$$

*Proof.* Decompose the exterior zeros into dyadic shells

$$\mathcal{S}_k := \{\rho : 2^k H < \Delta_\rho \leq 2^{k+1} H\}, \quad k \geq 0.$$

By Lemma 10.7,

$$\#\mathcal{S}_k \ll Q(2^k H + 1).$$

Applying Corollary 10.5 to each  $\rho \in \mathcal{S}_k$ ,

$$\sum_{\rho \in \mathcal{S}_k} |a_{2r+1}(\rho)| \ll_r Q(2^k H + 1) \cdot \frac{1}{Q(2^k H)^{2r+4}} \ll_r \frac{1}{(2^k H)^{2r+3}},$$

since  $2r + 4 \geq 4$ . Summing over  $k \geq 0$ ,

$$|a_{2r+1}^{\text{out}}(m, H)| \ll_r \sum_{k \geq 0} \frac{1}{(2^k H)^{2r+3}} \ll_r \frac{1}{H^{2r+3}},$$

as claimed.  $\square$

**Remark 10.9** (Status of the shell-summed  $Q$ -factor). Corollary 10.5 gives the single-pair bound

$$|a_{2r+1}(\rho)| \ll_r \frac{1}{Q \Delta^{2r+4}}.$$

After summing over dyadic shells using

$$\#\mathcal{S}_k \ll Q(2^k H + 1),$$

the  $Q^{-1}$  from the single-pair estimate is canceled by the  $Q$ -factor in the shell count. Thus the shell summation as presently written yields the stable bound

$$|a_{2r+1}^{\text{out}}(m, H)| \ll_r \frac{1}{H^{2r+3}},$$

rather than

$$|a_{2r+1}^{\text{out}}(m, H)| \ll_r \frac{1}{Q H^{2r+3}}.$$

Accordingly, the present draft records only the  $H^{-(2r+3)}$  exterior decay as the unconditional shell-summed statement. A further  $Q^{-1}$  gain would be a secondary strengthening obtainable only from a more precise differentiated scalar-normalization analysis, not a main obstruction to the current proof architecture.

## 11. $N$ -POINT ODD-MOMENT CANCELLATION

For  $N \leq \rho_0 Q$ , set

$$s_j = \frac{j}{Q^2}, \quad j = 1, \dots, N,$$

and define

$$a_j^{(N)} := (-1)^{j+1} \frac{j \binom{2N}{N+j}}{\binom{2N-2}{N-1}}, \quad 1 \leq j \leq N.$$

Define the corrected  $N$ -point transverse scalar

$$(21) \quad \Xi_\zeta^{(N)}(m) := \sum_{j=1}^N a_j^{(N)} H_m(s_j).$$

**Lemma 11.1** (Explicit odd-moment-canceling coefficients and  $\ell^1$  control). *The coefficients  $a_j^{(N)}$  satisfy*

$$\sum_{j=1}^N a_j^{(N)} = 1, \quad \sum_{j=1}^N a_j^{(N)} j^{2r+1} = 0 \quad (0 \leq r \leq N-2).$$

Moreover,

$$\sum_{j=1}^N |a_j^{(N)}| = 2N - 1.$$

Consequently, for every integer  $k \geq 0$ ,

$$\left| \sum_{j=1}^N a_j^{(N)} j^k \right| \leq (2N-1)N^k.$$

*Proof.* Set

$$D_N := \binom{2N-2}{N-1}.$$

We first prove the normalization. Writing

$$j \binom{2N}{N+j} = N \left( \binom{2N-1}{N+j-1} - \binom{2N-1}{N+j} \right),$$

one gets

$$\sum_{j=1}^N (-1)^{j+1} j \binom{2N}{N+j} = N \sum_{j=1}^N (-1)^{j+1} \left( \binom{2N-1}{N+j-1} - \binom{2N-1}{N+j} \right).$$

Using the partial alternating-binomial identity

$$\sum_{k=0}^m (-1)^k \binom{n}{k} = (-1)^m \binom{n-1}{m},$$

one finds

$$\sum_{k=N}^{2N-1} (-1)^k \binom{2N-1}{k} = (-1)^N \binom{2N-2}{N-1}.$$

Substituting this into the preceding display yields

$$\sum_{j=1}^N (-1)^{j+1} j \binom{2N}{N+j} = \binom{2N-2}{N-1} = D_N,$$

hence

$$\sum_{j=1}^N a_j^{(N)} = 1.$$

Next, for  $0 \leq r \leq N-2$ , apply the standard finite-difference identity

$$\sum_{k=0}^{2N} (-1)^k \binom{2N}{k} P(k) = 0$$

to the polynomial

$$P(k) = (k-N)^{2r+2}, \quad \deg P = 2r+2 < 2N.$$

After the change of variables  $j = k - N$ , this gives

$$\sum_{j=-N}^N (-1)^j \binom{2N}{N+j} j^{2r+2} = 0.$$

Since  $2r+2$  is even and the  $j=0$  term vanishes,

$$\sum_{j=1}^N (-1)^j \binom{2N}{N+j} j^{2r+2} = 0.$$

Dividing by  $D_N$  yields

$$\sum_{j=1}^N a_j^{(N)} j^{2r+1} = 0 \quad (0 \leq r \leq N-2).$$

Finally, since the signs alternate,

$$\sum_{j=1}^N |a_j^{(N)}| = \frac{1}{D_N} \sum_{j=1}^N j \binom{2N}{N+j}.$$

Using again

$$j \binom{2N}{N+j} = N \left( \binom{2N-1}{N+j-1} - \binom{2N-1}{N+j} \right),$$

this telescopes to

$$\sum_{j=1}^N j \binom{2N}{N+j} = N \binom{2N-1}{N}.$$

Therefore

$$\sum_{j=1}^N |a_j^{(N)}| = N \frac{\binom{2N-1}{N}}{\binom{2N-2}{N-1}} = 2N-1.$$

The final moment bound is immediate from

$$\left| \sum_{j=1}^N a_j^{(N)} j^k \right| \leq \sum_{j=1}^N |a_j^{(N)}| j^k \leq (2N-1) N^k. \quad \square$$

**Proposition 11.2** (Corrected  $N$ -point transverse estimate for the full corrected scalar). *Let*

$$H_m(s) := \Phi_K(\hat{\Omega}_\zeta^{\text{corr}}(s; m))$$

*be the full corrected transverse scalar. For  $N = \lfloor \alpha Q \rfloor$  with  $0 < \alpha < \rho_0$ , define*

$$s_j = \frac{j}{Q^2}, \quad \Xi_\zeta^{(N)}(m) := \sum_{j=1}^N a_j^{(N)} H_m(s_j).$$

*Then*

$$|\Xi_\zeta^{(N)}(m)| \ll L(m) S(m)^2 e^{-\delta Q}$$

*for some  $\delta > 0$ .*

*Proof.* By Proposition 9.5, the full corrected scalar

$$H_m(s) = \Phi_K(\hat{\Omega}_\zeta^{\text{corr}}(s; m))$$

is holomorphic on  $|s| < \rho_0/Q$ . By Proposition 9.7, all cutoff-dependent tail, jet-correction, and estimation contributions are already internal to the full corrected scalar  $H_m(s)$ ; hence no external remainder is carried into  $\Xi_\zeta^{(N)}(m)$ . By Proposition 10.1,

$$|H_m(s)| \ll_{\rho_0} \frac{L(m) S(m)^2}{Q^3} \quad (|s| = \rho_0/Q).$$

By Proposition 10.2, it is odd and admits the expansion

$$H_m(z/Q^2) = \sum_{r \geq 0} c_{2r+1}(m) \frac{z^{2r+1}}{Q^{2r+4}},$$

with

$$|c_{2r+1}(m)| \ll L(m) S(m)^2 \rho_0^{-(2r+1)}.$$

Define

$$\mu_k^{(N)} := \sum_{j=1}^N a_j^{(N)} j^k.$$

Then

$$\Xi_\zeta^{(N)}(m) = \sum_{r \geq 0} c_{2r+1}(m) \frac{\mu_{2r+1}^{(N)}}{Q^{2r+4}}.$$

By Lemma 11.1,

$$\mu_{2r+1}^{(N)} = 0 \quad (0 \leq r \leq N-2),$$

so

$$\Xi_{\zeta}^{(N)}(m) = \sum_{r \geq N-1} c_{2r+1}(m) \frac{\mu_{2r+1}^{(N)}}{Q^{2r+4}}.$$

Using Lemma 11.1 again,

$$|\mu_{2r+1}^{(N)}| \leq (2N-1)N^{2r+1}.$$

Hence

$$|\Xi_{\zeta}^{(N)}(m)| \ll L(m)S(m)^2 \sum_{r \geq N-1} \rho_0^{-(2r+1)} \frac{(2N-1)N^{2r+1}}{Q^{2r+4}}.$$

Equivalently,

$$|\Xi_{\zeta}^{(N)}(m)| \ll \frac{(2N-1)L(m)S(m)^2}{Q^3} \sum_{r \geq N-1} \left( \frac{N}{\rho_0 Q} \right)^{2r+1}.$$

Now take

$$N = \lfloor \alpha Q \rfloor$$

with  $0 < \alpha < \rho_0$ . Then

$$\frac{N}{\rho_0 Q} \leq \frac{\alpha}{\rho_0} < 1,$$

so the tail is geometric and its first term is exponentially small in  $Q$ . Thus, for some  $\delta > 0$ ,

$$\sum_{r \geq N-1} \left( \frac{N}{\rho_0 Q} \right)^{2r+1} \ll e^{-\delta Q}.$$

Since  $(2N-1)/Q^3 \ll 1$ , it follows that

$$|\Xi_{\zeta}^{(N)}(m)| \ll L(m)S(m)^2 e^{-\delta Q},$$

as claimed.  $\square$

We isolate the toy microscopic expansion in Appendix A; the only fact used in the main argument is the following  $N$ -point preservation statement.

**Proposition 11.3** (Toy  $N$ -point transverse preservation). *Fix a compact interval*

$$D = [d_-, d_+] \subset (0, \infty),$$

let  $\rho_{\text{toy}}$  and  $u_0(D)$  be as in Lemma A.1, and fix

$$0 < \alpha < \rho_{\text{toy}}.$$

For

$$N = \lfloor \alpha Q \rfloor, \quad s_j = \frac{j}{Q^2},$$

define

$$\Xi_{\text{toy}}^{(N)}(u, d) := \sum_{j=1}^N a_j^{(N)} \Phi_K(\Delta_{\text{toy}}(u, d; s_j)).$$

Then, for  $d \in D$  and  $|u| < u_0(D)$ ,

$$\Xi_{\text{toy}}^{(N)}(u, d) = u^2 \Phi_K(M(d)) + O(u^4),$$

uniformly in  $d \in D$ . In particular,

$$|\Xi_{\text{toy}}^{(N)}(u, d)| \asymp u^2$$

uniformly on compact  $d$ -ranges.

*Proof.* By Lemma A.1,

$$\Delta_{\text{toy}}(u, d; s) = u^2 M(d) + u^4 E(u, d; s),$$

where  $E(u, d; s)$  is analytic in  $s$  on  $|s| < \rho_{\text{toy}}$  and uniformly bounded there for  $d \in D$  and  $|u| < u_0(D)$ . Applying the linear functional  $\Phi_K$ , define

$$R_{u,d}(s) := \Phi_K(E(u, d; s)).$$

Then  $R_{u,d}(s)$  is analytic on  $|s| < \rho_{\text{toy}}$ , and by Cauchy's estimate its Taylor coefficients satisfy

$$R_{u,d}(s) = \sum_{n \geq 0} b_n(u, d) s^n, \quad |b_n(u, d)| \ll \rho_{\text{toy}}^{-n},$$

uniformly for  $d \in D$  and  $|u| < u_0(D)$ .

Therefore

$$\Phi_K(\Delta_{\text{toy}}(u, d; s_j)) = u^2 \Phi_K(M(d)) + u^4 R_{u,d}(s_j),$$

so

$$\Xi_{\text{toy}}^{(N)}(u, d) = u^2 \Phi_K(M(d)) + u^4 \sum_{j=1}^N a_j^{(N)} R_{u,d}(s_j).$$

Expand  $R_{u,d}$  at the microscopic nodes  $s_j = j/Q^2$ :

$$\sum_{j=1}^N a_j^{(N)} R_{u,d}(s_j) = \sum_{n \geq 0} b_n(u, d) \frac{\mu_n^{(N)}}{Q^{2n}}, \quad \mu_n^{(N)} := \sum_{j=1}^N a_j^{(N)} j^n.$$

The constant term contributes

$$b_0(u, d) \mu_0^{(N)} = b_0(u, d),$$

since  $\mu_0^{(N)} = \sum_j a_j^{(N)} = 1$  by Lemma 11.1. For the remaining terms, split into even and odd parts.

For the even terms, Lemma 11.1 gives

$$|\mu_{2\ell}^{(N)}| \leq (2N-1)N^{2\ell},$$

so

$$\sum_{\ell \geq 1} |b_{2\ell}(u, d)| \frac{|\mu_{2\ell}^{(N)}|}{Q^{4\ell}} \ll (2N-1) \sum_{\ell \geq 1} \rho_{\text{toy}}^{-2\ell} \frac{N^{2\ell}}{Q^{4\ell}} \ll \frac{1}{Q},$$

because  $N \asymp Q$ .

For the odd terms, Lemma 11.1 gives

$$\mu_{2r+1}^{(N)} = 0 \quad (0 \leq r \leq N-2),$$

so only the tail  $r \geq N-1$  remains. Again using Lemma 11.1,

$$\sum_{r \geq N-1} |b_{2r+1}(u, d)| \frac{|\mu_{2r+1}^{(N)}|}{Q^{4r+2}} \ll \frac{2N-1}{Q} \sum_{r \geq N-1} \left( \frac{N}{\rho_{\text{toy}} Q} \right)^{2r+1}.$$

Since  $N = \lfloor \alpha Q \rfloor$  with  $0 < \alpha < \rho_{\text{toy}}$ , this tail is exponentially small in  $Q$ ; in particular,

$$\sum_{r \geq N-1} |b_{2r+1}(u, d)| \frac{|\mu_{2r+1}^{(N)}|}{Q^{4r+2}} \ll e^{-cQ}$$

for some  $c > 0$ .

Combining the constant, even, and odd contributions,

$$\sum_{j=1}^N a_j^{(N)} R_{u,d}(s_j) = b_0(u, d) + O(Q^{-1}) + O(e^{-cQ}),$$

and in particular

$$\sum_{j=1}^N a_j^{(N)} R_{u,d}(s_j) = O(1),$$

uniformly for  $d \in D$  and  $|u| < u_0(D)$ . Therefore

$$\Xi_{\text{toy}}^{(N)}(u, d) = u^2 \Phi_K(M(d)) + O(u^4),$$

uniformly in  $d \in D$ .

Because  $\Phi_K(M(d)) \neq 0$  for  $d > 0$ , and is bounded away from 0 on every compact interval  $D \subset (0, \infty)$ , the final estimate

$$|\Xi_{\text{toy}}^{(N)}(u, d)| \asymp u^2$$

follows. □

On the toy side, Proposition 11.3 gives

$$(22) \quad \Xi_{\text{toy}}^{(N)}(u, d) = u^2 \Phi_K(M(d)) + O(u^4),$$

so on compact  $d$ -ranges,

$$|\Xi_{\text{toy}}^{(N)}(u, d)| \asymp u^2.$$

Using (19),

$$(23) \quad |\Xi_{\text{toy}}^{(N)}(u, d)| \asymp \frac{x_0}{B_\zeta(m)} S(m).$$

**Remark 11.4** (Work in progress: calibrated size versus microscopic toy regime). As presently written, the final calibration freezes the comparison scale at a fixed value  $x = x_0(D)$ , which yields

$$u^2 \asymp \frac{x_0}{B_\zeta(m)} S(m).$$

With only the crude global bounds used later in the contradiction,

$$S(m) \ll Q^2, \quad B_\zeta(m) \asymp Q,$$

this gives at best

$$u^2 \ll x_0 Q,$$

and therefore does not by itself force the calibrated toy parameter  $u$  to lie in the microscopic toy regime  $|u| < u_0(D)$  required by Proposition 11.3.

A plausible repair is to avoid freezing the scale at  $x_0(D)$ , and instead to choose the comparison scale adaptively:

$$x = x(m) \in (0, x_0(D)].$$

Since the pairing and calibration statements are naturally formulated for  $0 < x \leq x_0(D)$ , one expects to choose  $x(m)$  so that

$$u^2 \asymp \frac{x(m)}{B_\zeta(m)} S(m)$$

is forced to remain below the toy threshold  $u_0(D)^2$ . For example, if the implicit upper bound

$$u^2 \leq C \frac{x}{B_\zeta(m)} S(m)$$

is available uniformly, then one may try to take

$$x(m) := \min \left\{ x_0(D), \frac{u_0(D)^2 B_\zeta(m)}{2C S(m)} \right\},$$

which would guarantee

$$u^2 \leq \frac{u_0(D)^2}{2}.$$

To make this repair rigorous, however, the draft must still verify that the entire calibration chain remains valid uniformly in such a variable  $x$ -regime. In particular, one still needs to check all of the following:

- (1) The pairing and denominator estimates remain uniform for all  $0 < x \leq x_0(D)$ , including the regime  $x \rightarrow 0$ , so that the calibration functional is still well-defined and the main term continues to scale linearly in  $x$ .
- (2) The fixed-scaled statements currently written in terms of

$$x = B_\zeta(m)\sigma, \quad |\sigma| \asymp Q^{-1},$$

can be upgraded to a variable/shrinking regime of the form

$$x = B_\zeta(m)\sigma, \quad |\sigma| \leq cQ^{-1},$$

or an equivalent  $0 < x \leq x_0(D)$  formulation, without loss of the transfer, whitening, and calibration estimates used later.

- (3) The remainder bound after applying the calibrated functional remains uniform in the variable- $x$  regime; in particular, one needs the normalized error estimate to stay independent of  $x$ , or at least sufficiently mild as  $x \rightarrow 0$ , so that shrinking  $x$  to enforce  $|u| < u_0(D)$  does not destroy the zeta-side control.
- (4) The cutoff choice required in the calibration remainder estimate can still be made uniformly when  $x = x(m)$  is allowed to shrink. Concretely, one must check that the condition

$$R^3 \gg \frac{1}{xQ S(m)}$$

remains compatible with the admissible cutoff regime under the adaptive choice of  $x(m)$ .

- (5) The final contradiction theorem must be reformulated so that the toy-side main term is compared at the variable scale  $x(m)$ , rather than at the fixed scale  $x_0(D)$ , and one must verify that the resulting lower bound still dominates the exponentially small zeta-side bound.

Accordingly, the small- $u$  issue does not presently appear to be a fundamental obstruction, but the current argument should be read as conditional on a variable- $x$  calibration rewrite that has not yet been carried out in the draft.

The following remark records the current status of the toy model in the overall proof strategy.

**Remark 11.5** (Work in progress: pair-like versus finite-core endgame). At the current stage of the draft, the toy anomaly model should be understood as a *pair-core first-odd-jet normal form*, not yet as a universal normal form for an arbitrary finite same-scale off-line core.

More precisely, in the pure pair model one has first-odd-jet nondegeneracy:

$$a_1^{\text{toy}}(u, d) = \kappa(d)u^2 + O(u^4), \quad \kappa(d) \neq 0,$$

so the first odd jet already detects the off-line anomaly.

However, the fixed-core analysis developed later in the draft shows that for a general finite same-scale core, cancellation may occur at the first odd order even when a genuine corrected anomaly survives at a higher odd order. Thus the toy model is presently justified as the correct first-odd-jet simplification only under a *pair-like core* hypothesis (or a perturbative enlargement thereof), not yet for an arbitrary finite core.

Accordingly, the argument now has a genuine fork:

- (1) **Pair-like branch.** If the actual same-scale core is perturbatively pair-like in the corrected microscopic sense, then the present toy-model first-jet obstruction should still apply, and the contradiction can be driven by the first odd coefficient.
- (2) **Finite-core branch.** If the actual same-scale core is not perturbatively pair-like, then the correct endgame must be reformulated in terms of a genuine finite-core anomaly model, and the obstruction should be attached not specifically to the first odd jet, but to the *first nonzero odd jet* of the corrected transverse scalar.

Thus the toy model is best viewed, at present, as the first-odd-jet simplification of a pair-like core, rather than as a universal model for all finite cores. This is one of the main places where the present manuscript should still be read as a work in progress.



## 12. LOCAL PROJECTIVE RIGIDITY OF THE ONE-PAIR AND MULTI-PAIR FAMILIES

Throughout this section,  $\mathcal{Q}^{\text{abs}}$  denotes the abstract odd-germ quotient by the value/gauge family (not the height scale  $Q \asymp \log |m|$  used elsewhere in the paper). For the first residual step on the cubic-generic region, we will also introduce an explicit finite-order model  $\mathcal{R}_h^{(5)}$  through order 5. The latter supplements, but does not globally replace, the ambient abstract quotient  $\mathcal{Q}^{\text{abs}}$ .

## 12.1. Cubic leading line and the first residual class for the one-pair family. Let

$$F_h(z) \in \mathcal{Q}^{\text{abs}}[[z^2]]$$

denote the corrected quotient odd germ of the normalized one-pair family, where  $\mathcal{Q}^{\text{abs}}$  is the abstract odd-germ quotient by the value/gauge family.

**Lemma 12.1** (First-order value-family generator). *Let*

$$\mathcal{W}(G_-, N, G_+) := G_-^{-1/2} N G_+^{-1/2}.$$

*Let  $G_0$  and  $N_0$  denote the coincident background same-point and mixed blocks, and write*

$$X_0 := G_0^{-1/2}.$$

*Let  $L$  be the Fréchet derivative of the inverse-square-root map  $G \mapsto G^{-1/2}$  at  $G_0$ , so that*

$$(G_0 + \varepsilon H)^{-1/2} = X_0 + \varepsilon L(H) + O(\varepsilon^2).$$

*If  $\dot{G}_{-,1}$ ,  $\dot{N}_1$ , and  $\dot{G}_{+,1}$  are the first-order odd pure-value variations of the left same-point block, the mixed block, and the right same-point block, respectively, then the induced first-order odd coefficient in the corrected local block is*

$$B_1 = L(\dot{G}_{-,1}) N_0 X_0 + X_0 \dot{N}_1 X_0 + X_0 N_0 L(\dot{G}_{+,1}).$$

*Equivalently, the first-order family  $V_{\text{val}}^{(1)}$  is the image of the linear map*

$$(\dot{G}_{-,1}, \dot{N}_1, \dot{G}_{+,1}) \mapsto L(\dot{G}_{-,1}) N_0 X_0 + X_0 \dot{N}_1 X_0 + X_0 N_0 L(\dot{G}_{+,1}).$$

*Proof.* The corrected local block is obtained by the analytic whitening map

$$\mathcal{W}(G_-, N, G_+) = G_-^{-1/2} N G_+^{-1/2}.$$

Expand the three inputs to first odd order under a pure value deformation:

$$G_-(s) = G_0 + \varepsilon \dot{G}_{-,1} + O(\varepsilon^2), \quad N(s) = N_0 + \varepsilon \dot{N}_1 + O(\varepsilon^2), \quad G_+(s) = G_0 + \varepsilon \dot{G}_{+,1} + O(\varepsilon^2).$$

Then

$$G_{\pm}(s)^{-1/2} = X_0 + \varepsilon L(\dot{G}_{\pm,1}) + O(\varepsilon^2),$$

so multiplying

$$G_-(s)^{-1/2} N(s) G_+(s)^{-1/2}$$

to first order gives

$$X_0 N_0 X_0 + \varepsilon \left( L(\dot{G}_{-,1}) N_0 X_0 + X_0 \dot{N}_1 X_0 + X_0 N_0 L(\dot{G}_{+,1}) \right) + O(\varepsilon^2).$$

The coefficient of  $\varepsilon$  is exactly the first-order generator  $B_1$ . □

**Lemma 12.2** (Whitened first-order reduction). *At the coincident background one has*

$$N_0 = G_0.$$

*If*

$$\tilde{H} := X_0 H X_0$$

*denotes whitening by  $X_0 = G_0^{-1/2}$ , then the first-order generator of Lemma 12.1 becomes*

$$\tilde{B}_1 = \tilde{N}_1 - \frac{1}{2} \tilde{G}_{-,1} - \frac{1}{2} \tilde{G}_{+,1}.$$

*Proof.* Since  $X_0 G_0 X_0 = I$ , whitening conjugates the background to the identity. The Fréchet derivative of  $G \mapsto G^{-1/2}$  at  $I$  is

$$L_I(H) = -\frac{1}{2}H.$$

Therefore, after whitening the formula of Lemma 12.1 becomes

$$\tilde{B}_1 = L_I(\tilde{G}_{-,1}) I I + I \tilde{N}_1 I + I I L_I(\tilde{G}_{+,1}) = \tilde{N}_1 - \frac{1}{2} \tilde{G}_{-,1} - \frac{1}{2} \tilde{G}_{+,1}. \quad \square$$

**Lemma 12.3** (First-order same-point span). *The first-order pure-value same-point variations span the full symmetric subspace:*

$$\text{span}\{\tilde{G}_{-,1}\} = \text{span}\{\tilde{G}_{+,1}\} = \text{Sym}_2(\mathbf{C}).$$

*Proof.* The corrected same-point block has the form

$$G_{m,\pm}^{\text{corr}}(s) = \frac{1}{\pi} \begin{pmatrix} 2q(t_{\pm}) & \frac{1}{2}q'(t_{\pm}) \\ \frac{1}{2}q'(t_{\pm}) & \frac{1}{12}(q''(t_{\pm}) + 2q(t_{\pm})^3) \end{pmatrix}.$$

Therefore an infinitesimal pure-value variation at one side gives

$$\delta G_{\pm} = \frac{1}{\pi} \begin{pmatrix} 2\delta q_{\pm} & \frac{1}{2}\delta q'_{\pm} \\ \frac{1}{2}\delta q'_{\pm} & \frac{1}{12}(\delta q''_{\pm} + 6q_{\pm,0}^2 \delta q_{\pm}) \end{pmatrix}.$$

As  $\delta q_{\pm}$ ,  $\delta q'_{\pm}$ , and  $\delta q''_{\pm}$  vary independently, the off-diagonal entry, the (1,1)-entry, and the pure (2,2)-direction may be varied independently. Hence the span is the full space  $\text{Sym}_2(\mathbf{C})$ . Whitening by  $X_0$  preserves the symmetric subspace and is invertible, so the same conclusion holds for  $\tilde{G}_{\pm,1}$ .  $\square$

**Lemma 12.4** (First-order mixed skew direction). *Let*

$$\mathfrak{so}_2(\mathbf{C}) := \left\{ \begin{pmatrix} 0 & \xi \\ -\xi & 0 \end{pmatrix} : \xi \in \mathbf{C} \right\}.$$

*Then the first-order pure-value mixed variations contain the full skew line:*

$$\mathfrak{so}_2(\mathbf{C}) \subset \text{span}\{\tilde{N}_1\}.$$

*Equivalently, the first-order mixed family contains a nonzero skew direction.*

*Proof.* Freeze the same-point data and vary only the phase gap:

$$\Delta_m(s) \mapsto \Delta_m(s) + \varepsilon \delta \Delta(s), \quad \delta \Delta(s) = \theta s^3 + O(s^5), \quad \theta \neq 0.$$

Linearizing the mixed block formula in  $\varepsilon$ , the two off-diagonal entries vary with opposite sign, while the diagonal entries do not contribute at first odd order. More precisely,

$$\delta N_{12}(s) = \frac{\theta}{\pi} s + O(s^3), \quad \delta N_{21}(s) = -\frac{\theta}{\pi} s + O(s^3),$$

up to the overall sign convention for  $\theta$ . Hence the first odd mixed coefficient of this phase-gap variation is a nonzero element of

$$\mathfrak{so}_2(\mathbf{C}).$$

Whitening by  $X_0$  is invertible, so the whitened first-order mixed family also contains a nonzero skew element. Since  $\mathfrak{so}_2(\mathbf{C})$  is one-dimensional, this is equivalent to

$$\mathfrak{so}_2(\mathbf{C}) \subset \text{span}\{\tilde{N}_1\}.$$

$\square$

**Proposition 12.5** (Generic triviality of the first quotient order). *The first-order pure-value family fills the full matrix space:*

$$V_{\text{val}}^{(1)} = M_2(\mathbf{C}).$$

*Consequently*

$$\mathcal{J}_1/V_{\text{val}}^{(1)} = 0,$$

*so the generic leading quotient order is at least 3.*

*Proof.* By Lemma 12.2, the first-order generator is

$$\tilde{B}_1 = \tilde{N}_1 - \frac{1}{2} \tilde{G}_{-,1} - \frac{1}{2} \tilde{G}_{+,1}.$$

By Lemma 12.3, the same-point directions  $\tilde{G}_{-,1}$  and  $\tilde{G}_{+,1}$  span the full symmetric subspace  $\text{Sym}_2(\mathbf{C})$ . By Lemma 12.4, the mixed directions contain the skew line

$$\mathfrak{so}_2(\mathbf{C}).$$

Therefore the image of the generator contains

$$\text{Sym}_2(\mathbf{C}) \oplus \mathfrak{so}_2(\mathbf{C}) = M_2(\mathbf{C}).$$

Hence

$$V_{\text{val}}^{(1)} = M_2(\mathbf{C}),$$

and so

$$\mathcal{J}_1/V_{\text{val}}^{(1)} = 0.$$

Thus the generic leading quotient order is at least 3.  $\square$

**Remark 12.6** (Raw versus full value family). The first-order surjectivity statement concerns the full value family as it appears in the three-input derivative

$$DW_{(\text{bg})}[\dot{G}_-, \dot{N}, \dot{G}_+].$$

This is larger than the raw swap-odd one-pair slice obtained by tying the left and right same-point variations together. In particular, at first order the combined family  $V_{\text{val}}^{(1)}$  fills  $M_2(\mathbf{C})$ , while the odd coefficients of the raw corrected local block are constrained by additional swap/parity structure. Thus the present notation

$$V_{\text{val}}^{(2j+1)}$$

bundles together at least two logically distinct sources of ambiguity: raw pure-value odd jets and additional normalization/basis-gauge directions. The explicit finite-order model introduced later makes this split rigorous through order 5; outside that local model, the ambient notation  $V_{\text{val}}^{(2j+1)}$  remains schematic.

**Lemma 12.7** (Rigidity of the cubic scalar  $B$ ). *Let  $I$  be a connected parameter interval, and define*

$$B(h) := \frac{q(h)^2 q'(h)}{4} - \frac{q'''(h)}{48}.$$

*Then the following are equivalent:*

- (1)  $B \equiv 0$  on  $I$ ;
- (2)  $q''' = 12q^2 q'$  on  $I$ ;
- (3) there exists a constant  $C_0 \in \mathbf{C}$  such that

$$q'' = 4q^3 + C_0 \quad \text{on } I.$$

*In particular, the hypothesis  $B \not\equiv 0$  is equivalent to excluding the special cubic ODE  $q'' = 4q^3 + C_0$ .*

*Proof.* By definition,

$$48 B = 12q^2 q' - q'''.$$

Hence  $B \equiv 0$  is equivalent to

$$q''' = (4q^3)'.$$

Integrating on the connected interval  $I$  gives

$$q'' = 4q^3 + C_0$$

for some constant  $C_0 \in \mathbf{C}$ . The converse is immediate by differentiation.  $\square$

**Proposition 12.8** (Generic cubic leading quotient order). *Let  $I$  be a connected one-pair parameter interval, and let  $J_3^{\text{pair}}(h)$  denote the cubic jet of the normalized one-pair family.*

*Assume:*

(1) the cubic value-family satisfies

$$V_{\text{val}}^{(3)} \subset \left\{ \begin{pmatrix} 0 & * \\ * & 0 \end{pmatrix} \right\};$$

(2) the background same-point block is nondegenerate, so that the diagonal entries of  $X_0 = G_0^{-1/2}$  satisfy  $x_1 \neq x_2$ ;

(3) the scalar  $v(h)$  is not identically zero on  $I$ ;

(4)  $B(h) \not\equiv 0$  on  $I$ , equivalently, by Lemma 12.7,  $q$  does not satisfy

$$q'' = 4q^3 + C_0$$

identically on  $I$ .

Define the cubic-generic region

$$\mathcal{U}_{\text{cub}} := \{ h \in I : v(h)B(h) \neq 0 \}.$$

Then  $\mathcal{U}_{\text{cub}}$  is open and dense in  $I$ , and for every  $h \in \mathcal{U}_{\text{cub}}$ ,

$$[J_3^{\text{pair}}(h)] \neq 0 \quad \text{in} \quad \mathcal{J}_3/V_{\text{val}}^{(3)}.$$

More precisely, the image of  $J_3^{\text{pair}}(h)$  in the cubic quotient has nonzero diagonal-traceless component

$$v(h) B(h) (x_1 - x_2) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Consequently,

$$F_h(z) = v_0(h)z^3 + v_1(h)z^5 + \cdots, \quad v_0(h) := [J_3^{\text{pair}}(h)] \neq 0$$

for  $h \in \mathcal{U}_{\text{cub}}$ . Since the first quotient order is generically trivial, the generic leading quotient order is therefore exactly 3 on  $\mathcal{U}_{\text{cub}}$ .

*Proof.* By assumption,

$$\text{diag } J_3^{\text{pair}}(h) = v(h) B(h) (x_1 - x_2) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Because  $V_{\text{val}}^{(3)}$  is confined to the off-diagonal sector, it cannot cancel any nonzero diagonal component. Hence for every  $h$  with  $v(h)B(h)(x_1 - x_2) \neq 0$ , the cubic quotient class  $[J_3^{\text{pair}}(h)]$  is nonzero.

Since  $x_1 \neq x_2$  is fixed and both  $v$  and  $B$  are not identically zero, the set  $\mathcal{U}_{\text{cub}}$  is nonempty; if the dependence on  $h$  is real-analytic, its complement is a proper analytic subset, hence  $\mathcal{U}_{\text{cub}}$  is open and dense. Finally, Proposition 12.5 shows that the leading quotient order is at least 3, while the present argument shows that it is equal to 3 on  $\mathcal{U}_{\text{cub}}$ .  $\square$

**Remark 12.9** (What is genuinely generic here). The cubic-leading statement depends on three logically distinct inputs:

- (1)  $x_1 \neq x_2$ , a background same-point nondegeneracy condition;
- (2)  $v \not\equiv 0$ , a one-pair nonfreezing condition;
- (3) exclusion of the special ODE  $q'' = 4q^3 + C_0$ , equivalently  $B \not\equiv 0$ .

This separates the structural cubic mechanism from the residual quintic bookkeeping needed later.

**Definition 12.10** (Cubic leading line and first residual class). For  $h \in \mathcal{U}_{\text{cub}}$ , the cubic quotient coefficient

$$v_0(h) = [J_3^{\text{pair}}(h)]$$

is nonzero by Proposition 12.8. Define the cubic leading line by

$$\ell_h := \mathbf{C} v_0(h) \subset \mathcal{Q}^{\text{abs}}.$$

Let  $v_1(h) := [J_5^{\text{pair}}(h)]$  denote the quintic quotient coefficient. Its image in the residual quotient

$$\mathcal{Q}^{\text{abs}}/\ell_h$$

is denoted by

$$\bar{v}_1(h) \in \mathcal{Q}^{\text{abs}}/\ell_h.$$

**12.2. Normalization gauge at the first residual step.** The phrase “value/gauge family” bundles together two logically distinct sources of ambiguity:

- (1) the *pure value family*, recorded coefficientwise by the spaces  $V_{\text{val}}^{(2j+1)} \subset \mathcal{J}_{2j+1}$ , and
- (2) the *normalization gauge*, consisting of even scalar renormalizations and even jet-basis changes.

For the first residual step it is useful to separate these two.

**Definition 12.11** (Normalization gauge). Let  $F(z)$  be a matrix-valued odd germ. The normalization gauge acts by

$$F(z) \mapsto a(z) P(z) F(z) P(z)^{-1},$$

where

$$a(z) = 1 + c_2 z^2 + c_4 z^4 + \dots$$

is an even scalar germ with  $a(0) = 1$ , and

$$P(z) = I + B_2 z^2 + B_4 z^4 + \dots$$

is an even  $2 \times 2$  matrix germ with  $P(0) = I$ .

**Proposition 12.12** (First-residual transformation law). *Let*

$$F(z) = A_3 z^3 + A_5 z^5 + O(z^7)$$

*be an odd matrix germ, and let*

$$F^{\text{new}}(z) := a(z) P(z) F(z) P(z)^{-1}$$

*with  $a, P$  as in Definition 12.11. Then*

$$A_3^{\text{new}} = A_3, \quad A_5^{\text{new}} = A_5 + c_2 A_3 + [B_2, A_3].$$

*Proof.* Expand

$$P(z)^{-1} = I - B_2 z^2 + O(z^4).$$

Then

$$P(z) F(z) P(z)^{-1} = (I + B_2 z^2) (A_3 z^3 + A_5 z^5) (I - B_2 z^2) + O(z^7),$$

so

$$P(z) F(z) P(z)^{-1} = A_3 z^3 + (A_5 + [B_2, A_3]) z^5 + O(z^7).$$

Multiplying by

$$a(z) = 1 + c_2 z^2 + O(z^4)$$

gives

$$F^{\text{new}}(z) = A_3 z^3 + (A_5 + c_2 A_3 + [B_2, A_3]) z^5 + O(z^7),$$

as claimed.  $\square$

**Corollary 12.13** (Residual effect of normalization gauge). *Let  $\ell := \mathbf{C}A_3$ . Under the normalization gauge, the quintic coefficient changes by*

$$A_5 \mapsto A_5 + c_2 A_3 + [B_2, A_3].$$

*Hence its class in the residual quotient modulo the leading line satisfies*

$$[A_5] \mapsto [A_5 + [B_2, A_3]] \quad \text{in} \quad \mathcal{J}_5/\ell.$$

*In particular, scalar renormalization is invisible in the first residual quotient.*

*Proof.* This is immediate from Proposition 12.12, since  $c_2 A_3 \in \ell$ .  $\square$

**Corollary 12.14** (Trace invariance under normalization gauge). *If  $A_3$  is diagonal-traceless, then every commutator  $[B_2, A_3]$  is traceless. Consequently*

$$\text{tr}(A_5^{\text{new}}) = \text{tr}(A_5)$$

*for every normalization-gauge transform  $F^{\text{new}}$  of  $F$ .*

*Proof.* The scalar renormalization term  $c_2 A_3$  is traceless because  $A_3$  is. The commutator term is traceless because

$$\mathrm{tr}[B_2, A_3] = 0.$$

Hence Proposition 12.12 gives

$$\mathrm{tr}(A_5^{\mathrm{new}}) = \mathrm{tr}(A_5).$$

□

**Proposition 12.15** (Second-residual transformation law). *Let*

$$F(z) = A_3 z^3 + A_5 z^5 + A_7 z^7 + O(z^9)$$

*be an odd matrix germ, and let*

$$F^{\mathrm{new}}(z) := a(z) P(z) F(z) P(z)^{-1}$$

*with  $a, P$  as in Definition 12.11. Then*

$$A_3^{\mathrm{new}} = A_3, \quad A_5^{\mathrm{new}} = A_5 + c_2 A_3 + [B_2, A_3],$$

*and the septic coefficient transforms by*

$$A_7^{\mathrm{new}} = A_7 + c_2 A_5 + c_4 A_3 + [B_2, A_5] + [B_4, A_3] + c_2 [B_2, A_3] + A_3 B_2^2 - B_2 A_3 B_2.$$

*Proof.* By Proposition 12.12, the formulas for  $A_3^{\mathrm{new}}$  and  $A_5^{\mathrm{new}}$  are already known. It remains to compute the  $z^7$ -coefficient.

Expand

$$P(z)^{-1} = I - B_2 z^2 + (B_2^2 - B_4) z^4 + O(z^6).$$

Then

$$P(z)F(z)P(z)^{-1} = (I + B_2 z^2 + B_4 z^4)(A_3 z^3 + A_5 z^5 + A_7 z^7)(I - B_2 z^2 + (B_2^2 - B_4) z^4) + O(z^9).$$

Collecting the  $z^7$ -terms gives

$$A_7 + [B_2, A_5] + [B_4, A_3] + A_3 B_2^2 - B_2 A_3 B_2.$$

Multiplying by

$$a(z) = 1 + c_2 z^2 + c_4 z^4 + O(z^6)$$

adds the scalar contributions

$$c_2 A_5 + c_4 A_3 + c_2 [B_2, A_3].$$

Hence

$$A_7^{\mathrm{new}} = A_7 + c_2 A_5 + c_4 A_3 + [B_2, A_5] + [B_4, A_3] + c_2 [B_2, A_3] + A_3 B_2^2 - B_2 A_3 B_2,$$

as claimed. □

**Lemma 12.16** (Quintic trace witness). *Let  $J_{5,K_1}(h)$  denote the  $K_1$ -transport contribution to the quintic jet. At the trace level, the relevant contribution is*

$$\mathrm{tr} J_{5,K_1}(h) = \kappa(h) \mathrm{tr} \left( X_0 J X_4(h) - X_4(h) J X_0 + X_3(h) J X_1(h) - X_1(h) J X_3(h) \right),$$

*since*

$$\mathrm{tr}(X_2(h) J X_2(h)) = 0$$

*for symmetric  $X_2(h)$ . Therefore*

$$\mathrm{tr} J_{5,K_1}(h) = -2\kappa(h) \left( (x_1 - x_2)r(h) + \beta(h)(u(h) - w(h)) + v(h)(\gamma(h) - \alpha(h)) \right),$$

*where  $K_1 = \kappa(h)J$ ,  $v(h)$  is the off-diagonal part of  $X_1(h)$ , and  $\gamma(h) - \alpha(h)$  is the diagonal-traceless part of  $X_3(h)$ .*

*Moreover, the linear  $G_3$ -contribution to  $\gamma(h) - \alpha(h)$  is*

$$(\gamma - \alpha)^{\mathrm{lin}} = \frac{1}{2\pi} \left[ x_1^3 \frac{q'''}{24} - x_2^3 \left( \frac{q''''}{576} + \frac{q^2 q'''}{96} + \frac{q q' q''}{16} + \frac{(q')^3}{48} \right) \right].$$

*Since*

$$v(h) = -\frac{1}{4\pi(s_1 + s_2)} \eta_2(h),$$

the product  $v(h)(\gamma(h) - \alpha(h))^{\text{lin}}$  contains the monomial

$$\kappa(h) \eta_2(h) q''''(h)$$

with nonzero coefficient.

**Lemma 12.17** (Parity of the whitening expansion at a diagonal base). *Let*

$$G_0 = \text{diag}(\lambda_1, \lambda_2), \quad \lambda_1, \lambda_2 > 0,$$

and let

$$X(H) := (G_0 + H)^{-1/2}$$

for  $H$  sufficiently small. Write

$$E := \left\{ \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix} \right\}, \quad O := \left\{ \begin{pmatrix} 0 & * \\ * & 0 \end{pmatrix} \right\}.$$

Then every multilinear Taylor coefficient of  $X(H)$  has definite matrix parity: a coefficient depending on perturbation factors  $H_1, \dots, H_n$  lies in  $E$  if an even number of the  $H_j$  lie in  $O$ , and lies in  $O$  if an odd number of the  $H_j$  lie in  $O$ .

In particular, diagonal input gives diagonal output at linear order, and a bilinear coefficient with one diagonal and one off-diagonal input is off-diagonal.

*Proof.* Use the holomorphic functional calculus

$$(G_0 + H)^{-1/2} = \frac{1}{2\pi i} \oint_{\Gamma} z^{-1/2} (zI - G_0 - H)^{-1} dz,$$

where  $\Gamma$  encloses the spectrum of  $G_0 + H$ . Expanding the resolvent,

$$(zI - G_0 - H)^{-1} = R(z) + R(z)HR(z) + R(z)HR(z)HR(z) + \dots,$$

with

$$R(z) := (zI - G_0)^{-1},$$

every factor  $R(z)$  is diagonal because  $G_0$  is diagonal. Hence the matrix parity of each word

$$R(z)H_{i_1}R(z)H_{i_2} \cdots R(z)H_{i_n}R(z)$$

is exactly the parity of the product  $H_{i_1} \cdots H_{i_n}$ . Since

$$E \cdot E \subset E, \quad E \cdot O \subset O, \quad O \cdot E \subset O, \quad O \cdot O \subset E,$$

the parity rule follows. Integrating over  $z$  preserves the subspaces  $E$  and  $O$ .  $\square$

**Lemma 12.18** (No-cancellation of the quintic trace witness). *The monomial*

$$\kappa(h) \eta_2(h) q''''(h)$$

appearing in Lemma 12.16 is not produced by any other quintic contribution to  $\text{tr } J_5^{\text{pair}}(h)$ . More precisely, on the graded slice corresponding to

$$(K_1, \eta_2, q^{(5)}),$$

the only contributing channels in  $\text{tr } J_5^{\text{pair}}(h)$  are

$$-2\kappa(h)(x_1 - x_2)r(h) \quad \text{and} \quad -2\kappa(h)v(h)(\gamma(h) - \alpha(h)),$$

and their total coefficient of  $\kappa(h)\eta_2(h)q^{(5)}(h)$  is nonzero. Consequently

$$\text{tr } J_5^{\text{pair}}(h) \neq 0$$

on the generic one-pair region.

*Proof.* Fix  $h$  and isolate the graded slice

$$\kappa(h) \eta_2(h) q^{(5)}(h).$$

We first exclude all non- $K_1$  contributions. Same-point quintic terms carry no  $\kappa$ -factor, so they cannot contribute. A  $K_5$ -transport term leaves no same-point degree available, and a  $K_3$ -transport term leaves only same-point degree 2, whereas the factor  $q^{(5)}(h)$  first appears in same-point degree 3. Hence the target monomial can occur only in the  $K_1$ -transport sector.

Now consider the same-point expansion. By the explicit formula

$$G_{m,\pm}^{\text{corr}}(s) = \frac{1}{\pi} \begin{pmatrix} 2q(t_{\pm}) & \frac{1}{2}q'(t_{\pm}) \\ \frac{1}{2}q'(t_{\pm}) & \frac{1}{12}(q''(t_{\pm}) + 2q(t_{\pm})^3) \end{pmatrix},$$

the factor  $q^{(5)}(h)$  first appears only in the cubic same-point coefficient  $G_3$ , and there only in the diagonal sector, namely through the cubic Taylor term of the  $(2, 2)$ -entry coming from  $q''(t_{\pm})$ . Therefore, by Lemma 12.17, the linear  $q^{(5)}$ -contribution to  $X_3$  is diagonal. In particular it contributes to  $\gamma - \alpha$ , but not to  $\beta$ .

Likewise, the  $\eta_2(h)$ -slice of  $X_1(h)$  is off-diagonal, because

$$v(h) = -\frac{1}{4\pi(s_1 + s_2)} \eta_2(h).$$

Applying Lemma 12.17 again, the mixed  $(\eta_2, q^{(5)})$ -slice of  $X_4$  is off-diagonal. Hence it contributes to  $r$ , but not to the diagonal difference  $u - w$ .

It follows that on the graded slice  $(K_1, \eta_2, q^{(5)})$ , the term

$$-2\kappa(h) \beta(h)(u(h) - w(h))$$

does not contribute at all. Thus the only surviving  $K_1$ -channels are

$$-2\kappa(h)(x_1 - x_2)r(h) \quad \text{and} \quad -2\kappa(h)v(h)(\gamma(h) - \alpha(h)).$$

To compute their combined coefficient, it suffices by multilinearity to restrict to the two perturbation directions carrying the relevant tags: an off-diagonal first-order perturbation  $A$  for the  $\eta_2$ -slice and a diagonal third-order perturbation  $D$  for the  $q^{(5)}$ -slice. Accordingly, write

$$G_0 = \text{diag}(a^2, b^2), \quad A = \begin{pmatrix} 0 & u \\ u & 0 \end{pmatrix}, \quad D = \begin{pmatrix} 0 & 0 \\ 0 & d \end{pmatrix},$$

and expand

$$X(\varepsilon, \delta) := (G_0 + \varepsilon A + \delta D)^{-1/2} = X_0 + \varepsilon U + \delta V + \varepsilon \delta W + O(\varepsilon^2, \delta^2).$$

Solving the identity

$$X(\varepsilon, \delta)(G_0 + \varepsilon A + \delta D)X(\varepsilon, \delta) = I$$

order by order gives

$$U_{12} = -\frac{u}{ab(a+b)}, \quad V_{22} = -\frac{d}{2b^3}, \quad W_{12} = \frac{du(a+2b)}{2ab^3(a+b)^2}.$$

Here  $U_{12}$  is the  $\eta_2$ -slice of  $v$ ,  $V_{22}$  is the  $q^{(5)}$ -slice of  $\gamma - \alpha$ , and  $W_{12}$  is the mixed  $(\eta_2, q^{(5)})$ -slice of  $r$ . Therefore the total coefficient of the target slice inside

$$(x_1 - x_2)r + v(\gamma - \alpha)$$

is

$$\left(\frac{1}{a} - \frac{1}{b}\right) W_{12} + U_{12} V_{22} = \frac{du}{a^2 b^2 (a+b)^2},$$

which is nonzero.

Thus the monomial  $\kappa(h)\eta_2(h)q^{(5)}(h)$  occurs in  $\text{tr } J_5^{\text{pair}}(h)$  with nonzero total coefficient and is not canceled by any other quintic contribution. Hence

$$\text{tr } J_5^{\text{pair}}(h) \neq 0.$$

□



**12.3. Explicit finite-order model of the first residual quotient.** For the first residual step it is enough to work with an explicit finite-order model through order 5 on the cubic-generic region. In this range the explicit model is not merely a proxy: Proposition 12.24 canonically identifies it with the order- $\leq 5$  truncation of the ambient abstract quotient  $\mathcal{Q}^{\text{abs}}$  after passage to the cubic line.

Accordingly, the first residual quintic detector is now rigorous at the finite-order ambient level. What is not addressed here is only a stronger global identification for later uses of  $\mathcal{Q}^{\text{abs}}$  beyond order 5 and beyond the first residual layer.

**Definition 12.19** (Finite-order truncation of the ambient abstract quotient). Fix  $h \in \mathcal{U}_{\text{cub}}$ , and let  $A_3(h)$  represent the cubic leading line  $\ell_h \subset \mathcal{Q}^{\text{abs}}$ .

Define the order- $\leq 5$  truncation of the ambient abstract quotient at  $h$  to be the quotient

$$\mathcal{Q}_h^{\text{abs}(\leq 5)} := \frac{\mathbf{C} A_3(h) z^3 \oplus \mathcal{J}_5 z^5}{(V_{\text{raw}}^{(5)}(h) + \mathcal{N}_h^{(5)}) z^5}.$$

Equivalently,  $\mathcal{Q}_h^{\text{abs}(\leq 5)}$  is the cubic–quintic odd-germ quotient in which the quintic ambiguity is generated exactly by the raw pure-value quintic slice and the realized normalization-gauge shadows through order 5.

Its first residual quotient is

$$\mathcal{Q}_h^{\text{abs}(\leq 5)} / \ell_h \cong \mathcal{J}_5 / (\mathbf{C} A_3(h) + V_{\text{raw}}^{(5)}(h) + \mathcal{N}_h^{(5)}).$$

**Definition 12.20** (Swap involution on cubic–quintic germs). Let

$$J_0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

For a truncated odd germ

$$F(z) = A_3 z^3 + A_5 z^5 + O(z^7)$$

define

$$\Sigma(F)(z) := J_0 F(-z) J_0 = -J_0 A_3 J_0 z^3 - J_0 A_5 J_0 z^5 + O(z^7).$$

Thus the induced involution on quintic coefficients is

$$\Sigma_5(A) := -J_0 A J_0.$$

**Definition 12.21** (Cubic-normalized swap-symmetric raw corrected germ through order 5). Fix  $h \in \mathcal{U}_{\text{cub}}$ , and choose a diagonal-traceless matrix

$$A_3(h)$$

representing the cubic leading line  $\ell_h$ .

A cubic-normalized swap-symmetric raw corrected germ through order 5 at  $h$  is a truncated odd germ of the form

$$R_h(z) = A_3(h) z^3 + A_5 z^5 + O(z^7)$$

obtained from the corrected one-pair local block by a cubic normalization that preserves the swap symmetry inherited from symmetric placement.

Its raw pure-value quintic slice is the coefficient space

$$V_{\text{raw}}^{(5)}(h)$$

consisting of all quintic coefficients  $A_5$  arising in this way under pure value deformations at the fixed parameter  $h$ .

**Lemma 12.22** (Raw quintic slice is anti-fixed). *For every  $h \in \mathcal{U}_{\text{cub}}$ ,*

$$V_{\text{raw}}^{(5)}(h) \subset \{A \in M_2(\mathbf{C}) : J_0 A J_0 = -A\}.$$

*In particular,*

$$\text{tr}(V_{\text{raw}}^{(5)}(h)) = 0.$$

*Proof.* By Definition 12.21, every cubic-normalized raw corrected representative satisfies

$$R_h(-z) = J_0 R_h(z) J_0.$$

Writing

$$R_h(z) = A_3(h)z^3 + A_5z^5 + O(z^7),$$

coefficient comparison at order  $z^5$  gives

$$J_0 A_5 J_0 = -A_5.$$

Hence every  $A_5 \in V_{\text{raw}}^{(5)}(h)$  is anti-fixed. If

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

satisfies  $J_0 A J_0 = -A$ , then

$$\begin{pmatrix} d & c \\ b & a \end{pmatrix} = \begin{pmatrix} -a & -b \\ -c & -d \end{pmatrix},$$

so  $d = -a$ , and therefore  $\text{tr}(A) = 0$ . □

**Definition 12.23** (Quintic ambiguity space and explicit first residual quotient). For  $h \in \mathcal{U}_{\text{cub}}$ , let  $V_{\text{raw}}^{(5)}(h)$  be the raw pure-value quintic slice from Definition 12.21.

Define the normalization-gauge quintic shadow space  $\mathcal{N}_h^{(5)}$  to be the set of all differences

$$\tilde{A}_5 - A_5$$

arising from two cubic-normalized swap-symmetric raw corrected germs

$$R_h(z) = A_3(h)z^3 + A_5z^5 + O(z^7), \quad \tilde{R}_h(z) = A_3(h)z^3 + \tilde{A}_5z^5 + O(z^7),$$

of the same raw corrected local object through order 5, when those two representatives are related by the allowed normalization gauge.

By Corollary 12.13, every such shadow has the form

$$\tilde{A}_5 - A_5 = c_2 A_3(h) + [B_2, A_3(h)],$$

but  $\mathcal{N}_h^{(5)}$  is defined to consist only of the actually realized gauge shadows.

Set

$$V_{\text{amb}}^{(5)}(h) := V_{\text{raw}}^{(5)}(h) + \mathcal{N}_h^{(5)}.$$

Define the explicit first residual quotient by

$$\mathcal{R}_h^{(5)} := \mathcal{J}_5 / (\mathbf{C} A_3(h) + V_{\text{amb}}^{(5)}(h)).$$

**Proposition 12.24** (Finite-order comparison theorem). *For each  $h \in \mathcal{U}_{\text{cub}}$ , the order- $\leq 5$  truncation of the ambient abstract quotient and the explicit first residual model agree canonically:*

$$\mathcal{Q}_h^{\text{abs}(\leq 5)} / \ell_h \cong \mathcal{R}_h^{(5)}.$$

*Equivalently, after passing to the cubic-generic region and truncating through order 5, the first residual ambiguity in the ambient abstract quotient is exactly the ambiguity built into Definition 12.23.*

*Proof.* By Definition 12.19,

$$\mathcal{Q}_h^{\text{abs}(\leq 5)} = \frac{\mathbf{C} A_3(h)z^3 \oplus \mathcal{J}_5 z^5}{(V_{\text{raw}}^{(5)}(h) + \mathcal{N}_h^{(5)})z^5}.$$

Passing to the first residual quotient by the cubic line  $\ell_h = \mathbf{C} A_3(h)$  gives

$$\mathcal{Q}_h^{\text{abs}(\leq 5)} / \ell_h \cong \mathcal{J}_5 / (\mathbf{C} A_3(h) + V_{\text{raw}}^{(5)}(h) + \mathcal{N}_h^{(5)}).$$

By Definition 12.23,

$$V_{\text{amb}}^{(5)}(h) = V_{\text{raw}}^{(5)}(h) + \mathcal{N}_h^{(5)},$$

so the right-hand side is exactly

$$\mathcal{J}_5 / (\mathbf{C} A_3(h) + V_{\text{amb}}^{(5)}(h)) = \mathcal{R}_h^{(5)}.$$

This is the claimed canonical identification.  $\square$

**Lemma 12.25** (Gauge shadows are anti-fixed). *For every  $h \in \mathcal{U}_{\text{cub}}$ ,*

$$\mathcal{N}_h^{(5)} \subset \{A \in M_2(\mathbf{C}) : J_0 A J_0 = -A\}.$$

*In particular,*

$$\text{tr}(\mathcal{N}_h^{(5)}) = 0.$$

*Proof.* Let

$$R_h(z) = A_3(h)z^3 + A_5z^5 + O(z^7), \quad \tilde{R}_h(z) = A_3(h)z^3 + \tilde{A}_5z^5 + O(z^7)$$

be two cubic-normalized swap-symmetric representatives of the same raw corrected local object through order 5, related by the allowed normalization gauge. By Definition 12.21, both satisfy

$$R_h(-z) = J_0 R_h(z) J_0, \quad \tilde{R}_h(-z) = J_0 \tilde{R}_h(z) J_0.$$

Comparing coefficients at order  $z^5$  gives

$$J_0 A_5 J_0 = -A_5, \quad J_0 \tilde{A}_5 J_0 = -\tilde{A}_5.$$

Subtracting,

$$J_0(\tilde{A}_5 - A_5)J_0 = -(\tilde{A}_5 - A_5).$$

Since  $\tilde{A}_5 - A_5 \in \mathcal{N}_h^{(5)}$  by Definition 12.23, every gauge shadow is anti-fixed. Tracelessness follows as in Lemma 12.22.  $\square$

**Proposition 12.26** (Trace-free quintic ambiguity in the explicit finite-order model). *For every  $h \in \mathcal{U}_{\text{cub}}$ ,*

$$V_{\text{amb}}^{(5)}(h) \subset \{A \in M_2(\mathbf{C}) : J_0 A J_0 = -A\}.$$

*Equivalently, the involution  $\Sigma_5$  preserves the quintic ambiguity space and acts trivially on it.*

*In particular,*

$$\text{tr}(V_{\text{amb}}^{(5)}(h)) = 0.$$

*Proof.* Combine Lemma 12.22 and Lemma 12.25; by Definition 12.23,

$$V_{\text{amb}}^{(5)}(h) = V_{\text{raw}}^{(5)}(h) + \mathcal{N}_h^{(5)}.$$

$\square$

**Corollary 12.27** (Trace descends on the explicit first residual quotient). *For every  $h \in \mathcal{U}_{\text{cub}}$ , trace annihilates*

$$\mathbf{C} A_3(h) + V_{\text{amb}}^{(5)}(h).$$

*Hence trace descends to a well-defined linear functional*

$$\tau_h : \mathcal{R}_h^{(5)} \rightarrow \mathbf{C}.$$

*Proof.* By Proposition 12.26, trace vanishes on  $V_{\text{amb}}^{(5)}(h)$ . It also vanishes on  $\mathbf{C} A_3(h)$  because  $A_3(h)$  is diagonal-traceless.  $\square$

**Proposition 12.28** (Generic first residual nonvanishing in the explicit finite-order model). *There exists an open dense subset*

$$\mathcal{U}_{\text{res}} \subset \mathcal{U}_{\text{cub}}$$

*such that*

$$\bar{v}_1(h) \neq 0 \quad \text{in } \mathcal{R}_h^{(5)}$$

*for every  $h \in \mathcal{U}_{\text{res}}$ .*

*Hence the one-pair quotient odd germ is not projectively constant on  $\mathcal{U}_{\text{res}}$  at the explicit first residual level.*

*Proof.* By Corollary 12.27, trace defines a linear functional

$$\tau_h : \mathcal{R}_h^{(5)} \rightarrow \mathbf{C}.$$

By Lemma 12.18,

$$\mathrm{tr} J_5^{\mathrm{pair}}(h) \neq 0$$

on the cubic-generic region. Hence there is an open dense subset

$$\mathcal{U}_{\mathrm{res}} \subset \mathcal{U}_{\mathrm{cub}}$$

on which

$$\mathrm{tr} J_5^{\mathrm{pair}}(h) \neq 0.$$

For such  $h$ ,

$$\tau_h(\bar{v}_1(h)) = \mathrm{tr} J_5^{\mathrm{pair}}(h) \neq 0,$$

so  $\bar{v}_1(h) \neq 0$  in  $\mathcal{R}_h^{(5)}$ . □

**Remark 12.29** (Status of the explicit first residual model). The first residual step is now closed at the finite-order ambient level. Indeed, Proposition 12.24 identifies the explicit quotient  $\mathcal{R}_h^{(5)}$  canonically with the order- $\leq 5$  truncation of the ambient abstract quotient  $\mathcal{Q}^{\mathrm{abs}}$  modulo the cubic line on the cubic-generic region.

Thus the remaining issue is no longer a finite-order comparison theorem for the first residual layer. What remains, if one wants a stronger global ambient identification, is a separate theorem showing that later uses of  $\mathcal{Q}^{\mathrm{abs}}$  beyond order 5 do not introduce any new ambiguity relevant to the downstream multi-pair and global arguments.

**Remark 12.30** (Beyond the first residual trace). The quintic trace is the first scalar residual detector, but it is not the end of the one-pair local structure. The later fixed-sector refinement  $A_5^{\mathfrak{f}}, A_7^{\mathfrak{f}}, \Delta_7$  indicates that the residual germ carries further geometric information beyond the first trace channel.

**12.4. Fixed-sector refinement at quintic and septic order.** The trace witness at quintic order suggests a more structured refinement of the one-pair residual germ than the trace alone records. For bookkeeping purposes, it is convenient to decompose the quintic and septic coefficients into fixed and anti-fixed parts with respect to the involution

$$\iota(A) := J_0 A J_0, \quad J_0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Write

$$\pi_{\mathfrak{f}}(A) := \frac{1}{2}(A + J_0 A J_0), \quad \pi_{\mathfrak{a}}(A) := \frac{1}{2}(A - J_0 A J_0).$$

Then

$$\mathfrak{f} = \mathrm{span}\{I, S\}, \quad \mathfrak{a} = \mathrm{span}\{D, J\},$$

where

$$I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad S = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad D = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

**Definition 12.31** (Fixed-sector quintic and septic coefficients). Let

$$F_h(z) = A_3(h)z^3 + A_5(h)z^5 + A_7(h)z^7 + \cdots$$

denote the normalized one-pair quotient odd germ on the cubic-generic region. Its fixed-sector quintic and septic coefficients are defined by

$$A_5^{\mathfrak{f}}(h) := \pi_{\mathfrak{f}}(A_5(h)), \quad A_7^{\mathfrak{f}}(h) := \pi_{\mathfrak{f}}(A_7(h)).$$

Write

$$A_5^{\mathfrak{f}}(h) = u_5(h)I + v_5(h)S, \quad A_7^{\mathfrak{f}}(h) = u_7(h)I + v_7(h)S.$$

**Definition 12.32** (Second residual determinant). On the cubic-generic region, define

$$\Delta_7(h) := u_7(h)v_5(h) - u_5(h)v_7(h).$$

**Remark 12.33** (Interpretation). The quintic fixed-sector vector  $A_5^{\mathfrak{f}}(h)$  refines the first residual data beyond the trace detector alone, and the determinant  $\Delta_7(h)$  records the next residual quotient at septic order. In the working notes, these form the first two nontrivial layers of a one-pair residual package:

$$c(h), \quad A_5^{\mathfrak{f}}(h), \quad \Delta_7(h).$$

The current draft now records explicit coefficient formulas for the quintic commutator contribution and the septic  $K_1$ -transport contribution in the existing notation  $(u, v, w, \alpha, \beta, \gamma, r, \dots)$ . The canonical order-7 fixed-sector content is the quotient class

$$[A_7^{\mathfrak{f}}] \in \mathfrak{f}/\mathbf{C}A_5^{\mathfrak{f}},$$

equivalently the second residual determinant

$$\Delta_7 = u_7v_5 - u_5v_7.$$

What remains non-unique is only the raw representative pair  $(u_7, v_7)$ , which is defined only up to the one-dimensional septic gauge ambiguity along the quintic fixed-sector line  $\mathbf{C}A_5^{\mathfrak{f}}$ . Thus the later geometry should be formulated in terms of the canonical quotient-level order-7 data  $[A_7^{\mathfrak{f}}]$  and  $\Delta_7$ , rather than a uniquely chosen raw pair  $(u_7, v_7)$ . At the canonical level, these are the natural variables for the geometric reformulation of the multi-pair problem; raw representatives  $(u_7, v_7)$  may be chosen locally by gauge-fixing when convenient, but are not themselves canonical.

**Proposition 12.34** (Projected septic gauge law). *Assume the normalization gauge is symmetry-preserving in the sense that*

$$B_2, B_4 \in \mathfrak{f}, \quad A_3 \in \mathfrak{a},$$

with

$$\mathfrak{f} = \text{span}\{I, S\}, \quad \mathfrak{a} = \text{span}\{D, J\}.$$

Then the fixed-sector septic coefficient satisfies

$$(A_7^{\mathfrak{f}})^{\text{new}} = A_7^{\mathfrak{f}} + c_2A_5^{\mathfrak{f}}.$$

Equivalently, the septic fixed-sector ambiguity is one-dimensional and lies along the quintic fixed-sector line.

*Proof.* By Proposition 12.15,

$$A_7^{\text{new}} = A_7 + c_2A_5 + c_4A_3 + [B_2, A_5] + [B_4, A_3] + c_2[B_2, A_3] + A_3B_2^2 - B_2A_3B_2.$$

Apply the fixed-sector projection  $\pi_{\mathfrak{f}}$  term by term.

Since  $A_3 \in \mathfrak{a}$ , one has

$$\pi_{\mathfrak{f}}(c_4A_3) = 0.$$

Because  $B_4 \in \mathfrak{f}$  and  $A_3 \in \mathfrak{a}$ ,

$$[B_4, A_3] \in [\mathfrak{f}, \mathfrak{a}] \subset \mathfrak{a},$$

so

$$\pi_{\mathfrak{f}}[B_4, A_3] = 0.$$

Likewise,

$$[B_2, A_3] \in [\mathfrak{f}, \mathfrak{a}] \subset \mathfrak{a},$$

hence

$$\pi_{\mathfrak{f}}(c_2[B_2, A_3]) = 0.$$

Now decompose

$$A_5 = A_5^{\mathfrak{f}} + A_5^{\mathfrak{a}}.$$

Since  $\mathfrak{f} = \text{span}\{I, S\}$  is abelian,

$$[B_2, A_5^{\mathfrak{f}}] = 0.$$

And since  $B_2 \in \mathfrak{f}$  and  $A_5^{\mathfrak{a}} \in \mathfrak{a}$ ,

$$[B_2, A_5^{\mathfrak{a}}] \in [\mathfrak{f}, \mathfrak{a}] \subset \mathfrak{a},$$

so

$$\pi_{\mathfrak{f}}[B_2, A_5] = 0.$$

Finally,  $B_2 \in \mathfrak{f}$  implies  $B_2^2 \in \mathfrak{f}$ . Therefore

$$A_3 B_2^2 \in \mathfrak{af} \subset \mathfrak{a}, \quad B_2 A_3 B_2 \in \mathfrak{faf} \subset \mathfrak{a},$$

and hence

$$\pi_{\mathfrak{f}}(A_3 B_2^2 - B_2 A_3 B_2) = 0.$$

Thus the only term surviving fixed projection is

$$c_2 A_5^{\mathfrak{f}},$$

and we conclude

$$(A_7^{\mathfrak{f}})^{\text{new}} = A_7^{\mathfrak{f}} + c_2 A_5^{\mathfrak{f}}. \quad \square$$

**Corollary 12.35** (Gauge invariance of the second residual determinant). *Under a symmetry-preserving normalization gauge, the scalar*

$$\Delta_7 = u_7 v_5 - u_5 v_7$$

*is invariant.*

*Proof.* By Proposition 12.34,

$$(A_7^{\mathfrak{f}})^{\text{new}} = A_7^{\mathfrak{f}} + c_2 A_5^{\mathfrak{f}}.$$

Writing

$$A_5^{\mathfrak{f}} = u_5 I + v_5 S, \quad A_7^{\mathfrak{f}} = u_7 I + v_7 S,$$

the new septic pair is

$$(u_7^{\text{new}}, v_7^{\text{new}}) = (u_7, v_7) + c_2(u_5, v_5).$$

Therefore

$$\Delta_7^{\text{new}} = (u_7 + c_2 u_5) v_5 - u_5 (v_7 + c_2 v_5) = u_7 v_5 - u_5 v_7 = \Delta_7. \quad \square$$

**Definition 12.36** (Septic normalization-shadow sectors). In the notation of Proposition 12.15, define the order-2 and order-4 septic normalization-shadow sectors by

$$A_{7,K_5} := c_2 A_5 + [B_2, A_5],$$

and

$$A_{7,K_3} := c_4 A_3 + [B_4, A_3] + c_2 [B_2, A_3] + A_3 B_2^2 - B_2 A_3 B_2.$$

Then the exact septic transformation law may be written

$$A_7^{\text{new}} = A_7 + A_{7,K_5} + A_{7,K_3}.$$

When the normalization gauge is symmetry-preserving, these are the septic  $K_5$ - and  $K_3$ -transport shadow sectors.

**Proposition 12.37** (Projected septic  $K_5$ -shadow law). *Assume the normalization gauge is symmetry-preserving in the sense of Proposition 12.34. Then*

$$A_{7,K_5}^{\mathfrak{f}} = c_2 A_5^{\mathfrak{f}}.$$

*In particular,*

$$A_{7,K_5}^{\mathfrak{f}} \in \mathbf{C} A_5^{\mathfrak{f}}.$$

*Proof.* By Definition 12.36,

$$A_{7,K_5} = c_2 A_5 + [B_2, A_5].$$

Apply the fixed-sector projection  $\pi_{\mathfrak{f}}$ . Decompose

$$A_5 = A_5^{\mathfrak{f}} + A_5^{\mathfrak{a}}.$$

Since  $\mathfrak{f} = \text{span}\{I, S\}$  is abelian,

$$[B_2, A_5^{\mathfrak{f}}] = 0.$$

And since  $B_2 \in \mathfrak{f}$  and  $A_5^{\mathfrak{a}} \in \mathfrak{a}$ ,

$$[B_2, A_5^{\mathfrak{a}}] \in [\mathfrak{f}, \mathfrak{a}] \subset \mathfrak{a},$$

so its fixed-sector projection vanishes. Therefore

$$A_{7,K_5}^{\mathfrak{f}} = \pi_{\mathfrak{f}}(c_2 A_5 + [B_2, A_5]) = c_2 A_5^{\mathfrak{f}}. \quad \square$$

**Proposition 12.38** (Projected septic  $K_3$ -shadow vanishing). *Assume the normalization gauge is symmetry-preserving in the sense of Proposition 12.34. Then*

$$A_{7,K_3}^{\mathfrak{f}} = 0.$$

*Proof.* By Definition 12.36,

$$A_{7,K_3} = c_4 A_3 + [B_4, A_3] + c_2 [B_2, A_3] + A_3 B_2^2 - B_2 A_3 B_2.$$

Since  $A_3 \in \mathfrak{a}$ , one has

$$c_4 A_3 \in \mathfrak{a}.$$

Because  $B_4 \in \mathfrak{f}$  and  $A_3 \in \mathfrak{a}$ ,

$$[B_4, A_3] \in [\mathfrak{f}, \mathfrak{a}] \subset \mathfrak{a}.$$

Likewise,

$$[B_2, A_3] \in [\mathfrak{f}, \mathfrak{a}] \subset \mathfrak{a},$$

hence

$$c_2 [B_2, A_3] \in \mathfrak{a}.$$

Finally,  $B_2 \in \mathfrak{f}$  implies  $B_2^2 \in \mathfrak{f}$ , so

$$A_3 B_2^2 \in \mathfrak{a} \mathfrak{f} \subset \mathfrak{a}, \quad B_2 A_3 B_2 \in \mathfrak{f} \mathfrak{a} \mathfrak{f} \subset \mathfrak{a}.$$

Thus every term of  $A_{7,K_3}$  is anti-fixed, and therefore

$$A_{7,K_3}^{\mathfrak{f}} = \pi_{\mathfrak{f}}(A_{7,K_3}) = 0. \quad \square$$

**Remark 12.39** (Interpretation of the septic shadow sectors). Propositions 12.37 and 12.38 show that the lower-order normalization shadows at septic order contribute no new canonical fixed-sector information beyond the already-known quintic line  $\mathbf{C} A_5^{\mathfrak{f}}$ .

**Proposition 12.40** (Explicit fixed-sector formulas at quintic order). *Write*

$$X_1 = \begin{pmatrix} u & v \\ v & w \end{pmatrix}, \quad X_3 = \begin{pmatrix} \alpha & \beta \\ \beta & \gamma \end{pmatrix}.$$

*Then the fixed-sector part of the basic quintic commutator channel*

$$X_3 J X_1 - X_1 J X_3$$

*is*

$$\pi_{\mathfrak{f}}(X_3 J X_1 - X_1 J X_3) = (v(\gamma - \alpha) + \beta(u - w))I + (\gamma u - \alpha w)S.$$

*Consequently, for the quintic fixed-sector coefficient*

$$A_5^{\mathfrak{f}}(h) = u_5(h)I + v_5(h)S,$$

*the commutator-type contribution is explicitly*

$$u_5^{(\text{comm})} = v(\gamma - \alpha) + \beta(u - w), \quad v_5^{(\text{comm})} = \gamma u - \alpha w.$$

*Proof.* A direct computation gives

$$X_3 J X_1 - X_1 J X_3 = \begin{pmatrix} -2\alpha v + 2\beta u & -\alpha w + \gamma u \\ -\alpha w + \gamma u & -2\beta w + 2\gamma v \end{pmatrix}.$$

The fixed-sector projection of a symmetric matrix

$$\begin{pmatrix} A & B \\ B & C \end{pmatrix}$$

is

$$\pi_{\mathfrak{f}} \begin{pmatrix} A & B \\ B & C \end{pmatrix} = \frac{A+C}{2} I + B S.$$

Applying this to the displayed matrix gives

$$\pi_{\mathfrak{f}}(X_3 J X_1 - X_1 J X_3) = \frac{(-2\alpha v + 2\beta u) + (-2\beta w + 2\gamma v)}{2} I + (\gamma u - \alpha w) S,$$

which simplifies to

$$(v(\gamma - \alpha) + \beta(u - w))I + (\gamma u - \alpha w)S.$$

□

**Proposition 12.41** (Explicit septic  $K_1$ -transport fixed-sector formulas). *Write*

$$\begin{aligned} X_0 &= \text{diag}(x_1, x_2), & X_1 &= \begin{pmatrix} u & v \\ v & w \end{pmatrix}, & X_2 &= \begin{pmatrix} a & b \\ b & c \end{pmatrix}, \\ X_4 &= \begin{pmatrix} p & r \\ r & s \end{pmatrix}, & X_5 &= \begin{pmatrix} \alpha_5 & \beta_5 \\ \beta_5 & \gamma_5 \end{pmatrix}, & X_6 &= \begin{pmatrix} \mu_6 & \nu_6 \\ \nu_6 & \lambda_6 \end{pmatrix}. \end{aligned}$$

Then the fixed-sector parts of the three septic  $K_1$ -transport channels are

$$\pi_{\mathfrak{f}}(X_5 J X_1 - X_1 J X_5) = (\beta_5(u - w) + v(\gamma_5 - \alpha_5))I + (\gamma_5 u - \alpha_5 w)S,$$

$$\pi_{\mathfrak{f}}(X_2 J X_4 - X_4 J X_2) = (b(p - s) + r(c - a))I + (cp - as)S,$$

and

$$\pi_{\mathfrak{f}}(X_0 J X_6 - X_6 J X_0) = (x_2 - x_1)\nu_6 I + (x_2\mu_6 - x_1\lambda_6)S.$$

Consequently, if the septic  $K_1$ -transport fixed-sector coefficient is written

$$A_{7,K_1}^{\mathfrak{f}}(h) = u_{7,K_1}(h)I + v_{7,K_1}(h)S,$$

then

$$u_{7,K_1} = \kappa \left( (x_2 - x_1)\nu_6 + \beta_5(u - w) + v(\gamma_5 - \alpha_5) + b(p - s) + r(c - a) \right),$$

$$v_{7,K_1} = \kappa \left( x_2\mu_6 - x_1\lambda_6 + \gamma_5 u - \alpha_5 w + cp - as \right).$$

*Proof.* For two symmetric matrices

$$X = \begin{pmatrix} a & b \\ b & c \end{pmatrix}, \quad Y = \begin{pmatrix} p & q \\ q & r \end{pmatrix},$$

a direct calculation gives

$$X J Y - Y J X = \begin{pmatrix} -2aq + 2bp & -ar + cp \\ -ar + cp & -2br + 2cq \end{pmatrix}.$$

Taking fixed-sector projection yields

$$\pi_{\mathfrak{f}}(X J Y - Y J X) = (b(p - r) + q(c - a))I + (cp - ar)S.$$

Applying this formula to  $(X, Y) = (X_5, X_1)$  and  $(X, Y) = (X_2, X_4)$  gives the first two displayed identities.

For the  $X_0/X_6$  channel, a direct multiplication gives

$$X_0 J X_6 - X_6 J X_0 = \begin{pmatrix} -2x_1\nu_6 & -x_1\lambda_6 + x_2\mu_6 \\ -x_1\lambda_6 + x_2\mu_6 & 2x_2\nu_6 \end{pmatrix},$$

whose fixed-sector part is

$$(x_2 - x_1)\nu_6 I + (x_2\mu_6 - x_1\lambda_6)S.$$

Summing the three channels gives the stated formulas for  $u_{7,K_1}$  and  $v_{7,K_1}$ . □

**Definition 12.42** (Septic  $K_1$ -residual determinant). If

$$A_{7,K_1}^{\mathfrak{f}}(h) = u_{7,K_1}(h)I + v_{7,K_1}(h)S,$$

define the corresponding septic  $K_1$ -residual determinant by

$$\Delta_{7,K_1}(h) := u_{7,K_1}(h)v_5(h) - u_5(h)v_{7,K_1}(h).$$



**Corollary 12.43** (Direct  $q^{(7)}$ -driven septic slice). *On the direct linear  $q^{(7)}$ -driven slice, the septic  $K_1$ -transport fixed-sector contribution is  $I$ -directional. More precisely,*

$$v_{7,K_1}^{(q^{(7)})} = 0,$$

and

$$u_{7,K_1}^{(q^{(7)})} = \kappa \left( (x_2 - x_1) \nu_6^{(q^{(7)})} + v(\gamma_5 - \alpha_5)^{(q^{(7)})} \right),$$

so in particular

$$\Delta_7^{(q^{(7)})} = u_7^{(q^{(7)})} v_5.$$

*Proof.* On the direct  $q^{(7)}$ -driven slice, the linear  $X_5$ -input is diagonal, hence

$$\beta_5^{(q^{(7)})} = 0.$$

The mixed  $X_6$ -slice is off-diagonal on this grading, so

$$\mu_6^{(q^{(7)})} = \lambda_6^{(q^{(7)})} = 0.$$

Moreover the  $X_2/X_4$ -channel does not lie on the pure  $(K_1, \eta_2, q^{(7)})$ -slice. Therefore the  $S$ -component vanishes and the displayed formula for  $u_{7,K_1}^{(q^{(7)})}$  follows from Proposition 12.41. Since

$$\Delta_7 = u_7 v_5 - u_5 v_7,$$

the direct  $q^{(7)}$ -slice satisfies

$$\Delta_7^{(q^{(7)})} = u_7^{(q^{(7)})} v_5. \quad \square$$

**Proposition 12.44** (Noncancellation of the direct septic  $K_1$  witness). *On the graded slice*

$$(K_1, \eta_2, q^{(7)}),$$

*the only contributing channels to*

$$u_{7,K_1}^{(q^{(7)})}$$

*are*

$$\kappa(x_2 - x_1) \nu_6^{(q^{(7)})} \quad \text{and} \quad \kappa v(\gamma_5 - \alpha_5)^{(q^{(7)})},$$

*and their total coefficient of the monomial*

$$\kappa(h) \eta_2(h) q^{(7)}(h)$$

*is nonzero. Consequently,*

$$u_{7,K_1}^{(q^{(7)})}(h) \neq 0$$

*on the generic one-pair region.*

*Proof.* Fix  $h$  and isolate the graded slice

$$\kappa(h) \eta_2(h) q^{(7)}(h).$$

By Corollary 12.43, the direct  $q^{(7)}$ -driven slice is  $I$ -directional and is given by

$$u_{7,K_1}^{(q^{(7)})} = \kappa \left( (x_2 - x_1) \nu_6^{(q^{(7)})} + v(\gamma_5 - \alpha_5)^{(q^{(7)})} \right).$$

Thus it remains only to show that the two displayed channels do not cancel.

Now consider the same-point expansion. By the explicit formula

$$G_{m,\pm}^{\text{corr}}(s) = \frac{1}{\pi} \begin{pmatrix} 2q(t_{\pm}) & \frac{1}{2}q'(t_{\pm}) \\ \frac{1}{2}q'(t_{\pm}) & \frac{1}{12}(q''(t_{\pm}) + 2q(t_{\pm})^3) \end{pmatrix},$$

the factor  $q^{(7)}(h)$  first appears only in the quintic same-point coefficient  $G_5$ , and there only in the diagonal sector, namely through the quintic Taylor term of the  $(2, 2)$ -entry coming from  $q''(t_{\pm})$ . Therefore, by Lemma 12.17, the linear  $q^{(7)}$ -contribution to  $X_5$  is diagonal. In particular it contributes to  $\gamma_5 - \alpha_5$ , but not to  $\beta_5$ .

Likewise, the  $\eta_2(h)$ -slice of  $X_1(h)$  is off-diagonal, because

$$v(h) = -\frac{1}{4\pi(s_1 + s_2)} \eta_2(h).$$

Applying Lemma 12.17 again, the mixed  $(\eta_2, q^{(7)})$ -slice of  $X_6$  is off-diagonal. Hence it contributes to  $\nu_6$ , but not to  $\mu_6$  or  $\lambda_6$ .

Thus the only contributing channels on the graded slice

$$(K_1, \eta_2, q^{(7)})$$

are exactly

$$\kappa(x_2 - x_1)\nu_6^{(q^{(7)})} \quad \text{and} \quad \kappa v(\gamma_5 - \alpha_5)^{(q^{(7)})}.$$

To compute their combined coefficient, it suffices by multilinearity to restrict to the two perturbation directions carrying the relevant tags: an off-diagonal first-order perturbation  $A$  for the  $\eta_2$ -slice and a diagonal fifth-order perturbation  $D$  for the  $q^{(7)}$ -slice. Accordingly, write

$$G_0 = \text{diag}(a^2, b^2), \quad A = \begin{pmatrix} 0 & u \\ u & 0 \end{pmatrix}, \quad D = \begin{pmatrix} 0 & 0 \\ 0 & d \end{pmatrix},$$

and expand

$$X(\varepsilon, \delta) := (G_0 + \varepsilon A + \delta D)^{-1/2} = X_0 + \varepsilon U + \delta V + \varepsilon \delta W + O(\varepsilon^2, \delta^2).$$

Solving the identity

$$X(\varepsilon, \delta) (G_0 + \varepsilon A + \delta D) X(\varepsilon, \delta) = I$$

order by order gives

$$U_{12} = -\frac{u}{ab(a+b)}, \quad V_{22} = -\frac{d}{2b^3}, \quad W_{12} = \frac{du(a+2b)}{2ab^3(a+b)^2}.$$

Here  $U_{12}$  is the  $\eta_2$ -slice of  $v$ ,  $V_{22}$  is the  $q^{(7)}$ -slice of  $\gamma_5 - \alpha_5$ , and  $W_{12}$  is the mixed  $(\eta_2, q^{(7)})$ -slice of  $\nu_6$ . Therefore the total coefficient of the target slice inside

$$(x_2 - x_1)\nu_6 + v(\gamma_5 - \alpha_5)$$

is

$$\left(\frac{1}{a} - \frac{1}{b}\right) W_{12} + U_{12} V_{22} = \frac{du}{a^2 b^2 (a+b)^2},$$

which is nonzero.

Thus the monomial

$$\kappa(h) \eta_2(h) q^{(7)}(h)$$

occurs in  $u_{7, K_1}^{(q^{(7)})}(h)$  with nonzero total coefficient, so

$$u_{7, K_1}^{(q^{(7)})}(h) \neq 0.$$

□

**Remark 12.45** (Septic  $K_1$  trace target). At the graded level, the septic  $K_1$ -transport channel is the first odd transport sector capable of coupling the new same-point  $q^{(7)}$ -slice into the fixed-sector septic residual. In particular, the expected highest new contribution to the septic determinant is carried by the  $K_1$ -channel, in parallel with the earlier quintic trace witness at order 5.

**Lemma 12.46** (Parity elimination of the pure same-point septic sector). *Let  $t_{\pm} = h \pm s/2$ , and let the odd two-point germ be defined as the odd part under the involution*

$$s \longmapsto -s \quad (\text{equivalently } t_+ \leftrightarrow t_-).$$

*Let  $A_{7, \text{pure}}$  denote the pure same-point septic sector, i.e. the septic contribution built only from the same-point blocks and their whitening, with no transport factor coming from the ordered pair  $(t_+, t_-)$ .*

Then  $A_{7,\text{pure}}$  is even in  $s$ , hence vanishes in the odd two-point germ. In particular,

$$A_{7,\text{pure}} = 0 \quad \text{in the odd germ}, \quad A_{7,\text{pure}}^f = 0.$$

Consequently, pure same-point septic terms do not contribute to  $\Delta_7$ , and in particular do not contribute to the weight-one part  $[\Delta_7]^{[1]}$ .

*Proof.* A pure same-point septic term depends only on midpoint jet data at  $h$ , i.e. on the same-point expansion at the midpoint, and carries no transport factor coming from the ordered pair  $(t_+, t_-)$ . Therefore it is invariant under exchanging the two points:

$$t_+ \leftrightarrow t_-.$$

Since this exchange is exactly the involution  $s \mapsto -s$ , every pure same-point septic term is an even function of  $s$ .

By definition, the odd two-point germ is obtained by projecting to the odd part under  $s \mapsto -s$ . Hence every even contribution vanishes in that projection. Therefore

$$A_{7,\text{pure}} = 0 \quad \text{in the odd germ.}$$

Applying fixed-sector projection gives

$$A_{7,\text{pure}}^f = 0.$$

Since  $\Delta_7$  is extracted from the odd pair germ after fixed-sector projection and quotient normalization, pure same-point septic terms contribute nothing to  $\Delta_7$ , and therefore nothing to  $[\Delta_7]^{[1]}$ .  $\square$

**Corollary 12.47** (Weight-one septic reduction to the direct  $K_1$  slice). *Let  $[\cdot]^{[1]}$  denote the part linear in the first genuinely new same-point  $q^{(7)}$ -slice. Then*

$$[\Delta_7]^{[1]} = [\Delta_{7,K_1}]^{[1]}.$$

*Equivalently, the highest genuinely new  $q^{(7)}$ -coefficient of the septic residual determinant comes only from the direct septic  $K_1$ -transport slice.*

*Proof.* The first genuinely new  $q^{(7)}$ -slice enters the same-point expansion at order 5, so

$$X_j^{[1]} = 0 \quad (j < 5).$$

Hence every septic transport sector built only from lower same-point orders is weight zero; in particular, the  $K_3$ - and  $K_5$ -transport sectors do not contribute to  $[\Delta_7]^{[1]}$ .

By Lemma 12.46, pure same-point septic terms also do not contribute to  $[\Delta_7]^{[1]}$ , because they vanish already in the odd two-point germ.

Thus the only septic sector that can see the weight-one  $q^{(7)}$ -slice is the direct  $K_1$ -transport sector, and therefore

$$[\Delta_7]^{[1]} = [\Delta_{7,K_1}]^{[1]}. \quad \square$$

**Proposition 12.48** (Generic nonvanishing of the direct septic slice under nontrivial quintic fixed-sector coefficient). *Let  $J$  be a connected open interval contained in the quintic-generic region. Assume*

$$v_5 \not\equiv 0 \quad \text{on } J.$$

*Then the direct  $q^{(7)}$ -driven slice of the full total septic residual determinant is generically nonzero on  $J$ . Equivalently,*

$$\Delta_7^{(q^{(7)})} \not\equiv 0 \quad \text{on } J.$$

*Proof.* By Corollary 12.47,

$$[\Delta_7]^{[1]} = [\Delta_{7,K_1}]^{[1]}.$$

Thus the highest genuinely new direct  $q^{(7)}$ -coefficient of the total septic residual determinant is the same as that of the direct  $K_1$ -transport slice.

By Proposition 12.44, the direct septic  $K_1$  witness is generically nonzero on the quintic-generic region. In particular,

$$u_{7,K_1}^{(q^{(7)})} \not\equiv 0 \quad \text{on } J.$$

Moreover, by Corollary 12.43, on the direct  $q^{(7)}$ -driven slice one has

$$v_{7,K_1}^{(q^{(7)})} = 0, \quad \Delta_{7,K_1}^{(q^{(7)})} = u_{7,K_1}^{(q^{(7)})} v_5.$$

Since  $v_5 \not\equiv 0$  on  $J$ , it follows that

$$\Delta_{7,K_1}^{(q^{(7)})} \not\equiv 0 \quad \text{on } J.$$

Therefore

$$\Delta_7^{(q^{(7)})} \not\equiv 0 \quad \text{on } J,$$

as claimed.  $\square$

**Theorem 12.49** (Quotient-septic closure). *Let the septic odd-germ coefficient be decomposed as*

$$A_7 = A_{7,K_1} + A_{7,K_5} + A_{7,K_3} + A_{7,\text{pure}},$$

where  $A_{7,K_1}$  is the direct  $K_1$ -transport sector,  $A_{7,K_5}$  and  $A_{7,K_3}$  are the septic normalization-shadow sectors from Definition 12.36, and  $A_{7,\text{pure}}$  is the pure same-point septic sector from Lemma 12.46.

Then

$$A_7^{\mathfrak{f}} = A_{7,K_1}^{\mathfrak{f}} + A_{7,K_5}^{\mathfrak{f}},$$

with

$$A_{7,K_5}^{\mathfrak{f}} \in \mathbf{C} A_5^{\mathfrak{f}}.$$

Consequently,

$$[A_7^{\mathfrak{f}}] = [A_{7,K_1}^{\mathfrak{f}}] \quad \text{in } \mathfrak{f}/\mathbf{C}A_5^{\mathfrak{f}},$$

and

$$\Delta_7 = \Delta_{7,K_1}.$$

*Proof.* By Proposition 12.38,

$$A_{7,K_3}^{\mathfrak{f}} = 0.$$

By Lemma 12.46,

$$A_{7,\text{pure}}^{\mathfrak{f}} = 0.$$

Therefore

$$A_7^{\mathfrak{f}} = A_{7,K_1}^{\mathfrak{f}} + A_{7,K_5}^{\mathfrak{f}}.$$

By Proposition 12.37,

$$A_{7,K_5}^{\mathfrak{f}} \in \mathbf{C} A_5^{\mathfrak{f}}.$$

Hence  $A_7^{\mathfrak{f}}$  and  $A_{7,K_1}^{\mathfrak{f}}$  define the same class in the quotient

$$\mathfrak{f}/\mathbf{C}A_5^{\mathfrak{f}}.$$

Writing

$$A_5^{\mathfrak{f}} = u_5 I + v_5 S, \quad A_{7,K_1}^{\mathfrak{f}} = u_{7,K_1} I + v_{7,K_1} S,$$

this quotient equality means precisely that

$$A_7^{\mathfrak{f}} = A_{7,K_1}^{\mathfrak{f}} + \chi A_5^{\mathfrak{f}}$$

for some scalar  $\chi$ . Therefore

$$\Delta_7 = (u_{7,K_1} + \chi u_5) v_5 - u_5 (v_{7,K_1} + \chi v_5) = u_{7,K_1} v_5 - u_5 v_{7,K_1} = \Delta_{7,K_1},$$

where  $\Delta_{7,K_1}$  is as in Definition 12.42.  $\square$

**Remark 12.50** (Status of the order-7 fixed-sector package). The order-7 fixed-sector analysis is now closed at the canonical quotient level. By Theorem 12.49, the full total septic fixed-sector class satisfies

$$[A_7^f] = [A_{7,K_1}^f] \quad \text{in } \mathfrak{f}/\mathbf{C}A_5^f,$$

and the canonical second residual determinant is

$$\Delta_7 = \Delta_{7,K_1}.$$

What remains non-unique is only the raw representative of  $A_7^f$  inside the affine line

$$A_{7,K_1}^f + \mathbf{C}A_5^f,$$

as recorded by Proposition 12.34. Equivalently, the full lower/mixed septic completion of the raw total pair  $(u_7, v_7)$  is not canonically determined. This remaining gauge-dependent choice does not affect the quotient class  $[A_7^f]$  or the residual determinant  $\Delta_7$  used in the later geometric arguments.

**Remark 12.51** (Local gauge-fixing for raw septic representatives). Although the raw representative pair  $(u_7, v_7)$  is not canonical, local choices may be made by fixing a gauge section along the quintic fixed-sector line. Indeed, Proposition 12.34 gives

$$(u_7^{\text{new}}, v_7^{\text{new}}) = (u_7, v_7) + c_2(u_5, v_5).$$

Hence on any region where  $v_5 \neq 0$ , one may choose

$$c_2 = -\frac{v_7}{v_5},$$

so that

$$v_7^{\text{new}} = 0, \quad u_7^{\text{new}} = \frac{\Delta_7}{v_5}.$$

Similarly, on any region where  $u_5 \neq 0$ , one may choose

$$c_2 = -\frac{u_7}{u_5},$$

so that

$$u_7^{\text{new}} = 0, \quad v_7^{\text{new}} = -\frac{\Delta_7}{u_5}.$$

Thus raw septic representatives can be chosen locally by gauge-fixing, but such choices are auxiliary and are not canonical.

**12.5. Normalized planar curve and curvature bridge.** On the region where  $c(h) \neq 0$ , define the normalized scalars

$$\rho_1(h) := \frac{v_5(h)}{c(h)}, \quad \rho_2(h) := \frac{\Delta_7(h)}{c(h)},$$

and the associated planar invariant curve

$$\Gamma(h) := (\rho_1(h), \rho_2(h)).$$

**Proposition 12.52** (Non-affinity of the second normalized residual scalar). *Let  $J$  be a connected open interval contained in the quintic-generic region and in  $\{h : c(h) \neq 0\}$ . Assume*

$$v_5 \not\equiv 0 \quad \text{on } J.$$

*Then there do not exist constants  $A, B \in \mathbf{C}$  such that*

$$\rho_2(h) \equiv A\rho_1(h) + B$$

*on  $J$ .*

*Equivalently,*

$$\Delta_7(h) \not\equiv A v_5(h) + B c(h)$$

*on  $J$ .*

*Proof.* Suppose

$$\Delta_7(h) \equiv A v_5(h) + B c(h)$$

on  $J$ . Then the direct  $q^{(7)}$ -driven slice of the right-hand side vanishes, because  $v_5$  and  $c$  are governed by the cubic/quintic layers and carry no direct septic  $q^{(7)}$ -slice.

On the other hand, by Proposition 12.48, the direct  $q^{(7)}$ -driven slice of  $\Delta_7$  is generically nonzero on  $J$ . This is a contradiction. Dividing by  $c(h)$  yields

$$\rho_2(h) \neq A \rho_1(h) + B. \quad \square$$

**Definition 12.53** (Normalized derivative numerators). Define

$$M(h) := c(h)v_5'(h) - c'(h)v_5(h), \quad N(h) := c(h)\Delta_7'(h) - c'(h)\Delta_7(h).$$

Then

$$\rho_1'(h) = \frac{M(h)}{c(h)^2}, \quad \rho_2'(h) = \frac{N(h)}{c(h)^2}.$$

**Proposition 12.54** (Curvature bridge for the normalized planar curve). *Let*

$$K_\Gamma(h) := \rho_1'(h)\rho_2''(h) - \rho_2'(h)\rho_1''(h)$$

*denote the affine curvature numerator of the planar curve  $\Gamma(h)$ . Then*

$$K_\Gamma(h) = \frac{M(h)N'(h) - N(h)M'(h)}{c(h)^4}.$$

*In particular, if  $K_\Gamma \equiv 0$  on a connected open interval  $J$  on which*

$$c \neq 0, \quad M \neq 0, \quad v_5 \neq 0,$$

*then  $\rho_2$  is affine in  $\rho_1$  on  $J$ , contradicting Proposition 12.52. Hence on every such interval one has*

$$K_\Gamma \neq 0.$$

*Proof.* By Definition 12.53,

$$\rho_1' = \frac{M}{c^2}, \quad \rho_2' = \frac{N}{c^2}.$$

Differentiating gives

$$\rho_1'' = \frac{M'c - 2Mc'}{c^3}, \quad \rho_2'' = \frac{N'c - 2Nc'}{c^3}.$$

Therefore

$$K_\Gamma = \frac{M}{c^2} \frac{N'c - 2Nc'}{c^3} - \frac{N}{c^2} \frac{M'c - 2Mc'}{c^3} = \frac{MN' - NM'}{c^4}.$$

If  $K_\Gamma \equiv 0$  on a connected open set with  $M \neq 0$ , then

$$0 = MN' - NM' = M^2 \left( \frac{N}{M} \right)',$$

so  $N/M$  is constant there:

$$\frac{N}{M} = A.$$

Since

$$\frac{N}{M} = \frac{\rho_2'}{\rho_1'},$$

it follows that

$$\rho_2' = A \rho_1',$$

hence

$$\rho_2 = A \rho_1 + B$$

on that connected interval. If in addition

$$c \neq 0, \quad v_5 \neq 0,$$

there, this contradicts Proposition 12.52. The final claim follows.  $\square$

**Remark 12.55** (Geometric consequence). Proposition 12.54 is the local-to-global bridge from the order-7 fixed-sector analysis back to the secant/tangent program. On the open set where

$$c(h) \neq 0, \quad M(h) \neq 0, \quad M(h)N'(h) - N(h)M'(h) \neq 0,$$

the normalized planar curve  $\Gamma(h)$  decomposes into  $C^2$  branches in the affine chart  $(\rho_1, \rho_2)$ , each of which is strictly convex or strictly concave. Thus the remaining global obstruction is not same-branch secancy, but the possibility of a common four-secant line meeting two separated ECT branches twice each.

**Definition 12.56** (Projective Wronskian of the strengthened one-pair triple). Define

$$W(h) := \det \begin{pmatrix} c(h) & v_5(h) & \Delta_7(h) \\ c'(h) & v_5'(h) & \Delta_7'(h) \\ c''(h) & v_5''(h) & \Delta_7''(h) \end{pmatrix}.$$

**Proposition 12.57** (Projective-Wronskian bridge). *With  $M(h)$  and  $N(h)$  as in Definition 12.53, one has*

$$M(h)N'(h) - N(h)M'(h) = c(h)W(h).$$

Consequently,

$$K_\Gamma(h) = \frac{W(h)}{c(h)^3}$$

on the region where  $c(h) \neq 0$ .

*Proof.* Using

$$M = cv_5' - c'v_5, \quad N = c\Delta_7' - c'\Delta_7,$$

a direct expansion gives

$$MN' - NM' = (cv_5' - c'v_5)(c\Delta_7'' - c''\Delta_7) - (c\Delta_7' - c'\Delta_7)(cv_5'' - c''v_5).$$

Expanding and regrouping yields

$$MN' - NM' = c \det \begin{pmatrix} c & v_5 & \Delta_7 \\ c' & v_5' & \Delta_7' \\ c'' & v_5'' & \Delta_7'' \end{pmatrix} = cW.$$

The displayed formula for  $K_\Gamma$  now follows from Proposition 12.54. □

**Lemma 12.58** (Exact identities for  $W$  and  $W'$ ). *Let*

$$F(h) := (c(h), v_5(h), \Delta_7(h)), \quad M(h) := c(h)v_5'(h) - c'(h)v_5(h).$$

*Then*

$$W(h) = \det(F(h), F'(h), F''(h))$$

*admits the exact expansion*

$$W = M\Delta_7'' + (v_5c'' - cv_5'')\Delta_7' + (c'v_5'' - v_5'c'')\Delta_7,$$

*and*

$$W'(h) = \det(F(h), F'(h), F'''(h))$$

*admits the exact expansion*

$$W' = M\Delta_7''' + (v_5c''' - cv_5''')\Delta_7' + (c'v_5''' - v_5'c''')\Delta_7.$$

*Proof.* Expand the determinant  $\det(F, F', F'')$  along the third column. This gives

$$W = \det \begin{pmatrix} c & v_5 \\ c' & v_5' \end{pmatrix} \Delta_7'' - \det \begin{pmatrix} c & v_5 \\ c'' & v_5'' \end{pmatrix} \Delta_7' + \det \begin{pmatrix} c' & v_5' \\ c'' & v_5'' \end{pmatrix} \Delta_7,$$

which is exactly the first identity.

For  $W'$ , differentiate  $W = \det(F, F', F'')$ . The terms  $\det(F', F', F'')$  and  $\det(F, F'', F'')$  vanish by repeated columns, so

$$W' = \det(F, F', F''').$$

Expanding along the third column gives the second formula. □

**Proposition 12.59** (Exclusion of  $W$ -flat intervals on the  $c \neq 0$ ,  $M \neq 0$  region). *Let  $J$  be a connected open interval on which*

$$c(h) \neq 0, \quad M(h) \neq 0.$$

*Then*

$$W \not\equiv 0 \quad \text{on } J.$$

*Proof.* Suppose for contradiction that

$$W \equiv 0 \quad \text{on } J.$$

By Proposition 12.57,

$$K_\Gamma(h) = \frac{W(h)}{c(h)^3} \equiv 0 \quad \text{on } J.$$

Since

$$M = c v'_5 - c' v_5$$

and  $c \neq 0$  on  $J$ , the hypothesis  $M \neq 0$  on  $J$  rules out  $v_5 \equiv 0$  on  $J$ . Hence  $v_5 \not\equiv 0$  on  $J$ .

By Proposition 12.54, it follows that

$$\rho_2(h) = A \rho_1(h) + B$$

for some constants  $A, B \in \mathbf{C}$ , i.e.

$$\Delta_7(h) = A v_5(h) + B c(h) \quad \text{on } J.$$

This contradicts Proposition 12.52. Therefore

$$W \not\equiv 0 \quad \text{on } J. \quad \square$$

**Remark 12.60** (Interpretation of the  $\lambda$ -reduction). On the direct highest-new slice one has

$$\Delta_7^{[1]} = \lambda^{[1]} v_5$$

by Corollary 12.43 together with Corollary 12.47. Thus if  $W \equiv 0$  on a connected open interval with  $c \neq 0$  and  $M \neq 0$ , then

$$v_5 \not\equiv 0$$

on that interval, because  $M = c v'_5 - c' v_5$ . The affine dependence

$$\Delta_7 = A v_5 + B c$$

forced by Proposition 12.59 implies

$$\Delta_7^{[1]} = A v_5,$$

hence

$$\lambda^{[1]} \equiv A$$

on that interval. In this sense, the present argument excludes  $W$ -flatness on an  $M \neq 0$  region without requiring a separate differential equation for  $\lambda^{[1]}$ ; what remains open is not the canonical quotient-level order-7 invariant, but only the lower/mixed septic completion of a raw representative pair  $(u_7, v_7)$ , which is irrelevant to the quotient class  $[A_7^f]$  and to  $\Delta_7$ .

**Definition 12.61** (Projective graph and dual tangent-line variables). On the region where  $v_5 \neq 0$ , define

$$\mu := \frac{c}{v_5}, \quad \lambda := \frac{\Delta_7}{v_5}.$$

On the subregion where  $M \neq 0$ , define

$$L := v_5 \Delta'_7 - v'_5 \Delta_7, \quad \beta := \frac{d\lambda}{d\mu}, \quad m := \lambda - \mu \beta.$$



**Proposition 12.62** (Closed minor system and dual tangent-line curve). *Let*

$$F := (c, v_5, \Delta_7), \quad U := (L, -N, M).$$

*Then*

$$F \times F' = U, \quad U \times U' = W F.$$

*Equivalently,*

$$MN' - NM' = cW, \quad ML' - LM' = v_5W, \quad NL' - LN' = \Delta_7W.$$

*On the region where  $v_5M \neq 0$ ,*

$$\mu' = -\frac{M}{v_5^2}, \quad \beta = -\frac{L}{M}, \quad m = \frac{N}{M},$$

*and*

$$\beta' = -\frac{v_5W}{M^2}, \quad m' = \frac{cW}{M^2}.$$

*Hence the dual tangent-line curve*

$$T(h) := (\beta(h), m(h))$$

*satisfies*

$$T'(h) = \frac{W(h)}{M(h)^2} (-v_5(h), c(h)).$$

*Moreover,*

$$\det(T'(h), T''(h)) = \frac{W(h)^2}{M(h)^3}.$$

*Proof.* The identity

$$F \times F' = (L, -N, M)$$

is the direct expansion of the cross product. Differentiating gives

$$U' = F \times F''.$$

Applying the vector identity

$$(A \times B) \times (A \times C) = \det(A, B, C) A$$

with  $A = F$ ,  $B = F'$ ,  $C = F''$  yields

$$U \times U' = (F \times F') \times (F \times F'') = \det(F, F', F'') F = W F,$$

which is exactly the three displayed scalar identities.

Next,

$$\mu' = \left( \frac{c}{v_5} \right)' = \frac{c'v_5 - cv_5'}{v_5^2} = -\frac{M}{v_5^2},$$

and

$$\lambda' = \left( \frac{\Delta_7}{v_5} \right)' = \frac{v_5\Delta_7' - v_5'\Delta_7}{v_5^2} = \frac{L}{v_5^2},$$

so

$$\beta = \frac{\lambda'}{\mu'} = -\frac{L}{M}.$$

Also

$$\mu L + M\lambda = \frac{c}{v_5}(v_5\Delta_7' - v_5'\Delta_7) + (cv_5' - c'v_5)\frac{\Delta_7}{v_5} = c\Delta_7' - c'\Delta_7 = N,$$

hence

$$m = \lambda - \mu\beta = \lambda + \mu\frac{L}{M} = \frac{N}{M}.$$

The formulas for  $\beta'$  and  $m'$  follow immediately from

$$ML' - LM' = v_5W, \quad MN' - NM' = cW.$$

Finally, write

$$T' = \alpha V, \quad \alpha := \frac{W}{M^2}, \quad V := (-v_5, c).$$

Then

$$\det(T', T'') = \alpha^2 \det(V, V') = \frac{W^2}{M^4} \det \begin{pmatrix} -v_5 & -v'_5 \\ c & c' \end{pmatrix} = \frac{W^2}{M^3},$$

because

$$\det(V, V') = cv'_5 - c'v_5 = M.$$

□

**Corollary 12.63** (Graph curvature identities in  $(\mu, \lambda)$ -coordinates). *On the region where  $v_5 M \neq 0$ ,*

$$\lambda_{\mu\mu} = \frac{v_5^3}{M^3} W,$$

and

$$\lambda_{\mu\mu\mu} = -\frac{v_5^5}{M^4} \left( W' + 3 \left( \frac{v'_5}{v_5} - \frac{M'}{M} \right) W \right).$$

*In particular, at a point where  $W = 0$ ,  $M \neq 0$ , and  $v_5 \neq 0$ ,*

$$\lambda_{\mu\mu\mu} = -\frac{v_5^5}{M^4} W'.$$

*Proof.* Since

$$\frac{d}{d\mu} = \frac{1}{\mu'} \frac{d}{dh} = -\frac{v_5^2}{M} \frac{d}{dh},$$

the identity

$$\lambda_{\mu\mu} = \frac{v_5^3}{M^3} W$$

follows from

$$\lambda_\mu = \beta, \quad \beta' = -\frac{v_5 W}{M^2}.$$

Differentiating once more gives

$$\lambda_{\mu\mu\mu} = -\frac{v_5^2}{M} \frac{d}{dh} \left( \frac{v_5^3}{M^3} W \right),$$

which expands to the displayed formula. The last statement is the specialisation  $W = 0$ . □

**Remark 12.64** (Determinant interpretation on a fixed-sign  $W$ -interval). On any connected interval where  $c$ ,  $M$ , and  $W$  are nonzero and  $W$  keeps one sign, the graph  $\lambda = \lambda(\mu)$  is strictly convex or strictly concave. Consequently, for any ordered triple of distinct points on that interval, the determinant

$$D(s, t, u) = \det(F(s), F(t), F(u))$$

has the corresponding fixed Vandermonde sign. Thus the unresolved global issue is not interval-wise sign regularity, but propagation across turning seams and possible nonadjacent recurrence of tangent-slope / tangent-line data.

**Proposition 12.65** (Local determinant sign transport across a generic turning boundary). *Let  $h_b$  be a generic turning boundary:*

$$M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0.$$

*Then there exists a neighborhood  $U$  of  $h_b$  and a sign  $\sigma_{\text{turn}} \in \{\pm 1\}$  such that for any three distinct points  $h_1, h_2, h_3 \in U$ ,*

$$\text{sgn } D(h_1, h_2, h_3) = \sigma_{\text{turn}} \text{sgn}((h_2 - h_1)(h_3 - h_1)(h_3 - h_2)),$$

where

$$D(h_1, h_2, h_3) := \det(F(h_1), F(h_2), F(h_3)), \quad F = (c, v_5, \Delta_7).$$

*Equivalently, the triple  $(c, v_5, \Delta_7)$  is locally discretely sign-regular of order 3 across a generic turning boundary.*

*Proof.* Work in the normalized affine chart

$$x(h) := \frac{v_5(h)}{c(h)}, \quad y(h) := \frac{\Delta_7(h)}{c(h)}.$$

At a generic turning boundary one has

$$x'(h_b) = 0, \quad x''(h_b) \neq 0, \quad y'(h_b) \neq 0.$$

Hence  $y$  is a valid local coordinate near  $h_b$ , and the normalized curve may be written as

$$x = \chi(y)$$

with

$$\chi'(y_b) = 0, \quad \chi''(y_b) \neq 0.$$

After shrinking  $U$ , the sign of  $\chi''$  is constant on  $U$ .

Now for  $h_i \in U$ , write

$$x_i := x(h_i), \quad y_i := y(h_i).$$

Since

$$F(h_i) = c(h_i) (1, x_i, y_i),$$

we have

$$D(h_1, h_2, h_3) = c(h_1)c(h_2)c(h_3) \det \begin{pmatrix} 1 & x_1 & y_1 \\ 1 & x_2 & y_2 \\ 1 & x_3 & y_3 \end{pmatrix}.$$

Using  $x_i = \chi(y_i)$ , the  $3 \times 3$  determinant factors as

$$\det \begin{pmatrix} 1 & x_1 & y_1 \\ 1 & x_2 & y_2 \\ 1 & x_3 & y_3 \end{pmatrix} = (y_2 - y_1)(y_3 - y_1)(y_3 - y_2) \chi[y_1, y_2, y_3],$$

where  $\chi[y_1, y_2, y_3]$  is the second divided difference of  $\chi$ . By the scalar divided-difference mean-value theorem,

$$\chi[y_1, y_2, y_3] = \frac{1}{2} \chi''(\eta)$$

for some  $\eta$  between the  $y_i$ . Thus  $\chi[y_1, y_2, y_3]$  has fixed sign on  $U$ .

Also  $c$  has fixed sign on a sufficiently small neighborhood of  $h_b$ , and since  $y'(h_b) \neq 0$ , the map  $h \mapsto y(h)$  is monotone on  $U$ . Hence

$$\operatorname{sgn}((y_2 - y_1)(y_3 - y_1)(y_3 - y_2)) = \operatorname{sgn}(y')^3 \operatorname{sgn}((h_2 - h_1)(h_3 - h_1)(h_3 - h_2)),$$

with  $\operatorname{sgn}(y')$  constant on  $U$ .

Collecting the fixed signs of  $c$ ,  $y'$ , and  $\chi''$  gives the desired constant  $\sigma_{\text{turn}}$ .  $\square$

**Proposition 12.66** (Dual asymptote at a generic turning boundary). *Let  $h_b$  be a generic turning boundary:*

$$M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0.$$

*Assume  $v_5(h_b) \neq 0$ , and write*

$$\mu_b := \frac{c(h_b)}{v_5(h_b)}, \quad \lambda_b := \frac{\Delta_7(h_b)}{v_5(h_b)}.$$

*Let*

$$T(h) = (\beta(h), m(h)) = \left( -\frac{L(h)}{M(h)}, \frac{N(h)}{M(h)} \right)$$

*be the dual tangent-line curve on the neighboring  $M \neq 0$  branches. Then both one-sided dual arcs have the common asymptote*

$$m = -\mu_b \beta + \lambda_b,$$

*and, more precisely,*

$$m + \mu_b \beta - \lambda_b = \frac{1}{4a\beta} + o\left(\frac{1}{\beta}\right) \quad \text{as } |\beta| \rightarrow \infty,$$

where  $a \neq 0$  is the quadratic coefficient in the local turning-point normal form

$$\mu - \mu_b = a(\lambda - \lambda_b)^2 + O((\lambda - \lambda_b)^3).$$

In particular, the two adjacent dual arcs approach the same asymptote from opposite sides. This asymptotic separation is the dual-curve explanation of the local adjacent turning-boundary exclusion theorem proved later.

*Proof.* At a generic turning boundary one has  $\mu'(h_b) = 0$  and  $\lambda'(h_b) \neq 0$ , so  $\lambda$  is a valid local coordinate and

$$\mu - \mu_b = a(\lambda - \lambda_b)^2 + O((\lambda - \lambda_b)^3), \quad a \neq 0.$$

Differentiating gives

$$\frac{d\mu}{d\lambda} = 2a(\lambda - \lambda_b) + O((\lambda - \lambda_b)^2),$$

hence

$$\beta = \frac{d\lambda}{d\mu} = \frac{1}{2a(\lambda - \lambda_b)} + O(1).$$

Thus  $|\beta| \rightarrow \infty$  as  $h \rightarrow h_b$  from either side.

Now

$$m = \lambda - \mu\beta,$$

so

$$m + \mu_b\beta - \lambda_b = (\lambda - \lambda_b) - (\mu - \mu_b)\beta.$$

Using

$$\mu - \mu_b = a(\lambda - \lambda_b)^2 + O((\lambda - \lambda_b)^3)$$

and

$$\beta = \frac{1}{2a(\lambda - \lambda_b)} + O(1),$$

we obtain

$$(\mu - \mu_b)\beta = \frac{1}{2}(\lambda - \lambda_b) + O((\lambda - \lambda_b)^2).$$

Therefore

$$m + \mu_b\beta - \lambda_b = \frac{1}{2}(\lambda - \lambda_b) + O((\lambda - \lambda_b)^2).$$

Substituting

$$\lambda - \lambda_b = \frac{1}{2a\beta} + o\left(\frac{1}{\beta}\right)$$

gives

$$m + \mu_b\beta - \lambda_b = \frac{1}{4a\beta} + o\left(\frac{1}{\beta}\right).$$

Since  $\beta \rightarrow +\infty$  on one side and  $\beta \rightarrow -\infty$  on the other, the correction term changes sign, so the two one-sided dual arcs approach the same asymptote from opposite sides.  $\square$

**Proposition 12.67** (ECT intervals for the strengthened one-pair triple). *Let  $I$  be a connected interval on which*

$$c(h) \neq 0, \quad M(h) \neq 0, \quad W(h) \neq 0.$$

*Then the triple*

$$(c, v_5, \Delta_7)$$

*is an extended complete Chebyshev system on  $I$ . Equivalently, every nontrivial linear combination*

$$A c(h) + B v_5(h) + C \Delta_7(h)$$

*has at most two zeros on  $I$ , counting multiplicity.*

*In particular, in the affine chart*

$$x = \frac{v_5}{c}, \quad y = \frac{\Delta_7}{c},$$

*every affine line meets  $\Gamma(I)$  in at most two points, counting tangency with multiplicity.*

*Proof.* Define

$$x(h) := \frac{v_5(h)}{c(h)}, \quad y(h) := \frac{\Delta_7(h)}{c(h)}.$$

Since the one-pair family is real-analytic on the generic region, the functions  $x$  and  $y$  are  $C^2$  on  $I$ .

By Definition 12.53,

$$x'(h) = \frac{M(h)}{c(h)^2}, \quad y'(h) = \frac{N(h)}{c(h)^2}.$$

Because  $M(h) \neq 0$  and  $c(h) \neq 0$  on  $I$ , one has

$$x'(h) \neq 0 \quad (h \in I),$$

so  $x$  is strictly monotone on  $I$  and may be used as a parameter on  $\Gamma(I)$ .

Moreover,

$$\frac{dy}{dx} = \frac{y'(h)}{x'(h)} = \frac{N(h)}{M(h)}.$$

Differentiating with respect to  $x$  gives

$$\frac{d^2y}{dx^2} = \frac{\frac{d}{dh}(N/M)}{x'(h)} = \frac{(MN' - NM')/M^2}{M/c^2}.$$

Using Proposition 12.57,

$$MN' - NM' = cW,$$

hence

$$\frac{d^2y}{dx^2} = \frac{c(h)^3 W(h)}{M(h)^3}.$$

Since  $c(h) \neq 0$ ,  $M(h) \neq 0$ , and  $W(h) \neq 0$  on  $I$ , it follows that

$$\frac{d^2y}{dx^2} \neq 0 \quad (h \in I).$$

Therefore  $\Gamma(I)$  is the graph of a strictly convex or strictly concave  $C^2$  function  $y = y(x)$ .

Now let

$$L(h) := A c(h) + B v_5(h) + C \Delta_7(h)$$

be a nontrivial linear combination. Since  $c(h) \neq 0$  on  $I$ ,

$$L(h) = 0 \iff A + B x(h) + C y(h) = 0.$$

Thus zeros of  $L$  correspond exactly to intersections of  $\Gamma(I)$  with the affine line

$$A + B x + C y = 0.$$

If  $C = 0$ , then the equation is

$$A + B x = 0,$$

which has at most one solution because  $x$  is strictly monotone on  $I$ . If  $C \neq 0$ , the equation may be written as

$$y = -\frac{A}{C} - \frac{B}{C}x,$$

and a strictly convex or strictly concave graph meets an affine line in at most two points, counting tangency with multiplicity.

Hence every nontrivial linear combination

$$A c + B v_5 + C \Delta_7$$

has at most two zeros on  $I$ , counting multiplicity. This is exactly the ECT property for the triple  $(c, v_5, \Delta_7)$ . The affine-line statement is the same conclusion in the chart  $(x, y)$ .  $\square$

**Definition 12.68** (ECT interval). An ECT interval is a connected interval

$$I \subset \{h : c(h) \neq 0, M(h) \neq 0, W(h) \neq 0\}.$$

Equivalently, by Proposition 12.67, it is a connected interval on which the triple

$$(c, v_5, \Delta_7)$$

is an extended complete Chebyshev system.

**Corollary 12.69** (At most two points from one ECT interval). *Let  $I$  be an ECT interval. Then every affine line in the chart*

$$x = \frac{v_5}{c}, \quad y = \frac{\Delta_7}{c}$$

*meets  $\Gamma(I)$  in at most two points, counting tangency with multiplicity.*

*In particular, if a finite configuration is cut out by a single affine relation in the normalized chart, then its contribution from any fixed ECT interval has cardinality at most two.*

*Proof.* This is exactly the affine-line formulation of Proposition 12.67.  $\square$

**Remark 12.70** (Branch geometry on an ECT interval). On an ECT interval  $I$ , the quantities  $c$ ,  $M$ , and  $W$  have fixed nonzero sign. Hence

$$x = \frac{v_5}{c}$$

is a monotone parameter on  $I$ , and in the affine chart

$$\Gamma(I) = \left\{ \left( \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)} \right) : h \in I \right\}$$

is a single  $C^2$  branch which is strictly convex or strictly concave according to the sign of

$$\frac{c(h)^3 W(h)}{M(h)^3}.$$

The later tangent-separator analysis is therefore an adjacent-branch problem, not a same-branch secancy problem.

**Definition 12.71** (Tangent-separator kernel). Let

$$F(h) := (c(h), v_5(h), \Delta_7(h)).$$

For  $s, t$  in the generic region, define

$$K(s, t) := \det(F(s), F'(s), F(t)).$$

For fixed  $s$ , the zeros of  $t \mapsto K(s, t)$  are exactly the points  $\Gamma(t)$  lying on the affine tangent line to  $\Gamma$  at  $\Gamma(s)$ . Equivalently,  $K(s, t) = 0$  is the tangent-separator incidence condition for the tangent at  $s$  and the point at  $t$ .

**Lemma 12.72** (Affine form of the tangent-separator kernel). *On the affine chart  $c \neq 0$ , write*

$$x(h) := \frac{v_5(h)}{c(h)}, \quad y(h) := \frac{\Delta_7(h)}{c(h)}.$$

*Then for every  $s, t$  with  $c(s)c(t) \neq 0$ ,*

$$K(s, t) = c(s)^2 c(t) x'(s) \left( (y(t) - m(s)x(t)) - (y(s) - m(s)x(s)) \right),$$

*where*

$$m(s) = \frac{dy}{dx}(s) = \frac{N(s)}{M(s)}.$$

*Equivalently,*

$$K(s, t) = c(s)^2 c(t) x'(s) (\phi_{m(s)}(t) - \tau_s),$$

*with*

$$\phi_{m(s)}(t) := y(t) - m(s)x(t), \quad \tau_s := y(s) - m(s)x(s).$$

*Proof.* Write

$$F(h) = (c(h), c(h)x(h), c(h)y(h)).$$

Then

$$F'(s) = (c'(s), c'(s)x(s) + c(s)x'(s), c'(s)y(s) + c(s)y'(s)).$$

Subtracting  $x(s)$  times the first column from the second and  $y(s)$  times the first column from the third gives

$$K(s, t) = \det \begin{pmatrix} c(s) & 0 & 0 \\ c'(s) & c(s)x'(s) & c(s)y'(s) \\ c(t) & c(t)(x(t) - x(s)) & c(t)(y(t) - y(s)) \end{pmatrix}.$$

Expanding along the first row yields

$$K(s, t) = c(s)^2 c(t) (x'(s)(y(t) - y(s)) - y'(s)(x(t) - x(s))).$$

Since

$$m(s) = \frac{y'(s)}{x'(s)},$$

this becomes

$$K(s, t) = c(s)^2 c(t) x'(s) ((y(t) - m(s)x(t)) - (y(s) - m(s)x(s))),$$

as claimed.  $\square$

**Lemma 12.73** (Basic properties of the tangent-separator kernel). *For every  $s$  in the generic region,*

$$K(s, s) = 0, \quad \partial_t K(s, s) = 0, \quad \partial_t^2 K(s, s) = W(s).$$

*In particular, on any interval where  $W$  has fixed sign, the tangent line at  $\Gamma(s)$  is locally supporting.*

*Proof.* The first two identities follow because, at  $t = s$ , the third column of the determinant equals  $F(s)$ , and after differentiating once in  $t$  it equals  $F'(s)$ . Differentiating a second time gives

$$\partial_t^2 K(s, s) = \det(F(s), F'(s), F''(s)) = W(s). \quad \square$$

**Lemma 12.74** (Compact first-zero reduction on adjacent intervals). *Let  $I_1 < I_2$  be adjacent ECT intervals in the affine  $x = v_5/c$  ordering. Assume there are compact subintervals*

$$J_1 \subseteq I_1, \quad J_2 \subseteq I_2$$

*such that every zero of the tangent-separator kernel*

$$K(s, t) = \det(F(s), F'(s), F(t)), \quad F(h) = (c(h), v_5(h), \Delta_7(h)),$$

*with  $s \in I_2$ ,  $t \in I_1$ , lies in  $J_2 \times J_1$ .*

*If  $K$  has a zero on  $J_2 \times J_1$ , then there exists a zero*

$$(s_*, t_*) \in J_2 \times J_1$$

*such that  $s_*$  is minimal in the  $x = v_5/c$  ordering among all  $s$ -coordinates of zeros of  $K$  on  $J_2 \times J_1$ .*

*Proof.* Because  $K$  is continuous, its zero set in the compact set  $J_2 \times J_1$  is closed, hence compact. The projection of that zero set to the  $s$ -axis is therefore compact as well. Since  $x = v_5/c$  is monotone on  $I_2$ , the induced  $x$ -ordering is equivalent to the interval ordering on  $J_2$ . Thus the projected zero set has a minimal element  $s_*$ , and there exists  $t_* \in J_1$  such that  $(s_*, t_*)$  is a zero of  $K$ .  $\square$

**Lemma 12.75** (First adjacent zero implies tangency). *In the setting of Lemma 12.74, let*

$$(s_*, t_*) \in J_2 \times J_1$$

*be a zero with minimal  $s$ -coordinate in the  $x = v_5/c$  ordering. Then*

$$\partial_t K(s_*, t_*) = 0.$$

Consequently,

$$K(s_*, t_*) = 0, \quad \partial_t K(s_*, t_*) = 0$$

force the tangent line at  $\Gamma(s_*)$  to be tangent also at  $\Gamma(t_*)$ .

*Proof.* Assume for contradiction that

$$\partial_t K(s_*, t_*) \neq 0.$$

Since

$$(s_*, t_*) \in J_2 \times J_1 \subseteq I_2 \times I_1,$$

the implicit function theorem applies in the open set  $I_2 \times I_1$  and gives a local  $C^1$  zero curve

$$t = t(s)$$

through  $(s_*, t_*)$ , defined for  $s$  in a neighborhood of  $s_*$ , such that

$$K(s, t(s)) = 0.$$

For  $s$  sufficiently close to  $s_*$  with  $s < s_*$ , one still has

$$(s, t(s)) \in I_2 \times I_1$$

and

$$K(s, t(s)) = 0.$$

By the zero-capture hypothesis in Lemma 12.74, every zero of  $K$  with  $s \in I_2$  and  $t \in I_1$  lies in  $J_2 \times J_1$ . Hence these nearby zeros also lie in  $J_2 \times J_1$ , contradicting the minimality of  $s_*$  among all zeros of  $K$  on  $J_2 \times J_1$ .

Therefore

$$\partial_t K(s_*, t_*) = 0.$$

Finally, the identities

$$K(s, t) = \det(F(s), F'(s), F(t)), \quad \partial_t K(s, t) = \det(F(s), F'(s), F'(t))$$

show that  $K(s_*, t_*) = 0$  places  $\Gamma(t_*)$  on the tangent line at  $\Gamma(s_*)$ , while  $\partial_t K(s_*, t_*) = 0$  places the tangent direction at  $t_*$  on that same line. Hence the line is tangent to  $\Gamma$  at both  $\Gamma(s_*)$  and  $\Gamma(t_*)$ ; that is, it is a common tangent line.  $\square$

**Proposition 12.76** (Adjacent slope-range separation). *Let  $I_1 < I_2$  be adjacent ECT intervals, meeting at a boundary point  $h_b$ .*

(1) *If*

$$M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0,$$

*then the interior tangent-slope ranges of  $I_1$  and  $I_2$  are disjoint.*

(2) *If*

$$W(h_b) = 0, \quad M(h_b) \neq 0,$$

*then  $x = v_5/c$  remains a valid local parameter, but no adjacent tangent separation statement is asserted here. In that case the local affine jet is governed by*

$$\left. \frac{d^2 y}{dx^2} \right|_{h_b} = 0, \quad \left. \frac{d^3 y}{dx^3} \right|_{h_b} = \frac{c(h_b)^5}{M(h_b)^4} W'(h_b),$$

*so the generic behavior is an ordinary affine inflection unless further special structure forces  $W'(h_b) = 0$ .*

*Proof.* In case (1),  $M(h_b) = 0$  is a simple zero and  $N(h_b) \neq 0$ . Therefore

$$m(h) = \frac{N(h)}{M(h)}$$

has opposite one-sided blow-up:

$$m(h) \rightarrow \pm\infty$$

as  $h \rightarrow h_b^\pm$ . Hence the interior tangent-slope ranges on the two sides are disjoint.

Case (2) is the local affine-jet computation recorded in Remark 12.84; it identifies the cubic-inflection coefficient but does not produce a tangent-separation theorem.  $\square$



**Lemma 12.77** (Endpoint tangent exclusion at a generic turning boundary). *Let  $I_1 < I_2$  be adjacent ECT intervals meeting at a boundary point  $h_b$ , and assume*

$$M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0.$$

*Then the boundary tangent line at  $\Gamma(h_b)$  is vertical and meets neither adjacent interval in its interior.*

*Proof.* The boundary tangent line is

$$x = \rho_1(h_b), \quad \rho_1(h) = \frac{v_5(h)}{c(h)}.$$

Since  $M(h_b) = 0$  is a simple zero,

$$\rho_1'(h_b) = 0, \quad \rho_1''(h_b) \neq 0,$$

so  $\rho_1$  has a strict local extremum at  $h_b$ . Therefore

$$\rho_1(h) - \rho_1(h_b)$$

has fixed sign on each adjacent interval and vanishes only at the boundary point. Hence the vertical tangent line meets neither adjacent interval in its interior.  $\square$

**Theorem 12.78** (Conditional adjacent tangent-separator sign regularity across turning boundaries). *Let  $I_1 < I_2$  be adjacent ECT intervals, and assume their common boundary point  $h_b$  is a generic turning boundary:*

$$M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0.$$

*Assume in addition the zero-capture hypothesis of Lemma 12.74 on the adjacent pair  $I_2 \times I_1$ . Then*

$$K(s, t) \neq 0 \quad (s \in I_2, t \in I_1),$$

*where*

$$K(s, t) = \det(F(s), F'(s), F(t)), \quad F(h) = (c(h), v_5(h), \Delta_7(h)).$$

*Equivalently, tangent lines from the right interval do not meet the adjacent left interval across a generic turning boundary.*

*Proof.* By Lemma 12.77, for a generic turning boundary there is a neighborhood of the boundary pair on which  $K(s, t) \neq 0$  for  $s \in I_2, t \in I_1$ .

Assume for contradiction that some zero  $(s_0, t_0) \in I_2 \times I_1$  exists. By the zero-capture hypothesis, there are compact subintervals

$$J_1 \Subset I_1, \quad J_2 \Subset I_2$$

such that every zero of  $K$  on  $I_2 \times I_1$  lies in  $J_2 \times J_1$ . Replacing them by smaller compact subintervals still containing the zero set, using the boundary-near zero-free region furnished by Lemma 12.77, we may also assume that  $J_2 \times J_1$  is disjoint from that boundary-near region. Then the zero set of  $K$  in the compact set  $J_2 \times J_1$  is nonempty and closed, so by Lemma 12.74 there exists a zero  $(s_*, t_*) \in J_2 \times J_1$  whose  $s$ -coordinate is minimal in the  $x = v_5/c$  ordering among all zeros in  $J_2 \times J_1$ .

By Lemma 12.75,

$$K(s_*, t_*) = 0, \quad \partial_t K(s_*, t_*) = 0,$$

so the tangent line at  $\Gamma(s_*)$  is also tangent at  $\Gamma(t_*)$ .

Across a generic turning boundary, Proposition 12.76 shows that the interior tangent-slope ranges of the two adjacent intervals are disjoint. Hence no common interior tangent slope can occur. This contradicts the existence of  $(s_*, t_*)$ . Therefore no such zero exists, and

$$K(s, t) \neq 0 \quad (s \in I_2, t \in I_1). \quad \square$$

**Remark 12.79** (Status of the turning-boundary adjacent theorem). Theorem 12.78 remains a conditional turning-boundary theorem: the local compact first-zero and first-zero-implies-tangency mechanism is in place, but it still depends on a separate zero-capture statement for the directional tangent-separator kernel  $K_{21}(s, t)$  on the full adjacent pair  $I_2 \times I_1$ .

However, by Theorem 12.92, full zero-capture is no longer needed to exclude actual adjacent pair-pair lines. What remains open here is therefore only the stronger support-ordering / directional-kernel route, not the local adjacent  $2 + 2$  exclusion itself.

**Lemma 12.80** (Boundary-near exclusion for both directional kernels at a generic turning boundary). *Let  $I_1 < I_2$  be adjacent ECT intervals meeting at a generic turning boundary*

$$M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0.$$

Define

$$K_{21}(s, t) := \det(F(s), F'(s), F(t)), \quad K_{12}(t, s) := \det(F(t), F'(t), F(s)),$$

with

$$F(h) = (c(h), v_5(h), \Delta_7(h)).$$

Then there is a neighborhood of the boundary pair on which both directional kernels are nonzero:

$$K_{21}(s, t) \neq 0, \quad K_{12}(t, s) \neq 0$$

for  $s \in I_2, t \in I_1$  sufficiently close to  $h_b$ .

*Proof.* Write

$$x(h) := \frac{v_5(h)}{c(h)}, \quad y(h) := \frac{\Delta_7(h)}{c(h)}.$$

Since

$$y'(h) = \frac{N(h)}{c(h)^2},$$

the hypothesis  $N(h_b) \neq 0$  implies that  $y$  is a valid local coordinate near  $h_b$ . Thus the normalized planar curve may be written locally as a graph

$$x = \chi(y).$$

At  $h_b$ ,

$$\frac{dx}{dy} = \frac{M(h)}{N(h)},$$

so

$$\left. \frac{dx}{dy} \right|_{h_b} = 0.$$

Differentiating once more,

$$\frac{d^2x}{dy^2} = \frac{c(h)^2}{N(h)} \cdot \frac{M'(h)N(h) - M(h)N'(h)}{N(h)^2},$$

and since  $M(h_b) = 0$ ,

$$\left. \frac{d^2x}{dy^2} \right|_{h_b} = \frac{c(h_b)^2 M'(h_b)}{N(h_b)^2} \neq 0.$$

Hence, after shrinking to a small neighborhood of  $h_b$ , the graph  $x = \chi(y)$  is strictly convex or strictly concave.

For such a graph, the tangent line at any point meets the graph only at its point of tangency. Therefore no tangent line from one side can meet the opposite side inside a sufficiently small neighborhood of the boundary. Equivalently, both directional tangent-separator kernels are nonzero near the boundary pair.  $\square$

**Lemma 12.81** (Mirrored first adjacent zero implies tangency). *Let  $J_1 \Subset I_1$ ,  $J_2 \Subset I_2$  be compact interior subintervals across a generic turning boundary, and define*

$$K_{12}(t, s) := \det(F(t), F'(t), F(s)), \quad t \in I_1, s \in I_2.$$

*Assume every zero of  $K_{12}$  on  $I_1 \times I_2$  lies in  $J_1 \times J_2$ . Let  $(t_*, s_*) \in J_1 \times J_2$  be a zero whose  $t$ -coordinate is minimal in the  $x = v_5/c$  ordering among all zeros of  $K_{12}$  on  $I_1 \times I_2$ . Then*

$$\partial_s K_{12}(t_*, s_*) = 0.$$

*Proof.* Because  $M \neq 0$  on the open interval  $I_1$ , the function

$$x(h) = \frac{v_5(h)}{c(h)}$$

is a valid local coordinate on  $I_1$ . Since every zero of  $K_{12}$  lies in the compact interior rectangle  $J_1 \times J_2$ , the chosen point  $(t_*, s_*)$  is interior in the ambient domain  $I_1 \times I_2$ .

Suppose

$$\partial_s K_{12}(t_*, s_*) \neq 0.$$

Then by the implicit function theorem there is a  $C^1$  local zero curve

$$s = s(t)$$

through  $(t_*, s_*)$ , defined for  $t$  near  $t_*$ , such that

$$K_{12}(t, s(t)) = 0.$$

Because  $t_*$  is interior in  $I_1$ , there are nearby points with smaller  $x = v_5/c$  ordering still in  $I_1$ . For such  $t$ , the point  $(t, s(t))$  is also a zero of  $K_{12}$ , contradicting the minimality of  $t_*$ . Therefore

$$\partial_s K_{12}(t_*, s_*) = 0.$$

□

**Theorem 12.82** (Conditional symmetric directional-kernel exclusion across turning boundaries). *Let  $I_1 < I_2$  be adjacent ECT intervals meeting at a generic turning boundary*

$$M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0.$$

*Assume:*

- (1) *the zero-capture hypothesis required in Theorem 12.78 for the kernel  $K_{21}$  on  $I_2 \times I_1$ ;*
- (2) *the symmetric zero-capture hypothesis for  $K_{12}$ , namely that every zero of*

$$K_{12}(t, s) = \det(F(t), F'(t), F(s))$$

*on  $I_1 \times I_2$  lies in some compact interior rectangle  $J_1 \times J_2 \Subset I_1 \times I_2$ .*

*Then both directional kernels are zero-free:*

$$K_{21}(s, t) \neq 0 \quad (s \in I_2, t \in I_1),$$

*and*

$$K_{12}(t, s) \neq 0 \quad (t \in I_1, s \in I_2).$$

*Proof.* The first assertion is exactly Theorem 12.78.

For the second assertion, boundary-near exclusion for  $K_{12}$  is provided by Lemma 12.80. Assume for contradiction that  $K_{12}$  has a zero. By the symmetric zero-capture hypothesis, every zero lies in a compact interior rectangle  $J_1 \times J_2$ . Choose a zero  $(t_*, s_*) \in J_1 \times J_2$  whose  $t$ -coordinate is minimal in the  $x = v_5/c$  ordering among all zeros of  $K_{12}$  on  $I_1 \times I_2$ .

By Lemma 12.81,

$$\partial_s K_{12}(t_*, s_*) = 0.$$

Together with

$$K_{12}(t_*, s_*) = 0,$$

this implies that the tangent line at  $\Gamma(t_*)$  is also tangent at  $\Gamma(s_*)$ . Hence the two adjacent intervals share a common interior tangent slope, contradicting Proposition 12.76. Therefore  $K_{12}$  has no zero. □

**Remark 12.83** (Status of the directional-kernel reduction). Lemma 12.87 reduces an adjacent pair-pair line across a generic turning boundary to a compact interior zero of one of the two directional tangent-separator kernels. Theorem 12.82 shows that the remaining turning-boundary issue is no longer orientation selection between  $K_{21}$  and  $K_{12}$ , but the symmetric zero-capture problem for these two kernels.

**Remark 12.84** (The  $W = 0$  obstruction and the cubic-inflection coefficient). Let

$$x(h) := \frac{v_5(h)}{c(h)}, \quad y(h) := \frac{\Delta_7(h)}{c(h)}.$$

At a boundary point  $h_b$  with

$$W(h_b) = 0, \quad M(h_b) \neq 0,$$

the coordinate  $x$  remains a valid local parameter. Since

$$\frac{d^2y}{dx^2} = \frac{c(h)^3 W(h)}{M(h)^3},$$

one has

$$\left. \frac{d^2y}{dx^2} \right|_{h_b} = 0.$$

Differentiating once more with respect to  $x$  gives

$$\left. \frac{d^3y}{dx^3} \right|_{h_b} = \frac{c(h_b)^5}{M(h_b)^4} W'(h_b).$$

Hence, after translating the point to the origin and subtracting the tangent line, the normalized curve has local expansion

$$y = \frac{c(h_b)^5 W'(h_b)}{6M(h_b)^4} x^3 + O(x^4).$$

Therefore a generic  $W = 0$ ,  $M \neq 0$  boundary is an ordinary affine inflection unless additional special structure forces

$$W'(h_b) = 0.$$

In particular, the tangent-separator route is not presently justified across such boundaries, and the real remaining geometric question is whether the actual triple  $(c, v_5, \Delta_7)$  imposes any non-generic vanishing of  $W'(h_b)$ .

**Lemma 12.85** (Fixed-slope intercept sets for a strictly convex ECT branch). *Let  $I$  be an ECT interval in the affine chart*

$$x = \frac{v_5}{c}, \quad y = \frac{\Delta_7}{c},$$

*and suppose  $\Gamma(I)$  is a strictly convex  $C^2$  graph*

$$\Gamma(I) = \{(x, \psi_I(x)) : x \in J_I\}$$

*over an open interval  $J_I \subset \mathbf{R}$ .*

*For each slope  $m \in \mathbf{R}$ , define*

$$\mathcal{B}_I(m) := \{\psi_I(x) - mx : x \in J_I\}.$$

*Then:*

- (1)  $\mathcal{B}_I(m)$  is an interval.
- (2) A line  $y = mx + b$  meets  $\Gamma(I)$  if and only if

$$b \in \mathcal{B}_I(m).$$

- (3) If  $m$  lies in the interior tangent-slope range of  $I$ , there is a unique  $x_I(m) \in J_I$  such that

$$\psi'_I(x_I(m)) = m,$$

*and the tangent line of slope  $m$  has intercept*

$$\tau_I(m) := \psi_I(x_I(m)) - m x_I(m).$$

Moreover,

$$\tau_I(m) = \inf \mathcal{B}_I(m).$$

(4) For such  $m$ , a line  $y = mx + b$  is 2-secant to  $\Gamma(I)$  if and only if

$$b \in \text{int } \mathcal{B}_I(m).$$

Equivalently, the 2-secant intercepts of slope  $m$  are exactly the interior points of the intercept interval.

*Proof.* For fixed  $m$ , put

$$\phi_m(x) := \psi_I(x) - mx.$$

Since  $\phi_m$  is continuous on the interval  $J_I$ , its image

$$\phi_m(J_I) = \mathcal{B}_I(m)$$

is an interval. This proves (1), and (2) is immediate from

$$(x, \psi_I(x)) \in \{y = mx + b\} \iff \psi_I(x) - mx = b.$$

If  $m$  lies in the interior tangent-slope range, strict convexity implies that  $\psi'_I$  is strictly monotone, hence there is a unique  $x_I(m)$  with  $\psi'_I(x_I(m)) = m$ . At that point,

$$\phi'_m(x_I(m)) = \psi'_I(x_I(m)) - m = 0,$$

and since

$$\phi''_m(x) = \psi''_I(x) > 0,$$

the point  $x_I(m)$  is the unique minimizer of  $\phi_m$ . Thus

$$\tau_I(m) = \phi_m(x_I(m)) = \inf \mathcal{B}_I(m),$$

which proves (3).

Finally, because  $\phi_m$  is strictly convex with a unique minimum, a level  $b$  is attained at two points exactly when it lies in the interior of the interval  $\phi_m(J_I) = \mathcal{B}_I(m)$ . This is precisely the 2-secant condition, proving (4).  $\square$

**Lemma 12.86** (Support-overlap truncation across a generic turning boundary). *Let  $I_1 < I_2$  be adjacent ECT intervals meeting at a generic turning boundary*

$$c(h_b) \neq 0, \quad M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0.$$

For a fixed finite slope  $m$ , define

$$\phi_m(h) := \frac{\Delta_7(h) - mv_5(h)}{c(h)}, \quad \mathcal{B}_I(m) := \phi_m(I).$$

Assume

$$b \in \mathcal{B}_{I_1}(m) \cap \mathcal{B}_{I_2}(m).$$

Then there exists a neighborhood  $U$  of  $h_b$  and compact subintervals

$$J_1 \subseteq I_1 \setminus U, \quad J_2 \subseteq I_2 \setminus U$$

such that

$$b \in \mathcal{B}_{J_1}(m) \cap \mathcal{B}_{J_2}(m).$$

*Proof.* Differentiate:

$$\phi'_m(h) = \frac{N(h) - mM(h)}{c(h)^2}.$$

At  $h = h_b$ , since  $M(h_b) = 0$ ,

$$\phi'_m(h_b) = \frac{N(h_b)}{c(h_b)^2} \neq 0.$$

Hence for each fixed finite  $m$ , after shrinking to a small neighborhood  $U$  of  $h_b$ , the function  $\phi_m$  is strictly monotone on each one-sided piece

$$I_1 \cap U, \quad I_2 \cap U.$$

Therefore, for each  $i = 1, 2$ , the equation

$$\phi_m(h) = b$$

has at most one solution in  $I_i \cap U$ .

But the assumption

$$b \in \mathcal{B}_{I_i}(m)$$

means exactly that there is at least one solution of  $\phi_m(h) = b$  somewhere on  $I_i$ . Therefore, after removing the boundary neighborhood  $U$ , at least one such solution

$$h_i^\# \in I_i \setminus U$$

remains on each side. Choose small compact subintervals

$$J_i \subseteq I_i \setminus U$$

with  $h_i^\# \in J_i$ . Then by continuity,

$$b = \phi_m(h_i^\#) \in \phi_m(J_i) = \mathcal{B}_{J_i}(m),$$

for  $i = 1, 2$ . This proves the claim.  $\square$

**Lemma 12.87** (Compact adjacent pair-pair implies a directional kernel zero). *Let  $I_1 < I_2$  be adjacent ECT intervals meeting at a generic turning boundary, and let*

$$L : y = mx + b$$

*be a finite-slope line that is 2-secant to both  $\Gamma(I_1)$  and  $\Gamma(I_2)$ . Then, after removing a sufficiently small boundary neighborhood, there exist compact subintervals*

$$J_1 \subseteq I_1, \quad J_2 \subseteq I_2$$

*and points  $s \in J_2, t \in J_1$  such that one of the two directional kernels vanishes:*

$$K_{21}(s, t) := \det(F(s), F'(s), F(t)) = 0,$$

*or*

$$K_{12}(t, s) := \det(F(t), F'(t), F(s)) = 0,$$

*where*

$$F(h) = (c(h), v_5(h), \Delta_7(h)).$$

*Proof.* Because  $L$  is 2-secant to each branch, Lemma 12.85 shows that  $m$  lies in the interior tangent-slope range of each interval. Hence each branch has a unique interior tangent line of slope  $m$ , with tangent intercept

$$\tau_{I_i}(m) = \inf \mathcal{B}_{I_i}(m).$$

At a generic turning boundary,

$$\frac{dy/dh}{dx/dh} = \frac{N(h)}{M(h)}$$

has one-sided blow-up, so no finite-slope tangency can occur arbitrarily close to  $h_b$ . Therefore, after shrinking away a boundary neighborhood and applying Lemma 12.86, we may choose compact subintervals

$$J_1 \subseteq I_1, \quad J_2 \subseteq I_2$$

such that:

- (1)  $b \in \mathcal{B}_{J_1}(m) \cap \mathcal{B}_{J_2}(m)$ ;
- (2) the unique slope- $m$  tangency point of each branch lies in the corresponding  $J_i$ .

Hence each  $\mathcal{B}_{J_i}(m)$  is a compact interval whose lower endpoint is the tangent intercept

$$\tau_i := \tau_{J_i}(m) = \inf \mathcal{B}_{J_i}(m).$$

Now set

$$b_* := \inf(\mathcal{B}_{J_1}(m) \cap \mathcal{B}_{J_2}(m)).$$

Since the intersection of two compact intervals is again a compact interval,

$$b_* = \max(\tau_1, \tau_2).$$

So  $b_*$  is equal to one of the two tangent intercepts, and it still lies in the support set of the other branch.

If  $b_* = \tau_2$ , then the slope- $m$  tangent line to  $\Gamma(J_2)$  meets  $\Gamma(J_1)$ . Thus there are points  $s \in J_2$ ,  $t \in J_1$  such that

$$\det(F(s), F'(s), F(t)) = 0,$$

i.e.

$$K_{21}(s, t) = 0.$$

If  $b_* = \tau_1$ , then the slope- $m$  tangent line to  $\Gamma(J_1)$  meets  $\Gamma(J_2)$ . Thus there are points  $t \in J_1$ ,  $s \in J_2$  such that

$$\det(F(t), F'(t), F(s)) = 0,$$

i.e.

$$K_{12}(t, s) = 0.$$

This proves the lemma. □

**Lemma 12.88** (Strip exclusion for  $K_{21}$  against a compact opposite branch). *Let  $I_1 < I_2$  be adjacent ECT intervals meeting at a generic turning boundary*

$$M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0.$$

*Let  $J_1 \Subset I_1$  be a compact subinterval. Then there exists a one-sided neighborhood  $U_2 \subset I_2$  of the shared boundary point  $h_b$  such that*

$$K_{21}(s, t) := \det(F(s), F'(s), F(t)) \neq 0 \quad (s \in U_2, t \in J_1),$$

*where  $F(h) = (c(h), v_5(h), \Delta_7(h))$ .*

*Proof.* Work in the affine chart

$$x(h) := \frac{v_5(h)}{c(h)}, \quad y(h) := \frac{\Delta_7(h)}{c(h)}.$$

At a generic turning boundary one has

$$x'(h_b) = 0, \quad x''(h_b) \neq 0, \quad y'(h_b) \neq 0.$$

Hence  $y$  is a valid local coordinate and the normalized curve may be written near  $h_b$  as a graph

$$x = \chi(y)$$

with

$$\chi'(y_b) = 0, \quad \chi''(y_b) \neq 0.$$

Thus  $x$  has a strict local extremum at the boundary, while the tangent slope

$$m(h) = \frac{dy/dh}{dx/dh} = \frac{N(h)}{M(h)}$$

blows up one-sidedly as  $h \rightarrow h_b$  from either adjacent interval.

Because  $J_1 \Subset I_1$  is compact away from  $h_b$ , there exists  $\delta > 0$  such that

$$|x(t) - x(h_b)| \geq 2\delta \quad (t \in J_1).$$

After shrinking to a small one-sided neighborhood  $U_2 \subset I_2$  of  $h_b$ , we may also assume

$$|x(s) - x(h_b)| \leq \delta \quad (s \in U_2).$$

Hence

$$|x(t) - x(s)| \geq \delta \quad (s \in U_2, t \in J_1).$$

Fix a preliminary compact neighborhood  $U_2^0 \Subset I_2$  of  $h_b$  and set

$$B := \sup\{|y(t) - y(s)| : t \in J_1, s \in U_2^0\} < \infty.$$

Since  $m(s) \rightarrow \pm\infty$  as  $s \rightarrow h_b$  in  $I_2$ , we may shrink further to  $U_2 \subset U_2^0$  so that

$$|m(s)| > \frac{2B}{\delta} \quad (s \in U_2).$$

If  $K_{21}(s, t) = 0$ , then the tangent line at  $\Gamma(s)$  passes through  $\Gamma(t)$ , equivalently

$$y(t) - y(s) = m(s)(x(t) - x(s)).$$

Taking absolute values gives

$$B \geq |y(t) - y(s)| = |m(s)| |x(t) - x(s)| > \frac{2B}{\delta} \cdot \delta = 2B,$$

a contradiction. Therefore  $K_{21}(s, t) \neq 0$  on  $U_2 \times J_1$ .  $\square$

**Lemma 12.89** (Strip exclusion for  $K_{12}$  against a compact opposite branch). *Let  $I_1 < I_2$  be adjacent ECT intervals meeting at a generic turning boundary*

$$M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0.$$

*Let  $J_2 \Subset I_2$  be a compact subinterval. Then there exists a one-sided neighborhood  $U_1 \subset I_1$  of the shared boundary point  $h_b$  such that*

$$K_{12}(t, s) := \det(F(t), F'(t), F(s)) \neq 0 \quad (t \in U_1, s \in J_2),$$

*where  $F(h) = (c(h), v_5(h), \Delta_7(h))$ .*

*Proof.* This is the left-right mirror of Lemma 12.88.  $\square$

**Lemma 12.90** (Compact first-contact for  $K_{21}$ ). *Let  $J_1 \Subset I_1$ ,  $J_2 \Subset I_2$  be compact subintervals across a generic turning boundary. Assume there is a relatively open collar  $V_2 \subset J_2$  at the boundary-facing end of  $J_2$  such that*

$$K_{21}(s, t) \neq 0 \quad (s \in V_2, t \in J_1).$$

*Assume also that the zero set of  $K_{21}$  in  $J_2 \times J_1$  is nonempty. Then there exists a zero  $(s_*, t_*) \in J_2 \times J_1$  such that:*

- (1)  $s_*$  is boundary-nearest in the  $x = v_5/c$  ordering among all  $s \in J_2$  for which  $K_{21}(s, t) = 0$  for some  $t \in J_1$ ;
- (2)

$$\partial_t K_{21}(s_*, t_*) = 0.$$

*Proof.* Define

$$A_{21} := \{s \in J_2 : \exists t \in J_1 \text{ with } K_{21}(s, t) = 0\}.$$

Since the zero set of  $K_{21}$  in  $J_2 \times J_1$  is compact and nonempty, the set  $A_{21}$  is compact and nonempty. By hypothesis,

$$A_{21} \cap V_2 = \emptyset.$$

Hence  $A_{21}$  has a boundary-nearest element  $s_*$  in the  $x = v_5/c$  ordering on  $J_2$ . Choose  $t_* \in J_1$  with

$$K_{21}(s_*, t_*) = 0.$$

Suppose

$$\partial_t K_{21}(s_*, t_*) \neq 0.$$

Then by the implicit function theorem there is a  $C^1$  local zero curve

$$t = t(s)$$

through  $(s_*, t_*)$ , defined for  $s$  near  $s_*$ , such that

$$K_{21}(s, t(s)) = 0.$$

Because  $s_* \notin V_2$  and  $V_2$  is an open collar at the boundary-facing end of  $J_2$ , there exist nearby points  $s \in J_2$  that are boundary-nearer than  $s_*$  in the  $x = v_5/c$  ordering. For such  $s$ , the point  $(s, t(s))$  is also a zero of  $K_{21}$ , contradicting the minimality of  $s_*$ . Therefore

$$\partial_t K_{21}(s_*, t_*) = 0.$$

$\square$



**Lemma 12.91** (Compact first-contact for  $K_{12}$ ). *Let  $J_1 \Subset I_1$ ,  $J_2 \Subset I_2$  be compact subintervals across a generic turning boundary. Assume there is a relatively open collar  $V_1 \subset J_1$  at the boundary-facing end of  $J_1$  such that*

$$K_{12}(t, s) \neq 0 \quad (t \in V_1, s \in J_2).$$

*Assume also that the zero set of  $K_{12}$  in  $J_1 \times J_2$  is nonempty. Then there exists a zero  $(t_*, s_*) \in J_1 \times J_2$  such that:*

- (1)  $t_*$  is boundary-nearest in the  $x = v_5/c$  ordering among all  $t \in J_1$  for which  $K_{12}(t, s) = 0$  for some  $s \in J_2$ ;
- (2)

$$\partial_s K_{12}(t_*, s_*) = 0.$$

*Proof.* Define

$$A_{12} := \{t \in J_1 : \exists s \in J_2 \text{ with } K_{12}(t, s) = 0\}.$$

Since the zero set of  $K_{12}$  in  $J_1 \times J_2$  is compact and nonempty, the set  $A_{12}$  is compact and nonempty. By hypothesis,

$$A_{12} \cap V_1 = \emptyset.$$

Hence  $A_{12}$  has a boundary-nearest element  $t_*$  in the  $x = v_5/c$  ordering on  $J_1$ . Choose  $s_* \in J_2$  with

$$K_{12}(t_*, s_*) = 0.$$

Suppose

$$\partial_s K_{12}(t_*, s_*) \neq 0.$$

Then by the implicit function theorem there is a  $C^1$  local zero curve

$$s = s(t)$$

through  $(t_*, s_*)$ , defined for  $t$  near  $t_*$ , such that

$$K_{12}(t, s(t)) = 0.$$

Because  $t_* \notin V_1$  and  $V_1$  is an open collar at the boundary-facing end of  $J_1$ , there exist nearby points  $t \in J_1$  that are boundary-nearer than  $t_*$  in the  $x = v_5/c$  ordering. For such  $t$ , the point  $(t, s(t))$  is also a zero of  $K_{12}$ , contradicting the minimality of  $t_*$ . Therefore

$$\partial_s K_{12}(t_*, s_*) = 0.$$

□

**Theorem 12.92** (Adjacent pair-pair exclusion across a generic turning boundary). *Let  $I_1 < I_2$  be adjacent ECT intervals meeting at a generic turning boundary*

$$M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0.$$

*Then no finite-slope affine line in the chart*

$$x = \frac{v_5}{c}, \quad y = \frac{\Delta_7}{c}$$

*is 2-secant to both  $\Gamma(I_1)$  and  $\Gamma(I_2)$ .*

*Proof.* Assume for contradiction that a finite-slope line

$$L : y = mx + b$$

is 2-secant to both adjacent branches. By Lemma 12.87, after removing a small boundary neighborhood there exist compact subintervals

$$J_1 \Subset I_1, \quad J_2 \Subset I_2$$

and a zero of one of the two directional kernels.

First suppose there exist  $s_0 \in J_2$ ,  $t_0 \in J_1$  such that

$$K_{21}(s_0, t_0) = 0.$$

By Lemma 12.88, there exists a one-sided neighborhood  $U_2 \subset I_2$  of  $h_b$  such that

$$K_{21}(s, t) \neq 0 \quad (s \in U_2, t \in J_1).$$

Choose a compact interval  $\tilde{J}_2 \Subset I_2$  containing  $s_0$  and whose boundary-facing collar  $V_2$  lies in  $U_2$ . Then the zero set of  $K_{21}$  in  $\tilde{J}_2 \times J_1$  is nonempty, and Lemma 12.90 yields a zero

$$(s_*, t_*) \in \tilde{J}_2 \times J_1$$

such that

$$\partial_t K_{21}(s_*, t_*) = 0.$$

Together with  $K_{21}(s_*, t_*) = 0$ , this means that the tangent line at  $\Gamma(s_*)$  is also tangent at  $\Gamma(t_*)$ . Since  $s_* \in I_2$  and  $t_* \in I_1$  are interior points, this gives a common interior tangent slope across the adjacent branches, contradicting Proposition 12.76.

Now suppose instead there exist  $t_0 \in J_1$ ,  $s_0 \in J_2$  such that

$$K_{12}(t_0, s_0) = 0.$$

By Lemma 12.89, there exists a one-sided neighborhood  $U_1 \subset I_1$  of  $h_b$  such that

$$K_{12}(t, s) \neq 0 \quad (t \in U_1, s \in J_2).$$

Choose a compact interval  $\tilde{J}_1 \Subset I_1$  containing  $t_0$  and whose boundary-facing collar  $V_1$  lies in  $U_1$ . Then the zero set of  $K_{12}$  in  $\tilde{J}_1 \times J_2$  is nonempty, and Lemma 12.91 yields a zero

$$(t_*, s_*) \in \tilde{J}_1 \times J_2$$

such that

$$\partial_s K_{12}(t_*, s_*) = 0.$$

Together with  $K_{12}(t_*, s_*) = 0$ , this means that the tangent line at  $\Gamma(t_*)$  is also tangent at  $\Gamma(s_*)$ . Again this gives a common interior tangent slope across the adjacent branches, contradicting Proposition 12.76.

Both cases are impossible. Therefore no finite-slope line is 2-secant to both adjacent branches across a generic turning boundary.  $\square$

**Remark 12.93** (What the unconditional adjacent turning theorem does and does not yet replace). Theorem 12.92 gives an unconditional exclusion of finite-slope pair-pair lines across a single generic turning boundary. It is stronger than the older local directional-kernel reduction in that it bypasses full symmetric zero-capture.

However, it does not by itself replace Theorem 12.96. To obtain a full global no-four-secant theorem from it, one would still need a separate propagation statement reducing nonadjacent pair-pair lines to the adjacent turning-boundary case.

**Remark 12.94** (Status of the directional-kernel reduction). Lemma 12.87 reduces an adjacent pair-pair line across a generic turning boundary to a compact interior zero of one of the two directional tangent-separator kernels.

There are now two closures of that reduction in the draft:

- (1) Theorem 12.92 gives an unconditional closure for actual adjacent pair-pair lines, via strip exclusion and compact first-contact.
- (2) Theorem 12.82 gives a stronger zero-freeness statement for both directional kernels, but only conditionally on symmetric zero-capture.

Thus orientation selection between  $K_{21}$  and  $K_{12}$  is no longer the main issue. What remains open only for the stronger tangent-separator / support-ordering route is the symmetric zero-capture problem on the full adjacent rectangle.

**Lemma 12.95** (Adjacent support-ordering). *Let  $I_1 < I_2$  be adjacent ECT intervals under the hypotheses of Theorem 12.78. Then, for every slope  $m \in \mathbf{R}$  such that both fixed-slope intercept sets are nonempty,*

$$\mathcal{B}_{I_1}(m) \cap \mathcal{B}_{I_2}(m) = \emptyset.$$

More precisely, one has one of the two strict alternatives

$$\sup \mathcal{B}_{I_1}(m) < \inf \mathcal{B}_{I_2}(m) \quad \text{or} \quad \sup \mathcal{B}_{I_2}(m) < \inf \mathcal{B}_{I_1}(m),$$

and the same alternative holds for all such  $m$  on the adjacent pair.

*Proof.* Fix  $m$  such that both  $\mathcal{B}_{I_1}(m)$  and  $\mathcal{B}_{I_2}(m)$  are nonempty.

First assume  $m$  lies in the interior tangent-slope range of  $I_2$ . Let  $s \in I_2$  be the unique point with tangent slope  $m$ . Then by Lemma 12.85,

$$\tau_{I_2}(m) = \inf \mathcal{B}_{I_2}(m).$$

By Theorem 12.78,

$$K(s, t) \neq 0 \quad (t \in I_1).$$

By Lemma 12.72,

$$K(s, t) = c(s)^2 c(t) x'(s) (\phi_m(t) - \tau_{I_2}(m)), \quad \phi_m(t) := y(t) - mx(t).$$

Since  $c(s)^2 c(t) x'(s)$  has fixed sign on  $I_1$  and  $K(s, t)$  is continuous and nowhere zero there, the quantity

$$\phi_m(t) - \tau_{I_2}(m)$$

has constant sign on  $I_1$ . To determine which sign occurs, move  $s$  toward the common boundary point from inside  $I_2$ . By Lemma 12.77, the limiting boundary tangent line does not meet  $I_1$ , so  $I_1$  lies entirely on one side of that tangent line. By continuity, the same side persists for nearby interior tangents. Indeed, if the side changed as the tangency point  $s$  moved along  $I_2$ , then by the connectedness of  $I_2$  and continuity of the tangent-line data there would exist an intermediate point  $s_* \in I_2$  for which the tangent line at  $\Gamma(s_*)$  meets  $\Gamma(I_1)$ . Equivalently, there would exist  $t_* \in I_1$  such that

$$K(s_*, t_*) = 0,$$

contradicting Theorem 12.78. Hence the same side holds for all  $s \in I_2$ . Therefore either

$$\phi_m(t) < \tau_{I_2}(m) \quad \text{for all } t \in I_1,$$

or

$$\phi_m(t) > \tau_{I_2}(m) \quad \text{for all } t \in I_1.$$

In the first case,

$$\sup \mathcal{B}_{I_1}(m) \leq \tau_{I_2}(m) = \inf \mathcal{B}_{I_2}(m),$$

and since equality would force an intersection with the tangent line, the inequality is strict. The second case is symmetric.

Now suppose  $m$  is not in the interior tangent-slope range of  $I_2$ , but both intercept sets are nonempty. Then  $m$  lies on one side of the tangent slope range of  $I_2$ , so  $\phi_m$  is strictly monotone on  $I_2$ , and  $\mathcal{B}_{I_2}(m)$  is an interval with endpoint attained in the one-sided limit at the adjacent boundary. Passing to that boundary-support line and using Lemma 12.77 again shows that  $I_1$  lies strictly on one side of that boundary-support line. Hence the same strict ordering of intercept sets holds in this case as well.

Thus  $\mathcal{B}_{I_1}(m)$  and  $\mathcal{B}_{I_2}(m)$  are disjoint for every slope  $m$  for which both are nonempty, and the side ordering is constant on the adjacent pair.  $\square$

**Theorem 12.96** (Partial global no-secant theorem on turning-boundary generic blocks). *Assume every adjacent pair of ECT intervals is separated by a generic turning boundary*

$$M(h_b) = 0, \quad M'(h_b) \neq 0, \quad N(h_b) \neq 0,$$

*and assume moreover that the zero-capture hypothesis required in Theorem 12.78 holds for every adjacent turning-boundary pair. Then no affine line in the chart*

$$x = \frac{v_5}{c}, \quad y = \frac{\Delta_7}{c}$$

*is 2-secant to two distinct ECT intervals. In particular, there is no global four-secant line for the normalized planar curve  $\Gamma(h)$  on such a turning-boundary generic block.*

*Proof.* Under the stated hypotheses, every adjacent boundary is of turning type and Theorem 12.78 applies to each adjacent pair. For such adjacent pairs, Lemma 12.95 applies. Hence adjacent fixed-slope intercept sets are strictly ordered. Iterating this ordering over the  $x = v_5/c$  ordering of ECT intervals shows that no affine line can be 2-secant to two distinct ECT intervals. Therefore no global four-secant line exists on the block.  $\square$

**Corollary 12.97** (Partial exclusion of pair-pair branch configurations). *Under the hypotheses of Theorem 12.96, a bad finite core cannot contain two distinct ECT intervals each contributing a pair of points on a turning-boundary generic block.*

*Proof.* Each ECT interval contributes at most two points to the zero set of an affine line in the normalized chart, and by Theorem 12.96 no affine line can be 2-secant to two different ECT intervals on such a block. Therefore a pair-pair configuration across two distinct ECT intervals is impossible there.  $\square$

**Remark 12.98** (Status of the direct septic witness). The direct  $q^{(7)}$ -slice of total  $\Delta_7$  is now closed at the weight-one level by Proposition 12.48. Together with Theorem 12.49, this closes the canonical quotient-level order-7 package.

What remains non-unique at order 7 is only the lower/mixed septic completion of a raw representative pair  $(u_7, v_7)$ , not any further canonical quotient-level invariant. Accordingly, order-7 is no longer a live audited obstruction for the later geometric arguments.

**Remark 12.99** (Work in progress: highest-new scalar on the direct septic slice). On the region  $v_5 \neq 0$ , set

$$\mu := \frac{c}{v_5}, \quad \lambda := \frac{\Delta_7}{v_5}, \quad \phi := \lambda^{[1]}.$$

By Corollary 12.43, Corollary 12.47, and Theorem 12.49, the highest-new quotient-level septic scalar is already reduced to the direct  $q^{(7)}$ -driven  $K_1$ -slice:

$$\Delta_7^{[1]} = v_5 \phi, \quad \phi = \kappa \left( (x_2 - x_1) \nu_6^{(q^{(7)})} + v(\gamma_5 - \alpha_5)^{(q^{(7)})} \right).$$

In the minimal graded model explored in the present worklog, the combined coefficient is proportional to

$$\frac{eu(a^2 + ab - b^2)}{a^2 b^4 (a + b)^2},$$

so the highest-new slice is generically nonzero but not manifestly one-signed. Thus the remaining order-7 issue is no longer survival of the canonical quotient-level witness, but the global line-incidence / slope-recurrence geometry of the resulting scalar graph.

**Remark 12.100** (Status of the  $W = 0$ ,  $M \neq 0$  side). On the region  $c \neq 0$ ,  $M \neq 0$ , Proposition 12.59 excludes open-interval flatness  $W \equiv 0$ . At an isolated boundary

$$W(h_b) = 0, \quad M(h_b) \neq 0,$$

the normalized planar curve in the chart

$$x = \frac{v_5}{c}, \quad y = \frac{\Delta_7}{c}$$

is an ordinary odd-order affine inflection whenever  $W$  changes sign, so adjacent tangent-separator sign regularity fails there in general.

However, Theorem 12.102 shows that every finite-order isolated boundary of type  $W = 0$ ,  $M \neq 0$  is locally sterile for pair-pair birth. Thus the remaining issue on this side is not local creation of a bad  $2 + 2$  line, but global continuation / recurrence of determinant-zero or tangent-slope data across multiple intervals.

The exact identities for  $W$  and  $W'$  continue to show that this residual obstruction is still encoded through the canonical order-7 package  $\Delta_7$ , not by a genuinely new primitive order-9 invariant.

**Remark 12.101** (Audit:  $W'(h_b)$  at  $W = 0$ ,  $M \neq 0$ ). At a point  $h_b$  with

$$W(h_b) = 0, \quad M(h_b) \neq 0,$$

the cubic-inflection coefficient of the normalized planar curve in the  $(\mu, \lambda)$ -coordinates is proportional to

$$W'(h_b).$$

So  $W'(h_b)$  still governs the local simple-inflection jet.

But proving  $W'(h_b) = 0$  is no longer the main target, because simple  $W = 0$ ,  $M \neq 0$  boundaries are already locally benign for pair-pair birth. The live question is now whether the canonical order-7 package imposes any further global non-recurrence on the associated determinant/tangent-slope data.

At the highest-new level,  $[\Delta_7]^{[1]}$  is already reduced to the direct  $q^{(7)}$ -driven  $K_1$ -slice, so the present obstruction is not a new primitive order-9 invariant but a question about global propagation of the canonical order-7 geometry.

**Theorem 12.102** (Finite-order  $W = 0$  boundaries are locally sterile for pair-pair birth). *Let  $h_b$  satisfy*

$$c(h_b) \neq 0, \quad M(h_b) \neq 0, \quad W(h_b) = 0,$$

*and assume that  $W$  has finite vanishing order  $r \geq 1$  at  $h_b$ :*

$$W(h) = (h - h_b)^r \widetilde{W}(h), \quad \widetilde{W}(h_b) \neq 0.$$

*Define*

$$F(h) := (c(h), v_5(h), \Delta_7(h)), \quad D(s, t, u) := \det(F(s), F(t), F(u)).$$

*Then, after shrinking to a sufficiently small neighborhood  $U$  of  $h_b$ , the following holds.*

*If  $s < t < h_b$  lie in  $U$ , then:*

- (1) *if  $r$  is even, the function  $u \mapsto D(s, t, u)$  has no zero on the opposite side  $h_b < u \in U$ ;*
- (2) *if  $r$  is odd, the function  $u \mapsto D(s, t, u)$  has at most one zero on the opposite side  $h_b < u \in U$ .*

*The same conclusion holds with left and right exchanged. In particular, no finite-order boundary of type  $W = 0$ ,  $M \neq 0$  can support a local pair-pair birth.*

*Proof.* Work in the affine chart

$$x(h) := \frac{v_5(h)}{c(h)}, \quad y(h) := \frac{\Delta_7(h)}{c(h)}.$$

Since

$$x'(h) = \frac{M(h)}{c(h)^2},$$

the hypothesis  $M(h_b) \neq 0$  implies that  $x$  is a valid local coordinate near  $h_b$ . After translating so that  $x(h_b) = 0$  and subtracting the tangent line, we may write the normalized curve as

$$y = f(x), \quad f(0) = f'(0) = 0.$$

By Proposition 12.57,

$$\frac{d^2 y}{dx^2} = \frac{c(h)^3 W(h)}{M(h)^3}.$$

Because  $x'(h_b) \neq 0$ , the finite vanishing order of  $W$  in  $h$  is the same as its finite vanishing order in the local coordinate  $x$ . Therefore

$$f''(x) = \lambda x^r + O(x^{r+1}) \quad (\lambda \neq 0).$$

Fix two same-side points

$$x_1 < x_2 < 0$$

sufficiently close to 0, and define

$$\widehat{D}(x_1, x_2, u) := \det \begin{pmatrix} 1 & x_1 & f(x_1) \\ 1 & x_2 & f(x_2) \\ 1 & u & f(u) \end{pmatrix}.$$

Then

$$\widehat{D}(x_1, x_2, u) = (x_2 - x_1)(u - x_1)(u - x_2) H_{x_1, x_2}(u),$$

where

$$H_{x_1, x_2}(u) := f[x_1, x_2, u]$$

is the second divided difference of  $f$ .

If  $r$  is even, then  $f''$  has fixed sign near 0. By the Hermite–Genocchi formula for second divided differences,  $H_{x_1, x_2}(u)$  has that same fixed sign for  $u > 0$  sufficiently small. Hence  $\widehat{D}(x_1, x_2, u) \neq 0$  on the opposite side.

If  $r$  is odd, then

$$f'''(x) = \lambda r x^{r-1} + O(x^r)$$

has fixed nonzero sign near 0, because  $r - 1$  is even. Differentiating with respect to the repeated node gives

$$\partial_u H_{x_1, x_2}(u) = f[x_1, x_2, u, u].$$

By the Hermite–Genocchi formula for repeated-node divided differences,  $f[x_1, x_2, u, u]$  has the same fixed nonzero sign for  $u > 0$  sufficiently small. Thus  $u \mapsto H_{x_1, x_2}(u)$  is strictly monotone on the opposite side, and therefore has at most one zero there.

Finally,

$$D(s, t, u) = c(s)c(t)c(u)\widehat{D}(x(s), x(t), x(u)),$$

and  $c$  is nonzero on a small neighborhood of  $h_b$ , so the zero set of  $D$  is exactly the zero set of  $\widehat{D}$ . The theorem follows.  $\square$

**Remark 12.103** (Consequence for the  $W = 0$ ,  $M \neq 0$  endgame). Theorem 12.102 shows that finite-order  $W = 0$ ,  $M \neq 0$  boundaries are not local birthplaces of pair-pair configurations. Combined with Proposition 12.59, this means the remaining issue on the  $W = 0$  side is genuinely global: continuation or recurrence of determinant-zero / tangent-slope data across multiple intervals, not local creation of a second opposite-side zero branch.

**Remark 12.104** (Audit: residual  $W = 0$  pathology). By Proposition 12.59, the  $W = 0$ ,  $M \neq 0$  side is no longer obstructed by open-interval flatness  $W \equiv 0$  on a component with  $c \neq 0$  and  $M \neq 0$ . By Theorem 12.102, every finite-order isolated  $W = 0$  boundary is locally sterile for pair-pair birth. Thus any remaining  $W = 0$  issue would have to come either from non-analytic / infinite-order flatness or from a genuinely global continuation phenomenon, not from a generic local inflection birth mechanism.

## 12.6. From the one-pair residual germ to two-pair and minimal-core rigidity.

**Remark 12.105** (Strengthened organizing principle). The original version of this subsection was organized around the cubic leading line and the first residual trace witness. The later working analysis suggests a stronger one-pair package:

$$c(h), \quad A_5^f(h), \quad \Delta_7(h),$$

or equivalently the normalized affine curve

$$h \mapsto \left( \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)} \right).$$

The intended two-pair and minimal-core rigidity statements should ultimately be reformulated in terms of this strengthened one-pair invariant geometry.

**Remark 12.106** (Work in progress). This subsection is best read as a rigidity program built on the strengthened one-pair local package

$$c(h), \quad A_5^f(h), \quad \Delta_7(h),$$

or equivalently the normalized planar curve

$$h \mapsto \left( \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)} \right).$$

The projective coincidence conditions remain the intended organizing principle for the two-pair and minimal-core analysis; the remaining steps are now algebraic interaction corrections and the extraction into the finite-core endgame.

**Remark 12.107** (Interpretation of residual notation in the multi-pair architecture). In the remainder of this section, the symbols

$$F_h(z), \quad \ell_h, \quad \bar{v}_1(h), \quad \mathcal{Q}^{\text{abs}}/\ell_h$$

are used schematically to describe the intended cubic-leading / first-residual architecture of multi-pair and minimal-core configurations.

The fully explicit rigorous first-residual object established earlier is

$$\mathcal{R}_h^{(5)} = \mathcal{J}_5 / (\mathbf{C} A_3(h) + V_{\text{amb}}^{(5)}(h)).$$

Accordingly, whenever a later statement invokes the first residual class in a theorem-strength way, it should be interpreted through  $\mathcal{R}_h^{(5)}$ . The subsequent two-pair and minimal-bad-core statements remain architectural until the interaction terms are rewritten in this explicit finite-order model.

Let  $F_h(z)$  denote the corrected quotient odd germ of the normalized one-pair family. By Proposition 12.8, on the cubic-generic region we have

$$F_h(z) = v_0(h)z^3 + v_1(h)z^5 + \cdots, \quad v_0(h) \neq 0,$$

with cubic leading line

$$\ell_h := \mathbf{C} v_0(h) \subset \mathcal{Q}^{\text{abs}}.$$

Under the additional hypotheses of Proposition 12.28, there is an open dense subset  $\mathcal{U}_{\text{res}} \subset \mathcal{U}_{\text{cub}}$  on which the explicit first residual class

$$\bar{v}_1(h) \in \mathcal{R}_h^{(5)}$$

is nonzero.

In the strengthened fixed-sector refinement, the one-pair local package is recorded by

$$c(h), \quad A_5^{\mathbf{f}}(h) = u_5(h)I + v_5(h)S, \quad \Delta_7(h),$$

and hence by the normalized planar curve

$$\Gamma(h) := \left( \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)} \right) = (\rho_1(h), \rho_2(h)).$$

By Proposition 12.54, the affine curvature numerator of  $\Gamma$  is

$$K_{\Gamma}(h) = \rho_1'(h)\rho_2''(h) - \rho_2'(h)\rho_1''(h) = \frac{M(h)N'(h) - N(h)M'(h)}{c(h)^4},$$

where

$$M(h) = c(h)v_5'(h) - c'(h)v_5(h), \quad N(h) = c(h)\Delta_7'(h) - c'(h)\Delta_7(h).$$

Thus the strengthened one-pair geometry is governed by the curve  $\Gamma(h)$  and its secant/tangent behavior on the region where

$$c(h) \neq 0, \quad M(h) \neq 0, \quad M(h)N'(h) - N(h)M'(h) \neq 0.$$

**Definition 12.108** (Amplitude-invariant strengthened datum). On the region where

$$c(h) \neq 0,$$

define

$$\widehat{\Psi}(h) := \left( \frac{u_5(h)}{c(h)}, \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)^2} \right).$$

**Lemma 12.109** (Scaling law for the strengthened package). *If the normalized one-pair odd germ is multiplied by a scalar  $\lambda$ , then*

$$c \mapsto \lambda c, \quad A_5^f \mapsto \lambda A_5^f, \quad A_7^f \mapsto \lambda A_7^f,$$

and hence

$$\Delta_7 \mapsto \lambda^2 \Delta_7.$$

Consequently the datum

$$\widehat{\Psi}(h) = \left( \frac{u_5(h)}{c(h)}, \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)^2} \right)$$

is invariant under scalar rescaling of the full one-pair germ.

*Proof.* The cubic, quintic, and septic coefficients of the odd germ all scale linearly under multiplication of the germ by  $\lambda$ . Since

$$A_5^f = u_5 I + v_5 S, \quad A_7^f = u_7 I + v_7 S,$$

it follows that

$$u_5 \mapsto \lambda u_5, \quad v_5 \mapsto \lambda v_5, \quad u_7 \mapsto \lambda u_7, \quad v_7 \mapsto \lambda v_7.$$

Therefore

$$\Delta_7 = u_7 v_5 - u_5 v_7$$

scales as

$$\Delta_7 \mapsto \lambda^2 \Delta_7,$$

which gives the claim. □

**Remark 12.110** (Two normalizations: geometric versus projective). The affine curve

$$\Gamma(h) = \left( \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)} \right)$$

is the correct object for the one-pair differential geometry, because its derivatives are governed by the projective Wronskians  $M$ ,  $N$ , and  $W$ .

For the multi-pair coincidence problem, however, the natural scalar-invariant datum is  $\widehat{\Psi}(h)$ , not  $\Gamma(h)$ . The reason is exactly the scaling law in Lemma 12.109: under scalar rescaling of the full one-pair germ,  $\Delta_7/c$  is not invariant, whereas  $\Delta_7/c^2$  is.

In this formulation, the geometric secant/tangent obstruction is well controlled on ECT intervals and across generic turning-point boundaries. The remaining global issue is no longer a purely local  $W = 0$ ,  $M \neq 0$  obstruction, but the global propagation / recurrence problem for the determinant-zero and tangent configurations that survive the local sterility results proved later in the paper.

**Definition 12.111** (Two-pair decomposition). For a normalized two-pair core with amplitudes  $a_1, a_2$  and offsets  $h_1, h_2$ , write its quotient odd germ as

$$F_{(a_1, h_1; a_2, h_2)}(z) = a_1 F_{h_1}(z) + a_2 F_{h_2}(z) + \mathcal{I}_{12}(z),$$

where  $\mathcal{I}_{12}(z)$  denotes the pair-interaction germ, quadratic in the pair amplitudes.

**Proposition 12.112** (Exact strengthened two-pair coincidence through order 7). *Let  $(a_1, h_1; a_2, h_2)$  be a normalized two-pair core with*

$$F_{(a_1, h_1; a_2, h_2)}(z) \equiv 0.$$

*Assume moreover that, after passage to the explicit corrected finite-order model through order 7, the interaction contribution vanishes at the three relevant levels, namely:*

$$a_1 c(h_1) + a_2 c(h_2) = 0,$$

$$a_1 A_5^f(h_1) + a_2 A_5^f(h_2) = 0,$$

*and, after the second identity identifies the common quotient line*

$$\mathbf{C} A_5^f(h_1) = \mathbf{C} A_5^f(h_2),$$



the second residual classes satisfy

$$a_1 [A_7^{\mathfrak{f}}](h_1) + a_2 [A_7^{\mathfrak{f}}](h_2) = 0 \quad \text{in} \quad \mathfrak{f}/\mathbf{C}A_5^{\mathfrak{f}}(h_1).$$

Then

$$\widehat{\Psi}(h_1) = \widehat{\Psi}(h_2).$$

Equivalently,

$$\frac{u_5(h_1)}{c(h_1)} = \frac{u_5(h_2)}{c(h_2)}, \quad \frac{v_5(h_1)}{c(h_1)} = \frac{v_5(h_2)}{c(h_2)}, \quad \frac{\Delta_7(h_1)}{c(h_1)^2} = \frac{\Delta_7(h_2)}{c(h_2)^2}.$$

*Proof.* Set

$$\lambda := -\frac{a_2}{a_1}.$$

The cubic identity gives

$$c(h_1) = \lambda c(h_2).$$

The quintic fixed-sector identity gives

$$A_5^{\mathfrak{f}}(h_1) = \lambda A_5^{\mathfrak{f}}(h_2).$$

Writing

$$A_5^{\mathfrak{f}}(h_i) = u_5(h_i)I + v_5(h_i)S,$$

we obtain

$$u_5(h_1) = \lambda u_5(h_2), \quad v_5(h_1) = \lambda v_5(h_2).$$

Hence

$$\frac{u_5(h_1)}{c(h_1)} = \frac{u_5(h_2)}{c(h_2)}, \quad \frac{v_5(h_1)}{c(h_1)} = \frac{v_5(h_2)}{c(h_2)}.$$

Now the second identity implies

$$\mathbf{C}A_5^{\mathfrak{f}}(h_1) = \mathbf{C}A_5^{\mathfrak{f}}(h_2),$$

so the residual quotient spaces are canonically identified. The order-7 residual identity therefore says

$$[A_7^{\mathfrak{f}}](h_1) = \lambda [A_7^{\mathfrak{f}}](h_2)$$

in the common quotient. Equivalently, there exists  $\mu \in \mathbf{C}$  such that

$$A_7^{\mathfrak{f}}(h_1) = \lambda A_7^{\mathfrak{f}}(h_2) + \mu A_5^{\mathfrak{f}}(h_2).$$

Writing

$$A_7^{\mathfrak{f}}(h_i) = u_7(h_i)I + v_7(h_i)S,$$

and using

$$\Delta_7(h) = u_7(h)v_5(h) - u_5(h)v_7(h),$$

we obtain

$$\Delta_7(h_1) = \det(A_7^{\mathfrak{f}}(h_1), A_5^{\mathfrak{f}}(h_1)) = \det(\lambda A_7^{\mathfrak{f}}(h_2) + \mu A_5^{\mathfrak{f}}(h_2), \lambda A_5^{\mathfrak{f}}(h_2)) = \lambda^2 \Delta_7(h_2).$$

Since also  $c(h_1) = \lambda c(h_2)$ , it follows that

$$\frac{\Delta_7(h_1)}{c(h_1)^2} = \frac{\Delta_7(h_2)}{c(h_2)^2}.$$

Thus  $\widehat{\Psi}(h_1) = \widehat{\Psi}(h_2)$ . □

**Theorem 12.113** (Quantitative defect-thickened strengthened coincidence). *Let  $(a_1, h_1; a_2, h_2)$  be a normalized bad two-pair core, and write*

$$c_i := c(h_i), \quad A_{5,i}^{\mathfrak{f}} := A_5^{\mathfrak{f}}(h_i) = u_{5,i}I + v_{5,i}S, \quad \Delta_i := \Delta_7(h_i).$$

Assume

$$a_1 \neq 0, \quad c_1 c_2 \neq 0.$$

Suppose the cubic, quintic, and septic quotient defects are written as

$$a_1 c_1 + a_2 c_2 + E_{12}^{(3)} = 0,$$

$$a_1 A_{5,1}^f + a_2 A_{5,2}^f + E_{12}^{(5)} = 0,$$

and

$$a_1 [A_{7,1}^f] + a_2 [A_{7,2}^f] + \overline{E}_{12}^{(7;1)} = 0 \quad \text{in} \quad \mathfrak{f}/\mathbf{C}A_{5,1}^f.$$

Set

$$\lambda := -\frac{a_2}{a_1}, \quad e := \frac{E_{12}^{(3)}}{a_1}, \quad E := \frac{E_{12}^{(5)}}{a_1} = e_u I + e_v S.$$

Choose any representative  $R \in \mathfrak{f}$  of the quotient class

$$-\frac{1}{a_1} \overline{E}_{12}^{(7;1)},$$

so that

$$A_{7,1}^f = \lambda A_{7,2}^f + R + \eta A_{5,1}^f$$

for some  $\eta \in \mathbf{C}$ . Then the strengthened invariant

$$\widehat{\Psi}(h) = \left( \frac{u_5(h)}{c(h)}, \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)^2} \right)$$

satisfies the exact identities

$$\begin{aligned} \frac{u_{5,1}}{c_1} - \frac{u_{5,2}}{c_2} &= \frac{u_{5,2}e - c_2 e_u}{c_1 c_2}, \\ \frac{v_{5,1}}{c_1} - \frac{v_{5,2}}{c_2} &= \frac{v_{5,2}e - c_2 e_v}{c_1 c_2}, \end{aligned}$$

and

$$\frac{\Delta_1}{c_1^2} - \frac{\Delta_2}{c_2^2} = \frac{-\lambda c_2^2 \det(A_{7,2}^f, E) + c_2^2 \det(R, A_{5,1}^f) + \Delta_2(2\lambda c_2 e - e^2)}{c_1^2 c_2^2}.$$

*Proof.* The cubic defect identity gives

$$c_1 = \lambda c_2 - e.$$

The quintic defect identity gives

$$A_{5,1}^f = \lambda A_{5,2}^f - E,$$

hence

$$u_{5,1} = \lambda u_{5,2} - e_u, \quad v_{5,1} = \lambda v_{5,2} - e_v.$$

Therefore

$$\frac{u_{5,1}}{c_1} - \frac{u_{5,2}}{c_2} = \frac{u_{5,1}c_2 - u_{5,2}c_1}{c_1 c_2} = \frac{(\lambda u_{5,2} - e_u)c_2 - u_{5,2}(\lambda c_2 - e)}{c_1 c_2} = \frac{u_{5,2}e - c_2 e_u}{c_1 c_2},$$

and similarly

$$\frac{v_{5,1}}{c_1} - \frac{v_{5,2}}{c_2} = \frac{v_{5,2}e - c_2 e_v}{c_1 c_2}.$$

For the third coordinate, by the choice of  $R$  we have

$$A_{7,1}^f = \lambda A_{7,2}^f + R + \eta A_{5,1}^f$$

for some  $\eta \in \mathbf{C}$ . Using

$$\Delta_i = \det(A_{7,i}^f, A_{5,i}^f),$$

we obtain

$$\Delta_1 = \det(A_{7,1}^f, A_{5,1}^f) = \lambda \det(A_{7,2}^f, A_{5,1}^f) + \det(R, A_{5,1}^f),$$

since the  $\eta A_{5,1}^f$  term drops out. Substituting

$$A_{5,1}^f = \lambda A_{5,2}^f - E$$

gives

$$\det(A_{7,2}^f, A_{5,1}^f) = \lambda \Delta_2 - \det(A_{7,2}^f, E),$$

hence

$$\Delta_1 - \lambda^2 \Delta_2 = -\lambda \det(A_{7,2}^f, E) + \det(R, A_{5,1}^f).$$

Also,

$$c_1 = \lambda c_2 - e$$

implies

$$\lambda^2 c_2^2 - c_1^2 = 2\lambda c_2 e - e^2.$$

Therefore

$$\frac{\Delta_1}{c_1^2} - \frac{\Delta_2}{c_2^2} = \frac{c_2^2 \Delta_1 - c_1^2 \Delta_2}{c_1^2 c_2^2} = \frac{c_2^2 (\Delta_1 - \lambda^2 \Delta_2) + \Delta_2 (\lambda^2 c_2^2 - c_1^2)}{c_1^2 c_2^2},$$

which is exactly the claimed formula.

Finally, the expression

$$\det(R, A_{5,1}^f)$$

is independent of the chosen representative of the quotient defect, since replacing  $R$  by  $R + \xi A_{5,1}^f$  does not change the determinant.  $\square$

**Corollary 12.114** (Bad two-pair cores are defect-scale close on good  $M \neq 0$  patches). *Let  $J \subseteq \{c \neq 0, M \neq 0\}$  be a compact interval. Assume*

$$|a_1| \geq a_* > 0$$

*for the normalized bad two-pair core under consideration. Then there exists a constant  $C_J > 0$  such that whenever  $h_1, h_2 \in J$ ,*

$$|h_1 - h_2| \leq C_J \left( |E_{12}^{(3)}| + \|E_{12}^{(5)}\| \right).$$

*In particular, on good  $M \neq 0$  patches, bad two-pair cores force the two parameters to be quantitatively close once the cubic/quintic interaction defects are small.*

*Proof.* The second coordinate of  $\widehat{\Psi}$  is

$$x(h) := \frac{v_5(h)}{c(h)}, \quad x'(h) = \frac{M(h)}{c(h)^2}.$$

Since  $J \subseteq \{c \neq 0, M \neq 0\}$ , there exists  $\eta_J > 0$  such that

$$|x(h_1) - x(h_2)| \geq \eta_J |h_1 - h_2| \quad (h_1, h_2 \in J).$$

On the other hand, Theorem 12.113 gives the exact identity

$$x(h_1) - x(h_2) = \frac{v_{5,2}e - c_2 e_v}{c_1 c_2},$$

with

$$e = \frac{E_{12}^{(3)}}{a_1}, \quad e_v = S\text{-coefficient of } \frac{E_{12}^{(5)}}{a_1}.$$

Since  $J$  is compact and  $a_1$  is bounded away from 0, the right-hand side is bounded by

$$C'_J \left( |E_{12}^{(3)}| + \|E_{12}^{(5)}\| \right)$$

for some constant  $C'_J$ . Combining the two estimates yields the claim.  $\square$

**Proposition 12.115** (No sublinear two-pair defects on good patches). *Let*

$$J \subseteq \{c \neq 0, M \neq 0\}$$

*be a compact interval. Then there exists a constant  $C_J > 0$  such that every normalized bad two-pair core with  $h_1, h_2 \in J$  and  $h_1 \neq h_2$  satisfies*

$$|E_{12}^{(3)}| + \|E_{12}^{(5)}\| + |\overline{E}_{12}^{(7;1)}| \geq C_J^{-1} |h_1 - h_2|.$$

*In particular, there is no sequence of nontrivial normalized bad two-pair cores in  $J$  for which*

$$|E_{12}^{(3)}| + \|E_{12}^{(5)}\| + |\overline{E}_{12}^{(7;1)}| = o(|h_1 - h_2|).$$

*Proof.* By Corollary 12.114, there exists a constant  $C'_J > 0$  such that

$$|h_1 - h_2| \leq C'_J \left( |E_{12}^{(3)}| + \|E_{12}^{(5)}\| \right)$$

for every normalized bad two-pair core with  $h_1, h_2 \in J$ . Since

$$|E_{12}^{(3)}| + \|E_{12}^{(5)}\| \leq |E_{12}^{(3)}| + \|E_{12}^{(5)}\| + |\overline{E}_{12}^{(7;1)}|,$$

the same estimate yields

$$|h_1 - h_2| \leq C'_J \left( |E_{12}^{(3)}| + \|E_{12}^{(5)}\| + |\overline{E}_{12}^{(7;1)}| \right).$$

Renaming  $C'_J$  as  $C_J$  proves the first claim. The final statement is immediate: if a sequence satisfied

$$|E_{12}^{(3)}| + \|E_{12}^{(5)}\| + |\overline{E}_{12}^{(7;1)}| = o(|h_1 - h_2|),$$

then the displayed lower bound would force

$$|h_1 - h_2| = o(|h_1 - h_2|),$$

which is impossible for a nontrivial two-point core with  $h_1 \neq h_2$ .  $\square$

**Proposition 12.116** (Conditional source-level quadratic interaction remainder). *Assume there exists a finite-dimensional corrected quotient-germ target  $\mathcal{G}$  and an analytic map*

$$\mathfrak{Q} : \mu \longmapsto \mathfrak{Q}(\mu) \in \mathcal{G}$$

*defined on small finite atomic sources*

$$\mu = \sum_j a_j \delta_{h_j},$$

*such that the corrected one-pair and two-pair objects are given by*

$$\mathcal{F}_1(a, h) := \mathfrak{Q}(a \delta_h), \quad \mathcal{F}_2(a_1, h_1; a_2, h_2) := \mathfrak{Q}(a_1 \delta_{h_1} + a_2 \delta_{h_2}).$$

*Define the raw interaction remainder by*

$$\mathcal{I}(a_1, h_1; a_2, h_2) := \mathcal{F}_2(a_1, h_1; a_2, h_2) - \mathcal{F}_1(a_1, h_1) - \mathcal{F}_1(a_2, h_2).$$

*Then there exists an analytic germ-valued map  $\mathcal{J}$  such that*

$$\boxed{\mathcal{I}(a_1, h_1; a_2, h_2) = a_1 a_2 (h_1 - h_2)^2 \mathcal{J}(a_1, h_1; a_2, h_2)}.$$

*Proof.* The source-level definition of  $\mathcal{F}_2$  is symmetric in the two atoms, so

$$\mathcal{I}(a_1, h_1; a_2, h_2) = \mathcal{I}(a_2, h_2; a_1, h_1).$$

Also, if one amplitude is zero, then the two-pair source reduces to a one-pair source, hence

$$\mathcal{I}(0, h_1; a_2, h_2) = \mathcal{I}(a_1, h_1; 0, h_2) = 0.$$

Thus  $\mathcal{I}$  is divisible by  $a_1 a_2$ , so

$$\mathcal{I}(a_1, h_1; a_2, h_2) = a_1 a_2 G(h_1, h_2)$$

for some analytic  $\mathcal{G}$ -valued map  $G$ .

Now if  $h_1 = h_2 = h$ , then

$$a_1 \delta_h + a_2 \delta_h = (a_1 + a_2) \delta_h,$$

so

$$\mathcal{F}_2(a_1, h; a_2, h) = \mathcal{F}_1(a_1 + a_2, h),$$

and therefore

$$\mathcal{I}(a_1, h; a_2, h) = 0.$$

Equivalently,

$$G(h, h) = 0.$$

Passing to midpoint/separation variables

$$m = \frac{h_1 + h_2}{2}, \quad s = h_2 - h_1,$$

the symmetry of  $G$  means that

$$\tilde{G}(m, s) := G\left(m - \frac{s}{2}, m + \frac{s}{2}\right)$$

is an even analytic function of  $s$ , and the diagonal vanishing gives

$$\tilde{G}(m, 0) = 0.$$

Hence  $\tilde{G}(m, s)$  is divisible by  $s^2$ :

$$\tilde{G}(m, s) = s^2 H(m, s)$$

for analytic  $H$ . Rewriting in  $(h_1, h_2)$  gives the claimed factorization.  $\square$

**Corollary 12.117** (Conditional quadratic defect estimate and two-point exclusion). *Assume, in addition to Proposition 12.116, that the raw cubic, quintic, and septic quotient defects are obtained from the raw interaction remainder by linear coefficient-extraction maps*

$$\Pi_3 : \mathcal{G} \rightarrow \mathbf{C}, \quad \Pi_5 : \mathcal{G} \rightarrow \mathfrak{f}, \quad \bar{\Pi}_7 : \mathcal{G} \rightarrow \mathfrak{f}/\mathbf{C}A_5^{\mathfrak{f}}.$$

*Then on compacta, the raw defects satisfy*

$$\mathcal{E}^{(3)} = a_1 a_2 O(\delta^2), \quad \mathcal{E}^{(5)} = a_1 a_2 O(\delta^2), \quad \bar{\mathcal{E}}^{(7)} = a_1 a_2 O(\delta^2), \quad \delta := |h_1 - h_2|.$$

*If the normalized defects in the present two-pair section are obtained by dividing by  $a_1$ , then on normalized families with  $|a_2|$  bounded one has*

$$E_{12}^{(3)} = O(\delta^2), \quad E_{12}^{(5)} = O(\delta^2), \quad \bar{E}_{12}^{(7;1)} = O(\delta^2).$$

*Consequently, the two-point bad-core branch is excluded on compact good patches*

$$J \in \{c \neq 0, M \neq 0\}.$$

*Proof.* The first claim is immediate from the factorization in Proposition 12.116 and the linearity of the coefficient-extraction maps.

For the normalized defects, divide by  $a_1$  and use boundedness of  $|a_2|$  on normalized families. Then Theorem 12.113 gives

$$|\hat{\Psi}(h_1) - \hat{\Psi}(h_2)| \lesssim |E_{12}^{(3)}| + \|E_{12}^{(5)}\| + |\bar{E}_{12}^{(7;1)}| = O(\delta^2).$$

On the other hand, on every compact good patch

$$J \in \{c \neq 0, M \neq 0\},$$

the second coordinate

$$x(h) = \frac{v_5(h)}{c(h)}$$

is bi-Lipschitz, since

$$x'(h) = \frac{M(h)}{c(h)^2} \neq 0.$$

Hence

$$|\hat{\Psi}(h_1) - \hat{\Psi}(h_2)| \gtrsim \delta.$$

Combining the two inequalities gives

$$\delta \lesssim \delta^2,$$

which is impossible for sufficiently small nonzero  $\delta$ . Thus the two-point branch is excluded on compact good patches.  $\square$

**Remark 12.118** (Work in progress: source-level reduction for the two-point branch). The preceding proposition and corollary isolate the exact structural input needed to finish the two-point branch. At present, the draft does not yet define a source-level corrected quotient-germ map  $\mathfrak{Q}$  on finite atomic sources, so the factorization

$$\mathcal{I} = a_1 a_2 (h_1 - h_2)^2 \mathcal{J}$$

is not yet available formally from the present definitions. If such a source-level reformulation is supplied, then the two-point bad-core branch is finished on compact good patches.

**Remark 12.119** (What remains in the two-pair rewrite). Theorem 12.113 supplies the correct algebraic bridge from the two-pair core to the strengthened invariant  $\widehat{\Psi}$ . In particular, on good  $M \neq 0$  patches, bad two-pair cores are forced into a defect-scale close regime by Corollary 12.114, and Proposition 12.115 shows that the two-point branch is already excluded whenever the total defect is sublinear in the pair separation.

The strongest remaining algebraic target for the two-point branch is now the source-level structural reduction isolated in Proposition 12.116 and Corollary 12.117: if the corrected two-pair object is defined source-level and collapses on coincident atoms, then the normalized defects are automatically  $O(|h_1 - h_2|^2)$ , and the two-point branch is excluded on good patches.

What remains beyond that is the genuine multi-pair interaction bookkeeping through orders 3, 5, 7, together with the local lifted 3/4-point analysis for close-point branches on  $v_5 \neq 0$  patches.

**Remark 12.120** (Minimal-core reformulation). The proposition above shows that the correct projective datum for the downstream multi-pair and minimal-core extraction is the amplitude-invariant strengthened map

$$\widehat{\Psi}(h) = \left( \frac{u_5(h)}{c(h)}, \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)^2} \right),$$

not merely the pair  $(\ell_h, \bar{v}_1(h))$ .

By contrast, the affine curve

$$\Gamma(h) = \left( \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)} \right)$$

remains the correct object for the one-pair secant/tangent geometry. The finite-core endgame should therefore be viewed as a two-stage problem: first extract defect-corrected coincidence information for  $\widehat{\Psi}$ , then convert that information into affine-line incidence constraints for  $\Gamma$ .

**Definition 12.121** (Minimal bad core). A normalized finite core is called bad if its quotient odd germ vanishes identically. A minimal bad core is a bad normalized finite core with minimal number of constituent pairs.

**Theorem 12.122** (Defect-free quintic circuit reduction). *Let*

$$F_{\text{core}}(z) = \sum_{j=1}^k a_j F_{h_j}(z) + \mathcal{I}_{\geq 2}(z)$$

*be a normalized finite core, and assume that the multi-pair interaction germ  $\mathcal{I}_{\geq 2}(z)$  contributes nothing to the cubic and quintic coefficients. Write*

$$A_5^\dagger(h) = u_5(h)I + v_5(h)S, \quad V(h) := (c(h), u_5(h), v_5(h)) \in \mathbf{C}^3,$$

*and on  $\{c \neq 0\}$  define*

$$\Xi(h) := \left( \frac{u_5(h)}{c(h)}, \frac{v_5(h)}{c(h)} \right) \in \mathbf{C}^2.$$

*If*

$$F_{\text{core}}(z) \equiv 0,$$

*then*

$$\sum_{j=1}^k a_j V(h_j) = 0.$$

*Equivalently, with*

$$b_j := a_j c(h_j),$$

*one has*

$$\sum_{j=1}^k b_j = 0, \quad \sum_{j=1}^k b_j \Xi(h_j) = 0.$$

*Thus the points  $\Xi(h_j)$  are affinely dependent.*

*Proof.* Taking the cubic coefficient in the identity

$$F_{\text{core}}(z) \equiv 0$$

and using the hypothesis that  $\mathcal{I}_{\geq 2}$  contributes nothing at cubic order gives

$$\sum_{j=1}^k a_j c(h_j) = 0.$$

Taking the quintic fixed-sector coefficient and using the corresponding defect-free hypothesis at order 5 gives

$$\sum_{j=1}^k a_j A_5^{\dagger}(h_j) = 0.$$

Since

$$A_5^{\dagger}(h_j) = u_5(h_j)I + v_5(h_j)S$$

and  $I, S$  are linearly independent, this is equivalent to

$$\sum_{j=1}^k a_j u_5(h_j) = 0, \quad \sum_{j=1}^k a_j v_5(h_j) = 0.$$

Together these three identities are exactly

$$\sum_{j=1}^k a_j V(h_j) = 0.$$

If  $c(h_j) \neq 0$  for all  $j$ , dividing the last two identities by  $c(h_j)$  after setting  $b_j = a_j c(h_j)$  yields

$$\sum_{j=1}^k b_j = 0, \quad \sum_{j=1}^k b_j \Xi(h_j) = 0,$$

which is the affine dependence statement.  $\square$

**Corollary 12.123** (Defect-free minimal quintic circuits have size at most four). *In the defect-free order- $\leq 5$  model of Theorem 12.122, every minimal bad core has size at most 4. More precisely:*

- (1) if  $k = 2$ , then  $\Xi(h_1) = \Xi(h_2)$ ;
- (2) if  $k = 3$ , then the three points  $\Xi(h_1), \Xi(h_2), \Xi(h_3)$  are collinear;
- (3) if  $k = 4$ , then the four points  $\Xi(h_1), \dots, \Xi(h_4)$  form an affine circuit in  $\mathbf{C}^2$ .

*Proof.* By Theorem 12.122, a minimal defect-free bad core produces a minimal affine dependence among finitely many points of the affine plane  $\mathbf{C}^2$ . Such a minimal affine dependent set has size at most 4. The cases  $k = 2, 3, 4$  are the standard affine configurations in dimension 2: coincidence, collinear triple, or four-point affine circuit.  $\square$

**Definition 12.124** (Lifted  $v_5$ -patch coordinates). On a connected interval  $U \subset \{c \neq 0, v_5 \neq 0\}$ , define

$$x(h) := \frac{v_5(h)}{c(h)}, \quad y(h) := \frac{u_5(h)}{c(h)}, \quad \sigma_v(h) := \frac{\Delta_7(h)}{c(h)v_5(h)}.$$

The associated lifted planar and space curves are

$$\Lambda_v(h) := (x(h), \sigma_v(h)) \in \mathbf{C}^2, \quad Q_v(h) := (y(h), x(h), \sigma_v(h)) \in \mathbf{C}^3.$$

**Proposition 12.125** (Defect-free lifted affine dependence on a  $v_5$ -patch). *Let*

$$F_{\text{core}}(z) = \sum_{j=1}^k a_j F_{h_j}(z) + \mathcal{I}_{\geq 2}(z)$$

*be a normalized finite core, and assume:*

- (1) all parameters  $h_j$  lie in one connected interval  $U \subset \{c \neq 0, v_5 \neq 0\}$ ;

(2) the multi-pair interaction germ  $\mathcal{I}_{\geq 2}(z)$  contributes nothing to the cubic, quintic, or septic coefficients.

Then, after choosing on  $U$  the local septic gauge  $v_7 = 0$ , the vanishing identity

$$F_{\text{core}}(z) \equiv 0$$

implies

$$\sum_{j=1}^k b_j = 0, \quad \sum_{j=1}^k b_j y(h_j) = 0, \quad \sum_{j=1}^k b_j x(h_j) = 0, \quad \sum_{j=1}^k b_j \sigma_v(h_j) = 0,$$

where  $b_j := a_j c(h_j)$ . Equivalently, the points  $Q_v(h_j) \in \mathbf{C}^3$  are affinely dependent.

*Proof.* The cubic and quintic identities are exactly those in Theorem 12.122, so after setting  $b_j = a_j c(h_j)$  we obtain

$$\sum_{j=1}^k b_j = 0, \quad \sum_{j=1}^k b_j y(h_j) = 0, \quad \sum_{j=1}^k b_j x(h_j) = 0.$$

On the interval  $U$  we may choose the local septic gauge  $v_7 = 0$ . Then

$$u_7(h) = \frac{\Delta_7(h)}{v_5(h)} = c(h) \sigma_v(h).$$

Since the septic interaction is assumed absent, taking the septic fixed-sector coefficient of  $F_{\text{core}} \equiv 0$  yields

$$\sum_{j=1}^k a_j u_7(h_j) = 0.$$

Substituting  $u_7(h_j) = c(h_j) \sigma_v(h_j)$  gives

$$\sum_{j=1}^k b_j \sigma_v(h_j) = 0.$$

Hence the points  $Q_v(h_j)$  are affinely dependent. □

**Corollary 12.126** (Geometric meaning of defect-free lifted affine dependence). *Assume the hypotheses of Proposition 12.125. Then:*

- (1) if  $k = 2$ , the two points  $Q_v(h_1)$  and  $Q_v(h_2)$  coincide;
- (2) if  $k = 3$ , the three points  $Q_v(h_1), Q_v(h_2), Q_v(h_3)$  are collinear in  $\mathbf{C}^3$ ;
- (3) if  $k = 4$ , the four points  $Q_v(h_1), \dots, Q_v(h_4)$  are coplanar in  $\mathbf{C}^3$ .

*Proof.* By Proposition 12.125, the points  $Q_v(h_j)$  are affinely dependent. For  $k = 2$ , affine dependence of two points is exactly coincidence. For  $k = 3$ , affine dependence of three points is exactly collinearity. For  $k = 4$ , affine dependence of four points is exactly coplanarity. □

**Corollary 12.127** (Explicit four-point determinant form on a  $v_5$ -patch). *Assume the hypotheses of Proposition 12.125 with  $k = 4$ . Then the coplanarity of the four lifted points is equivalent to*

$$\det \begin{pmatrix} 1 & y(h_1) & x(h_1) & \sigma_v(h_1) \\ 1 & y(h_2) & x(h_2) & \sigma_v(h_2) \\ 1 & y(h_3) & x(h_3) & \sigma_v(h_3) \\ 1 & y(h_4) & x(h_4) & \sigma_v(h_4) \end{pmatrix} = 0.$$



Equivalently,

$$\det \begin{pmatrix} 1 & \frac{u_5(h_1)}{c(h_1)} & \frac{v_5(h_1)}{c(h_1)} & \frac{\Delta_7(h_1)}{c(h_1)v_5(h_1)} \\ 1 & \frac{u_5(h_2)}{c(h_2)} & \frac{v_5(h_2)}{c(h_2)} & \frac{\Delta_7(h_2)}{c(h_2)v_5(h_2)} \\ 1 & \frac{u_5(h_3)}{c(h_3)} & \frac{v_5(h_3)}{c(h_3)} & \frac{\Delta_7(h_3)}{c(h_3)v_5(h_3)} \\ 1 & \frac{u_5(h_4)}{c(h_4)} & \frac{v_5(h_4)}{c(h_4)} & \frac{\Delta_7(h_4)}{c(h_4)v_5(h_4)} \end{pmatrix} = 0.$$

Multiplying row  $j$  by  $v_5(h_j)$  gives the canonical determinant identity

$$\det \begin{pmatrix} v_5(h_1) & u_5(h_1) & \frac{v_5(h_1)^2}{c(h_1)} & \frac{\Delta_7(h_1)}{c(h_1)} \\ v_5(h_2) & u_5(h_2) & \frac{v_5(h_2)^2}{c(h_2)} & \frac{\Delta_7(h_2)}{c(h_2)} \\ v_5(h_3) & u_5(h_3) & \frac{v_5(h_3)^2}{c(h_3)} & \frac{\Delta_7(h_3)}{c(h_3)} \\ v_5(h_4) & u_5(h_4) & \frac{v_5(h_4)^2}{c(h_4)} & \frac{\Delta_7(h_4)}{c(h_4)} \end{pmatrix} = 0.$$

Equivalently, multiplying instead row  $j$  by  $c(h_j)v_5(h_j)$  gives the fully polynomial version

$$\det \begin{pmatrix} c(h_1)v_5(h_1) & u_5(h_1)v_5(h_1) & v_5(h_1)^2 & \Delta_7(h_1) \\ c(h_2)v_5(h_2) & u_5(h_2)v_5(h_2) & v_5(h_2)^2 & \Delta_7(h_2) \\ c(h_3)v_5(h_3) & u_5(h_3)v_5(h_3) & v_5(h_3)^2 & \Delta_7(h_3) \\ c(h_4)v_5(h_4) & u_5(h_4)v_5(h_4) & v_5(h_4)^2 & \Delta_7(h_4) \end{pmatrix} = 0.$$

*Proof.* For  $k = 4$ , Corollary 12.126 shows that the four lifted points

$$Q_v(h_j) = (y(h_j), x(h_j), \sigma_v(h_j))$$

are coplanar. This is equivalent to the vanishing of the affine  $4 \times 4$  determinant with first column 1. The displayed reformulations follow by substituting the definitions of  $x, y, \sigma_v$  and multiplying rows by the indicated nonzero factors.  $\square$

**Remark 12.128** (Parallel  $u_5$ -patch version). There is a completely parallel local story on any connected interval in  $\{c \neq 0, u_5 \neq 0\}$ , obtained by choosing the local septic gauge  $u_7 = 0$  instead of  $v_7 = 0$ . Then one defines

$$\tau_u(h) := -\frac{\Delta_7(h)}{c(h)u_5(h)}, \quad Q_u(h) := \left( \frac{u_5(h)}{c(h)}, \frac{v_5(h)}{c(h)}, \tau_u(h) \right),$$

and the same defect-free affine-dependence conclusions hold with  $Q_v$  replaced by  $Q_u$ . Since the present one-pair geometry already uses

$$x(h) = \frac{v_5(h)}{c(h)}$$

as the distinguished ECT coordinate, the  $v_5 \neq 0$  patch is the natural first setting and will be used below.

**Proposition 12.129** (Linear independence of  $1, u_5/c, v_5/c$ ). *Let  $J$  be a connected open interval on which*

$$c \neq 0, \quad M \neq 0.$$

*Then the three functions*

$$1, \quad \frac{u_5}{c}, \quad \frac{v_5}{c}$$

*are linearly independent on  $J$ .*

*Proof.* Assume there exist constants  $A, B, C \in \mathbf{C}$ , not all zero, such that

$$A + B \frac{u_5}{c} + C \frac{v_5}{c} \equiv 0 \quad \text{on } J.$$

Multiplying by  $c$  gives

$$A c + B u_5 + C v_5 \equiv 0.$$

If  $B = 0$ , then

$$A c + C v_5 \equiv 0,$$

so  $v_5/c$  is constant on  $J$ . But

$$\left(\frac{v_5}{c}\right)' = \frac{M}{c^2},$$

and  $M \neq 0$  on  $J$ . This is impossible.

Hence  $B \neq 0$ , and after renormalization we may write

$$u_5 = \alpha v_5 + \beta c$$

for constants  $\alpha, \beta$ .

Now isolate the graded slice corresponding to

$$(K_1, \eta_2, q^{(5)}).$$

By Lemmas 12.16 and 12.18, the tagged monomial

$$\kappa(h) \eta_2(h) q^{(5)}(h)$$

occurs with nonzero coefficient in  $\text{tr } J_5^{\text{pair}}(h)$ . Since

$$A_5^f(h) = u_5(h)I + v_5(h)S$$

and  $S$  is traceless, this means exactly that the same tagged monomial occurs in the  $I$ -coefficient  $u_5(h)$ .

It cannot occur in  $c(h)$ , since  $c$  is cubic and carries no quintic  $q^{(5)}$ -slice. It also cannot occur in  $v_5(h)$ . Indeed, on the tagged slice, the only surviving quintic  $K_1$ -transport channels are those already isolated in Lemma 12.18. For the channel

$$X_0 J X_4 - X_4 J X_0,$$

Lemma 12.17 shows that the relevant  $(\eta_2, q^{(5)})$ -slice is diagonal, hence contributes only to the  $I$ -component. For the commutator channel

$$X_3 J X_1 - X_1 J X_3,$$

Proposition 12.40 gives

$$v_5^{(\text{comm})} = \gamma u - \alpha w.$$

On the tagged slice the  $\eta_2$ -input in  $X_1$  is purely off-diagonal, so  $u = w = 0$ , while the  $q^{(5)}$ -input in  $X_3$  is diagonal. Hence

$$v_5^{(\text{comm})} = 0$$

on that tagged slice.

Thus the monomial  $\kappa \eta_2 q^{(5)}$  occurs in  $u_5$  but in neither  $v_5$  nor  $c$ , contradicting

$$u_5 = \alpha v_5 + \beta c.$$

This proves the claim. □

**Proposition 12.130** (Non-affinity of the lifted septic scalar). *Let  $J$  be a connected open interval contained in the quintic-generic region and in*

$$\{h : c(h) \neq 0, v_5(h) \neq 0\}.$$

*Then there do not exist constants  $A, B \in \mathbf{C}$  such that*

$$\sigma_v(h) \equiv A x(h) + B \quad \text{on } J.$$

*Equivalently,*

$$\Delta_7(h) \not\equiv A v_5(h)^2 + B c(h) v_5(h) \quad \text{on } J.$$

*Proof.* If

$$\sigma_v(h) \equiv A x(h) + B,$$

then multiplying by  $c(h)v_5(h)$  gives

$$\Delta_7(h) \equiv A v_5(h)^2 + B c(h)v_5(h).$$

The right-hand side is built only from the cubic and quintic layers and therefore has no direct septic  $q^{(7)}$ -slice. But by Proposition 12.48, the direct  $q^{(7)}$ -slice of  $\Delta_7$  is generically nonzero on  $J$ . This is a contradiction.  $\square$

**Definition 12.131** (Lifted derivative numerator and curvature numerator). On  $\{c \neq 0, v_5 \neq 0\}$ , define

$$P_v(h) := c(h)v_5(h)\Delta_7'(h) - \Delta_7(h)(c(h)v_5(h))'.$$

Then

$$\sigma_v'(h) = \frac{P_v(h)}{c(h)^2 v_5(h)^2}.$$

Define also the affine curvature numerator of the lifted planar curve

$$\Lambda_v(h) = (x(h), \sigma_v(h))$$

by

$$K_v(h) := x'(h)\sigma_v''(h) - \sigma_v'(h)x''(h).$$

**Proposition 12.132** (Curvature bridge for the lifted  $v_5$ -patch curve). On  $\{c \neq 0, M \neq 0, v_5 \neq 0\}$ ,

$$\frac{d\sigma_v}{dx} = \frac{P_v}{M v_5^2},$$

and

$$K_v(h) = \frac{M(h)^2}{c(h)^4} \frac{d}{dh} \left( \frac{P_v(h)}{M(h) v_5(h)^2} \right).$$

In particular, if  $K_v \equiv 0$  on a connected open interval  $J \subset \{c \neq 0, M \neq 0, v_5 \neq 0\}$ , then  $\sigma_v$  is affine in  $x$  on  $J$ , contradicting Proposition 12.130. Hence

$$K_v \not\equiv 0$$

on every such interval.

*Proof.* Since

$$x = \frac{v_5}{c}, \quad x' = \frac{M}{c^2}, \quad \sigma_v' = \frac{P_v}{c^2 v_5^2},$$

we obtain

$$\frac{d\sigma_v}{dx} = \frac{\sigma_v'}{x'} = \frac{P_v}{M v_5^2}.$$

Differentiating with respect to  $h$  and multiplying by

$$x' = \frac{M}{c^2}$$

gives

$$K_v = x'^2 \frac{d}{dx} \left( \frac{d\sigma_v}{dx} \right) = \frac{M^2}{c^4} \frac{d}{dh} \left( \frac{P_v}{M v_5^2} \right).$$

If  $K_v \equiv 0$  on a connected interval with  $M \neq 0$ , then

$$\frac{P_v}{M v_5^2}$$

is constant there, so  $d\sigma_v/dx$  is constant and  $\sigma_v$  is affine in  $x$ . This contradicts Proposition 12.130.  $\square$

**Corollary 12.133** (Local defect-free three-point exclusion on lifted ECT subintervals). *Let  $J \subset \{c \neq 0, M \neq 0, v_5 \neq 0\}$  be a connected interval on which  $K_v$  has fixed nonzero sign. Then the triple*

$$1, \quad x, \quad \sigma_v$$

*is an ECT system on  $J$ . In particular, no affine line meets the lifted planar curve*

$$\Lambda_v(h) = (x(h), \sigma_v(h))$$

*in three points of  $J$ . Hence the defect-free three-point branch is excluded on such a subinterval.*

*Proof.* The Wronskian of  $1, x, \sigma_v$  is exactly  $K_v$ . If  $K_v$  has fixed nonzero sign on  $J$ , the ECT criterion applies.  $\square$

**Corollary 12.134** (Lifted order-7 Wronskian is not identically zero). *Let  $J$  be a connected open interval contained in the quintic-generic region and in*

$$\{h : c(h) \neq 0, M(h) \neq 0, v_5(h) \neq 0\}.$$

*Then*

$$T_v(h) := W_h\left(1, \frac{u_5(h)}{c(h)}, \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)v_5(h)}\right)$$

*is not identically zero on  $J$ .*

*Proof.* Write

$$y := \frac{u_5}{c}, \quad x := \frac{v_5}{c}, \quad \sigma_v := \frac{\Delta_7}{c v_5}.$$

If  $T_v \equiv 0$  on  $J$ , then by analyticity the four functions

$$1, y, x, \sigma_v$$

are locally linearly dependent on  $J$ , hence linearly dependent on all of the connected interval  $J$ . By Proposition 12.129, the first three are linearly independent, so there exist constants  $A, B, C$  such that

$$\sigma_v = A + By + Cx.$$

Multiplying by  $c v_5$  gives

$$\Delta_7 = A c v_5 + B u_5 v_5 + C v_5^2.$$

The right-hand side is built entirely from the cubic/quintic layers and therefore carries no direct  $q^{(7)}$ -slice. This contradicts Proposition 12.48, which gives

$$\Delta_7^{(q^{(7)})} \neq 0$$

on  $J$ .  $\square$

**Proposition 12.135** (Exact  $x$ -coordinate identities for the lifted 4-point invariants). *On a good  $v_5$ -patch, write*

$$x = \frac{v_5}{c}, \quad Y = \frac{u_5}{c}, \quad S = \sigma_v = \frac{\Delta_7}{c v_5}.$$

*Then:*

(1) *the lifted planar curvature numerator satisfies*

$$K_v = (x')^3 S''(x);$$

(2) *the lifted 4-point Wronskian satisfies*

$$T_v = (x')^6 W_x(1, Y, x, S) = (x')^6 (S'' Y''' - S''' Y'');$$

(3) *equivalently,*

$$T_v = (x')^6 S''^2 \frac{d}{dx} \left( \frac{Y''}{S''} \right) = K_v^2 \frac{d}{dx} \left( \frac{Y''}{S''} \right).$$

*In particular,  $K_v$  and  $S''$  have the same zero set and the same vanishing order on every good  $v_5$ -patch.*

*Proof.* By definition,

$$K_v = x' \sigma_v'' - \sigma_v' x''.$$

Since  $S = \sigma_v$  as a function of the local coordinate  $x$ , we have

$$\sigma_v' = S' x', \quad \sigma_v'' = S'' (x')^2 + S' x'',$$

hence

$$K_v = x' (S'' (x')^2 + S' x'') - S' x' x'' = (x')^3 S''.$$

Next,

$$T_v = W_h(1, Y, x, S).$$

Under reparametrization  $h \mapsto x(h)$ , a  $4 \times 4$  Wronskian gains the factor  $(x')^{0+1+2+3} = (x')^6$ , so

$$T_v = (x')^6 W_x(1, Y, x, S).$$

Now

$$W_x(1, Y, x, S) = \det \begin{pmatrix} 1 & Y & x & S \\ 0 & Y' & 1 & S' \\ 0 & Y'' & 0 & S'' \\ 0 & Y''' & 0 & S''' \end{pmatrix} = S'' Y''' - S''' Y'',$$

which proves the second identity.

Finally,

$$S'' Y''' - S''' Y'' = S''^2 \frac{d}{dx} \left( \frac{Y''}{S''} \right),$$

and using  $K_v = (x')^3 S''$  gives

$$(x')^6 S''^2 = K_v^2.$$

This yields the third identity.  $\square$

**Corollary 12.136** (Local defect-free four-point exclusion on lifted ECT subintervals). *Let  $J \subset \{c \neq 0, M \neq 0, v_5 \neq 0\}$  be a connected interval on which  $T_v$  has fixed nonzero sign. Then the quadruple*

$$1, \quad \frac{u_5}{c}, \quad \frac{v_5}{c}, \quad \frac{\Delta_7}{c v_5}$$

*is an ECT system on  $J$ . Equivalently, no affine plane meets the lifted space curve*

$$Q_v(h) = \left( \frac{u_5(h)}{c(h)}, \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h) v_5(h)} \right)$$

*in four points of  $J$ . Hence the defect-free four-point branch is excluded on such a subinterval.*

*Proof.* The Wronskian of the displayed quadruple is exactly  $T_v$ . If  $T_v$  has fixed nonzero sign on  $J$ , the ECT criterion applies.  $\square$

**Proposition 12.137** (Shrinking exact lifted 3-point circuits concentrate into  $K_v = 0$ ). *Let  $P \subseteq \{c \neq 0, v_5 \neq 0, M \neq 0\}$  be compact, and use*

$$x = \frac{v_5}{c}, \quad S = \sigma_v = \frac{\Delta_7}{c v_5}.$$

*Suppose there is a sequence of distinct triples*

$$x_{1,n}, x_{2,n}, x_{3,n} \in x(P), \quad x_{1,n}, x_{2,n}, x_{3,n} \rightarrow x_*,$$

*such that*

$$\det \begin{pmatrix} 1 & x_{1,n} & S(x_{1,n}) \\ 1 & x_{2,n} & S(x_{2,n}) \\ 1 & x_{3,n} & S(x_{3,n}) \end{pmatrix} = 0 \quad \text{for every } n.$$

*Then*

$$K_v(x_*) = 0.$$

*Proof.* Set

$$\Delta_3(x_1, x_2, x_3) := \det \begin{pmatrix} 1 & x_1 & S(x_1) \\ 1 & x_2 & S(x_2) \\ 1 & x_3 & S(x_3) \end{pmatrix}.$$

By the standard divided-difference factorization,

$$\Delta_3(x_1, x_2, x_3) = V_3(x_1, x_2, x_3) S[x_1, x_2, x_3],$$

where

$$V_3(x_1, x_2, x_3) := (x_2 - x_1)(x_3 - x_1)(x_3 - x_2).$$

Since the nodes are distinct,  $V_3(x_{1,n}, x_{2,n}, x_{3,n}) \neq 0$ , so

$$S[x_{1,n}, x_{2,n}, x_{3,n}] = 0.$$

As  $x_{1,n}, x_{2,n}, x_{3,n} \rightarrow x_*$ , the second divided difference converges to

$$\frac{1}{2} S''(x_*).$$

Hence  $S''(x_*) = 0$ . Since on the  $x$ -patch the lifted curvature numerator  $K_v$  is  $S''$  up to a nowhere-zero factor, it follows that  $K_v(x_*) = 0$ .  $\square$

**Corollary 12.138** (Uniform small-diameter exclusion away from  $K_v = 0$ ). *Let  $P_0 \subseteq \{c \neq 0, v_5 \neq 0, M \neq 0\}$  be compact and assume*

$$P_0 \cap \{K_v = 0\} = \emptyset.$$

*Then there exists  $\rho_3 > 0$  such that no exact lifted 3-point circuit contained in  $P_0$  has  $x$ -diameter  $< \rho_3$ .*

*Proof.* Otherwise there would exist a sequence of exact lifted 3-point circuits in  $P_0$  with diameters tending to 0. By compactness, after passing to a subsequence the three nodes would converge to some  $x_* \in x(P_0)$ . This contradicts Proposition 12.137.  $\square$

**Proposition 12.139** (Shrinking exact lifted 4-point circuits concentrate into  $T_v = 0$ ). *Let  $P \subseteq \{c \neq 0, v_5 \neq 0, M \neq 0\}$  be compact, and use*

$$x = \frac{v_5}{c}, \quad Y = \frac{u_5}{c}, \quad S = \sigma_v = \frac{\Delta_7}{c v_5}.$$

*Suppose there is a sequence of distinct quadruples*

$$x_{1,n}, x_{2,n}, x_{3,n}, x_{4,n} \in x(P), \quad x_{1,n}, x_{2,n}, x_{3,n}, x_{4,n} \rightarrow x_*,$$

*such that*

$$\det \begin{pmatrix} 1 & Y(x_{1,n}) & x_{1,n} & S(x_{1,n}) \\ 1 & Y(x_{2,n}) & x_{2,n} & S(x_{2,n}) \\ 1 & Y(x_{3,n}) & x_{3,n} & S(x_{3,n}) \\ 1 & Y(x_{4,n}) & x_{4,n} & S(x_{4,n}) \end{pmatrix} = 0 \quad \text{for every } n.$$

*Then*

$$T_v(x_*) = 0.$$

*Proof.* Set

$$\Delta_4(x_1, x_2, x_3, x_4) := \det \begin{pmatrix} 1 & Y(x_1) & x_1 & S(x_1) \\ 1 & Y(x_2) & x_2 & S(x_2) \\ 1 & Y(x_3) & x_3 & S(x_3) \\ 1 & Y(x_4) & x_4 & S(x_4) \end{pmatrix}.$$

Since  $\Delta_4$  is alternating in the four nodes, it is divisible by the Vandermonde

$$V_4(x_1, x_2, x_3, x_4) := \prod_{1 \leq i < j \leq 4} (x_j - x_i).$$

Write

$$\Delta_4(x_1, x_2, x_3, x_4) = V_4(x_1, x_2, x_3, x_4) \mathcal{D}_4(x_1, x_2, x_3, x_4),$$

where  $\mathcal{D}_4$  is symmetric and analytic. By the standard Wronskian/divided-difference coalescence formula,

$$\mathcal{D}_4(x_1, x_2, x_3, x_4) \longrightarrow \frac{1}{1!2!3!} W_x(1, Y, x, S)(x_*) = \frac{1}{12} T_v(x_*),$$

as all four nodes tend to  $x_*$ . Since  $V_4 \neq 0$  for distinct nodes and  $\Delta_4 = 0$  along the sequence, one has  $\mathcal{D}_4 = 0$  along the sequence, so the limit forces  $T_v(x_*) = 0$ .  $\square$

**Corollary 12.140** (Uniform small-diameter exclusion away from  $T_v = 0$ ). *Let  $P_0 \subseteq \{c \neq 0, v_5 \neq 0, M \neq 0\}$  be compact and assume*

$$P_0 \cap \{T_v = 0\} = \emptyset.$$

*Then there exists  $\rho_4 > 0$  such that no exact lifted 4-point circuit contained in  $P_0$  has  $x$ -diameter  $< \rho_4$ .*

*Proof.* Otherwise there would exist a sequence of exact lifted 4-point circuits in  $P_0$  with diameters tending to 0. By compactness, after passing to a subsequence the four nodes would converge to some  $x_* \in x(P_0)$ . This contradicts Proposition 12.139.  $\square$

**Definition 12.141** (Repeated-node tangent-contact determinants on a  $v_5$ -patch). On a  $v_5 \neq 0$  patch with  $x = v_5/c$ ,  $Y = u_5/c$ , and  $S = \Delta_7/(c v_5)$ , define

$$H_3(a, u) := \det \begin{pmatrix} 1 & a & S(a) \\ 0 & 1 & S'(a) \\ 1 & u & S(u) \end{pmatrix} = S(u) - S(a) - S'(a)(u - a),$$

and

$$H_4(a, u, v) := \det \begin{pmatrix} 1 & Y(a) & a & S(a) \\ 0 & Y'(a) & 1 & S'(a) \\ 1 & Y(u) & u & S(u) \\ 1 & Y(v) & v & S(v) \end{pmatrix}.$$

Thus  $H_3(a, u) = 0$  means that  $(u, S(u))$  lies on the tangent line to  $\Lambda_v$  at  $a$ , while  $H_4(a, u, v) = 0$  means that  $Q_v(u)$  and  $Q_v(v)$  lie in the tangent plane to  $Q_v$  at  $a$ .

**Proposition 12.142** (Repeated-node limits of exact lifted determinants). *For the exact lifted determinants*

$$\Delta_3(a, b, u) := \det \begin{pmatrix} 1 & a & S(a) \\ 1 & b & S(b) \\ 1 & u & S(u) \end{pmatrix},$$

and

$$\Delta_4(a, b, u, v) := \det \begin{pmatrix} 1 & Y(a) & a & S(a) \\ 1 & Y(b) & b & S(b) \\ 1 & Y(u) & u & S(u) \\ 1 & Y(v) & v & S(v) \end{pmatrix},$$

one has

$$\frac{\Delta_3(a, b, u)}{b - a} \longrightarrow H_3(a, u) \quad (b \rightarrow a),$$

and

$$\frac{\Delta_4(a, b, u, v)}{b - a} \longrightarrow H_4(a, u, v) \quad (b \rightarrow a).$$

*Proof.* Subtract the first row from the second row in each determinant and divide by  $b - a$ . The resulting second row is the first difference quotient of the corresponding row entries, and the limit  $b \rightarrow a$  gives the derivative row.  $\square$

**Theorem 12.143** (Local classification of the lifted 3-point branch near a finite-order  $K_v$ -zero). *Let  $x = 0$  be an isolated finite-order zero of  $K_v$  on a good  $v_5$ -patch. After affine normalization of  $\Lambda_v = (x, S)$ , write*

$$S(x) = x^m + O(x^{m+1}), \quad m = r + 2 \geq 3,$$

where  $r = \text{ord}_0 K_v$ . Then the repeated-node tangent-contact equation

$$H_3(a, u) = 0$$

with  $a < 0 < u$  has the following local classification:

- (1) if  $r$  is even (equivalently  $m$  is even), there is no opposite-side tangent-contact branch near  $(a, u) = (0, 0)$ ;
- (2) if  $r$  is odd (equivalently  $m$  is odd), there exists a unique real-analytic opposite-side branch

$$u = \Upsilon(a), \quad a < 0, \quad |a| \ll 1,$$

with asymptotic expansion

$$\Upsilon(a) = \lambda_m(-a) + O(a^2),$$

where  $\lambda_m > 1$  is the unique positive root of

$$P_m(\lambda) := \lambda^m - m\lambda - (m-1).$$

*Proof.* Set  $a = -\alpha$  with  $\alpha > 0$ , and write  $u = \lambda\alpha$  with  $\lambda > 0$ . For the model germ  $S(x) = x^m$ , the tangent-contact equation is

$$H_3(-\alpha, \lambda\alpha) = \alpha^m \begin{cases} \lambda^m + m\lambda + (m-1), & m \text{ even}, \\ \lambda^m - m\lambda - (m-1), & m \text{ odd}. \end{cases}$$

If  $m$  is even, the displayed factor is strictly positive for every  $\lambda > 0$ , so no opposite-side branch exists.

If  $m$  is odd, define

$$P_m(\lambda) := \lambda^m - m\lambda - (m-1).$$

Then

$$P_m(1) = -2(m-1) < 0, \quad P_m(\lambda) \rightarrow +\infty \text{ as } \lambda \rightarrow +\infty,$$

and

$$P'_m(\lambda) = m(\lambda^{m-1} - 1)$$

is negative on  $(0, 1)$  and positive on  $(1, \infty)$ . Therefore  $P_m$  has a unique positive root  $\lambda_m > 1$ .

For the perturbed germ  $S(x) = x^m + O(x^{m+1})$ , divide  $H_3(-\alpha, \lambda\alpha)$  by  $\alpha^m$ . The resulting equation is

$$P_m(\lambda) + O(\alpha) = 0.$$

If  $m$  is even, positivity of the model factor persists for  $\alpha$  small, so there is still no branch. If  $m$  is odd, the root  $\lambda_m$  is simple, so the implicit function theorem yields a unique analytic branch

$$\lambda = \lambda(\alpha) = \lambda_m + O(\alpha),$$

equivalently

$$u = \lambda_m(-a) + O(a^2).$$

This proves the classification. □

**Remark 12.144** (Exact nonmixed jet reduction for the lifted 4-point branch). At a nonmixed lifted point

$$K_v(0) \neq 0, \quad T_v(0) = 0,$$

Proposition 12.135 gives

$$T_v = (x')^6 S''^2 \frac{d}{dx} \left( \frac{Y''}{S''} \right).$$

Since  $K_v(0) \neq 0$ , one has  $S''(0) \neq 0$ , so after analytic reparametrization and affine target normalization preserving coplanarity one may write the lifted germ as

$$Q_v(\xi) = (\tilde{Y}(\xi), \xi, \xi^2), \quad \tilde{Y}(0) = \tilde{Y}'(0) = \tilde{Y}''(0) = 0.$$

In these normalized coordinates,

$$W_\xi(1, \tilde{Y}, \xi, \xi^2) = 2\tilde{Y}'''.$$



Thus the nonmixed lifted 4-point problem is exactly the jet problem of excluding odd-order zeros of  $\tilde{Y}'''$ .

In particular, the low-order dangerous nonmixed cases are exactly:

$$\begin{aligned} \text{ord}_0 T_v = 1 &\iff \tilde{Y}'''(0) = 0, \tilde{Y}^{(4)}(0) \neq 0, \\ \text{ord}_0 T_v = 3 &\iff \tilde{Y}^{(3)}(0) = \tilde{Y}^{(4)}(0) = \tilde{Y}^{(5)}(0) = 0, \tilde{Y}^{(6)}(0) \neq 0, \\ \text{ord}_0 T_v = 5 &\iff \tilde{Y}^{(3)}(0) = \dots = \tilde{Y}^{(7)}(0) = 0, \tilde{Y}^{(8)}(0) \neq 0. \end{aligned}$$

So the nonmixed 4-point endgame has been reduced to the finite jet problem: rule out leading terms of  $\tilde{Y}$  in degrees 4, 6, 8.

This is strictly sharper than the earlier qualitative statement that nonmixed odd-order  $T_v$ -zeros can carry preexisting close branches.

**Remark 12.145** (Exact mixed jet reduction and low-order dangerous list). At a mixed lifted point

$$K_v(0) = 0, \quad T_v(0) = 0,$$

set

$$r := \text{ord}_0 K_v, \quad s := \text{ord}_0 T_v.$$

By Proposition 12.135,

$$K_v = (x')^3 S'',$$

so on a good  $v_5$ -patch one has

$$S''(x) = \kappa x^r + O(x^{r+1}), \quad \kappa \neq 0.$$

After removing affine terms and integrating twice,

$$S(x) = \frac{\kappa}{(r+1)(r+2)} x^{r+2} + O(x^{r+3}).$$

Thus the mixed exponent  $m$  of the monomial model

$$Q_v(x) \sim (x^n, x, x^m)$$

is exactly

$$m = r + 2.$$

Now let  $n$  be the first degree of  $Y(x)$  not absorbed by  $\text{span}\{1, x, S\}$ , so that

$$Y(x) = \alpha x^n + O(x^{n+1}), \quad \alpha \neq 0, \quad n > r + 2.$$

Using

$$T_v = (x')^6 (S''Y''' - S'''Y''),$$

one finds

$$T_v = (x')^6 \kappa \alpha n(n-1)(n-r-2)x^{n+r-3} + O(x^{n+r-2}),$$

and since  $n > r + 2$ , the leading coefficient is nonzero. Therefore

$$s = n + r - 3, \quad n = s - r + 3.$$

So the mixed exponents are exactly

$$m = r + 2, \quad n = s - r + 3, \quad s \geq 2r.$$

The dangerous low-order mixed models are therefore finite and explicit:

$$(m, n) = (3, 4), (3, 6), (4, 6),$$

equivalently

$$(r, s) = (1, 2), (1, 4), (2, 5).$$

Thus the mixed lifted 4-point problem has also been reduced to a finite jet obstruction list: rule out those three order pairs in the actual geometry.

**Proposition 12.146** (Realization lemma for lifted germs). *Let  $\tilde{Y}(\xi)$  and  $\tilde{S}(\xi)$  be arbitrary analytic germs at  $\xi = 0$ . Define*

$$c(\xi) \equiv 1, \quad v_5(\xi) := 1 + \xi, \quad u_5(\xi) := \tilde{Y}(\xi), \quad \Delta_7(\xi) := (1 + \xi)\tilde{S}(\xi).$$

*Then on a sufficiently small neighborhood of 0,*

$$c \neq 0, \quad v_5 \neq 0, \quad M = c v_5' - c' v_5 = 1 \neq 0,$$

*so this is a good  $v_5$ -patch. The normalized lifted coordinates are*

$$x = \frac{v_5}{c} = 1 + \xi, \quad Y = \frac{u_5}{c} = \tilde{Y}(\xi), \quad S = \frac{\Delta_7}{c v_5} = \tilde{S}(\xi).$$

*After the affine shift  $x \mapsto x - 1$ , the lifted germ becomes*

$$Q_v(\xi) = (\tilde{Y}(\xi), \xi, \tilde{S}(\xi)).$$

*Thus every analytic pair of germs  $(\tilde{Y}, \tilde{S})$  occurs, after affine normalization, as the lifted pair  $(Y, S)$  on some good  $v_5$ -patch.*

*Proof.* The displayed formulas are immediate from the definitions. The hypotheses  $c \neq 0$ ,  $v_5 \neq 0$ , and  $M = 1 \neq 0$  hold on a sufficiently small neighborhood of 0, and the formulas for  $x, Y, S$  are direct.  $\square$

**Corollary 12.147** (No new jet obstruction is available at the current lifted level). *No identity involving only*

$$c, \quad v_5, \quad u_5, \quad \Delta_7$$

*and their derivatives can exclude the low-order dangerous lifted 4-point patterns listed above.*

*In particular, the dangerous nonmixed models*

$$Q_v(\xi) = (\xi^n, \xi, \xi^2), \quad n = 4, 6, 8,$$

*and the dangerous mixed models*

$$Q_v(\xi) = (\xi^n, \xi, \xi^m), \quad (m, n) = (3, 4), (3, 6), (4, 6),$$

*are all realizable at the present lifted level.*

*Consequently, any RH-specific obstruction to these low-order 4-point patterns must use information not encoded in the current lifted variables*

$$(c, v_5, u_5, \Delta_7)$$

*alone. It must come from deeper source-level corrected quotient-germ structure, matrix-level identities behind  $(u_5, \Delta_7)$ , or an additional symmetry not yet visible in the present lifted formulation.*

*Proof.* Apply Proposition 12.146 with

$$(\tilde{Y}, \tilde{S}) = (\xi^n, \xi^2)$$

for the nonmixed cases, and with

$$(\tilde{Y}, \tilde{S}) = (\xi^n, \xi^m)$$

for the mixed cases.  $\square$

**Remark 12.148** (Work in progress: a dream parity identity). After nonmixed normalization

$$Q_v(\xi) = (\tilde{Y}(\xi), \xi, \xi^2),$$

a parity identity of the form

$$\tilde{Y}(-\xi) = -\tilde{Y}(\xi)$$

would immediately eliminate all currently dangerous nonmixed low-order jet patterns, since it would forbid leading terms of degrees 4, 6, 8, and would also remove the dangerous mixed sector with even  $n$ . The present lifted formalism does not supply such an identity; if it exists, it must come from a deeper RH-specific symmetry not yet visible in the current quotient-germ package.

**Remark 12.149** (What the four-point branch now reduces to). The lifted 4-point geometry is now sharply reduced.

On the nonmixed side, the exact invariant identity

$$T_v = (x')^6 S''^2 \frac{d}{dx} \left( \frac{Y''}{S''} \right)$$

shows that the low-order dangerous cases are precisely the odd-order zeros of  $\tilde{Y}'''$  of orders 1, 3, 5 after normalizing the lower triple to

$$(1, \xi, \xi^2).$$

On the mixed side, the dangerous low-order cases are precisely the three order pairs

$$(r, s) = (1, 2), (1, 4), (2, 5),$$

equivalently

$$(m, n) = (3, 4), (3, 6), (4, 6).$$

By Corollary 12.147, these patterns cannot be excluded by any identity involving only the present lifted variables

$$(c, v_5, u_5, \Delta_7)$$

and their derivatives.

Accordingly, the next serious four-point target is no longer another lifted jet computation. Any closure of the four-point branch must come from deeper source-level corrected quotient-germ structure, matrix-level identities behind  $(u_5, \Delta_7)$ , or a hidden RH-specific symmetry not yet encoded in the present lifted formulation.

**Remark 12.150** (Work in progress: what remains in the lifted order-7 geometry). The lifted  $v_5$ -patch geometry now splits more sharply than a blanket zero-capture problem.

First, Propositions 12.137 and 12.139 show that shrinking exact lifted three-point and four-point circuits can accumulate only at

$$K_v = 0 \quad \text{and} \quad T_v = 0,$$

respectively, and give uniform small-diameter exclusion away from those zero sets.

Second, Proposition 12.142 identifies the one-pair collapse limits as tangent-line and tangent-plane contact determinants for the lifted curves  $\Lambda_v$  and  $Q_v$ .

Third, the nonmixed and mixed four-point local pictures are genuinely different. On the nonmixed side, generic odd-order  $T_v$ -zeros can carry a preexisting close branch; on the mixed side  $K_v = T_v = 0$ , the order bookkeeping is

$$m = r + 2, \quad n = s - r + 3, \quad s \geq 2r,$$

and there is no generic parity obstruction beyond this order inequality.

Thus the missing global ingredient is no longer a single blanket first-contact lemma. It is a combination of:

- source-level control of the multi-pair interaction defects

$$E_{\geq 2}^{(3)}, \quad E_{\geq 2}^{(5)}, \quad \overline{E}_{\geq 2}^{(7)};$$

- extra algebraic compatibility conditions for true bad four-point cores coming from a deeper source-level corrected quotient-germ construction, not from the present lifted coordinates alone;
- and, on the mixed side, RH-specific restrictions excluding the three explicit low-order order pairs

$$(r, s) = (1, 2), (1, 4), (2, 5).$$

**Remark 12.151** (Work in progress: stratified lifted local picture). The lifted local obstruction now splits into three regimes.

First, exact shrinking lifted 3-point circuits concentrate into

$$K_v = 0,$$

and near a finite-order isolated  $K_v$ -zero the repeated-node tangent-contact branch is completely classified: even order gives no opposite-side branch, while odd order gives exactly one opposite-side branch.

Second, exact shrinking lifted 4-point circuits concentrate into

$$T_v = 0.$$

On the nonmixed side  $K_v \neq 0$ , finite-order  $T_v$ -zeros control local birth/crossing by parity, but generic models show that a preexisting close branch can still exist near certain odd-order nonmixed zeros. Thus the nonmixed 4-point side is not governed by a blanket local-exclusion theorem.

Third, the mixed case

$$K_v = T_v = 0$$

is genuinely different. After affine normalization with

$$x = \frac{v_5}{c}$$

as local coordinate, the lifted space curve is led to a monomial model

$$Q_v(x) \sim (x^n, x, x^m),$$

with order bookkeeping

$$m = r + 2, \quad n = s - r + 3, \quad s \geq 2r.$$

No generic parity obstruction is available at this level, so the mixed locus remains the genuinely new codimension-two local obstruction.

Thus the remaining lifted local issues are: the continuation / non-recurrence of the unique odd-order 3-point branch, the continuation / non-recurrence of nonmixed 4-point branches, possible infinite-order flatness, and the mixed codimension-two locus together with the extra bad-core compatibility conditions needed there.

**Remark 12.152** (Current geometric bottleneck). The remaining geometric bottleneck is now split into two layers.

On the one-pair geometric layer, the obstruction is no longer bulk  $W = 0$  geometry, and it is no longer local adjacent pair-pair birth across a generic turning boundary. On the  $W = 0$ ,  $M \neq 0$  side, Proposition 12.59 excludes open-interval flatness and Theorem 12.102 shows that every finite-order isolated  $W = 0$  boundary is locally sterile for pair-pair birth. On the turning-boundary side, Theorem 12.92 removes actual adjacent pair-pair lines unconditionally.

At the lifted order-7 level, the new local determinants

$$K_v \quad \text{and} \quad T_v$$

govern shrinking exact lifted 3- and 4-point circuits on  $v_5 \neq 0$  patches: exact shrinking 3-point circuits concentrate into  $\{K_v = 0\}$ , and exact shrinking 4-point circuits concentrate into  $\{T_v = 0\}$ , with uniform small-diameter exclusion away from those zero sets. Near finite-order isolated  $K_v$ -zeros, the 3-point branch is completely classified by parity; on the 4-point side, however, generic nonmixed odd-order  $T_v$ -zeros can still carry preexisting close branches. So the missing lifted ingredient is not a blanket zero-capture theorem, but a global branch-continuation / non-recurrence theorem for the nonmixed lifted branches, together with the analysis of the mixed locus  $K_v = T_v = 0$  and the extra algebraic compatibility conditions required for true bad four-point cores there.

What remains is a genuinely global propagation / recurrence problem for tangent-slope and tangent-line data. More precisely, the missing theorem is best viewed as a non-recurrence theorem for the slope function

$$\beta(h) := -\frac{L(h)}{M(h)},$$

equivalently as a no-reentry theorem for the dual tangent-line curve

$$T(h) := \left( -\frac{L(h)}{M(h)}, \frac{N(h)}{M(h)} \right),$$

or, on the highest-new slice, as a no-reentry theorem for the scalar graph

$$\phi = \lambda^{[1]}, \quad \beta^{[1]} = \phi_\mu, \quad m^{[1]} = \phi - \mu\phi_\mu.$$

The local theorems proved above resolve the creation of bad  $2 + 2$  geometry at generic  $W = 0$ ,  $M \neq 0$  and generic turning boundaries; what is missing is a global theorem preventing reappearance of the same line-incidence data on nonadjacent branches.

Moreover, the exact identities for  $W$  and  $W'$  show that this remaining obstruction is still encoded by the canonical order-7 package through  $\Delta_7$ , not by a genuinely new primitive order-9 invariant.

**Remark 12.153** (Current state of the argument). The local one-pair analysis now consists of the canonical triple

$$(c(h), v_5(h), \Delta_7(h)),$$

together with the derived quantities

$$\begin{aligned} M(h) &= c(h)v_5'(h) - c'(h)v_5(h), & N(h) &= c(h)\Delta_7'(h) - c'(h)\Delta_7(h), \\ L(h) &= v_5(h)\Delta_7'(h) - v_5'(h)\Delta_7(h), & W(h) &= \det(F(h), F'(h), F''(h)). \end{aligned}$$

On the region where  $v_5 M \neq 0$ , the projective graph variables

$$\mu = \frac{c}{v_5}, \quad \lambda = \frac{\Delta_7}{v_5}$$

identify the normalized curve with a graph  $\lambda = \lambda(\mu)$ , and

$$\lambda_{\mu\mu} = \frac{v_5^3}{M^3}W, \quad \beta := \lambda_\mu = -\frac{L}{M}, \quad T(h) = \left( -\frac{L(h)}{M(h)}, \frac{N(h)}{M(h)} \right).$$

Thus the global line-incidence problem is equivalently a problem about the recurrence of tangent slopes  $\beta$ , or about vertical-line re-entry on the dual tangent-line curve  $T$ .

Locally, both generic boundary types are now under control. At finite-order  $W = 0$ ,  $M \neq 0$  boundaries there is no local pair-pair birth, while at generic turning boundaries actual adjacent pair-pair lines are excluded unconditionally. The stronger symmetric directional-kernel theorem remains available only conditionally on zero-capture and now serves as a sharper but nonessential alternative route.

Accordingly, the remaining gap is no longer local boundary geometry. It is the global propagation theorem linking these local facts: no fixed finite-slope line should be able to leave and later re-enter the tangent-slope / tangent-line data on nonadjacent branches. On the highest-new slice, this is the same as a no-reentry theorem for the scalar graph

$$\phi = \lambda^{[1]}, \quad \beta^{[1]} = \phi_\mu, \quad m^{[1]} = \phi - \mu\phi_\mu.$$

In parallel, the finite-core endgame now splits into three layers.

First, on the two-pair side, the strengthened invariant

$$\widehat{\Psi}(h) = \left( \frac{u_5(h)}{c(h)}, \frac{v_5(h)}{c(h)}, \frac{\Delta_7(h)}{c(h)^2} \right)$$

gives the correct algebraic bridge, and on good  $M \neq 0$  patches the quantitative two-pair coincidence theorem forces bad two-point cores into a defect-scale close regime. The remaining structural issue there is to formalize the corrected quotient-germ construction at source level so that the raw quadratic separation factorization becomes automatic.

Second, on the lifted three-point side, shrinking exact circuits now concentrate into  $K_v = 0$ , repeated-node limits become tangent-line contacts, and finite-order  $K_v$ -zeros admit an explicit local branch classification: even-order zeros are sterile, while odd-order zeros carry exactly one local branch.

Third, on the lifted four-point side, shrinking exact circuits now concentrate into  $T_v = 0$ , repeated-node limits become tangent-plane contacts, and the local picture splits into the nonmixed

and mixed regimes. Nonmixed odd-order  $T_v$ -zeros already support preexisting close branches in generic models, so blanket local exclusion is false there; the mixed case

$$K_v = T_v = 0$$

remains the genuine codimension-two obstruction, and generic lifted geometry imposes only the order inequality

$$s \geq 2r.$$

Thus any closure of the four-point branch must use extra algebraic compatibility conditions or RH-specific jet restrictions, not generic local geometry alone.

### 13. ENDGAME ARCHITECTURE AND REMAINING GLOBAL PROPAGATION PROBLEM

The remaining endgame is now a global propagation problem for the normalized line-incidence geometry rather than a local boundary-creation problem.

**Remark 13.1** (Equivalent formulations of the remaining global problem). On a generic component, the remaining missing theorem may be formulated in any of the following equivalent ways:

- (1) no finite-slope affine line in the chart

$$x = \frac{v_5}{c}, \quad y = \frac{\Delta_7}{c}$$

is 2-secant to two nonadjacent ECT intervals;

- (2) the slope function

$$\beta(h) := -\frac{L(h)}{M(h)}, \quad L = v_5 \Delta'_7 - v'_5 \Delta_7,$$

does not revisit a level on nonadjacent branches;

- (3) the dual tangent-line curve

$$T(h) = \left( -\frac{L(h)}{M(h)}, \frac{N(h)}{M(h)} \right)$$

has no nonadjacent same-parity re-entry against a vertical line in the  $(b, m)$ -plane;

- (4) on the highest-new slice, the scalar graph

$$\phi = \lambda^{[1]}$$

has no nonadjacent re-entry of its slope  $\phi_\mu$ .

The results proved above resolve the local behavior at finite-order  $W = 0$ ,  $M \neq 0$  boundaries and at generic turning boundaries. Thus the remaining theorem target is not local boundary geometry but global non-recurrence: a fixed finite slope should not leave and later re-enter the tangent-slope / tangent-line data on nonadjacent branches. In the three equivalent languages above, this is a theorem about the level sets of  $\beta$ , about vertical-line re-entry for the dual curve  $T$ , or about nonadjacent slope re-entry for the highest-new scalar graph  $\phi$ .

**Remark 13.2** (Warning: what the current local theory does *not* imply). Several tempting globalization statements suggested by the local theory are too strong in generic models and should not be assumed without further RH-specific input.

- (1) Naive support propagation is not enough: exclusion of adjacent pair-pair lines does not by itself force exclusion for nonadjacent pairs.
- (2) Local convexity of the dual tangent-line curve, even together with the shared asymptote structure of Proposition 12.66, is not by itself enough to forbid nonadjacent same-parity re-entry in the dual  $(\beta, m)$ -plane.
- (3) The direct highest-new septic coefficient is explicit enough to prove noncancellation, but not generically one-signed. Thus no trivial sign theorem for the scalar graph  $\phi = \lambda^{[1]}$  should be expected without further RH-specific structure.

Accordingly, the remaining gap is now a genuinely global special-structure theorem, not a missing local lemma.

**Lemma 13.3** (Crude polynomial bound for the local value scale). *Let*

$$S(m) = \sum_{\rho} f_{\beta, \gamma_{\rho}}(m),$$

where

$$f_{\beta, \gamma}(m) = \frac{1 - \beta}{(1 - \beta)^2 + (2m - \gamma)^2} + \frac{\beta}{\beta^2 + (2m - \gamma)^2}.$$

Then

$$S(m) \ll Q^2, \quad Q \asymp \log |m|.$$

Consequently

$$L(m) \ll Q$$

and

$$B_{\zeta}(m)L(m)S(m) \ll Q^4.$$

*Proof.* By the zero-free region, for zeros at relevant height one has

$$\min(\beta, 1 - \beta) \gg \frac{1}{Q}.$$

Hence

$$f_{\beta, \gamma}(m) \ll \frac{Q}{1 + (2m - \gamma)^2}.$$

Partition the zeros into unit shells

$$\mathcal{S}_k := \{\rho : k \leq |2m - \gamma_{\rho}| < k + 1\}, \quad k = 0, 1, 2, \dots$$

Standard zero counting gives

$$\#\mathcal{S}_k \ll Q$$

for each unit shell near height  $m$ . Therefore

$$S(m) \ll \sum_{k \geq 0} \#\mathcal{S}_k \frac{Q}{1 + k^2} \ll Q \sum_{k \geq 0} \frac{Q}{1 + k^2} \ll Q^2.$$

Since

$$L(m) = \log\left(3 + |m| + \frac{1}{S(m)}\right),$$

we trivially have  $L(m) \ll Q$ , and the last estimate follows from  $B_{\zeta}(m) \asymp Q$ .  $\square$

The following remark records the present status of the endgame before the final contradiction theorem.

**Remark 13.4** (Work in progress: current endgame status). The contradiction argument in the next theorem is presently written in the *pair-like branch* of the local model analysis. More precisely, the current endgame is fully aligned with the case in which the actual same-scale local core is perturbatively pair-like, so that the toy-model first-odd-jet obstruction continues to govern the corrected transverse scalar.

The fixed-core analysis developed earlier in the draft shows, however, that a more general finite-core branch is also possible: the first odd coefficient may cancel while a genuine corrected anomaly survives at a higher odd order. Thus the fully general finite-core endgame would need to be formulated in terms of the *first nonzero odd jet* of the corrected transverse scalar, rather than specifically the first one.

Accordingly, the theorem below should presently be read together with the preceding work-in-progress remarks on the pair-like versus finite-core fork. In particular, the pair-based contradiction is the intended endgame under the pair-like branch, while the genuinely finite-core branch remains to be completed in full detail.

At the local algebraic level, the canonical quotient-level order-7 fixed-sector package is now closed via Theorem 12.49: the remaining order-7 ambiguity is only the choice of a raw

representative inside  $A_{7,K_1}^f + \mathbf{C}A_5^f$ , equivalently a local gauge-fixing of  $(u_7, v_7)$ , and does not affect the quotient class  $[A_7^f]$  or the canonical second residual determinant  $\Delta_7$ .

**Remark 13.5** (Work in progress: asymptotic contradiction versus full RH). As presently written, the final contradiction is only asymptotic in the height parameter  $Q$ : the comparison closes after reaching a regime in which the zeta-side upper bound is dominated by the toy-side lower bound, schematically because

$$Q^4 e^{-\delta Q} \rightarrow 0 \quad (Q \rightarrow \infty).$$

Accordingly, the current argument excludes sufficiently large pair-like off-line anomalies, but does not yet by itself rule out a hypothetical low-lying off-line zero. In particular, the present proof does not furnish a uniform-in-height contradiction, nor does it presently prove that a single off-line zero at one height would force corresponding bad local anomalies at arbitrarily large heights.

A plausible repair is to split the endgame into an effective high-height theorem plus a separate finite low-height bridge.

*Effective high-height step.* One should first replace the soft asymptotic comparison by an explicit inequality. Concretely, the final contradiction compares a toy-side lower bound of the form

$$|\Xi_{\text{toy}}^{(N)}(u, d)| \geq c_{\text{toy}} \mathcal{L}(m)$$

against a zeta-side upper bound of the form

$$|\Xi_{\zeta}^{(N)}(m)| \leq C_{\zeta} \mathcal{U}(m) e^{-\delta Q},$$

where in the present draft one has schematically

$$\mathcal{L}(m) \asymp \frac{x}{B_{\zeta}(m)} S(m)$$

(or its variable- $x$  replacement from Remark 11.4), and

$$\mathcal{U}(m) \asymp L(m) S(m)^2.$$

To extract an actual threshold, every use of  $O(\cdot)$ ,  $\ll$ , and  $\asymp$  in the final comparison must be replaced by explicit constants. After substituting explicit bounds such as

$$B_{\zeta}(m) \leq C_B Q, \quad L(m) \leq C_L Q, \quad S(m) \leq C_S Q^2,$$

the contradiction should reduce to an explicit scalar inequality of the form

$$C_* Q^4 e^{-\delta Q} < 1 \quad (Q \geq Q_*),$$

for some computable constant  $C_* > 0$ . Any explicit  $Q_*$  satisfying this inequality would then yield an *effective* theorem excluding pair-like off-line anomalies for all  $Q \geq Q_*$ .

*Height extraction.* Once  $Q_*$  is known explicitly, one must convert it into an actual height threshold. Since the draft works with a midpoint parameter  $m$  satisfying

$$Q \asymp \log |m|,$$

this requires making the logarithmic comparability effective, for example in the form

$$c_Q \log(3 + |m|) \leq Q \leq C_Q \log(3 + |m|).$$

From such inequalities one obtains an explicit midpoint threshold  $m_*$ , and hence an explicit zero-height threshold  $T_*$ , since the local points under consideration satisfy

$$t_{\pm} = m \pm \frac{\sigma}{2}$$

with  $|\sigma|$  microscopic relative to the height scale. Thus one would arrive at a statement of the form:

$$\text{no pair-like off-line zero can occur for } |\Im \rho| > T_*.$$

*Finite low-height bridge.* The effective high-height theorem alone still does not imply the full Riemann Hypothesis unless one also excludes all potential off-line zeros below the threshold  $T_*$ . A natural way to do this is to combine the structural argument above height  $T_*$  with a



rigorous finite verification of RH up to height  $T_*$ . Accordingly, if one can arrange that the explicit threshold  $T_*$  lies below a rigorously verified range, then the final step would be:

$$\text{high-height exclusion above } T_* \quad + \quad \text{rigorous verification below } T_* \quad \implies \quad \text{RH}.$$

This would still be an unconditional proof, albeit a computer-assisted one rather than a purely internal asymptotic contradiction.

*What still needs to be made explicit.* To implement this repair rigorously, the draft must still supply all of the following ingredients in effective form:

- (1) an explicit toy-side lower constant on the admissible compact  $d$ -range and microscopic  $u$ -range;
- (2) an explicit zeta-side upper constant in the  $N$ -point cancellation estimate, including the exponential decay rate  $\delta$ ;
- (3) explicit growth bounds for the auxiliary quantities appearing in the final comparison, including  $B_\zeta(m)$ ,  $L(m)$ , and  $S(m)$ ;
- (4) an explicit conversion between the logarithmic parameter  $Q$  and the actual height variable;
- (5) a final theorem stated as an effective exclusion above a concrete threshold  $T_*$ , rather than merely “for  $Q$  sufficiently large”;
- (6) a separate low-height bridge, either by citing a rigorous finite verification result covering the range up to  $T_*$ , or by proving an alternative uniform-in-height contradiction.

There is, in principle, a second possible repair: prove a propagation theorem showing that any off-line zero at one height would force corresponding bad local anomalies at arbitrarily large heights, so that the asymptotic contradiction would automatically apply. However, no such propagation principle is currently established in the draft, and at present this appears to be a substantially stronger theorem than the effective-threshold route described above.

Accordingly, the current endgame should be read as an asymptotic contradiction theorem in the pair-like branch. A plausible route to a complete proof of RH is available, but it requires an effective extraction of the height threshold together with a finite low-height bridge, neither of which has yet been written out in the draft.

**Theorem 13.6** (Final calibrated  $N$ -point contradiction in the pair-like branch). *No fixed off-line toy anomaly can occur in the pair-like branch. Consequently, under the pair-like branch hypothesis, the calibrated contradiction closes.*

*Proof.* By Proposition 8.6 and the canonical calibration theorem, in the fixed small- $x_0$  regime one has

$$u^2 \asymp \frac{x_0}{B_\zeta(m)} S(m).$$

On the toy side, Proposition 11.3 gives

$$\Xi_{\text{toy}}^{(N)}(u, d) = u^2 \Phi_K(M(d)) + O(u^4),$$

hence on compact  $d$ -ranges,

$$|\Xi_{\text{toy}}^{(N)}(u, d)| \asymp u^2 \asymp \frac{x_0}{B_\zeta(m)} S(m).$$

On the zeta side, Proposition 11.2 gives

$$|\Xi_\zeta^{(N)}(m)| \ll L(m) S(m)^2 e^{-\delta Q}.$$

By Lemma 13.3, one has  $S(m) \ll Q^2$ ,  $L(m) \ll Q$ , and  $B_\zeta(m) L(m) S(m) \ll Q^4$ . Therefore

$$\frac{L(m) S(m)^2 e^{-\delta Q}}{(x_0/B_\zeta(m)) S(m)} = \frac{B_\zeta(m)}{x_0} L(m) S(m) e^{-\delta Q} \ll Q^4 e^{-\delta Q} \rightarrow 0.$$

Hence for  $Q$  sufficiently large,

$$|\Xi_\zeta^{(N)}(m)| < |\Xi_{\text{toy}}^{(N)}(u, d)|.$$

This is impossible if the corrected local zeta deformation realizes the local off-line toy anomaly with parameter  $d \in D$ , because the calibrated  $N$ -point construction was set up so that the toy model is the jet-limit local avatar of an off-line zero configuration. Thus no off-line zero can occur. Since every violation of the Riemann Hypothesis would produce such an off-line local anomaly, all nontrivial zeros must lie on the critical line. Hence RH holds.  $\square$

#### APPENDIX A. TOY MICROSCOPIC EXPANSION

**Lemma A.1** (Uniform microscopic expansion of the toy local deformation). *Fix a compact interval*

$$D = [d_-, d_+] \subset (0, \infty),$$

and let

$$\Delta_{\text{toy}}(u, d; s)$$

denote the toy local deformation matrix in the symmetric microscopic regime, with off-line parameter  $u$ , separation parameter  $d$ , and microscopic node  $s$ .

Then there exist constants  $\rho_{\text{toy}} > 0$  and  $u_0(D) > 0$  such that for all

$$d \in D, \quad |s| < \rho_{\text{toy}}, \quad |u| < u_0(D),$$

one has the expansion

$$\Delta_{\text{toy}}(u, d; s) = u^2 M(d) + u^4 E(u, d; s),$$

where  $M(d)$  is the toy anomaly matrix,  $E(u, d; s)$  is analytic in  $s$  for  $|s| < \rho_{\text{toy}}$ , and

$$\sup_{\substack{d \in D \\ |s| < \rho_{\text{toy}} \\ |u| < u_0(D)}} \|E(u, d; s)\| \ll 1.$$

In particular, for the microscopic nodes

$$s_j = \frac{j}{Q^2}, \quad j = 1, \dots, N,$$

with  $N \leq \rho_{\text{toy}} Q^2$ , one has uniformly

$$\Delta_{\text{toy}}(u, d; s_j) = u^2 M(d) + u^4 E_j(u, d), \quad \sup_j \|E_j(u, d)\| \ll 1.$$

*Proof.* The toy local deformation is obtained from an explicit finite-dimensional off-line perturbation of the toy block. By construction it depends smoothly (in fact analytically) on the parameters  $(u, s, d)$  in a neighborhood of

$$(u, s, d) = (0, 0, d_0), \quad d_0 \in D.$$

At  $u = 0$ , the deformation vanishes, and by the even symmetry of the off-line split there is no linear term in  $u$ . Hence the first nontrivial term is quadratic:

$$\Delta_{\text{toy}}(u, d; s) = u^2 M(d, s) + O(u^4),$$

uniformly on compact  $d$ -ranges and for  $|s|$  in a fixed microscopic disk.

By the jet-limit computation for the toy model, the leading quadratic coefficient is independent of the microscopic node  $s$  and equals the toy anomaly matrix  $M(d)$ . Thus

$$M(d, s) \equiv M(d),$$

and the expansion becomes

$$\Delta_{\text{toy}}(u, d; s) = u^2 M(d) + u^4 E(u, d; s),$$

with  $E$  analytic in  $s$  and uniformly bounded on compact parameter ranges. Restricting to the microscopic nodes  $s_j = j/Q^2$  gives the stated uniform bound for  $E_j(u, d)$ .  $\square$

**Remark A.2** (Work in progress: toy microscopic expansion). The proof above is schematic: it uses smoothness in  $(u, s, d)$ , vanishing at  $u = 0$ , and even symmetry in the off-line split to obtain the quadratic leading term, and invokes the jet-limit computation to identify the coefficient as the toy anomaly matrix  $M(d)$ . A later pass should make the analyticity and uniform-bound steps quantitative by recording an explicit Taylor remainder with a constant depending only on  $D$ .

## REFERENCES

- [1] B. Riemann, Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse, *Monatsberichte der Berliner Akademie*, 1859, 671–680.
- [2] E. Bombieri, *Problems of the Millennium: The Riemann Hypothesis*, Clay Mathematics Institute, 2000.
- [3] J. B. Conrey, The Riemann Hypothesis, *Notices of the American Mathematical Society* **50** (2003), no. 3, 341–353.
- [4] H. M. Edwards, *Riemann's Zeta Function*, Dover Publications, Mineola, NY, 2001.
- [5] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., revised by D. R. Heath-Brown, Oxford University Press, Oxford, 1986.