A residual replacement for the linear incidence guide

15 October 2026

Abstract

Stochastic epidemic models of cumulative incidence are used when new infections are recorded in discrete windows and the latent path is not observed. A linear guide proposes the latent counts by freezing the hazard until the next observation and shifting it with a Gaussian approximation of the count. The importance weight keeps the discretised transition in the numerator. On repeated synthetic series the median effective sample size of this guide falls from 149.66 at the published spacing to 2.06 at a spacing five times coarser. A myopic proposal does not fall in the same way. We replace the Poisson means by the increment of a restarted linear-noise path, plus a conditional residual mean for infections. The denominator stays a product of Poisson probabilities for the same increments. The comparison fixed before the second run asked whether the residual median at the coarse spacing would exceed 2.06. The value is 169.45. That median still declines with the spacing, and at the coarse spacing the gain sits in the second window. In five of twenty replicates every draw in the first window has weight zero. We do not analyse a surveillance series.

Keywords. conditioned hazard, effective sample size, Markov jump process, particle filter, SIR model

1 Introduction

Cumulative incidence is a common observation in epidemic modelling when infection times are hidden and only counts of new cases are recorded in successive windows. Whitaker et al. (2025) take a time-discretised susceptible-exposed-infected-removed model, and its susceptible-infected-removed reduction, as the latent transmission process, and they assimilate noisy incidence with a particle filter. The same construction matters more widely for

partially observed jump processes, where a conditioned proposal is used because forward simulation ignores the next observation (Golightly and Wilkinson 2015; Golightly and Sherlock 2019).

The inferential obstacle is an intractable observed-data likelihood. The data are incomplete and discrete, the reporting model is noisy, and the latent incidence between two observation times is a path of counts. A bootstrap filter would draw that path from the unconditioned transition and weight by the observation alone. Whitaker et al. (2025, p. 8) give two reasons not to stop there. Resampling static parameters collapses them, and forward simulation of incidence ignores the next count.

Whitaker et al. (2025) meet the second difficulty with a conditioned hazard. Particles are propagated by replacing the Poisson rate with a hazard derived from a Gaussian approximation of the cumulative incidence between the current time and the next observation, given that observation. The particle is then reweighted by the discretised transition and the observation mass function. The limitations section of that paper states that a further assumption in the derivation is linearity of the latent process over an interval no longer than the observation interval. When the epidemic is sparsely observed, so that cumulative incidence does not evolve linearly, "the conditioned hazard is likely to result in trajectories that are inconsistent with the true conditioned process" (Whitaker et al. 2025, p. 25).

That paragraph does not give a numerical spacing at which the failure begins, and it does not say that the importance weight then targets a different distribution. It names recent work on sampling conditioned jump processes, in particular Golightly and Sherlock (2019) and Corstanje et al. (2023), as relevant and open to adaptation. It does not print either algorithm as a replacement inside the filter.

The contribution is in four parts. Firstly, we read the constant-hazard step in the 2025 derivation as the linearity assumption named on that page, and we read the inconsistency as a property of the proposal. Secondly, we report the effective sample size of the incidence weights on the synthetic susceptible-infected-removed mechanism of Whitaker et al. (2025), at the published spacing and at two coarser spacings fixed in advance. Thirdly, we record why equation (19) of Golightly and Sherlock (2019) and Theorem 4.2 of Corstanje et al. (2023) do not replace the proposal as the filter is written. Fourthly, we replace the Poisson means by the increment of a restarted incidence linear-noise path plus a conditional residual mean, and we compare the effective sample size of that proposal with the linear guide on the same mechanism. The density in the denominator remains a product of Poisson probabilities for those incidence increments.

The illustration is synthetic. Whitaker et al. (2025) also discuss weekly Ebola counts and weekly COVID-19 counts. Those counts are plotted and are not stored here as a data file, so we do not reanalyse them.

The remainder of this paper is organised as follows. Section 2 compares the guides that were read. Section 3 restates the 2025 filter and the linear incidence guide. Section 4 states what the diagnostic claims. Section 5 states the residual proposal. Section 6 reports both comparisons. Section 7 records the qualifications.

2 Related work

We compare guides by what they propagate, what they condition on, and whether an importance weight corrects the proposal. Table 1 keeps only rows whose cells were filled from a paper we opened. A paraphrase is not a substitute for that paper.

Equation (9) of Golightly and Wilkinson (2015) is the constant-hazard step that Whitaker et al. (2025) adapt. The observation in the 2015 paper is a Gaussian function of the state, not a binomial or negative binomial count of incidence. Equation (5) of Golightly and Sherlock (2019) restates that hazard. Their Section 3.2 says that normality of the remaining reaction counts, and a hazard that is constant on the whole remaining interval, are often unreasonable. Their Section 6 says that treating the time until the next observation as one discretisation makes the bridges less accurate as the inter-observation time increases. Equation (19) of the same paper is a different proposal: a jump-process bridge under a Gaussian observation of the state, with weight (3). It is not a product of Poisson probabilities on an incidence grid.

Theorem 4.2 of Corstanje et al. (2023) conditions an inhomogeneous Poisson process on an exact integer end-point. The likelihood ratio (16) contains $h(0, x_0)$, the transition probability of the original process. Their Section 4.1 does not treat a binomial observation of an increment on a grid. Section 5 does not put either construction into the incidence weight. The proposal stated there uses the incidence linear-noise equations of Golightly et al. (2023) and the residual covariance block of Whitaker et al. (2017, Section 3.1.2), and it evaluates a Poisson probability for the same increments as the numerator.

Lowe et al. (2023) accelerate particle Markov chain Monte Carlo for a chemical kinetic model. On the pages we read, the conditioned hazard is a jump hazard multiplied by a ratio of observation densities, with that density

replaced by a linear-noise Gaussian. That account points at Golightly and Sherlock (2019). It is not the incidence proposal in Whitaker et al. (2025). The same pages say that nonlinear behaviour between observations precludes a modified diffusion bridge that pushes particles towards the observations in a linear fashion. They name Durham and Gallant (2002) for that bridge, and they name Whitaker, Golightly, Boys and Sherlock (2017) for a residual bridge. Neither of those papers was read.

Jovanovski et al. (2024) build data-conditional proposals for approximate Bayesian computation on a diffusion. On the pages we read, the discussion of erratic paths is a conjecture about where their speed-up is largest. It is not a statement that the 2025 incidence hazard fails. Whitaker et al. (2025, pp. 2–3) also name Golightly and Kypraios (2018) beside the bridge. That paper was not read, and we do not use it as a source for the guide.

3 The 2025 filter and the linear guide

Consider the filter of Whitaker et al. (2025). Let X_t denote prevalence and let N_t denote cumulative incidence. The unconditioned hazard is h, and $H = \mathrm{diag}\{h\}$. A scalar observation y_{t_i} is a noisy count of one incidence component over $(t_{i-1}, t_i]$, selected by a vector P. In the susceptible-infected-removed applications, $P' = (1, 0)$. The contact rate enters on the log scale as $\tilde{\beta}_t$, and we use the precision form printed as equation (2) of Whitaker et al. (2025). The static parameter and the particle index are those of their Algorithm 2. We do not restate the gamma updates of their equations (9)–(11). The diagnostic below does not sample them.

Their Algorithm 2 is printed for the negative binomial observation. The synthetic study in their Section 4.1 uses the binomial observation. They do not print a second algorithm box, and we do not invent one. The observation factor in the weight is the mass function of whichever model is in use. In our simulation that factor is the binomial probability mass function.

Algorithm 1. Schematic of Algorithm 2 of Whitaker et al. (2025). The draw from q in step 1c is the object of the diagnostic. It is not replaced.

1. Propagate each particle. Jitter the observation parameter with the kernel printed for the negative binomial model. Draw the contact-rate path from its transition, which the paper calls a blind propagation. Draw the incidence path from the proposal q, which depends on the next observation.

2. Resample with the weight in Equation (1).

3. Update the sufficient statistic for the rate parameters.

4. Draw those rate parameters from their conditional posteriors.

The weight in step 2, for particle k on the window that ends at the next observation, is

$$w^{(k)} \propto \frac{\pi\big(\Delta n^{(k)} \mid \tilde{\beta}^{(k)}, \theta^{(k)}\big)\, \pi\big(y \mid \Delta n^{(k)}, \phi^{(k)}\big)}{q\big(\Delta n^{(k)} \mid \tilde{\beta}^{(k)}, \phi^{(k)}, \theta^{(k)}, y\big)}. \tag{1}$$

The numerator is the time-discretised transition and the observation mass function. It does not contain the conditioned hazard h^*. The denominator is q. We read this ratio as an importance weight for one propagate step. We do not add a convergence theorem for the particle filter.

The proposal q is built in their Section 3.4. At a time τ inside the observation window, equation (12) of Whitaker et al. (2025) approximates the remaining incidence by a Gaussian whose mean and variance are the current hazard times the remaining time. The observation is given a Gaussian approximation, and the mean inside the observation variance is replaced by the mean of that Gaussian. The display after their equation (18) is the conditioned hazard h^*. End-point paths are their forward recursion with h replaced by h^*. The density q is the product of Poisson probabilities with rates $h^*\Delta\tau$.

The printed correction multiplies by $\rho P' H$, a row when the observation is scalar. The cross-covariance block that enters the conditional mean of the remaining incidence is a column. For a diagonal H the row and the column have the same entries. On the susceptible-infected-removed model the shared entries correct the infection component only,

$$h^*_{\mathrm{inf}} = h_{\mathrm{inf}} + \rho h_{\mathrm{inf}} \frac{y - \hat{\mu}}{\rho^2 h_{\mathrm{inf}}(t_i - \tau) + \sigma^2}, \qquad h^*_{\mathrm{rem}} = h_{\mathrm{rem}}, \tag{2}$$

where $\hat{\mu} = \rho\big(n^{\mathrm{past}}_{\mathrm{inf}} + h_{\mathrm{inf}}(t_i - \tau)\big)$ and, for the binomial variance evaluated at $\hat{\mu}$, $\sigma^2 = (1-\rho)\hat{\mu}$. Equation (2) is that scalar form. It matches the entries of the printed display, and it matches equation (9) of Golightly and Wilkinson (2015) in the case where the stoichiometry is the identity, so that the correction is the column HP. It is not a new guide.

The study in Section 6 does not run the jitter, the blind contact-rate draw, the sufficient-statistic update or the conditional parameter draw. The contact-rate path and the prevalence at the left endpoint of the window are taken from the simulated path, and the removal rate and the reporting probability stay at the simulation values. A particle in Algorithm 1 does not have that information. The diagnostic is about h^* given those paths.

4 Diagnostic

Equation (12) of Whitaker et al. (2025) freezes the hazard at the left endpoint of the remaining interval. The mean of the remaining incidence is that frozen hazard times the remaining time. Golightly and Wilkinson (2015, Section 3.1) use the same constant-hazard Gaussian step for remaining reaction counts. Golightly and Sherlock (2019, Section 3.2) say that this step is often unreasonable, and their Section 6 says that accuracy falls as the inter-observation time grows. Whitaker et al. (2025, p. 25) call linearity of the latent process, over an interval no longer than the observation interval, a further assumption in the derivation of the conditioned hazard. Their Section 3.4 does not use the word linear, and it does not print a second condition beyond equation (12) and the display after (18). We do not separate those two passages. The constant-hazard step is the assumption the limitations paragraph is pointing at.

The same paragraph says the conditioned hazard is likely to propose trajectories inconsistent with the true conditioned process, when cumulative incidence does not evolve linearly. The law named on the page is the true conditioned process. The sentence is about the trajectories the hazard proposes. It does not say that the discretised transition in the numerator of Equation (1) is the wrong model, and it does not say that the weight targets a different distribution.

A poor h^* changes q and therefore the weights. On the printed ratio, the importance target of one propagate step stays the discretised transition times the observation mass function. This is a reading of the printed weight. It is not a proof that the particle filter converges.

In that sense, inconsistent is a property of the proposal. No paper we opened states a numerical spacing at which the 2025 guide fails for this epidemic model. Golightly and Sherlock (2019, Section 5.1) report that a linear form fails to describe exponential decay, and that effective sample size falls as the interval length grows, for a death process. We do not copy those figures. They are not results for the epidemic model.

The observable is the effective sample size of the unnormalised incidence weights in one window,

$$\mathrm{ESS}(\tilde{w}_{1:N}) = \frac{\left(\sum_{i=1}^{N} \tilde{w}_i\right)^2}{\sum_{i=1}^{N} \tilde{w}_i^2}, \tag{3}$$

as printed by Golightly and Sherlock (2019, Section 5.2). Zero weights are included. If every weight in a window is zero, Equation (3) is undefined and

the recorded value is 0, as a convention for the table. A draw has weight zero when the correction in Equation (2) is undefined, when a component of the hazard passed to the Poisson draw is negative, or when the updated counts would be negative. Whitaker et al. (2025) do not print a truncation. Golightly and Sherlock (2019, Section 3.2) truncate their equation (5) to keep it positive. We do not copy that truncation onto the 2025 display.

The prediction, fixed before the run, is that the median of this effective sample size across replicate data sets is smaller at the coarsest pre-registered spacing than at the published unit spacing. A myopic proposal, with h^* replaced by h, is a comparator. It is not a second target. The diagnostic will appear as this comparison. It is not a threshold, and it does not say that the filtering distribution changes.

Several claims are outside the paper. Section 5 states a different proposal. We do not replace q by equation (19) of Golightly and Sherlock (2019) or by Theorem 4.2 of Corstanje et al. (2023). We do not drop $h(0, x_0)$ from their ratio (16). We do not present a truncated negative hazard as the 2025 guide. We do not claim a spacing, beyond the three values fixed before the run, at which the guide fails. We do not claim that a poor h^* changes the filtering target. We do not use the death-model table of Golightly and Sherlock (2019) as evidence about the epidemic model. We do not compute a gold-standard posterior.

5 Residual incidence guide

The two constructions named in the limitations paragraph of Whitaker et al. (2025) do not supply the density in the denominator of Equation (1). Equation (19) of Golightly and Sherlock (2019) samples a Markov jump process, and the weight (3) is the jump-process likelihood ratio under a Gaussian observation of the state. Theorem 4.2 of Corstanje et al. (2023) conditions an inhomogeneous Poisson process on an exact integer end-point, and the ratio (16) contains the transition probability $h(0, x_0)$. Corollary 4.3.6 and Theorem 4.4.2 of Corstanje (2024) are path-space ratios for a guided chemical reaction process conditioned on an exact end constraint. Equations (15) and (25) of Whitaker et al. (2017) are Gaussian densities of a diffusion skeleton. We do not put any of those expressions in place of q.

The residual guide keeps the Poisson draw and keeps q as the product of Poisson probabilities on the same incidence increments. The numerator of Equation (1) is unchanged. The Poisson means change.

At a substep, the linear-noise approximation of cumulative incidence is

restarted at the current prevalence. The incidence coordinate η starts at zero, the fundamental matrix G starts at the identity, and the residual variance V starts at zero. The contact rate is held at its current value. We integrate equations (4), (8) and (9) of Golightly et al. (2023) over the next substep and over the time remaining until the observation. The Jacobian is the matrix printed after their equation (6). The restart is the one they describe after their equation (19). Fearnhead et al. (2014, Section 4.2) state that restart: the ordinary differential equation is re-initialised at the beginning of each interval. The integration uses a classical Runge–Kutta step of length 0.1. That length is a numerical choice, fixed before the comparison in Section 6. It is not a step in Golightly et al. (2023).

The mean of the binomial observation then uses the linear-noise mean of the remaining infections in place of the frozen product $h_{\mathrm{inf}}(t_i - \tau)$. The variance remains Equation (2)'s binomial variance, $\sigma^2 = (1 - \rho)\hat{\mu}$. For an observation of infections this is the variance in equation (22) of Golightly et al. (2023).

Let R denote the residual incidence about η, with variance V after the restart. The covariance between the residual at the next substep and the residual at the end of the remaining interval is $V_{\Delta\tau}(G_{t_i-\tau}G_{\Delta\tau}^{-1})'$. Whitaker et al. (2017, Section 3.1.2) write the same block as $P_t\psi_t P_T'$ after their display (24). The observation loads on the infection component by ρ. The conditional mean of the infection residual at the next substep is that infection covariance, times $\rho(y - \hat{\mu})$, divided by $\rho^2 V_{11}(t_i - \tau) + \sigma^2$. The removal component is not shifted, as in the scalar form of Equation (2).

The Poisson mean for the next infection increment is the ordinary-differential increment of η plus that conditional residual mean. The Poisson mean for the removal increment is the ordinary-differential increment of the removal coordinate alone. A negative mean is not replaced by zero. The weight is zero if the integration stops because a prevalence on the ordinary differential equation would be negative, if the determinant of $G_{\Delta\tau}$ is smaller than 10^{-12} in absolute value, if either Poisson mean is negative or is not finite, or if the updated counts would be negative. The truncation in Section 3.2 of Golightly and Sherlock (2019) is not used.

Equation (6) of Golightly and Kypraios (2018) is the constant-hazard correction that Whitaker et al. (2025) adapt. It is not this proposal. Fintzi et al. (2022) reparameterise a linear-noise approximation for incidence counts. The equations integrated here are those printed by Golightly et al. (2023), which cite that reparameterisation. We do not transcribe an unread display from Fintzi et al. (2022) into q.

6 Simulation

The study uses the synthetic susceptible-infected-removed mechanism of Section 4.1 of Whitaker et al. (2025), observed at spacings 1, 2 and 5. Spacing 1 is the published grid. Spacings 2 and 5 are coarser, and they were fixed before any weight was seen. All algorithms were coded in R. No processor is recorded for this run. The data are simulated. There is no gold-standard posterior: the diagnostic does not estimate a posterior distribution of the static parameters.

We drew 20 replicates, with 500 proposals in each window, from seed base 20251015. The seeds are a property of the study script, not of the guide. No fine-grid path was discarded. The run took 46.879 seconds. That time is a record of this run. It is not a comparison with the run times in Whitaker et al. (2025).

For each replicate, each spacing and each guide, the within-replicate summary is the median of the window values of Equation (3). The study summary is the median of those 20 numbers. The prediction uses the linear guide only. The myopic guide is computed on the same simulated paths and the same observations, with independent Poisson draws. Table 2 reports the linear guide. Figure 1 plots those three medians and the three myopic medians given below. The myopic series in the figure is a comparator and is not the prediction.

From Table 2, the linear median effective sample size is 149.660679297905 at spacing 1 and 2.05670742297318 at spacing 5. The value at spacing 5 is smaller, so the prediction holds. At spacing 2 the linear median is 8.09166818813341. At every linear row, the median that includes zero weights equals the median computed on the finite weights alone. The fall is present in the finite weights. The median proportion of invalid draws is 0.0329 at spacing 1, 0.009 at spacing 2 and 0 at spacing 5, so the fall at spacing 5 is not produced by the zero-weight rule.

At spacing 1 the myopic median is 139.525563697813. At spacing 2 it is 142.824313796391. At spacing 5 it is 195.181391518181. The myopic median does not fall as the spacing grows.

Two qualifications belong with the comparison. The guide was given the simulated prevalence and the simulated contact-rate path. The result is about h^* given those paths. At spacing 1 the linear median is 149.660679297905 out of 500 draws, so the guide is already uneven on the published grid. The prediction was the comparison across spacings, not a claim that the published grid is unaffected. The comparison does not measure a pathwise distance to the true conditioned process, and it does not say




Figure 1: Effective sample size against observation spacing. Filled circles: linear guide, the medians in Table 2. Open circles: myopic comparator. Triangles: residual guide, the medians in Table 3. The prediction for the residual guide is the comparison at spacing 5 with the linear median, not the slope of the triangle series.

whether a filtering distribution would change.

The residual guide of Section 5 was run on the same mechanism, the same seeds and the same observation draws. The linear proposals were not redrawn. Twenty replicates and 500 draws were used again. The ordinary differential equations were integrated with a Runge–Kutta step of length 0.1. No fine-grid path was discarded. The run took 1502.073 seconds. The prediction, fixed before this run, was that the residual median effective sample size at spacing 5 would be larger than 2.05670742297318. With two windows, the within-replicate median is the average of the two window values. That is the same functional as the linear summary at spacing 5.

From Table 3, the residual median is 281.708848707666 at spacing 1, 236.47088660943 at spacing 2 and 169.453475371897 at spacing 5. The spacing-5 value is larger than 2.05670742297318, so the prediction holds. The residual median declines as the spacing grows. It does not fall to the linear value. At spacing 5 the median on the finite weights is 173.333046076685, so the zero-weight rule shifts the summary by a small amount beside that gap. The median proportion of invalid draws is 0.0287 at spacing 1, 0.0096 at spacing 2 and 5×10^{-4} at spacing 5.

That last proportion is a median across replicates, and it hides a split be-

tween the two windows. In the stored window file, five of the twenty first windows at spacing 5 have every one of the 500 residual draws invalid. The median effective sample size of those twenty first windows is 5.87283517539277. The median of the twenty second windows is 331.985239567309. These two medians were not the pre-registered summary. They are descriptions of the stored windows. The linear guide, on the same spacing, has median effective sample size 1.36201529794476 on the first window and 1.61726241454208 on the second, and no linear window has effective sample size 0. The pre-registered residual number at spacing 5 therefore mixes a weak first window with a much stronger second window. The comparison with 2.05670742297318 uses the functional named before the run. Figure 1 adds the three residual medians from Table 3 to the linear and myopic series.

7 Discussion

We have given a reading of the linear incidence guide in Whitaker et al. (2025) and a diagnostic for its weights under coarser observation. The constant-hazard step is the linearity assumption named in their limitations section. On the printed importance weight, a poor conditioned hazard changes the proposal and the weights. We have shown, on the synthetic mechanism of their Section 4.1, that the median effective sample size of the linear guide is smaller at spacing 5 than at spacing 1, with the values in Table 2. The myopic comparator does not fall. The guide had the simulated state and the simulated contact-rate path, and even the published spacing already yields a median well below the number of draws.

We expect the diagnostic to be useful when an observation window is lengthened and the question is whether this proposal is the source of uneven weights. We do not expect it to certify a posterior, and we do not treat it as a threshold that divides adequate spacing from inadequate spacing. The three spacings were chosen before the run because they divide the published horizon. They are not estimated change-points.

The residual guide in Section 5 replaces the Poisson means inside that proposal. The new means are the increment of a restarted incidence linear-noise path, plus a conditional residual mean on the infection count. The denominator of Equation (1) remains the product of Poisson probabilities for those increments. On the pre-registered summary, the median effective sample size at spacing 5 is 169.453475371897, which is larger than the linear median 2.05670742297318. The same summary declines from 281.708848707666 at spacing 1. The gain at spacing 5 is not shared equally by the two windows.

The first window, which covers the rise of the outbreak, has median effective sample size 5.87283517539277, and in five replicates every draw in that window has weight zero. The second window has median 331.985239567309. A guide that repairs the second half of a sparsely observed outbreak and still fails on the first half is only a partial replacement.

Equation (19) of Golightly and Sherlock (2019) remains unused as q. It samples a Markov jump process, and the weight (3) is the jump-process likelihood ratio under a Gaussian observation of the state. Theorem 4.2 of Corstanje et al. (2023) remains unused as well. Its ratio (16) contains $h(0, x_0)$. We do not drop that factor.

The weekly Ebola series and the weekly COVID-19 series in Whitaker et al. (2025) are not analysed. The counts are not stored in this repository, and we do not read them off the published figures. A surveillance example would need those counts in a form that can be reproduced, and a claim that the simulations have not already settled.

Acknowledgements

This draft records a diagnostic study. No personal acknowledgements are attached.

Funding details

Funding details have not been supplied for this draft.

Disclosure statement

The authors report there are no competing interests to declare.

Data availability

All data used in this paper were simulated from the model stated in Section 6. The table and the figure are regenerated by the script named in the supplementary note.

Supplementary material

The supplement for this draft is the base R code that evaluates the linear guide and the residual guide, the two study scripts, the stored tables, and the reproduction script. No surveillance counts are included.

References

M. Corstanje, F. van der Meulen, and M. Schauer. Conditioning continuous-time Markov processes by guiding. *Stochastics*, 95(6):963–996, 2023. doi: 10.1080/17442508.2022.2150081.

M. A. Corstanje. *Guiding techniques for conditioning Markov processes*. PhD thesis, Vrije Universiteit Amsterdam, 2024.

P. Fearnhead, V. Giagos, and C. Sherlock. Inference for reaction networks using the linear noise approximation. *Biometrics*, 70:457–466, 2014. doi: 10.1111/biom.12152.

J. Fintzi, J. Wakefield, and V. N. Minin. A linear noise approximation for stochastic epidemic models fit to partially observed incidence counts. *Biometrics*, 78:1530–1541, 2022. doi: 10.1111/biom.13538.

A. Golightly and T. Kypraios. Efficient SMC^2 schemes for stochastic kinetic models. *Statistics and Computing*, 28:1215–1230, 2018. doi: 10.1007/s11222-017-9789-8.

A. Golightly and C. Sherlock. Efficient sampling of conditioned Markov jump processes. *Statistics and Computing*, 29:1149–1163, 2019. doi: 10.1007/s11222-019-09861-5.

A. Golightly and D. J. Wilkinson. Bayesian inference for Markov jump processes with informative observations. *Statistical Applications in Genetics and Molecular Biology*, 14(2):169–188, 2015. doi: 10.1515/sagmb-2014-0070.

A. Golightly, L. E. Wadkin, S. A. Whitaker, A. W. Baggaley, N. G. Parker, and T. Kypraios. Accelerating Bayesian inference for stochastic epidemic models using incidence data. *Statistics and Computing*, 33:134, 2023. doi: 10.1007/s11222-023-10311-6.

P. Jovanovski, A. Golightly, and U. Picchini. Towards data-conditional simulation for ABC inference in stochastic differential equations. *Bayesian Analysis*, pages 1–31, 2024. doi: 10.1214/24-BA1467.

T. E. Lowe, A. Golightly, and C. Sherlock. Accelerating inference for stochastic kinetic models. *Computational Statistics and Data Analysis*, 185:107760, 2023. doi: 10.1016/j.csda.2023.107760.

G. A. Whitaker, A. Golightly, R. J. Boys, and C. Sherlock. Improved bridge constructs for stochastic differential equations. *Statistics and Computing*, 27:885–900, 2017. doi: 10.1007/s11222-016-9660-3.

S. A. Whitaker, A. Golightly, C. S. Gillespie, and T. Kypraios. Sequential Bayesian inference for stochastic epidemic models of cumulative incidence. *Bayesian Analysis*, pages 1–30, 2025. doi: 10.1214/25-BA1547.

Table 1: Guides read for this diagnostic. The Whitaker row is the proposal under study. The other rows are not substitutes for it.

Guide	What is propagated	What the proposal conditions on	Weight
Whitaker et al. (2025), display after (18)	Poisson incidence increments on a grid, at rate $h^*\Delta\tau$	The next incidence observation, through a Gaussian approximation	Yes. The ratio uses the discretised transition and the observation mass function over q
Golightly and Wilkinson (2015), equation (9)	A Markov jump process, by exponential waits at the conditioned hazard	A Gaussian observation of the state. The hazard is frozen on the remaining interval	Yes. The complete-data likelihood ratio (10)
Golightly and Sherlock (2019), equation (5)	The same construction as Golightly and Wilkinson (2015), equation (9)	The same Gaussian observation. Section 3.2 says the constant-hazard step is often unreasonable	Yes, through the weight (3), when this hazard is the proposal
Golightly and Sherlock (2019), equation (19)	A Markov jump process, using the linear-noise conditioned hazard without restarting the ordinary differential equations	The same Gaussian observation, through the linear-noise transition density in their equation (6)	Yes. The weight is (3)
Corstanje et al. (2023), Theorem 4.2	An inhomogeneous Poisson process with a homogeneous-Poisson guide	An exact integer end-point	The ratio (16) is printed and contains the transition probability $h(0, x_0)$

Table 2: Linear incidence guide. Median effective sample size across 20 replicates. Finite-weight medians use the draws with a finite log weight. The myopic comparator is not in this table.

Spacing	ESS	ESS, finite weights	Invalid proportion
1	149.660679297905	149.660679297905	0.0329
2	8.09166818813341	8.09166818813341	0.009
5	2.05670742297318	2.05670742297318	0

Table 3: Residual incidence guide. Median effective sample size across 20 replicates. Finite-weight medians use the draws with a finite log weight. The spacing-5 prediction is the comparison with the linear median 2.05670742297318 in Table 2.

Spacing	ESS	ESS, finite weights	Invalid proportion
1	281.708848707666	281.708848707666	0.0287
2	236.47088660943	236.47088660943	0.0096
5	169.453475371897	173.333046076685	5×10^{-4}