

A first order universal growth chart on the $s = 0 + I \cdot t$ line for periodic 1st degree Dirichlet character L-functions.

John Martin

August 28, 2026

Executive Summary

A comparison of the extreme peak height behaviour of several periodic 1st degree L-functions along with some Davenport-Heilbronn functions is presented on the $s = 0 + I \cdot t$ line (and for the Ramanujan Tau L-function on the $s = 5.5 + I \cdot t$ line) i.e., the lower boundary of the L-function critical strip. It is empirically demonstrated (similar to observed critical line behaviour) that a first order universal growth chart on the critical line for periodic 1st degree L-functions can be constructed by (i) rescaling of the t co-ordinate and (ii) rescaling of the L-function height by the proportion of zero (periodic) dirichlet characters. In addition, a Box Cox transformation model is fitted to the known extreme peak heights $|Z|$ of the Riemann Zeta function on the $s = 0 + I \cdot t$ line to yield a extreme peak trend growth estimate for periodic L-functions with $|\chi| = 0, 1$ dirichlet characters on the lower boundary of the critical strip. The trend model performance is then assessed by mapping the model from $s = 0 + I \cdot t$ to $s = 1 + I \cdot t$ (using the Riemann Zeta functional equation) which helps highlight deficiencies in the trend model.

Introduction

Various authors [1-21] have identified large and/or extreme Riemann Siegel Z function peaks. Most recently, Martin [21] has reported extreme Riemann Siegel Z function peaks for a range of 1st degree L functions and some of their Davenport-Heilbronn counterparts on the critical line $s = 0.5 + I \cdot t$ (as well as the Ramanujan Tau L-function on its critical line $s = 6 + I \cdot t$) using for progressively higher t values (i) available high precision L-function calculations, (ii) high precision tapered L-function dirichlet series estimates at the second quiescent region $N_2 = 2 \cdot \left(\frac{t}{2\pi}\right)^d \cdot N_C$, (iii) multiple decimal place accuracy tapered first order Riemann Siegel formula L-function estimates at the first quiescent region $N_1 = \sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}$, (iv) multiple decimal place accuracy spectrally filtered Euler Product L-function analogues of the first order Riemann Siegel formula as well as (v) 10% accuracy (critical line), 1% accuracy ($s=0,1$ lines) preliminary peak height estimates for $t < 1e30$ using a spectrally filtered heavily truncated Euler Product method $N_{sub1} = \beta \cdot 0.5 \cdot \left(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}\right)^{0.5}$ where $\beta = N_C$ for 1st degree L-functions. Where overlap of adjoining methods was included to validate calculation performance.

In particular, Martin [21] noted that (i) universal growth chart behaviour of extreme peak behaviour of 1st degree L-functions with $\{0,1\}$ magnitude dirichlet characters (in first order agreement with the Riemann Zeta Riemann Siegel $|Z|$ function behaviour) can be observed by changing the x-axis (of such 1st degree L-functions) from $t \rightarrow \log \left[\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C} \right]$ and adjusting the y axis $\log(|Z|) \rightarrow \log(|Z| \cdot \frac{N_C}{N_{\chi \neq 0}})$ and (ii) an empirical trend growth estimate can be obtained for known Riemann Zeta Riemann Siegel $|Z|$ function 1st degree L-functions with $\{0,1\}$ magnitude dirichlet characters extreme peaks on the critical line for $t < 1e40$ via the fitted Box Cox

$$\text{transformation model } \log \left[|Z_L(k + I \cdot t)|_{\max} \cdot \frac{N_C}{N_{\chi \neq 0}} \right]_{\text{trend}} \lesssim 1.274014 + 0.866363 \cdot \left(\frac{\sqrt{\log \left[\sqrt{\left(\frac{t}{2\pi} \right)^d \cdot N_C} \right] - 1}}{0.5} \right).$$

In this paper, (i) the extreme peak height growth behaviour along the $s = 0 + I \cdot t$ line of the Riemann Siegel Z function analogues of

- $\zeta(0.0 + I \cdot t)$ [22-26],
- $L(\chi_3(2, \cdot), 0.0 + I \cdot t)$ [27],
- $L(\chi_4(3, \cdot), 0.0 + I \cdot t)$ [27],
- $L(\chi_5(4, \cdot), 0.0 + I \cdot t)$ [27],
- the dual pair $L(\chi_5(3, \cdot), 0.0 + I \cdot t)$ [27], $L(\chi_5(2, \cdot), 0.0 + I \cdot t)$ [27] and their Davenport-Heilbronn function counterparts $f_1(0.0 + I \cdot t)$ and $f_2(0.0 + I \cdot t)$ [28-30],
- $L(\chi_7(6, \cdot), 0.0 + I \cdot t)$ [27],
- the dual pair $L(\chi_7(5, \cdot), 0.0 + I \cdot t)$ [27], $L(\chi_7(3, \cdot), 0.0 + I \cdot t)$ [27], and their Davenport-Heilbronn function counterparts $f_3(0.0 + I \cdot t)$ and $f_4(0.0 + I \cdot t)$ [31],
- $L(\chi_{37}(36, \cdot), 0.0 + I \cdot t)$ [27],
- $L(\chi_{51}(50, \cdot), 0.0 + I \cdot t)$ [27],
- $L(\chi_{5704}(2851, \cdot), 0.0 + I \cdot t)$ [27],
- $L(\chi_{6389}(6388, \cdot), 0.0 + I \cdot t)$ [27]

and $s = 5.5 + I \cdot t$ line of the Riemann Siegel Z function analogue of

- the Ramanujan Tau L-function 2-1-1.1-c11-0-0 which has degree 2 and character $\chi_1(1, \cdot)$ [27,32].

are compared on the lower boundary of their critical strip to further understand the growth behaviour of L-functions and the Riemann Zeta Riemann Siegel Z function, (ii) Box Cox transformation modelling (and simpler $\log(|Z|)$ vs $\log(t)$ modelling) of the Riemann Zeta function extreme peak Riemann Siegel Z function growth is conducted to provide a trend growth prediction on the critical strip lower boundary (i.e. extreme peak heights occur either side of the trend growth line) and (iii) the fitted $s = 0 + I \cdot t$ trend growth estimate is used with the Riemann Zeta functional equation to estimate the $s = 1 + I \cdot t$ (critical strip upper boundary) extreme peak trend behaviour for $t < 1e40$.

In addition, the extreme peak height growth behaviour along the critical line $s = 0.5 + I \cdot t$ line of the Riemann Siegel Z function analogues of $L(\chi_{5704}(2851, \cdot), 0.0 + I \cdot t)$ [27], $L(\chi_{6389}(6388, \cdot), 0.0 + I \cdot t)$ [27] and the critical line $s = 6 + I \cdot t$ line (for a wider distribution of peaks) of the Riemann Siegel Z function analogue of the Ramanujan Tau L-function 2-1-1.1-c11-0-0 [27,32] are included to supplement the previous results contained in [21] on the critical line behaviour of L-function extreme peak height growth behaviour. In contrast to [21], where for the 2nd degree Ramanujan Tau L-function just methods (i-iii) were employed as the interval investigated was only $t < 6600$, in this paper, extreme peaks for the wider interval $t < 1e13$ were identified for the 2nd degree Ramanujan Tau L-function and it has been empirically observed when using the spectrally filtered heavily truncated Euler Product method that $\beta_{\text{TauL}} = 100, 10, 1$ hyperparameter settings for the 2nd degree Ramanujan Tau L-function extreme peak estimates were achieving $\sim 1\%$, $\sim 10\%$, weaker accuracy respectively on its critical line using method (iii) for benchmarking.

3



Combined comparison of results for 1st degree L-functions

In figures 1 and 2, the extreme peak growth behaviour of the above 1st degree L-functions (all with dirichlet character magnitudes $|\chi| = 0, 1$) are compared in the range $0 < t < 1e20$.

- When plotting $\log(|Z|)$ vs $\log(t)$ in figure 1 it can be seen that each L-function varies in their growth rates of the extreme peak heights as a function of $\log(t)$. They are similar but different.
- However in figure 2 by

1. changing the x-axis from $\log(t) \rightarrow \log \left[\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C} \right]$ AND

2a. adjusting the y axis $\log(|Z|) \rightarrow \log(|Z| \cdot \frac{N_C}{N_{\chi \neq 0}})$ for L-functions OR

2b. adjusting the y axis $\log(|Z|) \rightarrow \log \left(|Z| \cdot \left\{ \frac{N_C}{\sum_{k=1}^{N_C} |\chi_k|} \right\} \right)$ for N_C -periodic Davenport-Heilbronn functions

where N_C is the L-function conductor, $N_{\chi \neq 0}$ is the number of non-zero dirichlet characters Mod N_C and $|\chi_k|$ is the magnitude of the periodic dirichlet characters, it can be seen that under these rescaled xy axes the 1st degree L-functions growth rates coincide reasonably well with the (absolute value $|Z|$) Riemann Zeta Riemann Siegel Z function behaviour as $t \rightarrow \infty$.

Such behaviour (transformation of different L-functions to a universal reference frame) relies on the strong piecewise interval behaviour $\left(N_1 = \sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C} \right)$ of L-functions with $\{0,1\}$ magnitude dirichlet characters depending on the L-function conductor value N_C & degree d in the Riemann Siegel formula and the bounded (Dirac comb) nature occurring at extreme 1st-degree L-function peaks (i.e. $|\Im(\frac{\chi}{n^s})| \lesssim \pm|\chi| \lesssim \pm 1$ for $\chi \neq 0$ and $\frac{\chi}{n^s} = 0$ for $\chi = 0$ for the low lying Dirichlet series terms $\frac{\chi L}{n^s}$).

That is, for the subset of L-functions with dirichlet characters $|\chi| = 0, 1$ therefore each $|\Im(\frac{\chi}{n^s})| \lesssim \pm|\chi| \lesssim \pm 1$ for $\chi \neq 0$ and so each dirichlet term only contribute a maximum of 1 to $|Z|$ summing up to the dirichlet series truncation point $N_1 = \sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}$ (or $N_2 = 2 \cdot \left(\frac{t}{2\pi}\right)^d \cdot N_C$) depending on which approximation approach i.e., methods (iii), (v) (or (ii), (iii)) mentioned in the introduction are used. Hence the extreme Riemann Siegel function peak value for the subset of L-functions with dirichlet characters $|\chi| = 0, 1$ will differ in first order from the Riemann Zeta function behaviour by the proportion of dirichlet characters that are non-zero when compared on a common x-axis scale i.e. $\log(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C})$.

By close examination of the ordered source data or by magnifying figures 1 and 2 it can be seen that the $s = 0 + I \cdot t$ Riemann Siegel $|Z|$ function extreme peak heights are not strictly monotonically increasing with t but the trend growth component is clearly dominant. Model fitting results of the trend growth rate of the extreme peak height will be presented later in this paper.

Also present in figures 1 & 2 are some low height peaks (i.e., not extreme) for $L(\chi_5(3, \cdot), 0.0 + I \cdot t)$ & $L(\chi_5(2, \cdot), 0.0 + I \cdot t)$ and $(L(\chi_7(5, \cdot), 0.0 + I \cdot t) \& L(\chi_7(3, \cdot), 0.0 + I \cdot t))$ L-functions because the search method for predicting extreme peaks for these dual pair L-functions relied on pre-screen calculations of the behaviour of the Davenport-Heilbronn functions $f_1(0.0 + I \cdot t)$ and $(f_4(0.0 + I \cdot t))$ respectively. That is, half each of the $f_1(0.0 + I \cdot t)$ extreme peaks are due to $L(\chi_5(3, \cdot), 0.0 + I \cdot t)$ ($L(\chi_5(2, \cdot), 0.0 + I \cdot t)$) extreme peaks and likewise half each of the $f_4(0.0 + I \cdot t)$ extreme peaks are due to $L(\chi_7(5, \cdot), 0.0 + I \cdot t)$ ($L(\chi_7(3, \cdot), 0.0 + I \cdot t)$) extreme peaks. So when $L(\chi_5(3, \cdot), 0.0 + I \cdot t)$ ($L(\chi_5(2, \cdot), 0.0 + I \cdot t)$) grid search or euler product spectrum calculations are run over the $f_1(0.0 + I \cdot t)$ identified extreme peak locations only half of the $L(\chi_5(3, \cdot), 0.0 + I \cdot t)$ ($L(\chi_5(2, \cdot), 0.0 + I \cdot t)$) peaks are extreme (and likewise in figure 10 when $L(\chi_7(5, \cdot), 0.0 + I \cdot t)$ ($L(\chi_7(3, \cdot), 0.0 + I \cdot t)$) grid search or euler product spectrum calculations are run over the $f_4(0.0 + I \cdot t)$ extreme peaks locations).

Scaled growth behaviour of the $s=0+it$ largest Riemann Siegel $|Z|$ peaks for several 1st degree L-functions using a $\log(\sqrt{(t/2/\pi)^1 \cdot N_c})$ x-axis and (inversely) adjusting the y axis for the proportion of non-zero dirichlet characters. The y-axis adjustment for the Davenport-Heilbronn counterparts uses the sum of their non-zero dirichlet characters and N_c .




Figure 2: Rescaled XY axes overlay of the growth behaviour of extreme Riemann Siegel Z function peak heights $|Z|$ for different 1st degree L-functions in the interval $t < 1e20$. The $|Z|$ values on the Y axis are scaled by the ratio of the (L-function conductor value)/(number of non-zero dirichlet characters mod N_c) i.e. $\log(|Z| \cdot \frac{N_c}{N_{\chi \neq 0}})$. (For the 5-periodic Davenport-Heilbronn functions $\log(|Z| \cdot \sum_{k=1}^5 \frac{5}{|\chi_k|})$ is used for Y axis rescaling.) The t co-ordinate on the X axis is transformed to the first quiescent region value of the L-function i.e. $\log(\sqrt{(\frac{t}{2\pi})^1 \cdot N_c})$.

Individual comparison of results for 1st degree L-functions

Searching and identifying candidate locations of extreme Riemann Siegel $|Z|$ function peaks for L-functions

In similar manner to [19-21], for each given L-function (and Davenport-Heilbronn counterpart) extreme Riemann Siegel $|Z|$ function peaks in the range $0 < t < 1e30$ on the critical line using for progressively higher t values (i) available high precision L-function calculations, (ii) high precision tapered Riemann Zeta dirichlet series estimates at the second quiescent region $N_2 = 2 * (\frac{t}{2\pi})^d * N_C$ (iii) multiple decimal place accuracy tapered first order Riemann Siegel formula estimates at the first quiescent region $N_1 = \sqrt{(\frac{t}{2\pi})^d * N_C}$, (iv) multiple decimal place accuracy spectrally filtered Euler Product analogues of first order Riemann Siegel formula as well as (vi) preliminary peak height estimates using a spectrally filtered heavily truncated Euler Product method $N_{sub1} = \frac{N_C}{2} * \left(\sqrt{(\frac{t}{2\pi})^d * N_C} \right)^{0.5}$. Again with overlap of adjoining methods was included to validate calculation performance. Following [21], $\beta = N_C$ rather than $\beta = 1$ was used for the heavily truncated N_{sub1} method.

1. To identify extreme Riemann Siegel $|Z|$ function peak locations for each given L-function,

1. a grid search $\Delta = 0.05$ for $t < 26^2 \cdot (2 \cdot \pi)^d / N_C$ within each $t = [N^2 \cdot (2 \cdot \pi)^d / N_C, (N+1)^2 \cdot (2 \cdot \pi)^d / N_C)$ interval using estimation methods (i) is conducted with some manual cleanup for peaks on the boundary edges between piecewise intervals
2. a grid search $\Delta = 0.05$ for $26^2 \cdot (2 \cdot \pi)^d / N_C \leq t < 65^2 \cdot (2 \cdot \pi)^d / N_C$ within each $t = [N^2 \cdot (2 \cdot \pi)^d / N_C, (N+1)^2 \cdot (2 \cdot \pi)^d / N_C)$ interval using estimation method (ii) with a method (i) calculation for the final identified peak location
3. a grid search $\Delta = 0.1$ for $65^2 \cdot (2 \cdot \pi)^d / N_C \leq t < 1001^2 \cdot (2 \cdot \pi)^d / N_C$ within each $t = [N^2 \cdot (2 \cdot \pi)^d / N_C, (N+1)^2 \cdot (2 \cdot \pi)^d / N_C)$ interval using estimation method (iii) with a method (ii) calculation for the final identified peak location
4. using LLL algorithm predictions for extreme peak locations

(I) a short interval grid search $\Delta = 0.05$ over a $\delta t = \pm 0.25$ interval about the LLL predicted location using estimation method (iii) is conducted (with some use of method (ii) for lower t values) and/or

(II) a short 256 point fourier grid search $\Delta = 0.05$ over a $\delta t = [-6.35, 6.4]$ interval about the LLL predicted location using estimation method (iv) (spectral filtered Euler Product N_1 truncation) often conducted with a method (iii) calculation for the final identified peak location

(III) for $t \gtrsim 10^{15}$ a short 256 point fourier grid search $\Delta = 0.05$ over a $\delta t = [-6.35, 6.4]$ interval about the LLL predicted location using preliminary estimation method (v) (spectral filtered Euler Product with heavy N_{1sub} truncation)

2. Adjusting the LLL search algorithm [33] constraints for periodic 1st degree L-functions with non-zero dirichlet characters with magnitudes of order unity ($|\chi| = 1$)

The LLL search algorithm for each investigated periodic 1st degree L-function employed the $\frac{t_{peak} * \log(k \bmod N_C = \{0, \dots, N_C - 1\})}{2\pi}$ constraints identified by [21] which appears to be related to the denominator of the fractional exponents of the $(N_C - 1)$ th cyclotomic polynomial roots of unity plus the N_C th constraint $\frac{t_{peak} * \log(k \bmod N_C = 0)}{2\pi} = 0$ (where in practice, the last constraint drops particular integers from the LLL algorithm calculation).

The above prescription does successfully identify candidate locations of extreme peak heights. However, that does not mean the above prescription definitively describe all constraints for possible extreme peaks nor will an actual implemented LLL algorithm report all possible instances of extreme peak locations.

Box Cox transformation model of linear regression fit of trend growth in extreme $|Z|$ peak height for 1st degree periodic L-functions along $s = 0 + I \cdot t$

Box Cox transformation modelling [37,38] is conducted to approximate the non-linearity in the growth curve in figure 2. Under the suspicion that the non-linearity in figure 2 is approximately of the form $\log \left(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C} \right)$ the input linear regression model and the Box Cox transformation model output diagnostics in figure 3 both given below provides a reasonable fit of the trend growth behaviour to $\frac{(x^{0.9604378} - 1)}{0.9604378}$ Box Cox transformation behaviour. For example, the intercept and linear regression coefficients are highly statistically significant (and given in equation (1) below) and the mode of the Box Cox transformation fit in figure 3 is $\lambda = 1$ which is evidence for a linear polynomial fit (with respect to $\frac{(\left(\sqrt{\log(\sqrt{(\frac{t}{2\pi})^1 \cdot 1})})^{0.9604378} - 1)}{0.9604378}$) for the Riemann Zeta function extreme peaks shown in figure 2.

Yielding the following phenomenological model of trend growth in extreme $|Z_\zeta(0 + I \cdot t)|$ peak height on the critical line based on high quality data for $t < 1e31$,

$$\log \left[|Z_\zeta(0 + I \cdot t)|_{max} \right]_{trend} \approx 2.1656076 + 1.1459223 \cdot \left(\frac{\left(\log \left[\sqrt{\frac{t}{2\pi}} \right] \right)^{0.9604378} - 1}{0.9604378} \right) \quad (1)$$

which can be potentially generalised based on figure 2, for **1st degree periodic L-functions with non-zero dirichlet characters with magnitudes of order unity** ($|\chi| = 1$) to

$$\log \left[|Z_{L,d=1}(0 + I \cdot t)|_{max} \cdot \frac{N_C}{N_{\chi \neq 0}} \right]_{trend} \lesssim 2.1656076 + 1.1459223 \cdot \left(\frac{\left(\log \left[\sqrt{\left(\frac{t}{2\pi}\right)^1 \cdot N_C} \right] \right)^{0.9604378} - 1}{0.9604378} \right) \quad (2)$$

Extrapolation of the model into the region $1e31 < t < 1e40$ and beyond, however compared to the critical line case [21], really requires higher quality data and/or analytic results for $t > 1e31$ to properly test the veracity of the above trend growth model at higher t .

For instance a heuristic trend growth model that also performs qualitatively well in the region $t < 1e40$ is the simpler parametrisation

$$\log \left[|Z_\zeta(0 + I \cdot t)|_{max} \right]_{trend} \approx 2.1656 + \log \left(\sqrt{\frac{t}{2\pi}} \right) \quad (3)$$

as shown when comparing equations (1) (brown dot-dashed line) and (3) (green dashed line) in figure 4.

Of great interest with respect to the Box Cox model estimate result is that the confidence interval of magnitude of the intercept $2.1656076 \pm 1.96 * 0.0054552$ includes the known value of the exponential generating function for the harmonic numbers [39,40]

$$\sum_{n=1}^{\infty} H_n \frac{1^n}{n!} = e^1 \cdot (\gamma + \ln 1 + \Gamma(0, 1))$$

$$= 2.1653822153... \quad (4)$$

suggesting a measurable Euler-Mascheroni constant contribution to the trend growth rate of extreme $|Z|$ peaks on the Riemann Zeta function $s = 0 + I \cdot t$ line.

However, as presented later in figure 13 when vetting equation (1) or (3) based predictions co-opting the Riemann Zeta functional equation for the estimated trend growth rate against observed $s = 1 + I \cdot t$ extreme peak growth behaviour model equation (3) predictions are unphysical for $s = 1 + I \cdot t$ and model equation (1) is clearly insufficient for extrapolation beyond the fitted region $t < 1e31$.

Linear regression model fit for $\log \left[|Z_{\zeta}(0 + I \cdot t)|_{max} \right]_{trend} \sim \left(\frac{\left(\log \left[\sqrt{\frac{t}{2\pi}} \right] \right)^{0.9604378} - 1}{0.9604378} \right)$

```
## [1] 1
##
## Call:
## lm(formula = yt3 ~ xt3)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.47472 -0.03240  0.01243  0.03783  0.15306
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  2.1656076  0.0054552    397  <2e-16 ***
## xt3          1.1459223  0.0003616   3169  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.06545 on 316 degrees of freedom
## Multiple R-squared:  1, Adjusted R-squared:  1
## F-statistic: 1.004e+07 on 1 and 316 DF, p-value: < 2.2e-16
```

Summary of languages/IDEs used for calculations

The LLL search based algorithm, the high precision L-function calculations, tapered dirichlet series calculations, input discrete fourier spectra samples, the fourier transforms, the spectral filtering, the rescaled conjugate mirror reflection values and the inverse fourier transforms are calculated using Pari-GP [34] and the Box Cox transformation modelling and general graphing is performed using R [35] and RStudio IDE [36].




Figure 3: Box Cox modelling analysis using the MASS package on the growth behaviour of Riemann Zeta function extreme Riemann Siegel Z function peak heights $|Z|$ for in the interval $t < 1e31$. Modelling $Y \sim X$, the input data comprises $Y = \log(|Z|)$ and $X = \frac{(\log(\sqrt{(\frac{t}{2\pi})^d \cdot N_C})^{0.9604378} - 1)}{0.9604378}$. Top panel - wide view of the log-likelihood of the Box Cox model fit supporting $\lambda = 1$ (i.e. linear behaviour with respect to the input regression model), Bottom panel - magnified view about $\lambda = 1$

Individual L-function and Davenport-Heilbronn function results in comparison to $|Z_\zeta(0.5 + I \cdot t)|$

Figures 4-11 below, display for each investigated 1st degree 1 L-function or Davenport-Heilbronn function

1. left column - $\log(|Z_L|)$ vs $\log(t)$ of extreme L-function (or Davenport-Heilbronn function) peak height
2. right column - $\log\left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^{d=1} \cdot N_C}\right)$ of extreme L-function (or Davenport-Heilbronn function) peak height

Figure 4 displays a comparison of $|Z_\zeta|$ behaviour with located extreme Riemann Siegel Z function peaks for

- top row $L(\chi_1(1, .), 0.5 + I \cdot t)$ i.e. $\zeta(0.5 + I \cdot t)$ with $N_C = 1$ and $N_{\chi \neq 0} = 1$,
- middle row $L(\chi_3(2, .), 0.5 + I \cdot t)$ with $N_C = 3$ and $N_{\chi \neq 0} = 2$,

Figure 5 displays a comparison of $|Z_\zeta|$ behaviour with located extreme Riemann Siegel Z function peaks for

- top row $L(\chi_4(3, .), 0.5 + I \cdot t)$ with $N_C = 4$ and $N_{\chi \neq 0} = 2$
- bottom row $L(\chi_5(4, .), 0.5 + I \cdot t)$ with $N_C = 5$ and $N_{\chi \neq 0} = 4$,

Figure 6 displays a comparison of $|Z_\zeta|$ behaviour with located extreme Riemann Siegel Z function peaks for

- top row $L(\chi_5(3, .), 0.5 + I \cdot t)$ with $N_C = 5$ and $N_{\chi \neq 0} = 4$,
- bottom row $L(\chi_5(2, .), 0.5 + I \cdot t)$ with $N_C = 5$ and $N_{\chi \neq 0} = 4$

Figure 7 displays a comparison of $|Z_\zeta|$ behaviour with located extreme Riemann Siegel Z function peaks for

- top row $L(\chi_7(6, .), 0.5 + I \cdot t)$ with $N_C = 7$ and $N_{\chi \neq 0} = 6$,
- middle row $L(\chi_37(36, .), 0.5 + I \cdot t)$ with $N_C = 37$ and $N_{\chi \neq 0} = 36$,

Figure 8 displays a comparison of $|Z_\zeta|$ behaviour with located extreme Riemann Siegel Z function peaks for

- $L(\chi_{51}(50, .), 0.5 + I \cdot t)$ with $N_C = 51$ and $N_{\chi \neq 0} = 32$

Figure 9 displays a comparison of $|Z_\zeta|$ behaviour with located extreme Riemann Siegel Z function peaks for

- top row $f_2(0.5 + I \cdot t)$ a 5-periodic Davenport-Heilbronn function with $N_C = 5$ and $\sum_{k=1}^5 |\chi_k| \approx 9.04$,
- middle row $f_1(0.5 + I \cdot t)$ a 5-periodic Davenport-Heilbronn function with $N_C = 5$ and $\sum_{k=1}^5 |\chi_k| \approx 2.568$

Figure 10 displays a comparison of $|Z_\zeta|$ behaviour with located extreme Riemann Siegel Z function peaks for

- top row $L(\chi_7(5, .), 0.5 + I \cdot t)$ with $N_C = 7$ and $N_{\chi \neq 0} = 6$,
- bottom row $L(\chi_7(3, .), 0.5 + I \cdot t)$ with $N_C = 7$ and $N_{\chi \neq 0} = 6$


Figure 11 displays a comparison of $|Z_\zeta|$ behaviour with located extreme Riemann Siegel Z function peaks for

- top row $f_3(0.5 + I \cdot t)$ a 7-periodic Davenport-Heilbronn function with $N_C = 7$ and $\sum_{k=1}^7 |\chi_k| \approx 7.2077509$,
- bottom row $f_4(0.5 + I \cdot t)$ a 7-periodic Davenport-Heilbronn function with $N_C = 7$ and $\sum_{k=1}^7 |\chi_k| \approx 4.3042576$,


As mentioned earlier also displayed in figure 3 are two models of the trend growth rate given by equations (1) and (3) shown as a brown dot-dashed and green dashed lines respectively.

For equations 4-11, only the Box Cox transformation based trend growth rate model equation (1) is displayed in the right column panels.


**Growth behaviour of the $s=0+It$ line largest Riemann Siegel $|Z|$ peak:
standard $\log(|Z|)$ vs $\log(t)$ plot**



**Growth behaviour of the $s=0+It$ line largest Riemann Siegel $|Z|$ peak:
 $\log(|Z|)$ vs $\log(\sqrt{(t/2/\pi)^{1*1}})$ plot**



**Growth behaviour of the $s=0+It$ line largest L32 $|Z|$ peaks
standard $\log(|Z|)$ vs $\log(t)$ plot**



**Growth behaviour of the $s=0+It$ line largest L32 $|Z|$ peaks
 $\log(|Z|^{3/2})$ vs $\log(\sqrt{(t/2/\pi)^{1*3}})$ plot**






Figure 4: A graphical comparison for each investigated 1st degree L-function or Davenport-Heilbronn function of **left column** - $\log(|Z_L|)$ vs $\log(t)$, **right column** - $\log\left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}\right)$ for top row - $L(\chi_1(1, \cdot), 0.0 + I \cdot t)$ and bottom row - $L(\chi_3(2, \cdot), 0.0 + I \cdot t)$. With two (one) fitted models for extreme peak height trend growth rate shown in the top (bottom) row.


**Growth behaviour of the s=0+It line largest L43 |Z| peaks
standard log(|Z|) vs log(t) plot**



**Growth behaviour of the s=0+It line largest L43 |Z| peaks
log(|Z|*4/2) vs log(sqrt((t/2/pi)^1*4)) plot**



**Growth behaviour of the s=0+It line largest L54 |Z| peaks
standard log(|Z|) vs log(t) plot**



**Growth behaviour of the s=0+It line largest L54 |Z| peaks
log(|Z|*5/4) vs log(sqrt((t/2/pi)^1*5)) plot**






Figure 5: A graphical comparison for each investigated 1st degree L-function or Davenport-Heilbronn function of **left column** - $\log(|Z_L|)$ vs $\log(t)$, **right column** - $\log\left(|Z_L| \cdot \frac{N_C}{N_{2.1656+\log(seq(.1,50,by=.1))_{\chi \neq 0}}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}\right)$ for top row - $L(\chi_4(3, \cdot), 0.0 + I \cdot t)$ and bottom row - $L(\chi_5(4, \cdot), 0.0 + I \cdot t)$. With one conservative fitted model for extreme peak height trend growth rate.


**Growth behaviour of the s=0+It line largest L53 |Z| peaks
standard log(|Z|) vs log(t) plot**



**Growth behaviour of the s=0+It line largest L53 |Z| peaks
log(|Z|*5/4) vs log(sqrt((t/2/pi)^1*5)) plot**



**Growth behaviour of the s=0+It line largest L52 |Z| peaks
standard log(|Z|) vs log(t) plot**



**Growth behaviour of the s=0+It line largest L52 |Z| peaks
log(|Z|*5/4) vs log(sqrt((t/2/pi)^1*5)) plot**






Figure 6: A graphical comparison for each investigated 1st degree L-function or Davenport-Heilbronn function of **left column** - $\log(|Z_L|)$ vs $\log(t)$, **right column** - $\log\left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}\right)$ for top row - $L(\chi_5(3, \cdot), 0.0 + I \cdot t)$ and bottom row - $L(\chi_5(2, \cdot), 0.0 + I \cdot t)$. With one conservative fitted model for extreme peak height trend growth rate.


**Growth behaviour of the s=0+It line largest L76 |Z| peaks
standard log(|Z|) vs log(t) plot**



**Growth behaviour of the s=0+It line largest L76 |Z| peaks
log(|Z|*7/6) vs log(sqrt((t/2/pi)^1*7)) plot**



**Growth behaviour of the s=0+It line largest L37p36 |Z| peaks
standard log(|Z|) vs log(t) plot**



**Growth behaviour of the s=0+It line largest L37p36 |Z| peaks
log(|Z|*37/36) vs log(sqrt((t/2/pi)^1*37)) plot**





Figure 7: A graphical comparison for each investigated 1st degree L-function or Davenport-Heilbronn function of **left column** - $\log(|Z_L|)$ vs $\log(t)$, **right column** - $\log\left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}\right)$ for top row - $L(\chi_7(6, \cdot), 0.0 + I \cdot t)$ and bottom row - $L(\chi_{37}(36, \cdot), 0.0 + I \cdot t)$. With one conservative fitted model for extreme peak height trend growth rate.

**Growth behaviour of the s=0+It line largest L51p50 |Z| peaks
standard log(|Z|) vs log(t) plot**



**Growth behaviour of the s=0+It line largest L51p50 |Z| peaks
log(|Z|*51/32) vs log(sqrt((t/2/pi)^1*51)) plot**






Figure 8: A graphical comparison for each investigated 1st degree L-function or Davenport-Heilbronn function of **left column** - $\log(|Z_L|)$ vs $\log(t)$, **right column** - $\log\left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}\right)$ for $L(\chi_{51}(50, \cdot), 0.0 + I \cdot t)$. With one conservative fitted model for extreme peak height trend growth rate.


**Growth behaviour of the $s=0+it$ line largest DHf2 $|Z|$ peaks
standard $\log(|Z|)$ vs $\log(t)$ plot**



**Growth behaviour of the $s=0+it$ line largest DHf2 $|Z|$ peaks
 $\log(|Z|^5/9.04)$ vs $\log(\sqrt{t/2\pi})^{1*5}$ plot**



**Growth behaviour of the $s=0+it$ line largest DHf1 $|Z|$ peaks
standard $\log(|Z|)$ vs $\log(t)$ plot**



**Growth behaviour of the $s=0+it$ line largest DHf1 $|Z|$ peaks
 $\log(|Z|^5/2.568)$ vs $\log(\sqrt{t/2\pi})^{1*5}$ plot**






Figure 9: A graphical comparison for each 1st degree investigated 1st degree L-function or Davenport-Heilbronn function of **left column** - $\log(|Z_L|)$ vs $\log(t)$, **right column** - $\log\left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d} \cdot N_C\right)$ for top row - $f_2(0.0 + I \cdot t)$ and bottom row - $f_1(0.0 + I \cdot t)$. With one conservative fitted model for extreme peak height trend growth rate.


**Growth behaviour of the s=0+It line largest L75 |Z| peaks
standard log(|Z|) vs log(t) plot**



**Growth behaviour of the s=0+It line largest L75 |Z| peaks
log(|Z|*7/6) vs log(sqrt((t/2/pi)^1*7)) plot**



**Growth behaviour of the s=0+It line largest L73 |Z| peaks
standard log(|Z|) vs log(t) plot**



**Growth behaviour of the s=0+It line largest L73 |Z| peaks
log(|Z|*7/6) vs log(sqrt((t/2/pi)^1*7)) plot**






Figure 10: A graphical comparison for each investigated 1st degree L-function or Davenport-Heilbronn function of **left column** - $\log(|Z_L|)$ vs $\log(t)$, **right column** - $\log\left(|Z_L| \cdot \frac{N_C}{N_{x \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}\right)$ for top row - $L(\chi_7(5, \cdot), 0.0 + I \cdot t)$ and bottom row - $L(\chi_7(3, \cdot), 0.0 + I \cdot t)$. With one conservative fitted model for extreme peak height trend growth rate.


**Growth behaviour of the s=0+It line largest DHf3 |Z| peaks
standard log(|Z|) vs log(t) plot**



**Growth behaviour of the s=0+It line largest DHf3 |Z| peaks
log(|Z|*7/7.2077509) vs log(sqrt((t/2/pi)^1*7)) plot**



**Growth behaviour of the s=0+It line largest DHf4 |Z| peaks
standard log(|Z|) vs log(t) plot**



**Growth behaviour of the s=0+It line largest DHf4 |Z| peaks
log(|Z|*7/4.3042576) vs log(sqrt((t/2/pi)^1*7)) plot**




Figure 11: A graphical comparison for each investigated 1st degree L-function or Davenport-Heilbronn function of **left column** - $\log(|Z_L|)$ vs $\log(t)$, **right column** - $\log\left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}\right)$ for top row - $f_3(0.0 + I \cdot t)$, and bottom row - $f_4(0.0 + I \cdot t)$. With one conservative fitted model for extreme peak height trend growth rate.

Highlights part 1

For the L- functions with only dirichlet characters of unity $|\chi| = 1$ or zero $|\chi| = 0$: figures 4, 5, 6, 7, 8 and 10:

In the left column, there is divergence between the L-function and Riemann Zeta $|Z|$ extreme peak growth behaviour.

In the right column, except for the lowest t values the re-scaled L-function extreme growth behaviour to first order, tends to lie reasonably adjacent to the Riemann Zeta $|Z|$ extreme peak growth behaviour. While the LLL search results $t > 1e5$ do find extreme peaks it doesn't collect all of them as the grid search does for lower t so there is still room for re-scaled extreme L-function peaks to get closer to the Riemann Zeta $|Z|$ behaviour.

The re-scaling of the y-axis is a simple argument using the two bits of information (i) that the Riemann Zeta has only non-zero characters of unity ($\chi = 1$) while the other presented 1st degree L-functions have ($|\chi| = 0$ OR 1) and (ii) for extreme peak locations at high t , the **imaginary component** of each lowest lying dirichlet series term contributes approximately 1 to the dirichlet series sum.

The re-scaling of the x-axis is a recognition that each L-function has an intrinsic length $N_1 = \sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}$ and in properly comparing the growth behaviour of two L-functions the difference in their N_1 values must be taken into account. (The intrinsic length $N_2 = 2 \cdot \left(\frac{t}{2\pi}\right)^d \cdot N_C$ also exists and tapered Dirichlet series sums using this length can approach high precision L-function calculation performance for some L-functions.)

For the Davenport-Heilbronn functions in figures 9 and 11:

In the left column, there is divergence between the Davenport-Heilbronn and Riemann Zeta $|Z|$ extreme peak growth behaviour.

In the right columns (comparing to the pure L-function figures), except for the lowest t values there is weak but possible evidence that the re-scaled Davenport-Heilbronn extreme growth behaviour **may be lower than** the Riemann Zeta $|Z|$ extreme peak growth behaviour. In this case, the attempted y-axis re-scaling uses the $\sum_{k=1}^{N_C} |\chi_k|$ of the (absolute value of the) real coefficients of the known Davenport-Heilbronn dirichlet series.

Extreme peak height results displaying both (i) lower boundary of critical strip and (ii) critical line behaviour

Individual comparison of Ramanujan Tau L-function Riemann Siegel Z function extreme peak behaviour for $s = 5.5 + I \cdot t$ and $s = 6 + I \cdot t$ lines in the complex plane to the $|Z_\zeta(0 + I \cdot t)|$ and $|Z_\zeta(0.5 + I \cdot t)|$ extreme peak behaviour respectively.

To identify extreme Riemann Siegel $|Z|$ function peak locations for the Ramanujan Tau L-function,

1. a grid search $\Delta = 0.05$ for $t < 65 \cdot (2 \cdot \pi)/N_C$ within each $t = [N \cdot (2 \cdot \pi)/N_C, (N + 1) \cdot (2 \cdot \pi)/N_C)$ interval using estimation method (i) is conducted with some manual cleanup for peaks on the boundary edges between piecewise intervals
2. a grid search $\Delta = 0.05$ for $65 \cdot (2 \cdot \pi)/N_C \leq t < 650 \cdot (2 \cdot \pi)/N_C$ within each $t = [N \cdot (2 \cdot \pi)/N_C, (N + 1) \cdot (2 \cdot \pi)/N_C)$ interval using estimation method (iii) with a method (i) calculation for the final identified peak location. Method (ii) using 128 point tapering gave poorer results than method (iii) using method (i) as reference so method (ii) was dropped for this L-function. Some investigation suggested that greater tapering is required with method (ii) for the Ramanujan tau L-function due to the non-periodic nature of its dirichlet character magnitudes.

3. a grid search $\Delta = 0.05$ for $650 \cdot (2 \cdot \pi)/N_C \leq t < 10001 \cdot (2 \cdot \pi)/N_C$ within each $t = [N \cdot (2 \cdot \pi)/N_C, (N + 1) \cdot (2 \cdot \pi)/N_C)$ interval using estimation method (iii).

4. using LLL algorithm predictions for extreme peak locations

(I) a short interval grid search $\Delta = 0.05$ over a $\delta t = \pm 0.25$ interval about the LLL predicted location using estimation method (iii) is conducted

(II) a longer 32768 point fourier grid search $\Delta = 0.05$ over a $\delta t = [-819.15, 819.2]$ interval about the LLL predicted location using estimation method (iv) (spectral filtered Euler Product N_1 truncation) often conducted with a method (iii) calculation for the final identified peak location

(III) for $t \gtrsim 10^6$ a longer 32768 point fourier grid search $\Delta = 0.05$ over a $\delta t = [-819.15, 819.2]$ interval about the LLL predicted location using preliminary estimation method (v) (spectral filtered Euler Product with heavy N_{1sub} truncation) where $\beta_{tau} = 100, 10, 1$ hyperparameters values were used depending on the t value. Some peak heights were recalculated for $\beta_{tau} = 10$ results using $\beta_{tau} = 100$ (and $\beta_{tau} = 1$ results using $\beta_{tau} = 10$) as the higher accuracy method.

Adjusting the LLL search algorithm [33] constraints for the Ramanujan Tau L-function

Attempts at using the LLL search algorithm for the Ramanujan Tau L-function employed a variety of modifications to $\frac{t_{peak} \cdot \log(k)}{2\pi}$ constraints mainly based on known $\arccos\left(\frac{tau(k)}{k^{5.5}} \cdot \frac{1}{2}\right)$ behaviour arising from Ramanujan's third conjecture $|\tau(p)| \leq 2p^{\frac{11}{2}}$ proved by Deligne [41]. However, the performance at identifying extreme peaks was clearly not as successful as for periodic 1st degree L-functions and hence drove the dramatic increase in the analysed fourier spectrum width (from $\delta t = [-6.35, 6.4]$ with periodic 1st degree L-functions to $\delta t = [-819.15, 819.2]$ with the Ramanujan Tau L-function) which helped locate nearby larger peaks.

Ramanujan Tau L-function results

Figure 12 displays a comparison of $|Z_\zeta|$ behaviour with located extreme Riemann Siegel Z function peaks for top row - the lower boundary of the critical strip $s = 5.5 + I \cdot t$ and

bottom row - the critical line $s = 6 + I \cdot t$

versus the behaviour for the lower boundary of the Riemann Zeta function critical strip and its critical line respectively.

1. left column - $\log(|L_\tau(s)|)$ vs $\log(t)$

2. right column - $\log\left(|Z_L(s)|\tau \cdot \frac{N_C}{N_{\chi \neq 0}}\right)$ vs $\log\left(\frac{1}{d} \cdot \log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d}\right) + \log(\sqrt{N_C})\right)$

where $d=2$, $N_C=1$ and $\frac{N_C}{N_{\chi \neq 0}} := \frac{1}{N_{\chi \neq 0} := 1} := 1$ is used for the Ramanujan Tau L-function.

For the top row,

- the Box Cox transformation based trend growth rate model given in equation (1) of this current paper is displayed shown, as brown dot-dashed line.
- a further transformation $\frac{2.1656}{2} + \frac{\log\left(|Z_L(s)|\tau \cdot \frac{N_C}{N_{\chi \neq 0}}\right)}{2}$ is shown for some of the data points as green triangles.

For the bottom row,

- the Box Cox transformation based fitted trend growth rate model ($t < 1e31$) from [21] for the critical line of the Riemann Zeta function

$$\log \left[|Z_{\zeta}(0.5 + I \cdot t)|_{max} \right]_{trend} \approx 1.274014 + 0.866363 \cdot \left(\frac{\sqrt{\log \left[\sqrt{\frac{t}{2\pi}} \right] - 1}}{0.5} \right) \quad (5)$$

is displayed, shown as brown dot-dashed line.

The bottom row panels in figure 12 contain a much wider set of extreme peaks (and lesser peaks) than was reported in [21].

Highlights part 2

For the Ramanujan Tau L-function

Examining the top row panels of figure 12,

- the growth rate of extreme peaks on the lower boundary of the Ramanujan Tau L-function critical strip clearly exceeds the growth rate of extreme peaks on the lower boundary of the Riemann Zeta function critical strip.
- the green triangle behaviour in the righthand panel of the top row estimates the growth rate of $|Z|$ extreme peaks along $s = 5.5 + I \cdot t$ for the Ramanujan Tau L-function as roughly the square of the $|Z_{zeta}(s = 0 + I \cdot t)|$ extreme peak growth rate.

Examining the bottom row panels of figure 12,

- the wider range of extreme peaks investigated in this paper for the Ramanujan Tau L-function on its critical line continues to exhibit to first order the same extreme peak growth rate as $|Z_{zeta}(s = 0.5 + I \cdot t)|$







Figure 12: A graphical comparison for the 2nd degree Ramanujan's Tau L-function and the Riemann Zeta function extreme peaks of **left panel** - $\log(|Z_L|)$ vs $\log(t)$, **right panel** - $\log(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}})$ vs $\frac{1}{d} \cdot \log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d}\right) + \log(\sqrt{N_C})$ for **top row** - $s = 5.5 + I \cdot t$ (zero line), and **bottom row** - $s = 6 + I \cdot t$ (critical line). The x axis rescaling in the right panel is different in figure 12 compared to figures 4-11 because the two L-functions being compared in figure 12 are of different degrees. With one a different fitted model for extreme peak height trend growth rate depending on $\text{Re}(s)$. Riemann Zeta function behaviour shown in gray for known peak heights (and cyan for high t preliminary peak height estimates).

Large N_C value L-function extreme peak height behaviour

Figures 13 & 14 below, display for two investigated large N_C value 1st degree 1 L-functions

1. left column - $\log(|Z_L|)$ vs $\log(t)$
2. right column - $\log\left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^{d=1} \cdot N_C}\right)$

of extreme L-function peak height respectively.

Figure 13 displays a comparison of $|Z_\zeta|$ behaviour with located extreme Riemann Siegel Z function peaks for

- top row $L(\chi_{5704}(2851, .), 0 + I \cdot t)$,
- bottom row $L(\chi_{5704}(2851, .), 0.5 + I \cdot t)$

with $N_C = 5704$ and $N_{\chi \neq 0} = 2640$ [27,34].

Figure 14 displays a comparison of $|Z_\zeta|$ behaviour with located extreme Riemann Siegel Z function peaks for

- top row $L(\chi_{6389}(6388, .), 0 + I \cdot t)$,
- bottom row $L(\chi_{6389}(6388, .), 0.5 + I \cdot t)$

with $N_C = 6389$ and $N_{\chi \neq 0} = 6388$ [27,34].

For the top row,

- the Box Cox transformation based trend growth rate model given in equation (1) of this current paper is displayed shown, as brown dot-dashed line.

For the bottom row,

- the Box Cox transformation based fitted trend growth rate model ($t < 1e31$) from [21] for the critical line of the Riemann Zeta function

$$\log \left[|Z_\zeta(0.5 + I \cdot t)|_{max} \right]_{trend} \approx 1.274014 + 0.866363 \cdot \left(\frac{\sqrt{\log \left[\sqrt{\frac{t}{2\pi}} \right] - 1}}{0.5} \right) \quad (6)$$

is displayed, shown as brown dot-dashed line.

Highlights part 3

For the example large N_C L-functions

Examining the top row panels of figures 13 & 14,

- the growth rate of extreme peaks at low t , on the lower boundary of the large N_C L-function critical strip clearly exceeds the growth rate of extreme peaks on the lower boundary of the Riemann Zeta function critical strip.
- the xy scale transformation $\log\left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^{d=1} \cdot N_C}\right)$ in the right panel produces to first order a universal growth rate behaviour for the two large N_C L-functions.

It is important to remember that on the lower boundary of the critical strip the dirichlet series terms have no decay component to their magnitude since

$$\left| \frac{\chi_k}{k^s} \right| \rightarrow \left| \frac{\chi_k}{k^{I \cdot t}} \right| \rightarrow |\chi_k| \text{ when } s = 0 + I \cdot t \quad (7)$$

Examining the bottom row panels of figure 13 & 14

- the growth rate of extreme peaks at low t , on the lower boundary of the large N_C L-function critical strip clearly exceeds the growth rate of extreme peaks on the lower boundary of the Riemann Zeta function critical strip.
- for $L(\chi_{6389}(6388, \cdot), 0.5 + I \cdot t)$ in figure 14, the xy scale transformation $\log \left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}} \right)$ vs $\log \left(\sqrt{\left(\frac{t}{2\pi}\right)^{d=1} \cdot N_C} \right)$ in the right panel produces similar behaviour to the low N_C 1st degree L-functions whereby many extreme peaks lie very close to the Riemann Zeta extreme peak behaviour (in the transformed scale).
- for $L(\chi_{5704}(2851, \cdot), 0.5 + I \cdot t)$ in figure 13, the xy scale transformation $\log \left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}} \right)$ vs $\log \left(\sqrt{\left(\frac{t}{2\pi}\right)^{d=1} \cdot N_C} \right)$ in the right panel displays very few extreme peaks which lie very close to the Riemann Zeta extreme peak behaviour (in the transformed scale).

Since the $s = 0 + I \cdot t$ extreme peak behaviour growth rate behaviour (in the transformed universal growth chart) is similar for $L(\chi_{6389}(6388, \cdot), 0.5 + I \cdot t)$ and $L(\chi_{5704}(2851, \cdot), 0.5 + I \cdot t)$ the most likely reason for their different behaviour (in the transformed universal growth chart) for $s = 0.5 + I \cdot t$ is the combined impact of large number of dirichlet terms with $\chi_k = 0$ and decay in $|\chi_k|$ induced by the real component of s .

That is, on the critical line

$$\left| \frac{\chi_k}{k^s} \right| \rightarrow \left| \frac{\chi_k}{k^{0.5+I \cdot t}} \right| \rightarrow \frac{1}{\sqrt{k}} \cdot |\chi_k| \text{ when } s = 0.5 + I \cdot t \quad (8)$$

So for $L(\chi_{6389}(6388, \cdot), 0.5 + I \cdot t)$, $6388/6389 > 0.9998$ dirichlet series terms are of the magnitude $\frac{1}{\sqrt{k}} \cdot |\chi_k| = 1$ whereas for $L(\chi_{5704}(2851, \cdot), 0.5 + I \cdot t)$ only $2640/6389 \approx 0.4132$ dirichlet series terms are of the magnitude $\frac{1}{\sqrt{k}} \cdot |\chi_k| = 1$ the rest of the dirichlet series terms being zero. Where the simple y-axis universal growth scaling $\log \left(|Z_L| \cdot \frac{N_C}{N_{\chi \neq 0}} \right)$ in the right columns doesn't appear to smartly address decay factor effect also occurring on the critical line it is sufficient (to first order) on the lower boundary of the critical strip. The empirical evidence displayed however supports that (equation (2) of [21])

$$\log \left[|Z_L(k + I \cdot t)|_{max} \cdot \frac{N_C}{N_{\chi \neq 0}} \right]_{trend} \lesssim 1.274014 + 0.866363 \cdot \left(\frac{\sqrt{\log \left[\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C} \right] - 1}}{0.5} \right)$$

remains a useful approximate extreme peak height trend growth upper bound for large N_C 1st degree L-functions on the critical line.




Figure 13: A graphical comparison for the L5704p2851 1st degree L-functions and the Riemann Zeta function extreme peaks of **left panel** - $\log(|Z_L|)$ vs $\log(t)$, **right panel** - $\log\left(|Z_L| \cdot \frac{N_C}{N_{x \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}\right)$ for **top row** - $s = 0 + I \cdot t$ (zero line), and **bottom row** - $s = 0.5 + I \cdot t$ (critical line). The x axis rescaling in the right panel is the same in figure 14 compared to figures 4-11. With one a different fitted model for extreme peak height trend growth rate depending on $\text{Re}(s)$. Riemann Zeta function behaviour shown in gray for known peak heights (and cyan for high t preliminary peak height estimates).




Figure 14: A graphical comparison for the L6389p6388 1st degree L-functions and the Riemann Zeta function extreme peaks of **left panel** - $\log(|Z_L|)$ vs $\log(t)$, **right panel** - $\log\left(|Z_L| \cdot \frac{N_C}{N_{x \neq 0}}\right)$ vs $\log\left(\sqrt{\left(\frac{t}{2\pi}\right)^d \cdot N_C}\right)$ for **top row** - $s = 0 + I \cdot t$ (zero line), and **bottom row** - $s = 0.5 + I \cdot t$ (critical line). The x axis rescaling in the right panel is the same in figure 15 compared to figures 4-11. With one a different fitted model for extreme peak height trend growth rate depending on $\text{Re}(s)$. Riemann Zeta function behaviour shown in gray for known peak heights (and cyan for high t preliminary peak height estimates).

Testing equation (1) and (3) trend model estimates by assessing their performance at predicting $s = 1 + I \cdot t$ behaviour

A second available method of assessing equation (1) and (3) trend model performance is by co-opting the Riemann Zeta functional equation to extract $s = 1 + I \cdot t$ extreme peak trend behaviour predictions and compare such predictions to observed $s = 1 + I \cdot t$ behaviour. Figure 15 displays this comparison as well as mapping the exact $s = 0 + I \cdot t$ peak heights to the $s = 1 + I \cdot t$ line shown as gray dots.

Given the Riemann Zeta functional equation [22-26]

$$\zeta(s) = \chi(s)\zeta(1-s) \quad (9)$$

$$= 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s) \quad (10)$$

$$\equiv e^{\log[\chi(s)]} \zeta(1-s) \quad (11)$$

$$= e^{\left((s-1/2) \cdot \log[\pi i] + \log \Gamma\left[\left(\frac{1-s}{2}\right)\right] - \log\left[\Gamma\left(\frac{s}{2}\right)\right]\right)} \zeta(1-s) \quad (12)$$

$$\equiv e^{-2\theta_{ext}(s)} \zeta(1-s) \quad (13)$$

where

$$\chi(s) = e^{\left((s-1/2) \cdot \log[\pi i] + \log \Gamma\left[\left(\frac{1-s}{2}\right)\right] - \log\left[\Gamma\left(\frac{s}{2}\right)\right]\right)} \quad (14)$$

from equation (12) provides the most numerically stable version of calculating $\chi(z)$ for the Pari-GP [34] calculations performed in this paper.

Transformation of equations (1) and (3) to $s = 1 + I \cdot t$ predictions involves the mapping

$$\left[|Z_\zeta(1 + I \cdot t)|_{max} \right]_{trend} \sim |\chi(1 + I \cdot t)| \cdot \exp \left[2.1656076 + 1.1459223 \cdot \left(\frac{\left(\log \left[\sqrt{\frac{t}{2\pi}} \right] \right)^{0.9604378} - 1}{0.9604378} \right) \right] \quad (15)$$

and

$$\left[|Z_\zeta(1 + I \cdot t)|_{max} \right]_{trend} \sim |\chi(1 + I \cdot t)| \cdot \exp \left[2.1656 + \log \left(\sqrt{\frac{t}{2\pi}} \right) \right] \quad (16)$$

as shown when comparing equations (1) (brown dot-dashed line) and (3) (green dashed line) based estimates to predict $s = 1 + I \cdot t$ in figure 15.

Highlights part 4

1. The close agreement between the gray dots and the coloured data points shows that the $s = 0 + I \cdot t$ peak heights are accurately mapped to the $s = 1 + I \cdot t$ peak heights via the functional equation behaviour.
2. Equation (15) shows reasonable performance as a trend estimate for $t < 1e18$ but is inaccurate for $t > 1e18$

3. Equation (16) is an unphysical trend estimate predicting a constant extreme peak height for the $s = 1 + I \cdot t$ line

Numerically it is easy to see that

$$|\chi(1 + I \cdot t)| \rightarrow \frac{1}{\sqrt{\frac{t}{2\pi}}} \quad \text{for } t > \pi \quad (17)$$

with four decimal place accuracy and rapidly improving.

Hence re-examining equation (16)

$$\begin{aligned} \left[|Z_\zeta(1 + I \cdot t)|_{max} \right]_{\text{trend}} &\sim |\chi(1 + I \cdot t)| \cdot \exp \left[2.1656 + \log \left(\sqrt{\frac{t}{2\pi}} \right) \right] \\ &\rightarrow \frac{1}{\sqrt{\frac{t}{2\pi}}} \cdot \exp[2.1656] \cdot \sqrt{\frac{t}{2\pi}} \quad \text{for } t > \pi \quad (18) \\ &= \exp[2.1656] \quad \text{for } t > \pi \quad (19) \end{aligned}$$

shows that constant amplitude behaviour of extreme peaks on $s = 1 + I \cdot t$ is the predicted (unphysical) behaviour of equation (3) when mapped to the $s = 1 + I \cdot t$ line.

Assessing self-dual 1st degree L-function and Davenport-Heilbronn mapped extreme peak height $s = 1 + I \cdot t$ behaviour from $s = 0 + I \cdot t$ results using the L-function's functional equation

Finally figure 16 maps the $s = 0 + I \cdot t$ $|Z|$ data in figure 1 to the $s = 1 + I \cdot t$ line using the functional equations of the various **self-dual** L-functions (and Davenport-Heilbronn functions) overlayed by explicit $\zeta(1 + I \cdot t)$ data.

Using the functional equation of an L-function,

$$L(\chi_{N_c}(m, \cdot), s) = \chi_L(s) \cdot \bar{L}(\chi_{N_c}(m, \cdot), 2k - s) \quad (20)$$

with critical line $k=1/2$

$$L(\chi_{N_c}(m, \cdot), 1 + I \cdot t) = \chi_L(1 + I \cdot t) \cdot \bar{L}(\chi_{N_c}(m, \cdot), 1 - (1 + I \cdot t)) \quad (21)$$

$$= \chi_L(1 + I \cdot t) \cdot \bar{L}(\chi_{N_c}(m, \cdot), 0 - I \cdot t) \quad (22)$$

$$= e^{-2\theta_{ext}(1+I \cdot t)} \cdot \bar{L}(\chi_{N_c}(m, \cdot), 0 - I \cdot t) \quad (23)$$

The (unnormalised) Riemann Siegel Z function is therefore of the form for both the LHS and RHS of equations (21,23)

$$Z_L(\chi, 1 + I \cdot t) \equiv e^{I \cdot \text{Im}(\theta_{ext}(1+I \cdot t))} \cdot L(\chi_{N_c}(m, \cdot), 1 + I \cdot t) \quad (24)$$

$$= e^{I \cdot \text{Im}(\theta_{ext}(1+I \cdot t))} \cdot \chi_L(1 + I \cdot t) \cdot \bar{L}(\chi_{N_c}(m, \cdot), 1 - (1 + I \cdot t)) \quad (25)$$

$$= e^{I \cdot \text{Im}(\theta_{ext}(1+I \cdot t))} \cdot e^{-2\theta_{ext}(1+I \cdot t)} \cdot \bar{L}(\chi_{N_c}(m, \cdot), 0 - I \cdot t) \quad (26)$$




Figure 15: Observed $|Z|$ versus $\log(t)$ scatterplot of (i) good precision extreme Riemann Siegel Z function peak heights on the $s = 1 + I \cdot t$ line between $1e6 < t < 1e13$ (red solid circles) equation (16) and (ii) preliminary extreme Riemann Siegel Z function peak heights between $1e6 < t < 1e40$ based on spectral filtered heavily truncated Euler products based approximation method (v) overlayed by (I) dot-dashed brown line and green dashed line of two mapped trend estimates based on fitted $s = 0 + I \cdot t$ extreme peak behaviour (equations (1) and (3)) and the known functional equation and (II) shown as gray points $s = 0 + I \cdot t$ extreme peak heights transformed by the known functional equation

with the (unnormalised) property (shown using critical line $k=1/2$)

$$|Z_L(\chi, s)| \equiv |L(\chi, s)| = |\chi_L(s) \cdot \bar{L}(\chi_{N_c}(m, \cdot), 1-s)| \quad (27)$$

since estimating the extreme peak heights of $|L(\chi, s)|$ is the aim of the exercise. Using estimation methods (i-iii) in this paper, $|L(\chi, s)|$ is directly calculated off the critical line (and Z_L on the critical line), however when using the spectral filtering based methods (iv) and (v), $Z_L(\chi, s)$ is required for calculation.)

Using the multiplicative property of the absolute value on the field of complex numbers $|z_1 \cdot z_2| = |z_1| \cdot |z_2|$ [42-44]

$$|Z_L(\chi, 1 + I \cdot t)| = |e^{I \cdot \text{Im}(\theta_{ext}(1+I \cdot t))}| \cdot |L(\chi_{N_c}(m, \cdot), 1 + I \cdot t)| = |L(\chi_{N_c}(m, \cdot), 1 + I \cdot t)| \quad (28)$$

$$= |e^{I \cdot \text{Im}(\theta_{ext}(1+I \cdot t))}| \cdot \chi_L(1 + I \cdot t) \cdot \bar{L}(\chi_{N_c}(m, \cdot), 1 - (1 + I \cdot t))| \quad (29)$$

$$= |e^{I \cdot \text{Im}(\theta_{ext}(1+I \cdot t))}| \cdot |\chi_L(1 + I \cdot t)| \cdot |\bar{L}(\chi_{N_c}(m, \cdot), 1 - (1 + I \cdot t))| \quad (30)$$

$$= |\chi_L(1 + I \cdot t)| \cdot |\bar{L}(\chi_{N_c}(m, \cdot), 1 - (1 + I \cdot t))| \quad (31)$$

Using the complex conjugate symmetry of 1st degree self-dual L-functions (shown using critical line $k=1/2$)

$$\bar{L}_{\text{self-dual}}(\chi, 1-s) = L_{\text{self-dual}}(\chi, 1-s) \quad (32)$$

$$|L(\chi, 0 - I \cdot t)| = |L(\chi, 0 + I \cdot t)| \quad (33)$$

Therefore, the magnitude of the Riemann Siegel Z function for self-dual L-functions $|Z_{L\text{self-dual}}(s_{UB})|$ on the upper boundary of the critical strip can be obtained from the magnitude of the Riemann Siegel Z function $|Z_{L\text{self-dual}}(s_{LB})|$ using the functional equation, equation (31), the complex conjugate symmetry of 1st degree self-dual L-functions and the multiplicative property of the absolute value on the field of complex numbers $|z_1 \cdot z_2| = |z_1| \cdot |z_2|$

$$|Z_L(\chi, 1 + I \cdot t)|_{\text{self-dual}} = |L_{\text{self-dual}}(\chi_{N_c}(m, \cdot), 1 + I \cdot t)| \quad (34)$$

$$= |\chi_L(1 + I \cdot t)| \cdot |\bar{L}_{\text{self-dual}}(\chi_{N_c}(m, \cdot), 1 - (1 + I \cdot t))| \quad (35)$$

$$= |\chi_L(1 + I \cdot t)| \cdot |L_{\text{self-dual}}(\chi_{N_c}(m, \cdot), 1 - (1 + I \cdot t))|, \quad (36)$$

$$\begin{aligned} & \text{since } \bar{L}_{\text{self-dual}}(\chi, 1-s) = L_{\text{self-dual}}(\chi, 1-s) \\ & = |\chi_L(1 + I \cdot t)| \cdot |L_{\text{self-dual}}(\chi_{N_c}(m, \cdot), 0 - I \cdot t)| \end{aligned} \quad (37)$$

$$= |\chi_L(1 + I \cdot t)| \cdot |L_{\text{self-dual}}(\chi_{N_c}(m, \cdot), 0 + I \cdot t)|, \quad (38)$$

$$\text{since } |L_{\text{self-dual}}(\chi_{N_c}(m, \cdot), 0 - I \cdot t)| = |L_{\text{self-dual}}(\chi_{N_c}(m, \cdot), 0 + I \cdot t)|$$

Where furthermore numerically for $t > \pi$, for the investigated 1st degree L-functions (and their Davenport-Heilbronn functions)

$$|\chi_L(1 + I \cdot t)| \rightarrow \frac{1}{\sqrt{\left(\frac{t}{2\pi}\right)^1 \cdot N_C}}, \text{ for } t > \pi \quad (39)$$

so in the existing datasets of $s = 0 + I \cdot t$ results $|L_{\text{self-dual}}(\chi_{N_c}(m, \cdot), 0 + I \cdot t)|$ and $\sqrt{\frac{t}{2\pi} \cdot N_C}$ are already available for easy mapping to $s = 1 + I \cdot t$ behaviour via equation (38).

Thus in figure 16 it is shown

- in the lefthand panel - the mapped 1st degree L-function peak heights are compared to **explicit** $\zeta(1 + I \cdot t)$ data for extreme peak heights in the interval $t < 1e30$ using the first order universal growth chart plot

$$|\chi_L(1 + I \cdot t)| \cdot \left(|Z_L(0 + I \cdot t)| \cdot \frac{Nc}{N_{\chi \neq 0}} \right) \text{ vs } \log \left[\log \left(\sqrt{\left(\left(\frac{t}{2\pi} \right)^1 * Nc \right)} \right) \right]$$

- in the righthand panel - the mapped 5-periodic and 7-periodic Davenport-Heilbronn counterparts of 1st degree L-function peak heights are compared to **explicit** $\zeta(1 + I \cdot t)$ data for extreme peak heights in the interval $t < 1e30$ using the first order universal growth chart plot

$$|\chi_L(1 + I \cdot t)| \cdot \left(|Z_L(0 + I \cdot t)| \cdot \frac{Nc}{\sum_1^{Nc} |\chi_k|} \right) \text{ vs } \log \left[\log \left(\sqrt{\left(\left(\frac{t}{2\pi} \right)^1 * Nc \right)} \right) \right]$$

- in both panels - the mapped data from $|\zeta(0 + I \cdot t)|$ (large gray discs) to high numerical precision superimposes onto the explicit $|\zeta(1 + I \cdot t)|$ data (various coloured symbols).

Since both $s = 0 - I \cdot t$ and $s = 0 + I \cdot t$ data would be required to map the lower boundary of the critical strip to the upper boundary for the dual pairs $L(\chi_5(3, \cdot), s)$ & $L(\chi_5(2, \cdot), s)$ and $L(\chi_7(5, \cdot), s)$ [27], $L(\chi_7(3, \cdot), s)$ and only $s = 0 + I \cdot t$ data was calculated and stored, the above non self-dual L-functions were not included in the lefthand panel of figure 16. Their Davenport-Heilbronn (D-H) counterparts are shown in the righthand panel however since these D-H functions have real coefficient Dirichlet Series and are therefore self-dual in nature.

Highlights part 5

1. In the lefthand panel, there is good first order agreement between the **maximum** of the extreme peak heights of the mapped $s = 0 + I \cdot t$ peak heights of the L-function via the functional equation behaviour to the $s = 1 + I \cdot t$ peak heights explicitly calculated for $\zeta(1 + I \cdot t)$.
2. In the righthand panel, there is a gap between the **maximum** of the extreme peak heights of the mapped $s = 0 + I \cdot t$ peak heights of the examined 5-periodic and 7-periodic Davenport-Heilbronn 1st degree L-function counterparts via the functional equation behaviour to the $s = 1 + I \cdot t$ peak heights explicitly calculated for $\zeta(1 + I \cdot t)$. In particular,
 - (i) in the transformed universal growth chart reference frame the Davenport-Heilbronn counterpart functions have lower extreme peak heights on $s = 1 + I \cdot t$ and $s = 0 + I \cdot t$ for $t < 1e30$ than $\zeta(1 + I \cdot t)$ and $\zeta(0 + I \cdot t)$. (In figure 2 it was visually much harder to assess the qualitative difference in extreme peak height because of the much faster growth rate for $s = 0 + I \cdot t$ compared to the slow growth along $s = 1 + I \cdot t$ in figure 16.)
 - (ii) in the transformed universal growth chart frame the $f_2(1 + I \cdot t)$ and $f_3(1 + I \cdot t)$ for $t < 1e30$ have lower extreme peak heights than $f_1(1 + I \cdot t)$ and $f_4(1 + I \cdot t)$ respectively. (It is noted for the record that there is already empirical evidence for low t [29-31], that the density of $f_1(1 + I \cdot t)$ and $f_4(1 + I \cdot t)$ off-critical line non-trivial zeroes is lower than for $f_2(1 + I \cdot t)$ and $f_3(1 + I \cdot t)$ respectively.)
 - (iii) in figure 16 the graphical evidence indicates that adjusting for the gap between the $\zeta(1 + I \cdot t)$ behaviour and the Davenport-Heilbronn functions behaviour for extreme peak heights would require a functional dependence on t (not an additional constant numerical factor).




Figure 16: Rescaled XY axes overlay of the growth behaviour of extreme Riemann Siegel Z function peak heights $|Z|$ for different 1st degree **self-dual** L-functions in the interval $t < 1e30$ mapped from the $s = 0 + I \cdot t$ line to the $s = 1 + I \cdot t$ line and compared to $\zeta(1 + I \cdot t)$ extreme peak behaviour. The $|Z|$ values on the Y axis are scaled by the ratio of the (L-function conductor value)/(number of non-zero dirichlet characters mod N_C) i.e. $\log(|Z| \cdot \frac{N_C}{N_{\chi \neq 0}})$. (For the n-periodic Davenport-Heilbronn functions $\log(|Z| \cdot \sum_{k=1}^n \frac{1}{|X_k|})$ (where $n=5,7$ for the investigated self-dual D-H functions in this paper) is used for Y axis rescaling.) The X axis is transformed by the first quiescent region value of the L-function and hence the displayed scale is $\log(\log(\sqrt{(\frac{t}{2\pi})^1 \cdot N_C}))$.

Conclusions

A method for re-scaling the growth behaviour of extreme Riemann Siegel Z function of 1st degree L-functions with non-zero dirichlet characters of magnitude unity $|\chi| = 1$ is demonstrated to work on the $s = 0 + I \cdot t$ in addition to previously reported critical line behaviour. The re-scaling method is also shown to work for two large $N_C = \{5704, 6389\}$ self-dual L-functions. Using a mapping from the $s = 0 + I \cdot t$ line to the $s = 1 + I \cdot t$ line, two simple candidate trend growth formula for extreme Riemann Siegel Z function peaks of the Riemann Zeta function fitted to $s = 0 + I \cdot t$ data are shown to have obvious deficiencies.

Supplementary Material

A zip file of the located (extreme or otherwise) peak locations for the investigated L- functions and Davenport_Heilbronn counterparts is available on github https://github.com/johnpdmartin/sampling-investigations/blob/master/files_with_extreme_peaks_of_L_functions_on_zeroline.zip.

The typical format is 4-7 columns for files containing calculations by this author:

column 1 - t value (accurate to 0.05),

column 2 - N_1 value,

column 3 - residual of N_1 value (used for investigating multi-modal distribution of peaks within $[N, N+1)$ piecewise intervals.

column 4 - $|Z|$ value,

column 5 - Z value (sometimes calculated by more accurate method as crosscheck)

column 6 - text message describing method used for column 4 and 5

column 7 - may contain a number indicating element number of vector of estimates where peak occurred.

For files containing data from references there are typically column headers.

References

1. Brent, R.P. (1979) “On the zeros of the Riemann zeta function in the critical strip” *Math. Comp.* 33 , p1361-1372 DOI: <https://doi.org/10.1090/S0025-5718-1979-0537983-2>
2. Brent R.P., van de Lune J., te Riele H.J.J. and Winter D.T. (1982) “On the Zeros of the Riemann Zeta function in the Critical Strip. II” *Mathematics of Computation*, Volume 39, Number 160, October 1982, Pages 681-688
3. van de Lune J. and te Riele H.J.J. “On the Zeros of the Riemann Zeta Function in the Critical Strip.” III (1983) *MATHEMATICS OF COMPUTATION* VOLUME 41, NUMBER 164 OCTOBER PAGES 759-767
4. van de Lune J., te Riele H.J.J. and Winter D.T. (1986) “On the Zeros of the Riemann Zeta function in the Critical Strip. IV” *Mathematics of Computation*, Volume 46, Pages 667-681
5. A.M. Odlyzko, A. Schönhage, Fast algorithms for multiple evaluations of the Riemann zeta function, *Trans. Am. Math. Soc.*, 309, (1988), 797–809.
6. Odlyzko, A. M. “On the Distribution of Spacings Between Zeros of the Zeta Function”, (1987) *Mathematics of Computation* Vol 48, no. 177, 1987, pages 273-308
7. Odlyzko, A.M. “The 10^{20} -th zero of the Riemann Zeta function and 175 million of its neighbors”. (1988) <http://www.dtc.umn.edu/~odlyzko/unpublished/>
8. Odlyzko A.M. “The 10^{22} -nd zero of the Riemann zeta function, Dynamical, Spectral, and Arithmetic Zeta Functions” (2001) *Amer. Math. Soc., Contemporary Math. series*, no. 290, 139-144.
9. Gourdon X. and Sebah P., *Numbers, Constants and Computation*, <http://numbers.computation.free.fr/Constants/constants.html>, (1999) table of maximum Riemann Zeta peaks <http://numbers.computation.free.fr/Constants/Miscellaneous/MaxiZAll.txt>
10. Gourdon X. “The 10^{13} -th first zeros of the Riemann Zeta function and zeros computation at very large height” (2004) <http://numbers.computation.free.fr/Constants/Miscellaneous/zetazeros1e13-1e24.pdf>
11. Kotnick T., “COMPUTATIONAL ESTIMATION OF THE ORDER OF $\zeta(1/2+it)$ ” (2003) *MATHEMATICS OF COMPUTATION*, Volume 73, Number 246, Pages 949–956 S 0025-5718(03)01568-0
12. Hiary G.A., “Fast methods to compute the Riemann Zeta function” (2011) *Annals of Mathematics*, Volume 174, Pages 891-946 <https://doi.org/http://dx.doi.org/10.4007/annals.2011.174.2.4>
13. Hiary G.A., Table of high Riemann Zeta peaks (second table on the url page) <https://people.math.osu.edu/hiary.1/fastmethods.html>
14. Bober J.W. and Hiary G.A., “New computations of the Riemann zeta function on the critical line” (2018) *Experimental Mathematics*, 27(2), 125-137. <https://doi.org/10.1080/10586458.2016.1233083>
15. Tihanyi N., Kovács A. and Szűcs A. (2014) “Distributed computing of simultaneous Diophantine approximation problems” (2014) *Stud. Univ. Babeş-Bolyai Math.* 59, No. 4, 557–566
16. Tihanyi N., Kovacs A. and Kovacs J., “Computing Extremely Large Values of the Riemann Zeta Function” (2017) *J Grid Computing* 15:527–534 <https://doi.org/10.1007/s10723-017-9416-0>
17. Tihanyi N. and Kovacs A. “IMPROVEMENTS ON FINDING LARGE VALUES OF THE RIEMANN ZETA FUNCTION ON THE CRITICAL LINE” (2018) *Annales Univ. Sci. Budapest., Sect. Comp.*, 48, p53–64
18. Tihanyi N. “Numerical computing of extremely large values of the Riemann-Siegel Z-function” (2019) Doctoral Thesis, Eötvös Loránd University

19. Martin, J.P.D. “Preliminary estimates of extreme Riemann Zeta function peak heights with $\pm 10\%$ accuracy on the critical line using spectrally filtered heavily truncated partial Euler Product calculations” (2026) <http://dx.doi.org/10.6084/m9.figshare.31990179>
20. Martin, J.P.D. “Preliminary estimates of extreme Riemann Zeta function peak heights on three lines $s = \{0 + I \cdot t, 1 + I \cdot t, 1.5 + I \cdot t\}$ away from the real axis with $\pm 1\%, \pm 1\%, \pm 0.1\%$ accuracy for $10^{12} < t < 10^{40}$ using spectrally filtered heavily truncated partial Euler Product calculations.” (2026) <http://dx.doi.org/10.6084/m9.figshare.32129782>
21. Martin, J.P.D. “A first order universal growth chart on the critical line for periodic 1st degree Dirichlet character L-functions and a phenomenological model for trend growth of the known Riemann Zeta extreme peak heights.” (2026) <http://dx.doi.org/10.6084/m9.figshare.32736930>
22. Riemann, Bernhard (1859). “Über die Anzahl der Primzahlen unter einer gegebenen Grösse”. Monatsberichte der Berliner Akademie.. In *Gesammelte Werke*, Teubner, Leipzig (1892), Reprinted by Dover, New York (1953).
23. Titchmarsh, E.C. (1986) *The Theory of the Riemann Zeta Function*. 2nd Revised (Heath-Brown, D.R.) Edition, Oxford University Press, Oxford.
24. Edwards, H.M. (1974). *Riemann’s zeta function*. Pure and Applied Mathematics 58. New York-London: Academic Press. ISBN 0-12-232750-0. Zbl 0315.10035.
25. Berry, M. V. “The Riemann-Siegel Expansion for the Zeta Function: High Orders and Remainders.” *Proc. Roy. Soc. London A* 450, 439-462, 1995.
26. J. Arias De Reyna, “High precision computation of Riemann’s Zeta function by the Riemann-Siegel formula”, *Mathematics of Computation* Vol 80, no. 274, 2011, Pages 995–1009
27. The LMFDB Collaboration, *The L-functions and Modular Forms Database*, <http://www.lmfdb.org>, 2019, [Online; accessed January 2020].
28. Spira, R. *Mathematics of Computation*, Volume 63, Number 208, October 1994, Pages 747-748
29. Balanzario, E.P. and Sanchez-Ortiz, J. *Mathematics of Computation*, Volume 76, Number 260, October 2007, Pages 2045–2049
30. E. Bombieri, A. Ghosh, “Around the Davenport–Heilbronn function”, *Uspekhi Mat. Nauk*, 66:2(398) (2011), 15–66; *Russian Math. Surveys*, 66:2 (2011), 221–270 <https://doi.org/10.4213/rm9410> IAS lecture https://www.youtube.com/watch?v=-JUHypc2_9A
31. Martin, J.P.D. “7-, & 9-periodic dual Davenport Heilbronn counterexamples are derived for the corresponding Degree 1 dual L-functions pairs given in <https://www.lmfdb.org/L/degree1>.” (2020) <http://dx.doi.org/10.6084/m9.figshare.11675877>
32. Keiper, J. B. “On the Zeros of the Ramanujan τ -Dirichlet Series in the Critical Strip.” *Mathematics of Computation* 65, no. 216 (1996): 1613-619. Accessed February 10, 2021. <http://www.jstor.org/stable/2153727>.
33. Lenstra A.K., Lenstra Jr. H.W., Lovász L., “Factoring polynomials with rational coefficients” (1982) *Math. Ann.*, 261, (4), 515–534.
34. The PARI-Group, *PARI/GP* version 2.12.0, Univ. Bordeaux, 2018, <http://pari.math.u-bordeaux.fr/>.
35. R Core Team (2017). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
36. RStudio Team (2015). *RStudio: Integrated Development for R*. RStudio, Inc., Boston, MA <http://www.rstudio.com/>.
37. Box, G. E. P., & Cox, D. R. (1964). An analysis of transformations. *Journal of the Royal Statistical Society: Series B (Methodological)*, 26(2), 211–252. <https://www.jstor.org/stable/2984418>

38. Venables, W. N. & Ripley, B. D. (2002) *Modern Applied Statistics with S*. Fourth Edition. Springer, New York. ISBN 0-387-95457-0
39. Gosper, R. W. “harmonic summation and exponential gfs.” math-fun@cs-arizona.edu posting, Aug 2, 1996
40. Sondow, J. and Weisstein, E. W. “Harmonic Number.” From MathWorld—A Wolfram Resource. <https://mathworld.wolfram.com/HarmonicNumber.html>
41. Deligne, P. “La conjecture de Weil : I.” *Publications Mathématiques de l’IHÉS* 43 (1974): 273–307. <https://doi.org/10.1007/BF02684373>
42. Rudin, W. “Principles of Mathematical Analysis.” (1976) 3rd Edition, McGraw-Hill, New York, Chapter 1, pp.14–15, Theorem 1.33 part (c)
43. Brown J.W. and Churchill R.V. “Complex Variables and Applications” (2008) 8th Edition, McGraw-Hill Chapter 1
44. Wikipedia contributors. (2026, June 16). Brahmagupta–Fibonacci identity. Section “Multiplication of complex numbers”. In Wikipedia, The Free Encyclopedia. Retrieved 02:41, August 27, 2026, from https://en.wikipedia.org/w/index.php?title=Brahmagupta%E2%80%93Fibonacci_identity&oldid=1359602387