

STAT 221 Homework 1

John Rapp Farnes

15 januari 2020

1.3

a)


```
set.seed(2020)

w = rnorm(550,0,1) # 50 extra to avoid startup problems
x = filter(w, filter=c(0,-.9), method="recursive")[-(1:50)] # remove first 50


par(mfrow=c(2, 1))
plot.ts(x, main="Autoregression")

v = filter(w, sides=2, filter=rep(1/4,4)) # moving average
plot.ts(v, main="moving average")
```

Autoregression



moving average



b)


```

t = 1:100
x = cos(2*pi * t / 4)

par(mfrow=c(2, 1))
plot.ts(x, main="periodic")

v = rep(0, 100) #filter(x, sides=2, filter=rep(1/4,4)) # moving average
plot.ts(v, main="moving average")

```



c)

```


t = 1:100
x = cos(2*pi * t / 4) + rnorm(100,0,1)

par(mfrow=c(2, 1))
plot.ts(x, main="periodic with noise")


v = filter(x, sides=2, filter=rep(1/4,4)) # moving average
plot.ts(v, main="moving average")

```

periodic with noise



moving average



d) The moving average smooths out a) a lot, while the oscillations cancel in b). c) looks more periodic.

1.20

a)

```
w = rnorm(500,0,1)
(sample_acf <- acf(w, 20, plot=FALSE)$acf[20])
```

```
## [1] -0.06195539
```

Estimated ACF = -0.0619554, actual ACF = 0

```
w = rnorm(50,0,1)
(sample_acf <- acf(w, 20, plot=FALSE)$acf[20])
```

```
## [1] -0.006222671
```

Now estimated ACF = -0.0062227

The higher n , the more closer the value will tend to the actual 0 value. With 50 vs 500, the difference doesn't seem that big however, as $n = 50$ sometimes is closer to 0 depending on time series realization