

Modelling the effects of lymph node swelling on T-cell response.

Supplementary File 1.

Supplementary Methods.

1.1 TC recruitment

Under non-inflammatory conditions, it was assumed that TC entry and exit remain constant and TCs occupy a constant percentage (55%) of the total paracortical volume. The model represents a LN of 0.113-0.268 mm³, implying a LN mass of 0.18-0.44mg (based on collaborative unpublished measurements of murine popliteal LN mass versus volume). Reported lymphocyte recruitment for a popliteal LN of 1.15g is 4×10⁷ lymphocytes/hour and up to 40% of lymphocytes are B cells [1–3]. The model represents half a paracortex, therefore TC recruitment rate was estimated as 1950-9000 TCs/hour under non-antigenic conditions. Naive TC transit time through the LN (T_{res}) was estimated as 6-24 hours [4].

1.2 Agents and agent migration

The agents were designated as members of the TC or DC class (S Fig 1). Simulated DCs appear in the paracortex at a constant entry rate for 2 days, subsiding over the following 12 hours. Entry rate of DCs was scaled from counts of migrating DCs and initial TCs in a murine LN post-immunisation, with DCs totalling 4% of cells present [5]. TC migration was assumed to follow a random walk with pauses. Previous models have described TC migration with Brownian motion, a random-walk with persistence, run and tumble, and Lévy walks amongst other methods, partly due to differing reports of *in-vivo* migration. [6–9].

1.3 Agent interaction and signal integration

During TC-DC interaction, TCs remain stationary. Antigenic signal is presented by the DCs in the form of representative values of MHCI or MHCII, and decays with time (t) with the form :

$$MHCI(t) = MHCI_i(0.5)^{\frac{t}{MHCI_{1/2}}} \quad (1)$$

$MHCI_i$ or $MHCII_i$ is initial MHCI or initial MHCII presented with estimated MHC half-lives $MHCI_{1/2}$ and $MHCII_{1/2}$ respectively. During cognate TC-DC interaction, CD4⁺/CD8⁺ TCs gain (S) at a rate proportional to the MHCI or MHCII signal respectively (Fig 2.C). Accumulated stimulation decays at a constant rate. Total change in TC stimulation is therefore given according to the first order rate equation:

$$\frac{dS}{dt} = K_s MHCII(t) - \lambda_S S(t) \quad (2)$$

Where K_s is a rate constant and λ_S is a decay factor. Accumulated stimulation decays to a minimal value of S=1 to allow differentiation between cognate TCs that gain and lose stimulation (S=1) and those that never gain stimulation (S=0). Probability of cognate CD4⁺ activation (P_{a4+}) or cognate CD8⁺ activation (P_{a8+}) is calculated with a sigmoidal function given by:

$$P_{a4+} = \frac{1}{1 + e^{\frac{-S - Act\mu_4}{Actl_4}}} \quad (3)$$

Parameter $Act\mu_4$ determines the value of S required for 50% probability of activation and $Actl_4$ determines steepness of sigmoid inflection (Fig 2.D). Activation probability for $CD8^+$ TCs (P_{a8+}) is determined with a sigmoid curve using a lower inflection point, ($Act\mu_8$), than for $CD4^+$ TCs, unless the interacting agDC is licenced (see 2.4). Model parameters were estimated such that TC activation became apparent 8-15 hours post DC-arrival [10–12]. This method of signal integration and the subsequent progressive patterns of TC proliferation and differentiation is supported by *in-vivo* observations and modelling descriptions [12–15]. It is assumed that co-stimulatory requirements are met as agDCs are highly efficient antigen-presenting cells.

Post-activation, proliferation is possible every 11 ($CD4^+$) or 9 ($CD8^+$) ± 1 hr [16–19]. Differentiation into effector or memory cells is possible after ≥ 4 divisions, with differentiation probability determined with a second set of sigmoidal probability curves with midpoint $Dif\mu_{4+}$ and $Dif\mu_{8+}$ respectively [20,21]. Greater $CD4^+$ TCs dependence on continued stimulation for differentiation than $CD8^+$ TCs was implemented by using a higher minimum threshold of accumulated stimulation for $CD4^+$ differentiation than for $CD8^+$ TCs [22–26]. Effectors TCs that underwent < 8 proliferations were less likely to differentiate into memory cells than those that underwent > 8 proliferations. This was implemented by assigning differentiation ratios of dif_{early} and dif_{late} to the two different subsets (≤ 8 or > 8 divisions).

During TC activation and proliferation, $S1P_1R$ expression was estimated from several *in-silico* studies. Expression change was considered following TC migration velocity into areas of high $S1P$ concentration, changes in TC egress from the LN and changes in $S1P_1R$ expression relative to naive $S1P_1R$ expression following TC activation and with subsequent proliferation (Fig 2.F) [27–29].

References

- [1] Cahill R, Frost H, Trnka Z. The effects of antigen on the migration of recirculating lymphocytes through single lymph nodes. *J Exp Med*. 1976;143(4):870–888.
- [2] Habenicht LM, Albershardt TC, Iritani BM, Ruddell A. Distinct mechanisms of B and T lymphocyte accumulation generate tumor-draining lymph node hypertrophy. *Oncoimmunology*. 2016 08;5(8):e1204505.
- [3] Battaglia A, Ferrandina G, Buzzonetti A, Malinconico P, Legge F, Salutari V, et al. Lymphocyte populations in human lymph nodes. Alterations in $CD4(+) CD25(+)$ T regulatory cell phenotype and T-cell receptor Beta-repertoire. *Immunology*. 2003 11;110(3):304–312.
- [4] Tomura M, Yoshida N, Tanaka J. Monitoring cellular movement in vivo with photoconvertible fluorescence protein 'Kaede' transgenic mice. *PNAS*. 2008;105(31):10871–6.
- [5] Acton SE, Farrugia AJ, Astarita JL, Mourao-Sa D, Jenkins RP, Nye E, et al. Dendritic cells control fibroblastic reticular network tension and lymph node expansion. *Nature*. 2014 10;514(7523):498–502.
- [6] Celli S, Day M, Müller AJ, Molina-Paris C, Lythe G, Bousso P. How many dendritic cells are required to initiate a T-cell response? *Blood*. 2012;120(19):3945–3948. Available from: <http://www.bloodjournal.org/content/120/19/3945>.

- [7] Bogle G, Dunbar PR. Simulating T-cell motility in the lymph node paracortex with a packed lattice geometry. *Imm Cell Biol*. 2008 08;86(8):676–687.
- [8] Brown LV, Gaffney EA, Wagg J, Coles MC. An in silico model of cytotoxic T-lymphocyte activation in the lymph node following short peptide vaccination. *Journal of the Royal Society, Interface*. 2018 03;15(140):2018.0041.
- [9] Harris TH, Banigan EJ, Christian DA, Konradt C, Tait Wojno ED, Norose K, et al. Generalized Lévy walks and the role of chemokines in migration of effector CD8⁺ T cells. *Nature*. 2012 05;486:545–548.
- [10] Hugues S, Fetler L, Bonifaz L, Helft J, Amblard F, Amigorena S. Distinct T cell dynamics in lymph nodes during the induction of tolerance and immunity. *Nature Immunology*. 2004 10;5:1235–42.
- [11] Mempel TR, Henrickson SE, Von Andrian UH. T-cell priming by dendritic cells in lymph nodes occurs in three distinct phases. *Nature*. 2004;427(6970):154–9.
- [12] Stoll S, Delon J, Brotz T, Germain R. Dynamic imaging of T cell-dendritic cell interactions in lymph nodes. *Science*. 2002;296(5574):1873–6.
- [13] Germain RN, Stefanová I. THE DYNAMICS OF T CELL RECEPTOR SIGNALING: Complex Orchestration and the Key Roles of Tempo and Cooperation. *Annual Review of Immunology*. 1999 07;17(1):467–522. Available from: <https://doi.org/10.1146/annurev.immunol.17.1.467>.
- [14] Rachmilewitz J, Lanzavecchia A. A temporal and spatial summation model for T-cell activation: signal integration and antigen decoding. *Trends in Immunology*. 2002;23(12):592–595. Available from: <http://www.sciencedirect.com/science/article/pii/S1471490602023426>.
- [15] Lanzavecchia A, Sallusto F. Progressive differentiation and selection of the fittest in the immune response. *Nature Reviews Immunology*. 2002 12;2:982–7.
- [16] De Boer RJ, Homann D, Perelson AS. Different Dynamics of CD4⁺ and CD8⁺ T Cell Responses During and After Acute Lymphocytic Choriomeningitis Virus Infection. *The Journal of Immunology*. 2003;171(8):3928–3935. Available from: <http://www.jimmunol.org/content/171/8/3928>.
- [17] Foulds KE, Zenewicz LA, Shedlock DJ, Jiang J, Troy AE, Shen H. Cutting Edge: CD4 and CD8 T Cells Are Intrinsically Different in Their Proliferative Responses. *The Journal of Immunology*. 2002;168(4):1528–1532. Available from: <http://www.jimmunol.org/content/168/4/1528>.
- [18] Nelson RW, Beisang D, Tubo NJ, Dileepan T, Wiesner DL, Nielsen K, et al. T cell receptor cross-reactivity between similar foreign and self peptides influences naïve cell population size and autoimmunity. *Immunity*. 2015 01;42(1):95–107.
- [19] Tubo NJ, Pagán AJ, Taylor JJ, Nelson RW, Linehan JL, Ertelt JM, et al. Single Naive CD4⁺ T Cells from a Diverse Repertoire Produce Different Effector Cell Types during Infection. *Cell*. 2013 04;153(4):785–796.
- [20] Miller MJ, Wei SH, Parker I, Cahalan MD. Two-photon imaging of lymphocyte motility and antigen response in intact lymph node. *Science*. 2002;296(5574):1869–73.

- [21] Linderman JJ, Riggs T, Pande M, Miller M, Marino S, Kirschner DE. Characterizing the Dynamics of CD4+ T Cell Priming within a Lymph Node. *The Journal of Immunology*. 2010;184(6):2873–2885. Available from: <http://www.jimmunol.org/content/184/6/2873>.
- [22] Schrum AG, Palmer E, Turka LA. Distinct temporal programming of naive CD4(+) T cells for cell division versus TCR-dependent death susceptibility by antigen-presenting macrophages. *European journal of immunology*. 2005 02;35(2):449–459.
- [23] Wong P, Pamer EG. Cutting Edge: Antigen-Independent CD8 T Cell Proliferation. *The Journal of Immunology*. 2001;166(10):5864–5868. Available from: <http://www.jimmunol.org/content/166/10/5864>.
- [24] Kaech SM, Ahmed R. Memory CD8+ T cell differentiation: initial antigen encounter triggers a developmental program in naïve cells. *Nature Immunology*. 2001 05;2:415–22. Available from: <http://dx.doi.org/10.1038/87720>.
- [25] van Stipdonk MJB, Lemmens EE, Schoenberger SP. Naïve CTLs require a single brief period of antigenic stimulation for clonal expansion and differentiation. *Nature Immunology*. 2001 05;2:423–9. Available from: <http://dx.doi.org/10.1038/87730>.
- [26] Shaulov A, Murali-Krishna K. CD8 T Cell Expansion and Memory Differentiation Are Facilitated by Simultaneous and Sustained Exposure to Antigenic and Inflammatory Milieu. *The Journal of Immunology*. 2008;180(2):1131–1138. Available from: <http://www.jimmunol.org/content/180/2/1131>.
- [27] Pham T, Okada T, Matloubian M, Lo C, Cyster J. S1P 1 receptor signaling overrides retention mediated by Gi-coupled receptors to promote T cell egress. *Immunity*. 2008;28(1):122–133.
- [28] Matloubian M, Lo C, Cinamon G, Lesneski M, Xu Y. Lymphocyte egress from thymus and peripheral lymphoid organs is dependent on S1P receptor 1. *Nature*. 2004;427(6972):355–60.
- [29] Garriss CS, Blaho VA, Hla T, Han MH. Sphingosine-1-phosphate receptor 1 signalling in T cells: trafficking and beyond. *Immunology*. 2014 07;142(3):347–353.