

# LADR Done Right

johnsmith0x3f

August 31, 2024

# Contents

<b>1</b>	<b>Vector Spaces</b>	<b>2</b>
1.A	$\mathbf{R}^n$ and $\mathbf{C}^n$ . . . . .	2
1.B	Definition of Vector Space . . . . .	2
1.C	Subspaces . . . . .	2
<b>2</b>	<b>Inner Product Spaces</b>	<b>4</b>
2.A	Inner Products and Norms . . . . .	4
2.B	Orthonormal Bases . . . . .	6

# Chapter 1

## Vector Spaces

### 1.A $\mathbf{R}^n$ and $\mathbf{C}^n$

**Exercise 7** Show that for every  $\alpha \in \mathbf{C}$ , there exists a unique  $\beta \in \mathbf{C}$  such that  $\alpha + \beta = 0$ .

*Proof.* Suppose  $\alpha = a + bi$ . Let  $\beta = -a - bi$ . Then

$$\alpha + \beta = (a - a) + (b - b)i = 0,$$

proving existence.

Suppose there exists  $\gamma \in \mathbf{C}$  such that  $\alpha + \gamma = 0$ . Then

$$\gamma = \gamma + (\alpha + \beta) = (\gamma + \alpha) + \beta = \beta,$$

proving uniqueness. ■

### 1.B Definition of Vector Space

### 1.C Subspaces

**Exercise 12** Prove that the union of two subspaces of  $V$  is a subspace of  $V$  if and only if one of the subspaces is contained in the other.

*Proof.* Suppose  $U$  and  $W$  are subspaces of  $V$ , such that  $U \cup W$  is also a subspace and neither is contained in the other. In other words,  $U \not\subset W$  and  $W \not\subset U$ . Let  $u \in U - W$  and  $w \in W - U$ . Since  $U \cup W$  is a subspace of  $V$ , we have  $u + w \in U \cup W$ . However,

$$u + w \in U \Rightarrow w = (u + w) - u \in U,$$

$$u + w \in W \Rightarrow u = (u + w) - w \in W,$$

which implies that  $u + w$  is in neither  $U$  nor  $W$ , a contradiction.



## Chapter 2

# Inner Product Spaces

### 2.A Inner Products and Norms

**Exercise 5** Suppose  $T \in \mathcal{L}(V)$  is such that  $\|Tv\| \leq \|v\|$  for every  $v \in V$ . Prove that  $T - \sqrt{2}I$  is invertible.

*Proof.* Suppose  $V$  is finite-dimensional. For all  $u \in \text{null}(T - \sqrt{2}I)$ , we have  $Tu = \sqrt{2}u$ , and hence  $\|Tu\| = \|\sqrt{2}u\| = \sqrt{2}\|u\|$ . Since  $\|Tv\| \leq \|v\|$ , this implies that  $u = 0$ . ■

**Exercise 6** Suppose  $u, v \in \mathcal{L}(V)$ . Prove that  $\langle u, v \rangle = 0$  if and only if

$$\|u\| \leq \|u + av\|$$

for all  $a \in \mathbf{F}$ .

*Solution.* If  $\langle u, v \rangle = 0$ , then

$$\|u + av\|^2 - \|u\|^2 = \|av\|^2 \geq 0.$$

by the Pythagorean Theorem.

If  $\|u\| \leq \|u + av\|$  for all  $a \in \mathbf{F}$ , then

$$0 \leq \|u + av\|^2 - \|u\|^2 = \bar{a}\langle u, v \rangle + a\overline{\langle u, v \rangle} + |a|^2\|v\|^2.$$

Letting  $a = -\frac{\langle u, v \rangle}{\|v\|^2}$  yields  $\langle u, v \rangle = 0$ . ■

**Exercise 8** Suppose  $u, v \in V$  and  $\|u\| = \|v\| = 1$  and  $\langle u, v \rangle = 1$ . Prove that  $u = v$ .

*Proof.* Note that

$$\|u - v\|^2 = \langle u - v, u - v \rangle = \|u\|^2 - \langle u, v \rangle - \langle v, u \rangle + \|v\|^2 = 0.$$

Hence  $u - v = 0$  by definiteness. ■

**Exercise 11** Prove that

$$16 \leq (a + b + c + d) \left( \frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} \right)$$

for all positive numbers  $a, b, c, d$ .

*Proof.* By Exercise 6.17(a), we have

$$\begin{aligned} 16 &= \left| \sqrt{a \cdot \frac{1}{a}} + \sqrt{b \cdot \frac{1}{b}} + \sqrt{c \cdot \frac{1}{c}} + \sqrt{d \cdot \frac{1}{d}} \right|^2 \\ &\leq (a + b + c + d) \left( \frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} \right). \end{aligned}$$

■

**Exercise 12** Prove that

$$(x_1 + x_2 + \cdots + x_n)^2 \leq n(x_1^2 + x_2^2 + \cdots + x_n^2)$$

for all positive integers  $n$  and all real numbers  $x_1, \dots, x_n$ .

*Proof.* In Exercise 6.17(a), let  $y_1 = y_2 = \cdots = y_n = 1$ . ■

**Exercise 17** Prove or disprove: there is an inner product on  $\mathbf{R}^2$  such that the associated norm is given by

$$\|(x, y)\| = \max\{x, y\}$$

for all  $(x, y) \in \mathbf{R}^2$ .

*Counterexample.* Let  $u = (0, 1), v = (1, 0)$ . Then 6.22 fails. ■

## 2.B Orthonormal Bases

**Exercise F** Find a polynomial  $q \in \mathcal{P}_2(\mathbf{R})$  such that

$$p\left(\frac{1}{2}\right) = \int_0^1 p(x)q(x) \, dx$$

for every  $p \in \mathcal{P}_2(\mathbf{R})$ .

*Solution.* Let  $\varphi(p) = p(\frac{1}{2})$  and  $\langle p, q \rangle = \int_0^1 p(x)q(x) \, dx$ . Then with the orthonormal basis found in Exercise 5 and the formula in 6.43, we can find that

$$q(x) = -15x^2 + 15x - \frac{3}{2}. \quad \blacksquare$$