


satisfied. (It would be wise to find the confidence interval with the low score excluded so we can see its effect.) This check of requirements is Step 1 in the process of finding a confidence interval of  $\sigma$ , so we proceed with Step 2. ✓

#### MINITAB



**Step 2:** The confidence interval can be found using technology. If using Table A-4, we first use the sample size of  $n = 22$  to determine that the number of degrees of freedom is given by  $df = n - 1 = 21$ . If we use Table A-4, we refer to the row corresponding to 21 degrees of freedom, and we refer to the columns with areas of 0.975 and 0.025. (For a 95% confidence level, we divide  $\alpha = 0.05$  equally between the two tails of the chi-square distribution, and we refer to the values of 0.975 and 0.025 across the top row of Table A-4.) The critical values are  $\chi_L^2 = 10.283$  and  $\chi_R^2 = 35.479$  (as shown in Example 1).

**Step 3:** Using the critical values of 10.283 and 35.479, the sample standard deviation of  $s = 14.3$ , and the sample size of  $n = 22$ , we construct the 95% confidence interval by evaluating the following:

$$\frac{(n-1)s^2}{\chi_R^2} < \sigma^2 < \frac{(n-1)s^2}{\chi_L^2}$$

$$\frac{(22-1)(14.3)^2}{35.479} < \sigma^2 < \frac{(22-1)(14.3)^2}{10.283}$$

**Step 4:** Evaluating the expression above results in  $121.0 < \sigma^2 < 417.6$ . Finding the square root of each part (before rounding), then rounding to one decimal place, yields this 95% confidence interval estimate of the population standard deviation:  $11.0 < \sigma < 20.4$ .

#### Interpretation

Based on this result, we have 95% confidence that the limits of 11.0 and 20.4 contain the true value of  $\sigma$ . The confidence interval can also be expressed as (11.0, 20.4).

The IQ test was designed so that the population of IQ scores would have a standard deviation of 16, and the value of 16 is contained within the confidence interval, so the variation of the IQ scores does not appear to be unusual.