

Notes on Abstract Algebra

JOHN R. SMITH
email: jrsmith@ucdavis.edu

*Physics Department
University of California, Davis
Davis, CA 95616-8677, USA*

July 1, 2026

1 Logic

Definition If a statement A leads logically to another statement B we say that ‘if A , then B ’ and write $A \rightarrow B$ which can be thought of as A is *sufficient* for B . If logical space were analogous to sets, we could express ‘if A , then B ’ as $A \subseteq B$. Another way to think about implication is that $A \rightarrow B$ can be thought of as ‘if A is true, then B must be true’. Yet another way is (‘ B , if A ’) or that B is *necessary* for A . This can also be expressed as ‘ A only if B ’.

$A \rightarrow B$ implies that A being true is sufficient for B to be true. If the we have both $A \rightarrow B$ and $B \rightarrow A$, then we can say that A is *necessary and sufficient* for B – which can also be expressed as $A \iff B$ or by analogy with set notation as $A \subseteq B$ & $B \subseteq A$.

These logical dependences can be summarized in the statement “*If* is sufficient, *Only If* is necessary.”

Necessary Condition: If A is necessary for B , then B is a subset of A – every time B happens, A must happen.

Sufficient Condition: If A is sufficient for B , then A is a subset of B – whenever A happens, B follows.

Necessary and Sufficient: A and B are exactly the same set

2 Axioms

Definition The laws of addition for Numbers:

A1: The Commutative Law of addition: $x + y = y + x$ for all pairs of numbers x and y .

A2: The Associative Law of addition: $(x + y) + z = x + (y + z)$ for any three numbers x , y and z .

A3: The existence of a zero. There is a number, called the zero and denoted by 0 , which has the property that

$$x + 0 = x \text{ and } 0 + x = x \text{ for all numbers } x.$$

A4: The existence of negatives: Corresponding to each number x there is a number called the negative of x and written $-x$ which satisfies

$$x + -x = 0 \text{ and } -x + x = 0.$$

Definition Distributive Law

D1: The Distributive Law:

$$\begin{aligned}(x + y)z &= xz + yz \\ x(y + z) &= xy + xz\end{aligned}$$

Definition The laws of multiplication:

M1: The Commutative Law of multiplication : $xy = yx$ for all pairs of numbers x and y .

M2: The Associative Law of multiplication: $(xy)z = x(yz)$ for any three numbers x , y and z .

M3: The existence of a unity. There is a number, called the unity and denoted by 1, which has the property that

$$x1 = x \text{ and } 1x = x \text{ for all numbers } x.$$

M4: The existence of inverses: Corresponding to each number x there is a number called the inverse of x and written x^{-1} which satisfies

$$xx^{-1} = 1 \text{ and } x^{-1}x = 1.$$

Definition The Distinction of Numbers

X1: $1 \neq 0$.

$a \cdot 1 = a$ for all $a \in D$ where D represents a suitable domain of numbers.

$a \cdot 0 = 0$ for all $a \in D$.

3 Equivalence Relations

Theorem 3.1 *The equivalence classes theorem*

Suppose in a set S we have a relation R defined between certain pairs of elements, xRy meaning that x and y stand in the given relation R .

Suppose further that R has the following 3 properties:

(1) It is reflexive: i.e. xRx for all $x \in S$.

(2) It is symmetric: i.e. $xRy \Rightarrow yRx$.

(3) It is transitive: i.e. xRy and $yRz \Rightarrow xRz$.

Then R is an equivalence relations: i.e. it divides S into mutually exclusive subsets so that every element of S is in one and only one subset, and so that two elements are in the same subset if and only if they stand in the relation R to one another.

The subsets are called equivalence classes defined by R .

Given any element x in S consider all the elements y such that xRy . These form a subset of S : let us call it A_x . We show first that two elements are in the same subset A_x if and only if they stand in the relation R to one another. Suppose yRz and $y \in A_x$. Then xRy and so xRz since R is transitive. Hence $z \in A_x$. Conversely, if y and z are both in A_x we have xRy and xRz , i.e. yRx by the symmetric property and xRz : thus yRz by transitivity.

For each element x we now have a subset A_x , but these will not all be distinct. We will show that two such are either mutually exclusive or else identical. Suppose A_x and A_y both have an element z , Then xRz and yRz , and so zRy by the symmetric property; hence xRy by the transitive property. Now take any element w of A_x . Then xRw and since xRy we have yRx , giving yRw . So w is in A_y . Hence we have shown that if A_x and A_y have one element z in common, any element w of A_x is in A_y , i.e. $A_x \subset A_y$. Similarly $A_y \subset A_x$, and so the subsets A_x and A_y are identical.

Thus we have mutually exclusive subsets $A_{x_1}, A_{x_2}, \dots$. Finally by the reflexive property any element t is in one of the subsets, viz. A_t .

4 Groups, Subgroups and Cosets

Abstract Group

A set of elements S forms a group if to any two elements x and y of S taken in a particular order there is associated a unique third element of S , called their product and denoted by xy , which satisfies the following laws

(M2) For any three elements

$$x, y \text{ and } z: (xy)z = x(yz).$$

(M3) There is an element called the *neutral element* and denoted by e which has the property that $xe = ex = x$ for all elements x .

(M4) Corresponding to each element x there is an element x^{-1} called the *inverse* of x which has the property that

$$xx^{-1} = x^{-1}x = e.$$

Subgroups

A non-empty subset H of a group G is a subgroup of G if:

- (a) Given any ordered pair (h_1, h_2) of elements of H the product h_1h_2 as defined in G is in H .
- (b) The neutral element of G is in H .

(c) Given any element h in H , the inverse h^{-1} as defined in G is in H .

This a subset of elements of G is a subgroup, if and only if it forms a group, under the same group product.

Note that (b) is not needed, since if $h \in H, h^{-1} \in H$ by (c) and so $hh^{-1} = e \in H$ by (a).

Theorem 4.1 *A necessary and sufficient condition for a non-empty subset H to be a subgroup of a group G is that $gh^{-1} \in H$ for all $g, h \in H$.*

- (1) Necessary: If H is a subgroup $h^{-1} \in H$ and so $gh^{-1} \in H$.
- (2) Sufficient: If g is any element of H we have $gg^{-1} \in H$, i.e, $e \in H$. Hence $eg^{-1} = g^{-1} \in H$, i.e, H contains inverses. Thus if $g, h \in H$, so is h^{-1} and hence $g(h^{-1})^{-1} \in H$, i.e. $gh \in H$.

Given a group G and a subgroup H it is possible to decompose the whole group into subsets ‘parallel’ to H , two elements being in the same subset if their ‘difference’ is in H . This concept is better adapted in the product notation where the difference between g and k is expressed as $k^{-1}g$. We therefore decompose group G into subsets such that g and k are in the same subset if $k^{-1}g \in H$. These subsets are known as cosets and we have a true decomposition into mutually exclusive subsets because $k^{-1}g \in H$ sets up an equivalence relation gRk if $k^{-1}g \in H$.

Theorem 4.2 *Equivalence Classes of G relative to H The relation defined by gRk above is an equivalence relation.*

- (1) Reflexive: For any $g \in H$ we have gRg because $g^{-1}g = e \in H$ since H is a subgroup.
- (2) Symmetric: If gRk then $k^{-1}g \in H$. Hence its inverse is in H , i.e. $g^{-1}k \in H$ and so kRg .
- (3) Transitive: If gRk and kRl then $k^{-1}g \in H$ and $l^{-1}k \in H$. Hence $(l^{-1}k)(k^{-1}g) = l^{-1}g \in H$ and so gRl .

These mutual exclusive classes are called left cosets of G relative to H . So if gRk then kRg and $g^{-1}k \in H$ or $k = gh$ for some $h \in H$. Hence the coset containing g is precisely the complex gH , the set $\{gh\}$ for all $h \in H$.

Theorem 4.3 *Left Cosets The left cosets of G relative to H are the complexes gH , $g \in G$. Any left coset may be expressed in this form for any g in it. Any two cosets are either the same or have no element in common.*

We may similarly define *right cosets* of G relative to H as the equivalence classes under the relation given by $gk^{-1} \in H$. They are the complexes Hg .

We normally will use left cosets and therefore will just refer to them as cosets.

5 Mappings and Homomorphisms

Definition An injective function $f : X \rightarrow Y$ (also known as injection, or one-to-one function) is a function f that maps distinct elements of its domain to distinct elements; that is, $f(x_1) = f(x_2)$ implies $x_1 = x_2$. Equivalently, $x_1 \neq x_2$ implies $f(x_1) \neq f(x_2)$ in the equivalent contrapositive statement.

Definition A surjective function $f : X \rightarrow Y$ (also known as surjection, or onto function) is a function f that every element y can be mapped from some element x so that $f(x) = y$. In other words, every element of the function's codomain is the image of at least one element of its domain. It is not required that x be unique; the function f may map one or more elements of the set X to the same element of the set Y .

Definition Homomorphisms and Mappings:

Monomorphism: 1-1, Epimorphism: onto, Isomorphism: 1-1 and onto.

Homomorphisms preserve structure (i.e. multiplication and addition in Groups and Rings).

Homomorphisms that have the same object and image space are called Endomorphisms.

And Endomorphism that is also an Isomorphism is called an Automorphism.

In a Group G , the mappings of G into itself, given by $g \rightarrow x^{-1}gx$, where x is any fixed element of G , is an Automorphism, known as an Inner Automorphism.

Suppose $f : A \rightarrow B$ and $g : B \rightarrow C$. Hence $(g \circ f) : A \rightarrow C$ exists. Show

(a) If f and g are 1-1 (monomorphisms), then $g \circ f$ is 1-1:

Suppose $(g \circ f)(x) = (g \circ f)(y)$. Then $g(f(x)) = g(f(y))$. Because g is 1-1, $f(x) = f(y)$ and because f is 1-1, $x = y$. Therefore $(g \circ f)(x) = (g \circ f)(y)$, implies $x = y$; hence $g \circ f$ is 1-1.

(b) If f and g are onto (epimorphisms), then $g \circ f$ is onto.

Suppose $c \in C$. Because g is onto, there exists $b \in B$ for which $g(b) = c$. Because f is onto, there exists $a \in A$ for which $f(a) = b$. Thus $(g \circ f)(a) = g(f(a)) = g(b) = c$, Hence $g \circ f$ is onto.

(c) If $g \circ f$ is 1-1, then f is 1-1.

Prove contrapositive, Suppose f is not 1-1., Then there exists distinct elements $x, y \in A$ for which $f(x) = f(y)$. Then $(g \circ f)(x) = g(f(x)) = g(f(y)) = (g \circ f)(y)$. Hence $g \circ f$ is not 1-1. Therefore if $g \circ f$ is 1-1, then f must be 1-1.

(d) If $g \circ f$ is onto, then g is onto.

Prove contrapositive, If $a \in A$, then $(g \circ f)(a) = g(f(a)) \in g(B)$. Hence $(g \circ f)(A) \subseteq g(B)$. Suppose g is not onto. Then $g(B)$ is properly contained in C and so $(g \circ f)(A)$ is properly contained in C ; thus $g \circ f$ is not onto. Therefore if $g \circ f$ is onto, then g must be onto.

Definition A is called the object space of the mapping, B is the image space, a is the object whose image is b and we say that we have a mapping of A into B . If the mapping is denoted by θ , we write $\theta : A \rightarrow B$, and we say that $a\theta = b$ or $a \rightarrow b$.

In any homomorphism between a group G and another group H the neutral element e of G maps into the neutral element f of H . The set of elements which form a subset of G which map into f is called the *kernel* of the homomorphism.

Theorem 5.1 *The kernel of any homomorphism is a subgroup of G .*

Theorem 5.2 *The image $G\theta$ is a subgroup of H .*

6 Rings

A set R or elements forms a ring if to any two elements x and y of R taken in a particular order there are associated elements of R , called their sum and product and denoted by $x + y$ and xy respectively, both being unique, and such that axioms $A1, A2, A3, A4$ and $M2$ and $D1$ above are satisfied.

Basic types of rings

Commutative rings

The most obvious additional axiom is that of commutativity of multiplication $M1$, that $xy = yx$ for all x and y in the ring. Such a ring is called a *commutative ring* (addition is always commutative).

Rings with unity A ring admits of multiplication but not necessarily of division. For the latter to be possible we must have a neutral element, and also inverses, which are defined in terms of the neutral element. If both these are present, the structure becomes a *field*. However, a ring may have a neutral element and yet not possess inverses (example – the ring of integers). Such a ring is called a *ring with unity*.

7 Ideals

Definition A subring I of a ring R is a two-sided ideal if for all $x \in R$ and $i \in I$, xi and ix are both in I .

Note that because R may not be a commutative ring we must have both xi and ix in I .

Theorem 7.1 If R has a unity then

$$\{xi : x \in R, i \in I\} \subseteq I \implies \{xi\} = I \text{ and similarly for } \{ix\}.$$

Proof $\{xi\} \supseteq \{1i\} = I$

Quotient Rings

Theorem 7.2 If I is an ideal of a ring R the cosets $x + I$ form a ring under the sum $(x + I) + (y + I) = (x + y) + I$ and products $(x + I)(y + I) = xy + I$.

The ring of cosets is called the *Quotient Ring of I in R* and it is written R/I . The mapping $a \rightarrow a' = a + I$ for $a \in R$, which carries each element of R into the coset containing it is a homomorphism of R into the quotient ring R/I and the elements mapped on zero in this homomorphism are exactly the elements of the ideal I . In the homomorphic ring, the zero is the coset $0 + I$, hence the elements of I are mapped onto the zero coset.

Theorem 7.3 If a homomorphism H maps R on R' and exactly the elements of I on zero, then R' is isomorphic to the quotient ring R/I .

If b is an element of a commutative ring with unity, the set $[b]$ of all multiples xb of b , for any $x \in R$, is an ideal known as a *principal ideal*; it is the smallest ideal of R containing b .

Inclusion between ideals is closely related to divisibility between numbers. In the ring J of integers $n|m$ means that $m = an$, for some integer a (n divides m). Hence, every multiple of m is a multiple of n . The multiples of n constitute

the principle ideal $[n]$, so the result means that $[m]$ is contained in $[n]$. Conversely $[m] \subseteq [n]$ means in particular that m is in $[n]$, and hence $m = an$. Therefore

$$[m] \subseteq [n] \text{ if and only if } n|m.$$

The “bigger” number corresponds to the “smaller” ideal; for instance the ideal $[6]$ of all multiples of 6 is properly contained in the ideal $[2]$ of all even integers.

8 Fields

Theorem 8.1 (*Fields have no non-trivial Ideals*): *If a field F has an ideal I , then $I = (0)$ or $I = (1) = F$.*

Proof Let $a \in I$ and $a \neq 0$. Then $aa^{-1} = 1 \in I$ and so $f \cdot 1 = f \in I$ for any $f \in F$. Therefore $F \subset I$. But $I \subset F$ by definition of Ideals. Therefore $I = F$. The only other possibility is $a = 0$ in which case we have the trivial Ideal $I = (0) = 0$.

If F is any field, then the smallest subfield of F that contains the identity element 1 is called the *prime subfield* of F . If F is a finite field, then its prime subfield is isomorphic to $\mathbb{Z}_p$, where $p = \text{char}(F)$ for some prime p . When we want to emphasize $\mathbb{Z}/p\mathbb{Z}$ with its field structure, we denote it by $\mathbb{F}_p$. The prime subfield of $\mathbb{Z}_p$ which is isomorphic to $\mathbb{Z}/p\mathbb{Z}$ is $\mathbb{F}_p$. The only prime fields are the fields of residues modulo p and the field of rationals.

The prime subfield of a field F is the intersection of all subfields of F . The intersection of any two subfields is a field and similarly for all the other subfields. Also the intersection of all subfields has no proper subfields, since any such would be subfields of F and hence must contain the intersection which is impossible. Hence the intersection is a prime subfield. F cannot contain two distinct prime subfields, since their intersection would also be a subfield and hence must be identical with both.

Theorem 8.2 *a Field has no proper homomorphic images*

Proof It is sufficient to show that a field, considered as a ring, can have no proper ideals.

Theorem 8.3 *Any non-empty subset I of polynomials $F[x]$ which contains the sum and difference of any two, and all the multiples of any one of its members, consists either: 1) of 0 alone; or 2) of the set of multiples $q(x)a(x)$ of any non-zero member $a(x)$ of the least degree.*

Proof Unless $I = 0$, it contains a non-zero polynomial $a(x)$ of least degree $d(a)$ and, with $a(x)$, all its multiples $q(x)a(x)$. In this case, if $b(x)$ is any polynomial in I , using the division algorithm there exists some polynomial $r(x) = b(x) - q(x)a(x)$ which has degree less than $d(a)$. But by hypothesis I contains $r(x)$ and by construction no non-zero polynomial of degree less than $d(a)$. Hence $r(x)$ is the zero polynomial and $b(x) = q(x)a(x)$ ■

Using the above result of Thm. (8.3) we have in the domain $F[x]$ of polynomials $f(x)$ over F every ideal is a principal ideal.

9 Integral Domain

An integral domain is a commutative ring with a unity that has no zero divisors: $xy = 0 \rightarrow x = 0$ or $y = 0$.

Theorem 9.1 *The absence of zero divisors is equivalent to the Cancellation Law.*

If $xy = 0 \rightarrow x = 0$ or $y = 0$, then

$$xy = xz \rightarrow y = z \text{ or } x = 0,$$

and conversely.

Proof Suppose first that there are no zero divisors, Then if

$$xy = xz, \quad xy - xz = 0 \text{ and so } x(y - z) = 0.$$

Hence either $x = 0$ or $y - z = 0$ by hypothesis, i.e., $x = 0$ or $y = z$.

Conversely let the Cancellation Law be true and suppose that $xy = 0$. Then since $x0 = 0, xy = x0$ and so by the Cancellation Law $x = 0$ or $y = 0$.

10 Invariant Subgroups

Invariant Subgroups as also known as Normal Subgroups.

Consider a subgroup H of G . For $g, g' \in G$ let $x = gH$ and $y = g'H$ be the corresponding cosets of g and g' . For H to be an Invariant Subgroup of G we require that xy to be in the same coset as $gg'H$ or

$$(gH)(g'H) \in gg'H.$$

To obtain the basic relation defining Invariant Subgroups, premultiply both sides by g^{-1} :

$$\begin{aligned} g^{-1}(gH)(g'H) &\in g^{-1}gg'H, \\ H(g'H) &\in g'H. \end{aligned}$$

In the right most H choose the neutral element as e

$$\begin{aligned} Hg'e &\in g'H \\ Hg' &\in g'H \end{aligned}$$

If $Hg' \in g'H$ for any $g' \in G$ we also must have that (changing g' to g) that $Hg \in gH$. Since $g^{-1} \in G$ we also have

$$Hg^{-1} \in g^{-1}H.$$

and therefore

$$\begin{aligned} (gH)g^{-1}g &\in gg^{-1}Hg, \\ gH &\in Hg, \end{aligned}$$

Therefore we have $gH \in Hg$ and $Hg \in gH$ and so

$$gH = Hg,$$

leading up to the following definition of an Invariant or Normal Subgroup:

Definition H is an invariant subgroup of G if it is a subgroup and if the left and right cosets of H in G coincide; i.e., if

$$gH = Hg \quad \forall g \in G.$$

Invariant subgroups are often called normal subgroups or self-conjugate subgroups.

If the left coset gH is also a right coset, then it contains $g(=eg)$ and so must be the right coset Hg . Hence if every left coset is a right coset (or vice versa) H is invariant.

Let us show that this condition is sufficient.

Theorem 10.1 *If H is invariant in G then $(gH)(g'H) = gg'H$ for all $g, g' \in G$.*

Proof

$$\begin{aligned} gHg'H &= g(Hg')H = g(g'H)H \text{ since } Hg' = g'H \text{ as } H \text{ is invariant,} \\ &= gg'H^2 = gg'H \text{ since } H^2 = H \text{ as } H \text{ is a subgroup.} \end{aligned}$$

Note we have already shown that it is sufficient to prove that

$$(gH)(g'H) \subseteq gg'H,$$

in order to show that H is an Invariant Subgroup.

Theorem 10.2 *A subgroup H is an invariant subgroup of G if and only if $g^{-1}Hg = H$ for all $g \in G$.*

11 Conjugacy

Definition The elements x and $g^{-1}xg$, where x and g are elements of a group G , are called conjugate elements in G : in other words x and y are conjugate if and only if there exists an element $g \in G$ such that $y = g^{-1}xg$.

Theorem 11.1 *The relation of conjugacy between elements of any group G is an equivalence relation.*

Reflexive. Since $x = e^{-1}xe$, x is conjugate to itself.

Symmetric. If $y = g^{-1}xg$ we have $x = gyg^{-1} = (g^{-1})^{-1}yg^{-1}$.

Transitive. If $y = g^{-1}xg$ and $z = h^{-1}yh$ we have

$$z = h^{-1}g^{-1}xgh = (gh)^{-1}x(gh).$$

It follows that the relation divides the elements of G into mutually exclusive classes; these are called *conjugacy classes* in G .

Theorem 11.2 *If H is a subgroup of G then $g^{-1}Hg$ is a subgroup for any element $g \in G$.*

Proof For

$$(g^{-1}Hg)^2 = (g^{-1}Hgg^{-1}Hg) = (g^{-1}HHg) = (g^{-1}Hg) \text{ since } H^2 = H$$

as H is a subgroup. Also

$$(g^{-1}Hg)^{-1} = g^{-1}H^{-1}g = g^{-1}Hg \text{ since } H^{-1} = H.$$

We say that H and $g^{-1}Hg$ are *conjugate subgroups* in G .

Theorem 11.3 *The relation of conjugacy between subgroups of G is an equivalence relation.*

Theorem 11.4 *Normal subgroups of G are composed of all elements of G which are common to the set of conjugate subgroups of G .*

Proof Suppose the set of conjugate subgroups of G are

$$H_1, H_2, \dots, H_i.$$

The elements which are common to these subgroups form a group, since with any two of them contained in H_i their product is also contained in H_i . Also, the transformations of H_1 by elements in G are the same conjugate subgroups

$$H_1, g_2^{-1}H_1g_2, \dots, g_i^{-1}H_1g_i,$$

the group of common elements, which is part of any one of them as it is of H_1 , is identical with its transforms and hence a normal subgroup of G .

A group of all elements which are common to conjugate subgroups, is called their **greatest common subgroup** and a greatest common subgroup is normal in the group.

Theorem 11.5 *H is an invariant subgroup of G if the conjugacy class of subgroups of G which contains H consists of just H itself: i.e., if the only subgroup conjugate to H is H .*

12 Quotient groups

Theorem 12.1 *If H is an invariant subgroup of G , the cosets of H in G form a group under the group product given by*

$$(gH)(g'H) = gg'H.$$

The product is well defined and unique, and is also a coset.

It is associative, since

$$((gH)(g'H))(g''H) = gg'g''H = (gH)((g'H)(g''H)).$$

The coset $H = (eH)$ is a unity since

$$(gH)(eH) = geH = gH \text{ and } (eH)(gH) = egH = gH.$$

The coset gH has an inverse $g^{-1}H$, since

$$(gH)(g^{-1}H) = gg^{-1}H = eH = H \text{ and } (g^{-1}H)(gH) = H$$

similarly.

The quotient group formed by the cosets of H in G is called the *Quotient Group of H in G* , or sometimes the *Factor Group of G relative to H* .

13 Quotient Group Associated with a Homomorphism

As we have already hinted, the subjects of invariant subgroups and homomorphisms of groups are closely connected. There is in fact a 1-1 correspondence between subgroups of a given group G and the essentially different homomorphisms which have G as the object space, so that the study of either topic is similar to that of the other.

Suppose that H is an invariant subgroup of G , so that the quotient group G/H is defined. Consider the mapping $\theta : G \rightarrow G/H$ defined by $g\theta = gH$, the coset containing g , which is of course an element of G/H .

Theorem 13.1 *The mapping $\theta : G \rightarrow G/H$ defined by $g\theta = gH$ is a homomorphism.*

Proof

$$\begin{aligned}(gg')\theta &= gg'H = gg'HH \\ &= gHg'H \\ &= (g\theta)(g'\theta)\end{aligned}$$

Corollary 13.2 *The kernel of θ is H and the image is G/H , so that θ is an epimorphism.*

The homomorphism θ is called the Natural Homomorphism or the Canonical Homomorphism associated with H . It is essential that H be an invariant subgroup, for otherwise G/H is not defined and we cannot speak of the product of the cosets $(gH)(g'H)$.

Any homomorphism $\theta : G \rightarrow G'$ is closely related to the natural homomorphism associated with an invariant subgroup of G .

Theorem 13.3 *The kernel of $\theta : G \rightarrow G'$ is an invariant subgroup of G .*

If h is in the kernel H and g is any element of G , we have

$$\begin{aligned}(g^{-1}hg)\theta &= (g^{-1}\theta)(h\theta)(g\theta) \\ &= (g\theta)^{-1}f(g\theta), \text{ where } f \text{ is the neutral element of } G' \\ &= (g\theta)^{-1}(g\theta) = f\end{aligned}$$

Theorem 13.4 *The mapping defined by $gH \rightarrow g\theta$ is a well-defined mapping of G/H onto $G\theta$.*

If g_1 and g_2 are in the same coset, then $g_1\theta = g_2\theta$. If g_1 and g_2 are in the same coset, then $g_2^{-1}g_1 \in H$. Hence $(g_2\theta)^{-1}(g_1\theta) = f$, the neutral element in G' , since H is in the kernel of θ . Therefore $g_2\theta = g_1\theta$.

Also any element of $G\theta$ is the image of some $g \in G$ and so the mapping is onto $G\theta$, although not necessarily onto G' .

Theorem 13.5 *The mapping defined by $gH \rightarrow g\theta$ is an isomorphism between G/H and $G\theta$.*

We have that

$$\begin{aligned}(g_1H)(g_2H) &= g_1g_2H, \\ (g_1\theta)(g_2\theta) &= (g_1g_2)\theta,\end{aligned}$$

because θ is a homomorphism and $g_1g_2H \rightarrow (g_1g_2)\theta$.

Since this mapping is onto $G\theta$, it is also an isomorphism provided it is 1-1, i.e., provided the kernel is the neutral element of G/H . Suppose $gH \rightarrow f$. Then $g\theta \rightarrow f$ and so $g \in H$, so that $gH = H$, the neutral element of G/H .

Notice that the sets of elements of G which map into the particular elements of $G\theta$ are the cosets of G relative to the kernel H .

We have proved that to any invariant subgroup H of G there is a natural homomorphism from G onto G/H , and conversely that any homomorphism of G is isomorphic to a natural homomorphism associated with its kernel. Therefore there is a 1-1 correspondence between the invariant subgroups and the essentially different homomorphisms of G . By essentially different homomorphisms we mean those having different kernels: two homomorphisms having the same kernel are both equivalent to the same natural homomorphism and therefore exhibit the same structure.

The image space G' is largely irrelevant : the subgroup $G\theta$ of G' is isomorphic to G/H but G' may be larger. Also there is no reason why $G\theta$ should be invariant in G' .

14 Relationship between Normal Subgroups and Their Images

Consider a group G that has a normal subgroup N and H and that $H \subseteq N$.

Theorem 14.1 *Because $N \supseteq H$, both N and H are normal subgroups of G . Thus G/N is the homomorphic image of G/H under the natural homomorphism $\pi : G/H \rightarrow G/N$ defined by $(xH)\pi = xN$.*

Proof 1) Define a function $\phi : G \rightarrow G/N$ by

$$x\phi = xN.$$

This is the standard canonical projection map. It is a surjective group homomorphism with $\ker(\phi) = N$.

2) We want to construct a map $\bar{\phi} : G/H \rightarrow G/N$. Define $\bar{\phi}$ by:

$$(xH)\bar{\phi} = x\phi = xN.$$

3) We must show that if $xH = yH$, then $(xH)\bar{\phi} = (yH)\bar{\phi}$.

$$\begin{aligned} xH &= yH \rightarrow y^{-1}x \in H. \\ H &\subseteq N \text{ given} \\ y^{-1}x &\in N \\ xN &= yN \\ (xH)\bar{\phi} &= (yH)\bar{\phi} \end{aligned}$$

The map is well-defined because $H \subseteq \ker(\phi)$.

4) $\bar{\phi}$ is a Homomorphism:

$$(xH)(yH)\bar{\phi} = (xyH)\bar{\phi} = xyN = (xN)(yN) = (xH)\bar{\phi}(yH)\bar{\phi}.$$

5) $\bar{\phi}$ is Surjective: Let $yN \in G/N$. Since $y \in G$, we can form the coset $yH \in G/H$:

$$(yH)\bar{\phi} = yN.$$

Every element in the co-domain has a pre-image

6) Apply the First Isomorphism Theorem By the First Isomorphism Theorem, the domain modulo the kernel is isomorphic to the image

$$(G/H)/\ker(\bar{\phi}) \cong \text{Im}(\bar{\phi}) = G/N.$$

Since G/N is isomorphic to a quotient group of G/H , it is by definition a homomorphic image of G/H . ■

Lattice Isomorphism Theorems

Theorem 14.2 (First Isomorphism Theorem:) *Let ϕ be a group homomorphism from G to G' . Then the mapping from $G/\text{Ker}(\phi)$ to $G\phi$ given by $g\text{Ker}(\phi) \rightarrow g\phi$ is an isomorphism.*

Example Consider the homomorphism $\phi : \mathbb{Z} \rightarrow \mathbb{Z}_4$ such that $x\phi = x \pmod{4}$. In this case let G be the group of integers (under addition)

$$G = \{\dots, -4, -3, -2, -1, 0, 1, 2, 3, 4, \dots\}$$

and

$$G' = \mathbb{Z}_4 = \{0, 1, 2, 3\}.$$

Also

$$\text{Ker}(\phi) = \{\dots - 8, -4, 0, 4, 8, \dots\} = \langle 4 \rangle \text{ (the cyclic subgroup generated by 4)}$$

So $G\phi = \{0, 1, 2, 3\}$. Note that $G\phi$ doesn't have to be the entire group G' . In this example it is the case that $G\phi = G'$. Then

$$G/\text{Ker}(\phi) = \{0 + \langle 4 \rangle, 1 + \langle 4 \rangle, 2 + \langle 4 \rangle, 3 + \langle 4 \rangle\} \text{ (four cosets)}$$

Notice that we can create a mapping $\bar{\phi}$ that maps $G/\text{Ker}(\phi)$ into $G\phi$ as follows:

$$\bar{\phi} : G/\text{Ker}(\phi) \rightarrow G\phi,$$

where

$$\begin{aligned} \bar{\phi}(0 + \langle 4 \rangle) &= (0)\phi = 0 \\ \bar{\phi}(1 + \langle 4 \rangle) &= (1)\phi = 1 \\ \bar{\phi}(2 + \langle 4 \rangle) &= (2)\phi = 2 \\ \bar{\phi}(3 + \langle 4 \rangle) &= (3)\phi = 3 \end{aligned}$$

Theorem 14.3 (Second Isomorphism Theorem:) *The Second Isomorphism Theorem: If S is a subgroup of G and N is a normal subgroup of G , then the product SN is a subgroup, and the quotient SN/N is isomorphic to $S/(S \cap N)$.*

Theorem 14.4 (Third Isomorphism Theorem:) *Normal Subgroups of Quotients: If N and H are normal subgroups of G (where $N \subseteq H$), then H/N is a normal subgroup of G/N . Furthermore, the second quotient is isomorphic to the primary quotient:*

$$(G/N)/(H/N) \cong G/H.$$

The Third Isomorphism Theorem is analogous to fractions:

$$\frac{G/N}{H/N} \cong \frac{G}{H}.$$

Theorem 14.5 (Fourth Isomorphism Theorem:) *There is a direct, structure-preserving correspondence between the subgroups of a quotient group G/N and the subgroups of G that contain N .*

Theorem 14.6 *The mapping $\phi : \mathbb{Z} \rightarrow \mathbb{Z}_n$ defined by $m\phi = m \bmod n$ is a homomorphism with kernel $\langle n \rangle$. By the First Isomorphism Theorem, $\mathbb{Z}/\langle n \rangle \cong \mathbb{Z}_n$.*

If ϕ is a homomorphism from a finite group G to G' , then $\|G\phi\|$ divides $\|G\|$ and $\|G'\|$. This follows from the First Isomorphism Theorem and the fact that $G\phi$ is a subgroup of G' and Lagrange's Theorem.

Lemma 14.7 *The inner automorphism θ_x given by $g \rightarrow x^{-1}gx$ is the identity automorphism if and only if $x \in C$ where C is the center.*

If $x \in C$, $x^{-1}gx = x^{-1}xg = g$ and so $\theta_x = I$, the identity.

If $\theta_x = I$, $x^{-1}gx = g \forall g$ and so $x \in C$.

Lemma 14.8 *The mapping $x \rightarrow \theta_x$ of G into the group of automorphisms of G is a homomorphism*

We already have that $\theta_{xy} = \theta_x\theta_y$ and so the result follows.

By Lemma(14.7) the kernel of this homomorphism is the center C . The image is the subgroup of inner automorphisms. Therefore we obtain

Theorem 14.9 *The group of inner automorphisms of a group $G \cong G/C$, where C is the center of G .*

15 Vector Spaces

Definition Vector Spaces:

For F a field of scalars, let $\lambda \in F$. We talk about a vector space *over* F : the specification of the field F is an integral part of the definition.

A set V of elements forms a vector space over a given field F if to any two elements $\mathbf{x}$ and $\mathbf{y}$ of V there is associated a unique third element of V , called their sum and denoted by $\mathbf{x} + \mathbf{y}$, and if to any element $\mathbf{x}$ of V and any element λ of F there is associated an element V called their product and denoted by $\lambda\mathbf{x}$, where the sum satisfied A1, A2, A3 and A4 (above) and their product satisfies the following laws:

V1: $\lambda(\mathbf{x} + \mathbf{y}) = \lambda\mathbf{x} + \lambda\mathbf{y}$ for all $\lambda \in F$ and $\mathbf{x}, \mathbf{y} \in V$.

V2: $(\lambda + \mu)\mathbf{x} = \lambda\mathbf{x} + \mu\mathbf{x}$ for all $\lambda, \mu \in F$ and $\mathbf{x} \in V$.

V3: $\lambda(\mu\mathbf{x}) = (\lambda\mu)\mathbf{x}$ for all $\lambda, \mu \in F$ and $\mathbf{x} \in V$.

V4: $1.\mathbf{x} = \mathbf{x}$ where 1 is the unity of F and $\mathbf{x}$ is any element of V

Definition n -dimensional Affine Space $\mathbb{R}_n$. An ordered n -tuple of real numbers, i.e., a system of $(a_1, a_2, \dots, a_n)$ of n real numbers, called components, in a definite order, is called a point (n a positive integer). The numbers $a_1, a_2, \dots, a_n$ are called the coordinates of the point; in particular a_1 is called its first, a_2 its second, ..., a_n its n -th coordinate. Two points $P = (a_1, a_2, \dots, a_n)$ and $Q = (b_1, b_2, \dots, b_n)$ are said to be the *same* or to coincide if and only if $a_1 = b_1, a_2 = b_2, \dots, a_n = b_n$. The totality of all n -tuples of real numbers is called n -dimensional Affine Space and is denoted by $\mathbb{R}_n$.

Vector Notation An element, $\mathbf{x} \in \mathbb{R}_n$ is defined by its n components $(x_1, x_2, \dots, x_n)$, as follows

$$\mathbf{x} = (x_1, x_2, \dots, x_n)$$

Linear Dependence/Independence In the abstract vector space V the finite set of vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ is said to be linear dependent if scalars $\lambda_1, \lambda_2, \dots, \lambda_n$ exist, not all zero, such that

$$\lambda_1 \mathbf{x}_1 + \lambda_2 \mathbf{x}_2 + \dots + \lambda_n \mathbf{x}_n = \mathbf{0}.$$

If no such scalars exist, i.e., if

$$\lambda_1 \mathbf{x}_1 + \lambda_2 \mathbf{x}_2 + \dots + \lambda_n \mathbf{x}_n = \mathbf{0} \implies \lambda_i = 0 \forall i = 1, 2, \dots, n$$

then the set is linearly independent.

Conventions: We adopt the Range and Summation Conventions:

Definition Range Convention. When a small Latin suffix (superscript or subscript) occurs unrepeated in a term, it is understood to take all the values $1, 2, \dots, n$, where n is the number of dimensions of the space.

Definition Summation Convention. When a small Latin suffix is repeated in a term, summation with respect to that suffix is understood, the range of summation being $1, 2, \dots, n$.

Kronecker delta:

$$\delta_r^s = \begin{cases} 1 & \text{if } s = r \\ 0 & \text{if } r \neq s \end{cases}$$

Also, the another form of the Kronecker delta is δ_{rs} which satisfies the same cases as above.

Permutation Symbol: Consider a set σ of $\sigma_1, \sigma_2, \dots, \sigma_n$ numbers taken from the set $\{1, 2, \dots, n\}$. Let

$$\epsilon_{\sigma_1 \sigma_2 \dots \sigma_n} = \begin{cases} +1 & \text{if } \sigma \text{ is an even permutation of } \{1, 2, \dots, n\} \\ -1 & \text{if } \sigma \text{ is an odd permutation of } \{1, 2, \dots, n\} \\ 0 & \text{if } \sigma \text{ has any duplicates} \end{cases}$$

Span The set of vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ are said to span or generate, the space V if any vector $\mathbf{x}$ of V is expressible as a linear combination of them, i.e. if

$$\mathbf{x} = \lambda_1 \mathbf{x}_1 + \lambda_2 \mathbf{x}_2 + \dots + \lambda_n \mathbf{x}_n$$

for some scalars $\lambda_1, \lambda_2, \dots, \lambda_n$ not necessarily unique.

Basis The set $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ is a Basis of V : 1) if they are linearly independent; and (2) they span V .

Theorem 15.1 If $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ span V then they form a basis if and only if $\mathbf{x}$ in V is expressible uniquely as a linear combination of $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$.

Theorem 15.2 If $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m$ span V , then there is a subset of these which is a basis of V .

Proof If $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m$ are linearly independent, they form a basis and the theorem is proved. If they are linearly dependent then some $\mathbf{x}_{k+1}$ is a linear combination of

$$\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \text{ say } \mathbf{x}_{k+1} = \sum_{i=1}^k \mu_i \mathbf{x}_i,$$

Consider the set $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k, \mathbf{x}_{k+2}, \dots, \mathbf{x}_m$. Any vector $\mathbf{x}$ can be expressed as $\mathbf{x} = \sum_{i=1}^m \lambda_i \mathbf{x}_i$, since the $\mathbf{x}_i$'s also span V . Hence

$$\mathbf{x} = \sum_{\substack{i=1 \\ i \neq k+1}}^k \lambda_i \mathbf{x}_i + \lambda_{k+1} \sum_{i=1}^k \mu_i \mathbf{x}_i$$

and is a linear combination of $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k, \mathbf{x}_{k+2}, \dots, \mathbf{x}_m$ so that these also span V .

If this set is now linearly independent the theorem is proved as they form a basis. If not repeat the above argument and obtain a spanning set of $m - 2$ vectors, which may or may not be linearly independent. Repeat until after a most $m - 1$ steps we do obtain a basis, which is a subset of the original set. This must occur since after $m - 1$ steps we are left with a single vector which spans V , and this must be non-zero unless V is the trivial space $\{\mathbf{0}\}$ and hence is linearly independent, since $\lambda_1 \mathbf{x}_1 = \mathbf{0}$ with $\mathbf{x}_1 \neq \mathbf{0} \implies \lambda_1 = 0$.

Corollary 15.3 *If V possesses a finite spanning set then V is finite dimensional.*

Theorem 15.4 *If $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ are linearly independent and V is finite dimensional then there exists a basis containing*

$$\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p.$$

Proof Let $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ be a basis of V . Consider the set $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p, \mathbf{e}_1, \dots, \mathbf{e}_n$. This set certainly spans V . Using the argument of Thm. (15.2) on this set, but notice that at each stage we obtain one of the $\mathbf{e}_i$'s as a linear combination of the preceding vectors, since no $\mathbf{x}$ can be a linear combination of the preceding $\mathbf{x}_i$'s as the $\mathbf{x}_i$'s form a linearly independent set.

Hence we are left with the $\mathbf{x}_i$'s and the process must stop before or exactly when the $\mathbf{e}_i$'s are exhausted. We thus obtain a basis which contains $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p$ together with some of the $\mathbf{e}_i$'s.

Theorem 15.5 (Steinitz Exchange Lemma) *Let U and W be finite subsets of a vector space V and let U be a set of linearly independent vectors and let W span V . If $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p\} \in U$ is a linearly independent set, and $\{\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m\} \in W$ spans V , then $p \leq m$.*

Proof Consider the set $\mathbf{x}_p, \mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m$. Since the $\mathbf{y}_i$'s span V , $\mathbf{x}_p$ is a linear combination of them and so the set is linearly dependent. Hence some $\mathbf{y}_{k+1}$ is a linear combination of

$$\mathbf{x}_p, \mathbf{y}_1, \dots, \mathbf{y}_k.$$

This means that $\mathbf{x}_p, \mathbf{y}_1, \dots, \mathbf{y}_k, \mathbf{y}_{k+2}, \dots, \mathbf{y}_m$ span V .

Continuing similarly, adding $\mathbf{x}_{p-1}$ to this set. We obtain a linearly dependent set and some $\mathbf{y}_{l_1}$ is expressible as a linear combination of the others, so that we have a spanning set comprising $\mathbf{x}_{p-1}, \mathbf{x}_p$, and all the $\mathbf{y}_i$'s except $\mathbf{y}_{k+1}$

and $\mathbf{y}_{l+1}$. By continuing this process we obtain a spanning set consisting of one more $\mathbf{x}$ and one less $\mathbf{y}$. We always obtain a $\mathbf{y}_i$ since the $\mathbf{x}_i$'s are linearly independent and so no $\mathbf{x}_{i+1}$ is a linear combination of the preceding $\mathbf{x}_i$'s that have been introduced.

The process must be possible until all the $\mathbf{x}_i$'s have been exhausted and we must at each stage have some $\mathbf{y}_i$'s left, otherwise the set consisting of the $\mathbf{x}_i$'s along could not be linearly independent. Hence we finally obtain a spanning set $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p$ together with $(m-p)\mathbf{y}_i$'s and thus $m-p \geq 0$. Therefore $p \leq m$.

Theorem 15.6 *If $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ and $\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_m$ are both basis for V , then $m = n$.*

Proof Both sets are linearly independent and both span V . Using Thm. (15.5), since the $\mathbf{e}_i$'s are linearly independent and the $\mathbf{f}_i$'s span V , we have $n \leq m$. Since the $\mathbf{e}_i$'s span V and the $\mathbf{f}_i$'s are linearly independent, $m \leq n$. Therefore $m = n$.

If V possesses a basis, then all the same number of elements. We say that V is *finite dimensional* and the number of elements in the basis is called the dimension of V and is denoted by $\dim V$. If V possesses no finite basis, then we say that V is *infinite dimensional*.

Theorem 15.7 *If V has dimension n , there cannot exist a set of more than n linearly independent vectors in V . Any set of more than n linear combinations of n given vectors is always linearly dependent.*

Theorem 15.8 *The maximal number of linearly independent vectors in $\mathbb{R}_n$ is n*

Theorem 15.9 *If V has dimension n , there cannot exist a set of less than n vectors which span V .*

Definition Vectors are directed segments. The vector in $\mathbb{R}_n$ whose components are $a_1, a_2, \dots, a_n$ is denoted by

$$\mathbf{a} = \{a_1, a_2, \dots, a_n\} \text{ (braces).}$$

The zero vector in $\mathbb{R}_n$ is denoted as

$$\mathbf{0} = \{0, 0, \dots, 0\},$$

with n components each 0.

Definition We call p vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ **linearly dependent** if there are p numbers $\lambda_1, \lambda_2, \dots, \lambda_p$ not all 0, such that

$$\lambda_1 \mathbf{a}_1 + \lambda_2 \mathbf{a}_2 + \dots + \lambda_p \mathbf{a}_p = \mathbf{0} \text{ (the zero vector).}$$

If no such p numbers exist, then the p vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ are called **linearly independent**.

Theorem 15.10 *If a subset of the p vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ is linearly dependent, then the set of p vectors is itself linearly dependent.*

Suppose the first r vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_r$, ($r < p$) are linearly dependent. Then there are r numbers $\lambda_1, \lambda_2, \dots, \lambda_r$ not all zero such that

$$\lambda_1 \mathbf{a}_1 + \lambda_2 \mathbf{a}_2 + \dots + \lambda_r \mathbf{a}_r = \mathbf{0}.$$

Setting $\lambda_{r+1}, \lambda_{r+2}, \dots, \lambda_p$ to zero we have

$$\lambda_1 \mathbf{a}_1 + \lambda_2 \mathbf{a}_2 + \dots + \lambda_r \mathbf{a}_r + \lambda_{r+1} \mathbf{a}_{r+1} + \dots + \lambda_p \mathbf{a}_p = \mathbf{0}.$$

where the numbers $\lambda_1, \lambda_2, \dots, \lambda_p$ are not all zero. Therefore the p vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ are linearly dependent.

Theorem 15.11 *If the p vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ are linearly independent, then so is every subset of these p vectors.*

For otherwise, by Theorem 15.10, the p vectors would be linearly dependent.

Theorem 15.12 *If the p vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ are linearly dependent and if $p > 1$, then at least one of these vectors is a linear combination of the others.*

For, there are numbers λ_i , which do not all vanish such that

$$\lambda_1 \mathbf{a}_1 + \lambda_2 \mathbf{a}_2 + \dots + \lambda_p \mathbf{a}_p = \mathbf{0}.$$

Suppose $\lambda_p \neq 0$. Then we can solve for $\mathbf{a}_p$,

$$\mathbf{a}_p = -\frac{\lambda_1}{\lambda_p} \mathbf{a}_1 - \frac{\lambda_2}{\lambda_p} \mathbf{a}_2 - \dots - \frac{\lambda_{p-1}}{\lambda_p} \mathbf{a}_{p-1},$$

i.e. $\mathbf{a}_p$ is a linear combination of the vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_{p-1}$.

Theorem 15.13 *If one of the vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ is a linear combination of the others, then vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ are linearly dependent.*

Suppose

$$\mathbf{a}_1 = \lambda_2 \mathbf{a}_2 + \lambda_3 \mathbf{a}_3 + \dots + \lambda_p \mathbf{a}_p.$$

Then we have that

$$\mathbf{a}_1 - \lambda_2 \mathbf{a}_2 - \lambda_3 \mathbf{a}_3 - \dots - \lambda_p \mathbf{a}_p = \mathbf{0},$$

and since the coefficient of $\mathbf{a}_1$ is not zero, the vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ are linearly dependent.

Theorem 15.14 *If the vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ are linearly independent, and if the vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p, \mathbf{b}$ are linearly dependent, then $\mathbf{b}$ is a linear combination of $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$.*

We have a relation of the form

$$\lambda_1 \mathbf{a}_1 + \lambda_2 \mathbf{a}_2 + \dots + \lambda_p \mathbf{a}_p + \lambda_{p+1} \mathbf{b} = \mathbf{0},$$

where not all the λ_i vanish. Now λ_{p+1} cannot be 0 because the last term would drop out and then all the other λ_i would vanish because of the linear independence of the $\mathbf{a}_i$ vectors. Therefore $\lambda_{p+1} \neq 0$ and so we have

$$\mathbf{b} = -\frac{\lambda_1}{\lambda_{p+1}} \mathbf{a}_1 - \frac{\lambda_2}{\lambda_{p+1}} \mathbf{a}_2 - \dots - \frac{\lambda_p}{\lambda_{p+1}} \mathbf{a}_p.$$

Theorem 15.15 *If the vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ are linearly independent, and if $\mathbf{b}$ is not a linear combination of $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$, then the vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p, \mathbf{b}$ are linearly independent.*

Theorem 15.15 follows from Theorem 15.14,

Let L be a *non-empty* set of vectors of $\mathbb{R}_n$ with the properties:

1. If $\mathbf{a}$ is a vector of the set L , then $\lambda \mathbf{a}$ belongs to L for every real λ .
2. If $\mathbf{a}$ and $\mathbf{b}$ are two (not necessarily distinct) vectors of the set L , then the vector $\mathbf{a} + \mathbf{b}$ also belongs to L .

Such a set of vectors, satisfying 1. and 2., is called a **vector space** in $\mathbb{R}_n$.

Definition Dimension and Basis

Let L be any vector space in $\mathbb{R}_n$. The maximal number of linearly independent vectors is $\leq n$. Let p be this maximal number of independent vectors in L . We call p the **dimension** of L . Thus,

$$0 \leq p \leq n.$$

We call any system of p linearly independent vectors of L , such as $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ a **basis** of L .

Theorem 15.16 *Every vector of L can be represented in exactly one way as a linear combination of the basis vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$.*

Theorem 15.17 *The dimension of the vector space spanned by the vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ is equal to the maximal number of linearly independent vectors among $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$.*

Theorem 15.18 *Any k ($k \leq p$) linearly independent vectors*

$$\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_k$$

of L can be extended to form a basis of L by suitably adjoining to them $p - k$ other vectors $\mathbf{a}_{k+1}, \mathbf{a}_{k+2}, \dots, \mathbf{a}_p$.

Let L' and L'' be any two vector spaces in $\mathbb{R}_n$. By the **intersection** D of L' and L'' we mean the set of all those vectors belonging to both L' and L'' .

If we form the totality S of all vectors of the form $\mathbf{a}' + \mathbf{a}''$, where $\mathbf{a}'$ belongs to L' and $\mathbf{a}''$ belongs to L'' . S is called the **sum** of L' and L'' .

Theorem 15.19 *If the dimensions of the vector spaces L', L'', D , and S are respectively p', p'', d and s , then*

$$p' + p'' = d + s.$$

Subspaces

We define a *vector subspace* of M of a vector space V to be a subset of V which is closed under addition and scalar multiplication : if $\mathbf{x}$ and $\mathbf{y}$ are in M then so is $\mathbf{x} + \mathbf{y}$ and so is $\lambda\mathbf{x}$ for any scalar λ .

Theorem 15.20 *If V has a finite dimension n and M is a non-trivial subspace of V , then M has finite dimension m where $0 < m < n$.*

Theorem 15.21 (Basis Reduction Theorem) *If $V = \text{span}(\mathbf{v}_1, \dots, \mathbf{v}_m)$, then either $(\mathbf{v}_1, \dots, \mathbf{v}_m)$ is a basis of V or some $\mathbf{v}_i$ can be removed to obtain a basis of V .*

Proof Suppose $V = \text{span}(\mathbf{v}_1, \dots, \mathbf{v}_m)$. We start with the list $S = (\mathbf{v}_1, \dots, \mathbf{v}_m)$, and sequentially run through all vectors $\mathbf{v}_k$ for $k = 1, 2, \dots, m$ to determine whether to keep or remove them from S :

Step 1. If $\mathbf{v}_1 = \mathbf{0}$, then remove $\mathbf{v}_1$ from S . Otherwise, leave S unchanged.

Step k. If $\mathbf{v}_k \in \text{span}(\mathbf{v}_1, \dots, \mathbf{v}_{k-1})$, then remove $\mathbf{v}_k$ from S . Otherwise, leave S unchanged.

The final list S still spans V since, at each step, a vector was only discarded if it was already in the span of the previous vectors. The process also ensures that no vector is in the span of the previous vectors. Hence by Linear Dependence [see Definition (15)] the final list is linearly independent. Therefore, the final list S is a basis of V .

Theorem 15.22 (Basis Extension Theorem) *If M is a subspace of V , where V is finite dimensional, then any basis of M is contained in a basis of V .*

Proof Suppose V is finite dimensional and that $(\mathbf{v}_1, \dots, \mathbf{v}_m)$ is linearly independent. Since V is finite dimensional, there exists a finite set of vectors $(\mathbf{w}_1, \dots, \mathbf{w}_n)$ that span V . We want to adjoin some of the $\mathbf{w}_k$ to $(\mathbf{v}_1, \dots, \mathbf{v}_m)$ in order to create a basis of V .

Step 1. If $\mathbf{w}_l \in \text{span}(\mathbf{v}_1, \dots, \mathbf{v}_m)$, then $S = (\mathbf{v}_1, \dots, \mathbf{v}_m)$. Otherwise, $S = (\mathbf{v}_1, \dots, \mathbf{v}_m, \mathbf{w}_l)$.

Step k. If $\mathbf{w}_k \in \text{span}(S)$, then leave S unchanged. Otherwise adjoin $\mathbf{w}_k$ to S .

After each step, the list S is still linearly independent because we only adjoined $\mathbf{w}_k$ if it was not in the span of the previous vectors. After n steps, $\mathbf{w}_k \in \text{span}(S)$ for all $k = 1, 2, \dots, n$. Since $(\mathbf{w}_1, \dots, \mathbf{w}_n)$ was a spanning set, S spans V so that S is indeed a basis of V .

The range and null space of a linear transformation

If $\theta : V \rightarrow W$ is a linear transformation, where V and W are not necessarily finite dimensional, the image of θ is called the *range* of θ and the kernel is called the *null space*. The range and null space image and kernel of θ considered as a group homomorphism, are subgroups of W and V respectively.

Theorem 15.23 *The range R of $\theta : V \rightarrow W$ is a vector subspace of W .*

Proof By Thm (5.2) the range is a subgroup of W . If $\mathbf{y} \in R$, so that $\mathbf{y} = \theta\mathbf{x}$ for some $\mathbf{x} \in V$, $\theta(\lambda\mathbf{x}) = \lambda\mathbf{y}$ and so $\lambda\mathbf{y} \in R$ for any scalar λ . Hence R is a subspace.

Theorem 15.24 *The null space N of $\theta : V \rightarrow W$ is a vector subspace of V .*

Proof By Thm. (5.1) the null space is a subgroup. If $\mathbf{x} \in N$, so that $\theta\mathbf{x} = \mathbf{0}$, then $\theta(\lambda\mathbf{x}) = \lambda\mathbf{0} = \mathbf{0}$, and hence $\lambda\mathbf{x} \in N$. Therefore N is a subspace.

If V and W are finite dimensional, and

$$\dim V = n, \quad \dim W = m,$$

we see by Thm (15.20) that the range R has finite dimension $\leq m$, and the null space N finite dimension $\leq n$. The dimension of R is called the *rank* of θ and that of N is called the *nullity* of θ . These are denoted by $r(\theta)$ and $\nu(\theta)$ respectively.

Therefore $0 \leq r(\theta) \leq m$ and $0 \leq \nu(\theta) \leq n$. If θ is the zero transformation, $r(\theta) = 0$ and $\nu(\theta) = n$, whereas if θ is an isomorphism $r(\theta) = n$ and $\nu(\theta) = 0$.

In both cases rank plus nullity is n . and this result is generally true.

Theorem 15.25 (Rank / Nullity Theorem) *If $\theta : V \rightarrow W$ is a linear transformation between finite dimensional vector spaces with $\dim V = n$, then*

$$r(\theta) + \nu(\theta) = n.$$

Theorem 15.26 (Grassmann dimension formula) *If M and N are subspaces of a finite dimensional vector space V , then*

$$\dim(M + N) = \dim M + \dim N - \dim(M \cap N).$$

Proof Let $\dim M = \mu$, $\dim N = \nu$, $\dim(M \cap N) = \rho$, which are all finite. Take $\mathbf{z}_1, \dots, \mathbf{z}_\rho$ as the basis of $M \cap N$. Since $M \cap N$ is a subspace of both M and N , it can be extended as follows to a basis for both M and N :

$$\begin{array}{lll} \mathbf{z}_1, \dots, \mathbf{z}_\rho, & \mathbf{x}_1, \dots, \mathbf{x}_{\mu-\rho} & \text{of } M \text{ and} \\ \mathbf{z}_1, \dots, \mathbf{z}_\rho, & \mathbf{y}_1, \dots, \mathbf{y}_{\nu-\rho} & \text{of } N \end{array}$$

We now prove that

$$\mathbf{z}_1, \dots, \mathbf{z}_\rho, \quad \mathbf{x}_1, \dots, \mathbf{x}_{\mu-\rho}, \quad \mathbf{y}_1, \dots, \mathbf{y}_{\nu-\rho}$$

is a basis of $M + N$.

Any element of M is a linear combination of $\mathbf{z}_1, \dots, \mathbf{z}_\rho, \mathbf{x}_1, \dots, \mathbf{x}_{\mu-\rho}$ and any element of N is a linear combination of $\mathbf{z}_1, \dots, \mathbf{z}_\rho, \mathbf{y}_1, \dots, \mathbf{y}_{\nu-\rho}$. Therefore, any element of $M + N$ is a linear combination of the $\mathbf{z}$'s, $\mathbf{x}$'s and $\mathbf{y}$'s, and so this set spans $M + N$.

Now suppose that

$$\sum_{i=1}^{\rho} \alpha_i \mathbf{z}_i + \sum_{j=1}^{\mu-\rho} \beta_j \mathbf{x}_j + \sum_{k=1}^{\nu-\rho} \gamma_k \mathbf{y}_k = \mathbf{0}.$$

Then

$$\sum_{k=1}^{\nu-\rho} \gamma_k \mathbf{y}_k = - \sum_{i=1}^{\rho} \alpha_i \mathbf{z}_i - \sum_{j=1}^{\mu-\rho} \beta_j \mathbf{x}_j \in M,$$

and also in N , and so is in $M \cap N$. Therefore $\sum_{k=1}^{\nu-\rho} \gamma_k \mathbf{y}_k = \sum_{i=1}^{\rho} \delta_i \mathbf{z}_i$ since $\mathbf{z}_1, \dots, \mathbf{z}_\rho$ are a basis for $M \cap N$. But the $\mathbf{y}_i$'s and $\mathbf{z}_i$'s together form a linearly independent set, being a basis of N , and therefore all the γ_i 's are zero. Similarly all the β_i 's are zero, and so $\sum_{i=1}^{\rho} \alpha_i \mathbf{z}_i = \mathbf{0}$ and all the α_i 's are zero because $\mathbf{z}_1, \dots, \mathbf{z}_\rho$ are linearly independent.

Therefore $\mathbf{z}_1, \dots, \mathbf{z}_\rho, \quad \mathbf{x}_1, \dots, \mathbf{x}_{\mu-\rho}, \quad \mathbf{y}_1, \dots, \mathbf{y}_{\nu-\rho}$ are linearly independent and hence form a basis of $M + N$ and we have

$$\dim(M + N) = \rho + \mu - \rho + \nu - \rho = \mu + \nu - \rho, \text{ or}$$

$$\dim(M + N) = \dim M + \dim N - \dim(M \cap N). \quad \blacksquare$$

Thm (15.26) is basically Inclusion-Exclusion for vector spaces. The main reason this is the case is that it is possible to build a sequence of vector spaces where the Euler characteristic (alternating sum of dimensions) vanishes, which is directly related to the Inclusion-Exclusion formula. The two principles are the same idea showing up in different contexts: counting elements versus measuring linear independence.

16 Fundamental Theorem of Algebra

Axiom I - Axiom of inequality for $\mathbb{R}$. Among the real numbers, $\mathbb{R}$, there is a set called the positive numbers which satisfies the conditions: (i) for any number $a \in \mathbb{R}$ exactly one of the three alternatives holds; a is positive,

$a = 0$, or $-a$ is positive; (ii) any finite sum or produce of positive numbers is positive.

Axiom C - Axiom of continuity. Suppose that an infinite sequence $x_1, x_2, \dots, x_n \dots$ is such that $x_{n+1} \geq x_n$ for all n , and there is a number $M \in \mathbb{R}$ such that $x_n \leq M$ for all n . Then there is a number $L \leq M$ such that $x_n \rightarrow L$ as $n \rightarrow \infty$ and $x_n \leq L$ for all n .

Theorem 16.1 (*Nested Intervals Theorem*): Suppose that

$$I_n = \{x : a_n \leq x \leq b_n\}, \quad n = 1, 2, \dots,$$

is a sequence of closed intervals such that $I_{n+1} \subset I_n$, for each n . If $\lim_{n \rightarrow \infty} (b_n - a_n) = 0$, then there is one and only one number x_0 which is in every I_n .

Proof By hypothesis, we have $a_n \leq a_{n+1}$ and $b_{n+1} \leq b_n$. Since $a_n < b_n$ for every n , the sequence $\{a_n\}$ is nondecreasing and bounded from above by b_1 . Similarly, the sequence b_n is non-increasing and bounded from below by a_1 . Using Axiom C, we conclude that there are numbers x_0 and x'_0 such that $a_n \rightarrow x_0, a_n \leq x_0$, and $b_n \rightarrow x'_0, b_n \geq x'_0$ as $n \rightarrow \infty$. Using the fact that $b_n - a_n \rightarrow 0$ as $n \rightarrow \infty$, we find that $x_0 = x'_0$ and

$$a_n \leq x_0 \leq b_n$$

for every n . Thus x_0 is in every I_n .

To show that x_0 is the only number in every I_n , suppose there is another number x_1 also in every I_n . Define $\epsilon = |x_1 - x_0|$. Now, since $a_n \rightarrow x_0$ and $b_n \rightarrow x_0$, there is an integer N_1 , such that $a_{N_1} > x_0 - \epsilon$; also there is an integer N_2 such that $b_{N_2} < x_0 + \epsilon$. These inequalities imply that I_n cannot contain x_1 for any n beyond N_1 and N_2 . Thus x_1 is not in every I_n . ■

Theorem 16.2 Suppose f is continuous on $[a, b]$, $x_n \in [a, b]$ for each n , and $x_n \rightarrow x_0$. Then $x_0 \in [a, b]$ and $f(x_n) \rightarrow f(x_0)$.

Proof Since $x_n \geq a$ for each n , it follows from the theorem on the limit of inequalities for sequences that $x_0 \geq a$. Likewise $x_0 \leq b$ so $x_0 \in [a, b]$. If $a < x_0 < b$, then $f(x_n) \rightarrow f(x_0)$ on account of the composite function theorem. The same conclusion holds if $x_0 = a$ or b .

Theorem 16.3 (*Intermediate-value theorem*): Suppose f is continuous on $[a, b]$, $c \in \mathbb{R}$, $f(a) < c$, and $f(b) > c$. Then there is at least one number x_0 on $[a, b]$ such that $f(x_0) = c$.

There may be more than one such point!

Proof Define $a_1 = a$ and $b_1 = b$. Then observe that $f((a_1 + b_1)/2)$ is either equal to c , greater than c , or less than c . If it equals c , then choose $x_0 = (a_1 + b_1)/2$ and the result is proved. If $f((a_1 + b_1)/2) > c$, then define $a_2 = a_1$ and $b_2 = (a_1 + b_1)/2$. If $f((a_1 + b_1)/2) < c$, then define $a_2 = (a_1 + b_1)/2$ and $b_2 = b_1$.

In each of the last two cases we have $f(a_2) < c$ and $f(b_2) > c$. Again compute $f((a_2 + b_2)/2)$. If this value equals c , the result is proved. If $f((a_2 + b_2)/2) > c$, set $a_3 = a_2$ and $b_3 = (a_2 + b_2)/2$. If $f((a_2 + b_2)/2) < c$, then define $a_3 = (a_2 + b_2)/2$ and $b_3 = b_2$.

Continuing in this way, we either find a solution in a finite number of steps or we find a sequence $\{[a_n, b_n]\}$ of closed intervals each of which is one of the two halves of the preceding one, and for which we have

$$b_n - a_n = 2^{1-n}(b_1 - a_1),$$

$$f(a_n) < c, \quad f(b_n) > c, \text{ for each } n,$$

From the Nested Intervals Theorem it follows that there is a unique point x_0 in all these intervals and

$$\lim_{n \rightarrow \infty} a_n = x_0 \text{ and } \lim_{n \rightarrow \infty} b_n = x_0.$$

From the theorem of continuous function on a sequence of points in a closed interval, we conclude that

$$f(a_n) \rightarrow f(x_0) \text{ and } f(b_n) \rightarrow f(x_0).$$

From the inequalities about and the limit of inequalities it follows that $f(x_0) \leq c$ and $f(x_0) \geq c$ so that $f(x_0) = c$.

Theorem 16.4 (Extreme Value Theorem for real-valued functions of two real variables:) *Let D be the closed disk of radius $r > 0$ and centered at the origin, i.e.,*

$$D = \{(x_1, x_2) \in \mathbb{R}^2 | x_1^2 + x_2^2 \leq r^2\}.$$

Let $f : D \rightarrow \mathbb{R}$ be a continuous function on the closed disk $D \subset \mathbb{R}^2$. Then f is bounded and attains its minimum and maximum values on D . In other words, there exists points $x_m, x_M \in D$ such that

$$f(x_m) \leq f(x) \leq f(x_M)$$

for every possible choice of point $x \in D$.

Definition A polynomial over a field F is a *rational* and *integral* expression of the form

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0, \tag{1}$$

where the coefficients $a_i \in F, i = 1, 2, \dots, n$ and *rational* means free from fractional powers and *integral* means never in the denominator of a fraction. A polynomial function is a function which can be written in the form of Eq. (1).

Theorem 16.5 (Eisenstein's Irreducibility Criterion) *For a given prime p , let $a(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0$ be a polynomial with integral coefficients, such that $a_n \not\equiv 0 \pmod{p}$, $a_{n-1} \equiv a_{n-2} \equiv \dots \equiv a_0 \equiv 0 \pmod{p}$, $a_0 \not\equiv 0 \pmod{p^2}$. Then $a(x)$ is irreducible over the field of rational.*

Proof 1 Suppose

$$a(x) = gh,$$

where

$$\begin{aligned} g &= g_0 x^h + g_1 x^{h-1} + \dots + g_{k-1} x + g_k \\ h &= h_0 x^l + h_1 x^{l-1} + \dots + h_{l-1} x + h_l \end{aligned}$$

Then g and h have integral coefficients since f has. Comparing coefficients, we set

$$\begin{aligned} a_0 &= g_k h_l \\ a_1 &= g_k h_{l-1} + g_{k-1} h_l \\ a_2 &= g_k h_{l-2} + g_{k-1} h_{l-1} + \dots + g_{k-2} h_l \\ \dots &\quad \dots \\ a_l &= g_k h_{l-k} + g_{k-1} h_{l-k+1} + \dots + g_1 h_{l-1} + h_l \\ \dots &\quad \dots, \end{aligned}$$

where the h_i with possible negative subscripts are 0 and where

$$h_0 = 1.$$

if now every a_i , other than a_0 is divisible by p and a_n not by p^2 , we conclude from the first equation that either g_k or h_l is divisible by p and the other is not. But when g_k is divisible by p , we conclude from the following equations that $g_{k-1}, g_{k-2}, \dots, g_1$ are divisible by p , and from the last equation that h_l is also divisible by p which is a contradiction from the assumption that a_n is not divisible by p^2 .

Proof 2 In any possible factorization ($n = m + k$)

$$a(x) = bc = (b_n x^n + b_{n-1} x^{n-1} + \dots + b_1 x + b_0)(c_k x^k + c_{k-1} x^{k-1} + \dots + c_1 x + c_0)$$

we may assume that both factors have integral coefficients. Since $a_0 = b_0 c_0$ the hypothesis $a_0 \not\equiv 0 \pmod{p^2}$ means that not both b_0 and c_0 are divisible by p . Suppose $b_0 \not\equiv 0 \pmod{p}$, while $c_0 \equiv 0 \pmod{p}$. But $b_m c_k = a_n \not\equiv 0 \pmod{p}$, so $c_k \not\equiv 0 \pmod{p}$. Pick the smallest $r \leq k$ for which $c_r \not\equiv 0 \pmod{p}$, with $c_{r-1} \equiv \dots \equiv c_0 \equiv 0 \pmod{p}$. Then

$$a_r = b_0 c_r + b_1 c_{r-1} + \dots + b_r c_0 = b_0 c_r \pmod{p}.$$

But $b_0 \not\equiv 0 \pmod{p}$ and $c_r \not\equiv 0 \pmod{p}$ give $a_r \not\equiv 0 \pmod{p}$, for p is a prime. By the hypothesis, the only coefficient a_r for which this is possible is a_n , therefore $r = n$: the degree of the second of the proposed factors must be n (the same as degree of $a(x)$, so that $a(x)$ is irreducible. ■

This criterion can be applied to the cyclotomic polynomial

$$\phi(x) = \frac{x^p - 1}{x - 1} = x^{p-1} + x^{p-2} + \dots + x + 1. \quad (2)$$

This is the polynomial whose roots are the primitive p -th roots of unity. The Eisenstein criterion does not apply to Eq. (2) as it stands, but a simple change of variables $y = x - 1$ does the trick, for the binomial expansion gives

$$\frac{x^p - 1}{x - 1} = \frac{(y + 1)^p - 1}{y} = y^{p-1} + p y^{p-2} + \frac{p(p-1)}{1 \cdot 2} y^{p-3} + \dots + p.$$

The binomial coefficients which appear on the right are all integers divisible by the prime p , for p occurs in each numerator as a factor and can never be cancelled out by the smaller integers in the denominator. The polynomial in y satisfies the hypothesis of the Eisenstein criterion and hence is irreducible. Therefore the original cyclotomic polynomial, Eq. (2), $\phi(x)$ is irreducible.

Theorem 16.6 (Euler-Gauss: Fundamental Theorem of Algebra) *Every polynomial $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$ of positive degree with complex coefficients has a complex root.*

We can regard $(x, y) \rightarrow |f(x + iy)|$ as a function $\mathbb{R}^2 \rightarrow \mathbb{R}$. We denote this function by $|f(\cdot)|$ or $|f|$. Since polynomials and the square root function are continuous, then $|f|$ is also continuous.

Lemma 16.7 *Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be any polynomial function. Then there exists a point $z_0 \in \mathbb{C}$ where the function $|f|$ attains a minimum value in $\mathbb{R}$.*

Proof If f is a constant polynomial function, then the statement of Lemma (16.7) is trivial, so we may take $z_0 = 0$. If f is not constant, then the degree of the polynomial defining f is at least one. In this case, we can denote f explicitly as Eq. (1) with $a_n \neq 0$. Now assume that $z \neq 0$, and set $A = \max\{|a_0|, |a_1|, \dots, |a_{n-1}|\}$. We can obtain a lower bound for $|f(z)|$ as follows:

$$\begin{aligned} |f(z)| &= |a_n||z^n| \left| 1 + \frac{a_{n-1}}{a_n z} + \dots + \frac{a_0}{a_n z^n} \right| \\ &\geq |a_n||z^n| \left(1 - \frac{A}{|a_n|} \sum_{k=1}^{\infty} \frac{1}{|z|^k} \right) = |a_n||z^n| \left(1 - \frac{A}{|a_n|} \frac{1}{|z| - 1} \right) \end{aligned}$$

For all $z \in \mathbb{C}$ such that $|z| \geq 2$, we can further simplify this expression to

$$|f(z)| \geq |a_n||z^n| \left(1 - \frac{2A}{|a_n||z|} \right).$$

It follows from this inequality that there is an $r > 0$ such that $|f(z)| > |f(0)|$, for all $z \in \mathbb{C}$ satisfying $|z| > r$. Let $D \subset \mathbb{R}^2$ be the disk of radius r centered at 0, and define a function $g : D \rightarrow \mathbb{R}$, by

$$g(x, y) = |f(x + iy)|.$$

Since g is continuous, we can apply Thm (16.4) in order to obtain a point $(x_0, y_0) \in D$ such that g attains its minimum at (x_0, y_0) . By the choice of r we have that for $z \in D$, $|f(z)| > |g(0, 0)| \geq |g(x_0, y_0)|$. Therefore $|f|$ attains its minimum at $z = x_0 + iy_0$.

Proof of Thm (16.6): We rely on the fact that the function $|f(x)|$ attains its minimum value by Lemma (16.7). Let $z_0 \in \mathbb{C}$ be a point where the minimum is attained. We will show that if $f(z_0) \neq 0$, then z_0 is not a minimum value, thus proving that the minimum value of $|f(z)|$ is zero and therefore $f(z_0) = 0$.

If $f(z_0) \neq 0$, then we can define a new function $g : \mathbb{C} \rightarrow \mathbb{C}$ by setting

$$g(z) = \frac{f(z + z_0)}{f(z_0)}, \text{ for all } z \in \mathbb{C}.$$

Not that g is a polynomial of degree n , and that the minimum of $|f|$ is attained at z_0 if and only if the minimum of $|g|$ is attained at $z = 0$. Moreover, it is clear that $g(0) = 1$. Therefore g is a polynomial of the form

$$g(z) = b_n z^n + \dots + b_k z^k + 1,$$

with $n \geq 1$ and $b_k \neq 0$ for some $1 \leq k \leq n$. Let $b_k = |b_k|e^{i\theta}$, and consider z of the form

$$z = r|b_k|^{-1/k} e^{i(\pi - \theta)/k}, \quad (3)$$

with $r > 0$. For z of this form we have

$$g(z) = 1 - r^k + r^{k+1}h(r),$$

with h a polynomial. Then for $r < 1$, we have by the triangle inequality that

$$|g(z)| \leq 1 - r^k + r^{k+1}|h(r)|.$$

For $r > 0$ sufficiently small we have $r|h(r)| < 1$, by the continuity of the function $rh(r)$ and the fact that it vanishes at $r = 0$. Hence we find that

$$|g(z)| \leq 1 - r^k(1 - r|h(r)|) < 1,$$

for some z having the form given in Eq. (3) with $r \in (0, r_0)$ and $r > 0$ sufficiently small. But then the minimum of $|g| : \mathbb{C} \rightarrow \mathbb{R}$ cannot possibly be equal to 1. ■

17 Synthetic Division

Let $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$ and consider dividing this polynomial by a linear factor $(x - h)$. This division can certainly be performed by long division, but a more compact method is to use Synthetic Division which is outlined below. Let the quotient when

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$$

is divided by $(x - h)$ be given by

$$Q(x) = b_{n-1} x^{n-1} + b_{n-2} x^{n-2} + \dots + b_1 x + b_0.$$

We then can represent $f(x)$ with the following identity:

$$f(x) = (x - h)Q(x) + R,$$

with quotient $Q(x)$ and remainder R . The right-hand side of this identity can be grouped according to terms which multiply the same power of x as follows:

$$\begin{array}{c|c|c|c|c|c|c} b_{n-1}x^n & b_{n-2}x^{n-1} & b_{n-3}x^{n-2} & b_{n-4}x^{n-3} & \dots & b_0x & R \\ -hb_{n-1}x^{n-1} & -hb_{n-2}x^{n-2} & -hb_{n-3}x^{n-3} & \dots & -hb_1x & -hb_0 & \end{array}$$

Equating the coefficients of x on both sides of the we get the following recursive relationships:

$$\begin{aligned} b_{n-1} &= a_n \\ b_{n-2} &= hb_{n-1} + a_{n-1} \\ b_{n-3} &= hb_{n-2} + a_{n-2} \\ &\dots = \dots \\ b_1 &= hb_2 + a_2 \\ b_0 &= hb_1 + a_1 \\ R &= hb_0 + a_0 \end{aligned}$$

These recursive relations can be represented in the following table (known as Synthetic Division):

$$\begin{array}{c|cccccc} h & a_n & a_{n-1} & \dots & a_2 & a_1 & a_0 \\ & & hb_{n-1} & \dots & hb_2 & hb_1 & hb_0 \\ \hline & b_{n-1} & b_{n-2} & \dots & b_1 & b_0 & R \end{array}$$

As an example let us use Synthetic division to perform the following division:

$$\frac{2x^3 - 7x^2 + 5}{x - 3}$$

Synthetic division give us

$$3 \left| \begin{array}{cccc} 2 & -7 & 0 & 5 \\ & 6 & -3 & -9 \\ \hline 2 & -1 & -3 & -4 \end{array} \right.$$

We can read the terms in the Quotient and Remainder from the terms below the horizontal line in the above.

$$\begin{aligned} 2x^3 - 7x^2 + 5 &= (x - 3)(2x^2 - x - 3) - 4 \text{ with quotient } Q(x) \text{ and remainder } R \text{ given by:} \\ Q(x) &= 2x^2 - x - 3 \\ R &= -4 \end{aligned}$$

18 Rational Roots

19 Elementary Symmetric Functions

If $a_n = 1$ we say the polynomial is in *monic* form. It will often be convenient to adopt this convention; and in such cases we shall write the polynomial as

$$f(x) = x^n + p_1x^{n-1} + p_2x^{n-2} + \dots + p_{n-1}x + p_n.$$

The roots $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ of a polynomial are the values of x such that $f(x = \alpha_i) = 0$. The roots are in general complex numbers. According to a theorem of Viète, the relationship between the roots and the coefficients for a monic polynomial is given by the

$$\begin{aligned} f(x) &= x^n + p_1x^{n-1} + \dots + p_{n-1}x + p_n \\ &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \dots (x - \alpha_n). \end{aligned}$$

Expanding the product on the right-hand side we find the explicit relationships:

$$\begin{aligned} p_1 &= -(\alpha_1 + \alpha_2 + \dots + \alpha_n) \\ p_2 &= (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \dots + \alpha_2\alpha_3 + \dots + \alpha_{n-1}\alpha_n) \\ p_3 &= -(\alpha_1\alpha_2\alpha_3 + \alpha_1\alpha_2\alpha_4 + \dots + \alpha_2\alpha_3\alpha_4 + \dots + \alpha_{n-2}\alpha_{n-1}\alpha_n) \\ &\vdots \\ p_n &= -(1)^n(\alpha_1\alpha_2\alpha_3\alpha_4 \dots \alpha_{n-1}\alpha_n) \end{aligned}$$

These relationships are encapsulated in the following theorem

Theorem 19.1 *In every algebraic equation, the coefficient of whose highest term is unity, the coefficient p_1 of the second term with its sign changed is equal to the sum of the roots.*

The coefficient p_2 of the third terms is equal to the sum of the products of the roots taken two by two.

The coefficient of the fourth term with its sign changed is equal to the sum of the products of the roots taken three by three; and so on, the signs of the coefficients being taken alternately negative and positive, and the number of roots multiplied together in each term of the corresponding function of the roots increasing by unity till finally that function is reached with consists of the produces of the n roots.

Definition The Elementary Symmetric functions $c_1, c_2, c_3, \dots, c_n$ are defined as functions of n indeterminants $x_1, x_2, x_3, \dots, x_n$ as follows

$$\begin{array}{lll}
s(x_1) & = x_1 + x_2 + \dots + x_n & = c_1 \\
s(x_1x_2) & = x_1x_2 + x_1x_3 + \dots + x_2x_3 + \dots + x_{n-1}x_n & = c_2 \\
s(x_1x_2, x_3) & = x_1x_2x_3 + x_1x_2x_4 + \dots + x_2x_3x_4 + \dots + x_{n-2}x_{n-1}x_n & = c_3 \\
\vdots & & \vdots \\
s(x_1x_2 \dots x_n) & = x_1x_2x_3x_4 \dots x_{n-1}x_n & = c_n
\end{array} \tag{4}$$

Theorem 19.2 The sums of the similare powers of the roots of an equation can be expressed in terms of the coefficients (the elementary symmetric functions).

Proof Consider the polynomial

$$\begin{aligned}
f(x) &= x^n + p_1x^{n-1} + p_2x^{n-2} \dots + p_{n-1}x + p_n, \\
&= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \dots (x - \alpha_n) = 0.
\end{aligned}$$

Define the following terms for the sums of the powers of the roots:

$$s_i = \sum_{j=1}^n \alpha_j^i, \quad i = 0, 1, 2, \dots$$

Given that $f(x)$ is a polynomial we can write the derivative of $f(x)$ as

$$\begin{aligned}
f'(x) &= \frac{f(x)}{x - \alpha_1} + \frac{f(x)}{x - \alpha_2} + \dots + \frac{f(x)}{x - \alpha_n} \\
&= nx^{n-1} + (n-1)p_1x^{n-2} + (n-2)p_2x^{n-3} + \dots + 2p_{n-2}x + p_{n-1}
\end{aligned}$$

Using the method of Synthetic Division to divide $f(x)$ by the monomial $(x - a)$ we write out the sequence of computations as follows;

$$\begin{array}{c|c|c|c}
\begin{array}{l} \frac{f(x)}{x-a} \\ \\ \\ \end{array} & \begin{array}{l} x^{n-1} + a \\ +p_1 \end{array} & \begin{array}{l} x^{n-2} + a^2 \\ +ap_1 \\ +p_2 \end{array} & \begin{array}{l} x^{n-3} + a^3 \\ +a^2p_1 \\ +ap_2 \\ +p_3 \end{array} & \begin{array}{l} x^{n-4} + \dots + a^{n-2} \\ +a^{n-2}p_1 \\ +a^{n-3}p_2 \\ +\dots \\ +ap_{n-2} \\ +p_{n-1} \end{array}
\end{array}$$

By replacing a by each of the quantities $a_1, a_2, \dots, a_n$ in succession and adding and using $s_p = \sum a^p$ we obtain the following expression for the derivative of $f(x)$:

$$\begin{array}{c|c|c|c}
f'(x) & \begin{array}{l} x^{n-1} + s_1 \\ +np_1 \end{array} & \begin{array}{l} x^{n-2} + s_2 \\ +s_1p_1 \\ +np_2 \end{array} & \begin{array}{l} x^{n-3} + s_3 \\ +s_2p_1 \\ +sp_2 \\ +np_3 \end{array} & \begin{array}{l} x^{n-4} + \dots + s_{n-2} \\ +s_{n-2}p_1 \\ +s_{n-3}p_2 \\ +\dots \\ +sp_{n-2} \\ +np_{n-1} \end{array}
\end{array}$$

Therefore we obtain the following relationships between the s_i 's and the coefficients p_i :

$$\begin{aligned}
s_1 + p &= 0 \\
s_2 + p_1s_1 + 2p_2 &= 0
\end{aligned}$$

$$\begin{aligned}
s_3 + p_1 s_2 + p_2 s_1 + 3p_3 &= 0 \\
s_4 + p_1 s_3 + p_2 s_2 + p_3 s_1 + 4p_4 &= 0 \\
&\dots\dots\dots \\
s_{n-1} + p_1 s_{n-2} + p_2 s_{n-3} + \dots + p_{n-2} s_1 + (n-1)p_{n-1} &= 0
\end{aligned}$$

We now proceed to extend our results to the sums of all positive powers of the roots, $s_n, s_{n+1}, \dots, s_m$ by multiplying the polynomial by x^{m-n} and using the $s_0 = n$, we obtain

$$x^{m-n} f(x) = x^m + p_1 x^{m-1} + p_2 x^{m-2} + \dots + p_n x^{m-n},$$

By replacing x by the roots $a_1, a_2, \dots, a_n$ in succession and adding we have,

$$s_m + p_1 s_{m-1} + p_2 s_{m-2} + \dots + p_n s_{m-n} = 0.$$

Letting m have the values $m = n, n+1, n+2, \dots$ successively and noting that $s_0 = n$, we obtain the following equations

$$\begin{aligned}
s_n + p_1 s_{n-1} + p_2 s_{n-2} + \dots + n p_n &= 0 \\
s_{n+1} + p_1 s_n + p_2 s_{n-1} + \dots + p_n s_1 &= 0 \\
s_{n+2} + p_1 s_{n+1} + p_2 s_n + \dots + p_n s_2 &= 0 \\
&\dots\dots\dots 0
\end{aligned}$$

By transforming the equation $f(x) = 0$ into one whose roots are the reciprocals of $a_1, a_2, a_3, \dots, a_n$, and applying the above formulas we can also express all the negative powers of the roots.

Theorem 19.3 *Every rational symmetric function of $x_1, x_2, \dots, x_n$ is rationally expressible in terms of the elementary symmetric functions.*

Proof It is sufficient to prove this theorem for integral functions only, since fractional symmetric functions can be reduced to a single fraction whose numerator and denominator are both integral symmetric functions. Every integral function of $a_1, a_2, a_3, \dots, a_n$ is the sum of a number of terms of the form $N a_1^p a_2^q a_3^r \dots$, where N is a numerical constant; and if this function be symmetrical we can write it under the form $N \sum a_1^p a_2^q a_3^r \dots$, all the terms being of the same type. Therefore if we can prove that this quantity be expressed rationally in terms of the coefficients, the theorem will be demonstrated.

Let us begin with the symmetric function

$$\sum a_1^p a_2^q = s_p s_q - s_{p+q}. \quad (5)$$

Which can be proven by multiplying s_p and s_q , where

$$\begin{aligned}
s_p &= a_1^p + a_2^p + a_3^p + \dots + a_n^p, \\
s_q &= a_1^q + a_2^q + a_3^q + \dots + a_n^q,
\end{aligned}$$

Therefore

$$s_p s_q = a_1^{p+q} + a_2^{p+q} + a_3^{p+q} + \dots + a_n^{p+q} + a_1^p a + 2^q + a_1^p a + 3^q + \dots, \text{ or}$$

$$s_p s_q = s_{p+q} + \sum a_1^q a_2^q,$$

and finally

$$\sum a_1^p a_2^p = s_p s_q - s_{p+q}.$$

We proceed now to prove a similar expression for the triple function,

$$\sum a_1^p a_2^q a_3^r = s_p s_q s_r - s_{q+r} s_p - s_{r+p} s_q - s_{p+q} s_r + 2s_{p+q+r}. \quad (6)$$

Multiplying together $\sum a_1^p a_2^q$ and s_r , where

$$\begin{aligned} \sum a_1^p a_2^q &= a_1^p a_2^q + a_1^q a_2^p + a_1^p a_3^q + \dots, \\ s_r &= a_1^r + a_2^r + a_3^r + \dots + a_n^r \end{aligned}$$

we obtain an expression consisting of three different parts of the forms $\sum a_1^{p+r} a_2^q$, $\sum a_1^{q+r} a_2^p$, and $\sum a_1^p a_2^q a_3^r$, a formula connecting double and triple symmetric functions. However, by Eq(5):

$$\begin{aligned} \sum a_1^{p+r} a_2^q &= s_{p+r} s_q - s_{p+q+r}, \\ \sum a_1^{q+r} a_2^p &= s_{q+r} s_p - s_{p+q+r}, \\ \sum a_1^p a_2^q &= s_p s_q - s_{p+q}, \end{aligned}$$

Substituting these values we find the triple function expressed as in Eq(6).

In the same manner the quadruple function $\sum a_1^p a_2^q a_3^r a_4^s$, can be made to depend on the triple function $\sum a_1^p a_2^q a_3^r$, and ultimately on $s_1, s_2, s_3, \dots$. Finally every rational symmetric function of the roots may be expressed in terms of the elementary symmetric functions (the coefficients of a polynomial).

The formulas Eq.(5) and Eq.(6) require modification when any of the exponents become equal.

Thus if $p = q$, $a_1^p a_2^q = a_2^p a_1^q$, and the terms in Eq.(5) become equal to two and two; therefore $\sum a_1^p a_2^q = 2 \sum a_1^p a_2^p$ and

$$\sum a_1^p a_2^p = \frac{1}{2}(s_p^2 - s_{2p}).$$

Similarly, if $p = q = r$, the six terms obtained by interchanging the roots in $a_1^p a_2^q a_3^r$ become equal and therefore

$$\sum a_1^p a_2^p a_3^p = \frac{1}{2 \cdot 3} (s_p^3 - 3s_p s_{2p} + 2s_{3p}).$$

And in general, if m exponents become equal, each term is repeated $m!$ times.

Theorem 19.4 (Theorems related to Symmetric Functions) *The sum of the exponents of all the roots in any term of any symmetric function of the roots is equal to the sum of the suffices in each term of the corresponding value in terms of the coefficients.*

The sum of the exponents is of course the same for every term of the symmetric function, and may be called the degree in all the roots of that function. In other words the suffix for each coefficient is equal to the degree in the roots of the corresponding function of the roots; hence in any product of any powers of the coefficients the sum of the suffices must equal to the degree in all the roots of the corresponding function of the roots.

Theorem 19.5 *When an equation is written with binomial coefficients, the expression in terms of the coefficients for any symmetric function of the roots, which is a function of their differences only, is such that the algebraic sum of the numerical factors of all the terms in it is equal to zero.*

Theorem 19.6 (Division Algorithm for Polynomials) *If F is any field, and $a(x) \neq 0$ and $b(x)$ are any two polynomials over F , then we can find polynomials $q(x)$ and $r(x)$ such that*

$$b(x) = q(x)a(x) + r(x),$$

where $r(x)$ is either zero or has a degree less than that of $a(x)$.

Theorem 19.7 *In $F[x]$, any two polynomials a and b have a greatest common divisor (GCD) d satisfying (i) $d|a$ and $d|b$, (ii) $c|a$ and $c|b$ imply that $c|d$. Moreover, (iii) d is a linear combination $d = sa + tb$ of a and b .*

The procedure for determining the GCD of $a(x)$ and $b(x)$ is given in Algorithm (1):

Algorithm 1 Polynomial Euclidean Algorithm

```

1: procedure POLYGCD( $a(x), b(x)$ )
2:   while  $b(x) \neq 0$  do
3:      $r(x) \leftarrow a(x) \bmod b(x)$ 
4:      $a(x) \leftarrow b(x)$ 
5:      $b(x) \leftarrow r(x)$ 
6:   end while
7:   return  $a(x)$ 
8: end procedure

```

20 Cyclotomic Polynomials

Definition The cyclotomic polynomial $\phi(x)$ for non-trivial p -th roots of unity (other than $x = 1$) is given by

$$\phi(x) = \frac{(x^p - 1)}{x - 1} = x^{p-1} + x^{p-2} + \dots + x + 1.$$

Theorem 20.1 $\phi(x)$ is an irreducible polynomial.

Proof As it stands, the Eisenstein criterion for irreducibility doesn't apply. However a change of variable $y = x - 1$ and the binomial expansion changes the form into

$$\frac{(x^p - 1)}{x - 1} = \frac{((y + 1)^p - 1)}{y} = y^{p-1} + py^{p-2} + \frac{p(p-1)}{1 \cdot 2}y^{p-3} + \dots + p.$$

The binomial coefficients which appear on the right hand side are all integers divisible by the prime p , for p occurs in each numerator as a factor, and can never be cancelled out by the smaller integers in the denominator. The polynomial in y therefore satisfies the Eisenstein criterion and is therefore irreducible – which means that the original equation for $\phi(x)$ is irreducible. ■

Theorem 20.2 (Irreducibility of $x^p - a = 0$) A polynomial $f(x) = x^p - a = 0$ of prime degree p over a field K is either irreducible over K or has a root in K .

Proof 1 based on the Rational Root Theorem Let α be a root of $f(x) = x^p - a$ (assume irreducible or not). Let $K = \mathbb{Q}(\alpha)$. (Assume we're working over $\mathbb{Q}$ or $\mathbb{Z}$ by Gauss's lemma for monic polynomials).

Since p prime, $[K : \mathbb{Q}] = 1$ or p .

If $=1$, $\alpha \in \mathbb{Q}$, so rational root.

Hence if no rational root, $[K : \mathbb{Q}] = p$, so minimum polynomial has degree p , so f (monic) is the minimum polynomial, hence irreducible.

Proof 2 Adjoin to K a primitive p -th root of unity ξ and the a root u of $x^p - a$. The resulting extension $K(\xi, u)$ contains the p roots $u, \xi u, \xi^2 u, \dots, \xi^{p-1} u$ of the polynomial $x^p - a$, hence is the root field of this polynomial, which has the factorization

$$x^p - a = (x - u)(x - \xi u)(x - \xi^2 u) \dots (x - \xi^{p-1} u).$$

Suppose that $x^p - a$ has a proper factor $g(x)$ over K of positive degree $m < p$. This factor $g(x)$ is then a product of m of the linear factors of $x^p - a$ over $K(\xi, u)$, so that the constant term b in $g(x)$ is a product of the roots $\xi^i u$. Therefore $b = \xi^k u^m$, for some integer k , and

$$b^p = (\xi^k u^m)^p = (\xi^p)^k (u^p)^m = (u^p)^m = a^m.$$

From this we can find in K a p -th root of a , for $m < p$ is relatively prime to p and there exists integers s and t with $sm + tp = 1$, so that

$$b^p = a^m = a^{1-tp} = a/a^{tp},$$

and $a = (b^s a^t)^p$. The assumed reducibility of $x^p - a$ over K therefore yields a root $(b^s a^t)$ of $x^p - a$ in K . ■

Theorem 20.3 (Abel's Version of the Irreducibility of $x^p - F$) If the equations

$$\begin{aligned} f_1 x^{p-1} + f_2 x^{p-2} + \dots + f_p &= 0 \\ x^p - F &= 0, \end{aligned} \tag{7}$$

where p is a prime number, are simultaneously satisfied, either $f_1, f_2, f_3, \dots, f_p$ all vanish, or else one of the roots of $x^p - F = 0$ can be expressed rationally in terms of $f_1, f_2, \dots, f_p$ and F .

Proof For, suppose the coefficients $f_1, f_2, \dots, f_p$ don't vanish. The the two equations will have a greatest common divisor (GCD) of the form

$$g(x) = x^k + g_1 x^{k-1} + g_2 x^{k-2} + \dots + g_k = 0,$$

the coefficients of which are rational functions of $F, f_1, f_2, \dots, f_p$. Now If x_1 be any one of the roots common to equations (7), the other roots will be of the form

$$\omega^\alpha x_1, \omega^\beta x_1, \dots, \text{ where } \omega^p = 1.$$

and hence

$$g_k = x_1^k \omega^{\alpha+\beta} \dots = \omega^\pi x_1^k.$$

Since p is a prime number we can find two numbers m and n of which one is negative, such that $mp + nk = 1$. Also

$$g_k^n = \omega^{n\pi} x_1^{\pi k} = \omega^{n\pi} x_1^{1-nk},$$

and therefore

$$\omega^{n\pi} x_1 = g_k^n F^m,$$

therefore $\omega^{n\pi} x_1$, which is a root of Eq. (7) is expressed rationally in terms of $F, f_1, f_2, \dots, f_k$.

21 Roots of Unity Filters

To see exactly why this cancellation happens, we can analyze the sum using the geometric series formula. Let the sum be $S = \sum_{k=0}^{n-1} (\omega^m)^k$, where $\omega = e^{\frac{2\pi i}{n}}$.

Case 1: When m is a multiple of n

If m is a multiple of n (so $m = c \cdot n$ for some integer c):

Each term becomes $(\omega^n)^c$. Since $\omega^n = 1$, each term equals $1^c = 1$. The sum adds 1 exactly n times. Result: $S = 1 + 1 + \dots + 1 = n$.

Case 2: When m is not a multiple of n

When m is not a multiple of n , the common ratio $r = \omega^m$ is not equal to 1. We apply the finite geometric series formula, $S = \frac{1-r^n}{1-r}$: [1]

Substitute the ratio: $S = \frac{1-(\omega^m)^n}{1-\omega^m}$ Swap the exponents: $S = \frac{1-(\omega^n)^m}{1-\omega^m}$ Evaluate the numerator: Since $\omega^n = 1$, the numerator becomes $1 - 1^m = 0$. Ensure non-zero denominator: Because m is not a multiple of n , $\omega^m \neq 1$, so the denominator is not zero. Result: $S = \frac{0}{\text{non-zero}} = 0$.

Visualizing the Cancellation Geometrically, when m is not a multiple of n , the terms ω^{km} represent vectors pointing symmetrically in all directions around the unit circle. Because of this perfect rotational symmetry, their vectors balance each other out completely, pulling the net sum directly to the origin $(0,0)$. [1]

The roots of unity filter (or roots of unity trick) is a mathematical technique used in generating functions and combinatorics to isolate and sum specific polynomial coefficients, such as every n -th term. When applied to a product of polynomials, it allows you to easily extract terms whose exponents are periodic. [1, 2]

How It Works

The core of the filter relies on the following summation property for an n -th root of unity ω (where $\omega^n = 1$): [1]

$$\sum_{k=0}^{n-1} \omega^{mk} = \begin{cases} n, & \text{if } m \text{ is a multiple of } n \\ 0, & \text{otherwise} \end{cases}$$

If you have a generating function $P(x)$, you can isolate the sum of every n -th coefficient (e.g., $a_0, a_n, a_{2n}, \dots$) by averaging $P(x)$ evaluated at all n -th roots of unity: [1, 2] $\sum_{k=0}^{\infty} a_{kn} = \frac{P(1) + P(\omega) + P(\omega^2) + \dots + P(\omega^{n-1})}{n}$

Applying It to Products In combinatorial problems (such as finding the number of subsets whose sum is divisible by a certain integer), your polynomial is often expressed as a product of factors, $P(x) = \prod_{i=1}^m F_i(x)$. [1, 2] Applying

the filter to this product involves substituting each root of unity ω^j into the product individually:

Construct the Generating Function: Represent the event or choice as a polynomial, where powers indicate the "value" and coefficients indicate the number of combinations. **Evaluate at Roots of Unity:** Calculate $P(\omega^j)$ for $j = 0, 1, \dots, n-1$. **Exploit the Product Rule:** Because $P(x)$ is a product, $P(\omega^j)$ becomes the product of the individual evaluated factors: $P(\omega^j) = F_1(\omega^j) \cdot F_2(\omega^j) \cdot \dots \cdot F_m(\omega^j)$. **Average the Results:** Sum these products and divide by n to isolate the terms matching your modulo condition. [1, 2, 3]

Example: Divisibility of Subset Sums Suppose you want to find the number of subsets of the set $\{1, 2, 3\}$ whose sum of elements is divisible by 2.

Generating Function: Each element can be either included (x) or not included (1). The polynomial for the whole set is the product: $P(x) = (1+x)(1+x)(1+x)$ [1] **Apply the Filter (mod 2):** Since we want the sum to be divisible by 2, we use the 2nd roots of unity, which are 1 and -1. [1] **Evaluate and Add:** $P(1) = (1+1)(1+1)(1+1) = 2 \cdot 2 \cdot 2 = 8$ $P(-1) = (1-1)(1+(-1))(1+(-1)) = 0 \cdot 2 \cdot 0 = 0$

Average: $\frac{P(1)+P(-1)}{2} = \frac{8+0}{2} = 4$

There are 4 valid subsets whose sum is divisible by 2: $\{\emptyset\}, \{2\}, \{1, 3\}, \{1, 2, 3\}$.

22 Descartes Rule of Signs

Theorem 22.1 (*Descartes Rule of Signs*): The number of variations of sign in the subsequent terms of a polynomial, s , minus the number of positive roots, p , is a non-negative even integer.

$$s - p = 2r, \quad r \geq 0$$

a) $s - p$ is an even integer.

Let $f(x)$ be a polynomial with a positive leading coefficient .

$$f(x) = a_0x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_{n-1}x + a_n, \quad \text{and} \quad a_0 > 0.$$

Then since the sequence of signs starts and ends with a $+$ we must have an even number of variations $+-$ and $-+$ of sign in the sequence

$$+ \dots + .$$

to get back eventually to find the final $+$ sign so s is even.

Similarly p is even because the function must start at $y = a_0$ and can only cross the x -axis an even number of times to eventually again be in the $y > 0$ region. If there are multiple roots of the form $(x - \alpha)^k = 0$, then these are counted with their respective multiplicity. If k is even, then the function does not cross the x -axis at the multiple root. If k is odd, then the function crosses the x axis and goes into the $y < 0$ region. In either case, p is even. Hence if $a_0 > 0$, we have that $s - p$ is even.

Similar arguments hold for the case when $a_0 < 0$. In that case both s and p are odd numbers and therefore $s - p$ is still even.

b) $s - p$ is non-negative.

Use induction on the degree of the polynomial.

$n = 1$. $x + a_n = 0$. Zero variations if a_n is positive and zero positive roots so $s - p = 0$ is even. If $a_n < 0$ then there is one variation in sign and one positive root, $s = 1$ and $p = 1$ and $s - p = 0$.

Induct on $n = k$.

$$f(x) = a_0x^k + a_1x^{k-1} + a_2x^{k-2} + \dots + a_{k-1}x + a_k.$$

Take the derivative with respect to x ,

$$f'(x) = ka_0x^{k-1} + (k-1)a_1x^{k-2} + (k-2)a_2x^{k-3} + \dots + a_{k-1}.$$

Now, since the k terms that multiply the various x powers are all positive then

$$s = \text{sign variation}[a_0, a_1, a_2, \dots, a_{k-1}, a_k]$$

and

$$s' = \text{sign variation}[a_0, a_1, a_2, \dots, a_{k-1}]$$

Therefore

$$s \geq s'.$$

Since the roots p' are interspersed between the roots of $f(x)$ by Rolle's theorem, the number of positive roots can be $p - 1$ which occur between the roots of $f(x)$ plus possibly one more between $x = 0$ and the first root of α_1 of $f(x)$, so:

$$p' \geq p - 1.$$

Therefore we have

$$s \geq s' \geq p' \geq p - 1$$

or

$$s - p \geq -1.$$

Part a) shows that $s - p$ must be an even number in all cases, so $s - p \geq -1 \implies s - p \geq 0$. Hence

$$s - p = 2r, \quad r \geq 0.$$

23 Determinants and Their Properties

Consider a set of vectors in an n -dimensional Vector Space: $V = \{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n\}$ in $\mathbb{R}^n$.

Determinant The Determinant $D(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n)$ has the following properties

- i) Invariance property: $D(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n)$ is to remain unchanged if some $\mathbf{a}_i$ is replaced by $\mathbf{a}_i + \mathbf{a}_j$ ($i \neq j$);
- ii) Homogeneity property: $D(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n)$ is to go over into $\lambda D(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n)$ if $\mathbf{a}_i$ is replaced by $\lambda \mathbf{a}_i$;
- iii) the Normalization: $D(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n) = 1$.

The determinant is the unique function of vectors $\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n\}$ that satisfy the above properties [3].

Expansion of Determinant Let the S_n be the set of all permutation of the numbers $\{1, 2, \dots, n\}$. In the case of n -by- n matrix A we obtain the complete development of the determinant of A as follows:

$$D(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n) = \det |A| = \sum_{\sigma \in S_n} \epsilon_{\sigma_1 \sigma_2 \dots \sigma_n} a_{1\sigma_1} a_{2\sigma_2} \dots a_{n\sigma_n} = \begin{vmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{i1} & a_{i2} & \dots & a_{in} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{vmatrix} \quad (8)$$

If the rows are permuted by $\rho \in S_n$, we find

$$\epsilon_{\rho_1 \rho_2 \dots \rho_n} \det |A| = \sum_{\sigma \in S_n} \epsilon_{\sigma_1 \sigma_2 \dots \sigma_n} a_{\rho_1 \sigma_1} a_{\rho_2 \sigma_2} \dots a_{\rho_n \sigma_n} \quad (9)$$

Cofactor of an element a_{rs} ,

$$M_{rs} = \text{cofactor}(a_{rs}) = \sum_{\sigma \in S_n, \sigma_r = s} \epsilon_{\sigma_1 \sigma_2 \dots \sigma_n} a_{1\sigma_1} a_{2\sigma_2} \dots a_{r-1\sigma_{r-1}} a_{r+1\sigma_{r+1}} \dots a_{n\sigma_n} \quad (10)$$

Relation between cofactors and minors obtained by taking the determinant obtained from A by striking out row r and column s from A . We denote such minor determinants by A_{rs} :

$$M_{rs} = (-1)^{r+s} A_{rs}$$

For every r , $1 \leq r \leq n$ we can write $\det |A|$ in terms of a cofactor expansion

$$\det |A| = \sum_{s=1}^n a_{rs} M_{rs} = \sum_{s=1}^n (-1)^{r+s} a_{rs} A_{rs} \quad (11)$$

Also, if there is a mismatch between the row indices between the matrix element and the cofactor or minor, i.e.,

$$\sum_{s=1}^n a_{ks} M_{rs} = \sum_{s=1}^n (-1)^{r+s} a_{ks} A_{rs} = 0, \text{ for } k \neq r \quad (12)$$

The case $k \neq r$ corresponds to a determinant with two identical rows and hence vanishes because the rows are linearly dependent. Putting both of the above equations together we find

$$\sum_{s=1}^n a_{ks} M_{rs} = \sum_{s=1}^n (-1)^{r+s} a_{ks} A_{rs} = \delta_{kr} \det |A| \quad (13)$$

Classical Adjoint and Inverse Matrix We define the Classical Adjoint, also known as the Adjugate, of the matrix A as

$$\mathbf{adj}(A) = M^\dagger,$$

where $M^\dagger$ is the transpose of the cofactor matrix, M . Using Eq.(13), the inverse of A is given by

$$A^{-1} = \frac{1}{|A|} \mathbf{adj}(A) = \frac{1}{|A|} M^\dagger.$$

Since $A^{-1}A = I$ we see that the determinant of A^{-1} is $|A|^{-1}$. Therefore the determinant of the Classical Adjoint matrix is

$$|\mathbf{adj}(A)| = |A|^n |A|^{-1} = |A|^{n-1},$$

for an n -by- n matrix A .

Left Inverse and Right Inverse Suppose we have a left inverse X and a right inverse Y of a matrix A . This means that

$$XA = AY = I.$$

Therefore we have

$$\begin{aligned} XAY &= X, \quad \text{and} \\ XAY &= Y. \end{aligned}$$

Therefore

$$X = XAY = Y, \text{ or } X = Y.$$

Therefore left and right inverses of a non-singular matrix must be equal.

The inverse of a Symmetric Matrix is Symmetric A symmetric matrix is defined by $A = A^\dagger$. The inverse of A satisfies

$$A^{-1}A = I$$

Therefore, by taking the transpose of both sides we have

$$A^T(A^{-1})^\dagger = A(A^{-1})^\dagger = I.$$

Therefore

$$A^{-1}A(A^{-1})^\dagger = A^{-1}I = A^{-1},$$

or

$$(A^{-1})^\dagger = A^{-1}.$$

Therefore the inverse of a symmetric matrix is also symmetric.

Special Cases – Inverting a Triangular Matrix

Theorem 23.1 (Upper Triangular Matrices) *Let U be an upper triangular matrix. If the inverse U^{-1} exists, then it is upper triangular.*

Proof Recall the $(n-1) \times (n-1)$ cofactor matrix, M_{rs} that results from omitting row r and column s of $U = (u_{ij})$. When it exists,

$$U^{-1} = \frac{1}{|U|} \mathbf{adj}(U),$$

so it is enough to prove that the $n \times n$ matrix $(|C_{rs}|)$ of cofactor determinants whose transpose is $(|C_{rs}|)^\dagger$ is the adjugate, is lower triangular.

In the case $r < s$, every element below the diagonal of the matrix C_{rs} is also below the diagonal of U , so must be equal to 0.

Hence C_{rs} is upper triangular, with the determinant equal to the product of its diagonal elements.

However, $s - r$ of these diagonal elements are $u_{i+1,j}$ for $i = r, \dots, s - 1$. These elements are from below the diagonal of U , and so are equal to 0.

Hence $r < s$ implies that $|C_{rs}| = 0$, so the $n \times n$ matrix $(|C_{rs}|)$ of cofactor determinants is indeed lower triangular. Hence the adjugate $(|C_{rs}|)^\dagger$ is upper triangular. Hence the inverse of an upper triangular matrix is upper triangular. By similar reasoning we can show that the inverse of a lower triangular matrix is lower triangular. ■

Reciprocal Vector Space in n -Dimensions Consider a set of n vectors $\{\mathbf{a}'_1, \mathbf{a}'_2, \dots, \mathbf{a}'_n\}$, given by

$$\mathbf{a}'_j = \frac{1}{|A|} \{M_{j1}, M_{j2}, \dots, M_{jn}\}.$$

We will refer to this set of vectors as being reciprocal to the set of vectors forming the columns of A . From the definition of inverse of matrix A we immediately have

$$\mathbf{a}_k \cdot \mathbf{a}'_j = \delta_{kj}.$$

Also the determinant of the n -by- n matrix formed from the set of $\mathbf{a}'_j, j = 1, 2, \dots, n$ vectors arranged in columns

$$A' = [\mathbf{a}'_1 \quad \mathbf{a}'_2 \quad \dots \quad \mathbf{a}'_n] \quad (14)$$

had a determinant equal to

$$|A'| = |A|^{-n} |\det(M)| = |A|^{-n} |A|^{n-1} = |A|^{-1}. \quad (15)$$

Therefore the volume of the set of reciprocal vectors $\text{volume}\{\mathbf{a}'_1, \mathbf{a}'_2, \dots, \mathbf{a}'_n\} = \frac{1}{|A|} = \frac{1}{V}$ where $V = |A|$ is the volume of the set of the vectors $\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n\}$. This is the n -dimensional Euclidean space version of Reciprocal Vectors in 3-dimensional Euclidean space.

Schur's complement: Suppose p, q are nonnegative integers, and suppose A, B, C, D are respectively p -by- p , p -by- q , q -by- p , and q -by- q matrices of complex numbers. Let

$$M = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \quad (16)$$

so that M is a $(p + q)$ -by- $(p + q)$ matrix.

If D is invertible, then the Schur complement of the block D of the matrix M is the p -by- p matrix defined by

$$M/D := A - BD^{-1}C. \quad (17)$$

and

$$\det M = \det D \det(A - BD^{-1}C).$$

If A is invertible, the Schur complement of the block A of the matrix M is the q -by- q matrix defined by

$$M/A := D - CA^{-1}B. \quad (18)$$

and

$$\det M = \det A \det(D - CA^{-1}B).$$

In the case that A or D is singular, substituting a generalized inverse for the inverses on M/A and M/D yields the generalized Schur complement.

Laplace's General Expansion Theorem: Divide the set of integers $1, 2, \dots, n$ into two complementary sets $\{\alpha_1, \alpha_2, \dots, \alpha_p\}$ and $\{\beta_1, \beta_2, \dots, \beta_q\}$ where $n = p + q$ and

$$\alpha_1 < \alpha_2 < \dots < \alpha_p \text{ and } \beta_1 < \beta_2 < \dots < \beta_q.$$

These two sets remained fixed throughout the following:

$$\det A = \sum_{\substack{\rho_1, \rho_2, \dots, \rho_p \\ \rho_1 < \rho_2 < \dots < \rho_p}} (-1)^{(\alpha_1 + \alpha_2 + \dots + \alpha_p + \rho_1 + \rho_2 + \dots + \rho_p)} \begin{vmatrix} a_{\alpha_1 \rho_1} & a_{\alpha_1 \rho_2} & \dots & a_{\alpha_1 \rho_p} \\ a_{\alpha_2 \rho_1} & a_{\alpha_2 \rho_2} & \dots & a_{\alpha_2 \rho_p} \\ \dots & \dots & \dots & \dots \\ a_{\alpha_i \rho_1} & a_{\alpha_i \rho_2} & \dots & a_{\alpha_i \rho_p} \\ \dots & \dots & \dots & \dots \\ a_{\alpha_p \rho_1} & a_{\alpha_p \rho_2} & \dots & a_{\alpha_p \rho_p} \end{vmatrix} \begin{vmatrix} a_{\beta_1 \sigma_1} & a_{\beta_1 \sigma_2} & \dots & a_{\beta_1 \sigma_q} \\ a_{\beta_2 \sigma_1} & a_{\beta_2 \sigma_2} & \dots & a_{\beta_2 \sigma_q} \\ \dots & \dots & \dots & \dots \\ a_{\beta_i \sigma_1} & a_{\beta_i \sigma_2} & \dots & a_{\beta_i \sigma_q} \\ \dots & \dots & \dots & \dots \\ a_{\beta_q \sigma_1} & a_{\beta_q \sigma_2} & \dots & a_{\beta_q \sigma_q} \end{vmatrix} \quad (19)$$

The sets $\{\alpha_1, \alpha_2, \dots, \alpha_p\}$ and $\{\beta_1, \beta_2, \dots, \beta_q\}$ were defined to be fixed. The summation is taken over all p -tuples $(\rho_1, \rho_2, \dots, \rho_p)$ from among the numbers $1, 2, \dots, n$ for which $\rho_1 < \rho_2 < \dots < \rho_p$ and where

$$\sigma_1 < \sigma_2 < \dots < \sigma_q$$

are the remaining q integers.

24 Cauchy-Binet Formula

Application 1: **Example Partitioned Matrix and the Cauchy-Binet Formula**

$$M = \left[\begin{array}{cccc|cccc} 0 & 0 & \dots & 0 & b_{11} & b_{12} & \dots & b_{1n} \\ 0 & 0 & \dots & 0 & b_{21} & b_{22} & \dots & b_{2n} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 & b_{m1} & b_{m2} & \dots & b_{mn} \\ \hline c_{11} & c_{12} & \dots & c_{1m} & 1 & 0 & \dots & 0 \\ c_{21} & c_{22} & \dots & c_{2m} & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & 0 \\ c_{n1} & c_{n2} & \dots & c_{nm} & 0 & 0 & \dots & 1 \end{array} \right] \quad (20)$$

In the above M is an $(m + n)$ -by- $(m + n)$ matrix which is partitioned into: A an m -by- m matrix of 0's, B an m -by- n matrix, C an n -by- m matrix and D is the n -by- n Identity matrix. Then using the determinant in the Schur's Complement form we have

$$\det M = \det D \det(A - BD^{-1}C) = \det(-BC) = (-1)^m \det(BC).$$

We can obtain a similar result by row operations. Denote the $(m + i)$ row by $\mathbf{a}_i$. Adding a multiple of row i to row j does not change the value of $\det M$. We successively subtract the $\sum_{i=1}^n b_{ki} \mathbf{a}_i$ from the k -th row of M , for $k = 1, 2, \dots, m$ changing M into a different matrix with the same determinant, viz.,

$$\tilde{M} = \left[\begin{array}{cccc|cccc} -d_{11} & -d_{12} & \dots & -d_{1m} & 0 & 0 & \dots & 0 \\ -d_{21} & -d_{22} & \dots & -d_{2m} & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ -d_{m1} & -d_{m2} & \dots & -d_{mm} & 0 & 0 & \dots & 0 \\ \hline c_{11} & c_{12} & \dots & c_{1m} & 1 & 0 & \dots & 0 \\ c_{21} & c_{22} & \dots & c_{2m} & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & 0 \\ c_{n1} & c_{n2} & \dots & c_{nm} & 0 & 0 & \dots & 1 \end{array} \right], \quad (21)$$

where $d_{kj} = \sum_{i=1}^n b_{ki} c_{ij}$, for $k, j = 1, \dots, m$. Since $\det M = \det \tilde{M}$ we have another demonstration of

$$\det M = (-1)^m \det(BC) \quad (22)$$

where B and C are the matrices corresponding to b_{kj} and c_{ij} respectively.

Another representation of $\det M$ can be obtained by applying Laplace's Generalized Theorem successively. First swap the bottom n rows with the top m rows (by moving each of the bottom n rows up m positions and changing the sign of $\det M$ accordingly

$$\det M = (-1)^{m^2} \det \left[\begin{array}{cccc|cccc} c_{11} & c_{12} & \dots & c_{1m} & 1 & 0 & \dots & 0 \\ c_{21} & c_{22} & \dots & c_{2m} & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & 0 \\ c_{n1} & c_{n2} & \dots & c_{nm} & 0 & 0 & \dots & 1 \\ \hline 0 & 0 & \dots & 0 & b_{11} & b_{12} & \dots & b_{1n} \\ 0 & 0 & \dots & 0 & b_{21} & b_{22} & \dots & b_{2n} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 & b_{m1} & b_{m2} & \dots & b_{mn} \end{array} \right] \quad (23)$$

Using Laplace's Generalized Theorem the first time across the bottom m rows and taking advantage of the m -by- m block of zeros in the lower left hand corner to drop some of the minors in the expansion we have only a sum over the m -by- m minors in the b_{ij} elements starting in columns $m + 1$ through $m + n$ of M . After forming any one of these minors we have to select m columns to remove to form the algebraic complement to use in the Laplace Theorem. The n -by- n complementary minor consists of the n -by- m c_{ij} matrix elements with an additional $n - m$ columns which remain in from the former Identity matrix in the upper right hand side of the working matrix.

We now apply Laplace's Generalized Theorem a second time, now expanding over the first m columns of the complementary n -by- n minor. We must again sum over the choice of m rows from the first n rows. The complementary minor to this second set of minors is obtained from the original n -by- n Identity matrix, I_n , with m columns removed by forming the first set of minors in the b_{ij} and removing the m rows to form the second set of minors. We are left with an $n - m$ -by- $n - m$ matrix from I_n . If the rows that were struck out do not match exactly the columns that were struck out then there will appear in the the reduced matrix rows and columns with all zero entries and the corresponding determinant vanishes. So in the second application of Laplace's Generalized theorem there is only one term surviving in the sum, for any given choice of m numbers $1 \leq \rho_1 \leq \rho_2 \leq \dots \leq \rho_m \leq n$, the only term that

survive is

$$\begin{vmatrix} c_{1\rho_1} & c_{1\rho_2} & \cdots & c_{1\rho_m} \\ c_{2\rho_1} & c_{2\rho_2} & \cdots & c_{2\rho_m} \\ \cdots & \cdots & \cdots & \cdots \\ c_{m\rho_1} & c_{m\rho_2} & \cdots & c_{m\rho_m} \end{vmatrix} \begin{vmatrix} b_{1\rho_1} & b_{1\rho_2} & \cdots & b_{1\rho_m} \\ b_{2\rho_1} & b_{2\rho_2} & \cdots & b_{2\rho_m} \\ \cdots & \cdots & \cdots & \cdots \\ b_{m\rho_1} & b_{m\rho_2} & \cdots & b_{m\rho_m} \end{vmatrix} \quad (24)$$

Putting both results together and using the fact that both methods give an expression for $\det M$ we obtain the Cauchy-Binet Formula for matrices B (n -by- m) and C (m -by- n) with $m \leq n$:

$$\det(BC) = \sum_{\substack{\rho_1, \rho_2, \dots, \rho_m=1 \\ \rho_1 < \rho_2 < \dots < \rho_m}}^n \begin{vmatrix} c_{1\rho_1} & c_{1\rho_2} & \cdots & c_{1\rho_m} \\ c_{2\rho_1} & c_{2\rho_2} & \cdots & c_{2\rho_m} \\ \cdots & \cdots & \cdots & \cdots \\ c_{m\rho_1} & c_{m\rho_2} & \cdots & c_{m\rho_m} \end{vmatrix} \begin{vmatrix} b_{1\rho_1} & b_{1\rho_2} & \cdots & b_{1\rho_m} \\ b_{2\rho_1} & b_{2\rho_2} & \cdots & b_{2\rho_m} \\ \cdots & \cdots & \cdots & \cdots \\ b_{m\rho_1} & b_{m\rho_2} & \cdots & b_{m\rho_m} \end{vmatrix} \quad (25)$$

1: For rectangular matrices, when the number of columns exceeds the number of rows. The overall determinant is formed summing the product of all the sub-determinants be formed from one array (*by taking a number of columns equal to the number of rows*) multiplied by the corresponding determinants from the other array.

2: When the number of rows exceeds the number of columns, the resulting determinant vanishes. It will be observed that the resulting determinant is the same as would arise if an appropriate number of zero-columns were added to each of the given arrays and then forming the determinant of the product.

Application 2: Alternate Derivation of the Cauchy-Binet Formula

Let A be an m -by- n matrix and let B be an n -by- m matrix. Let $1 \leq j_1, j_2, \dots, j_m \leq n$. Let $A_{j_1 j_2 \dots j_m}$ denote the m -by- m matrix consisting of the m columns $j_1, j_2, \dots, j_m$ of A . Let $B_{j_1 j_2 \dots j_m}$ denote the m -by- m matrix consisting of the m rows $j_1, j_2, \dots, j_m$ of B . Let $\{k_1, k_2, \dots, k_m\}$ be an ordered m -tuple of integers and let $\{j_1, j_2, \dots, j_m\}$ be the same set of integers but arranged into non-decreasing order:

$$j_1 \leq j_2 \leq \dots \leq j_m.$$

Then we have the relationships that

$$\begin{aligned} \det(B_{k_1 k_2 \dots k_m}) &= \epsilon_{k_1 k_2 \dots k_m} \det(B_{j_1 j_2 \dots j_m}), \\ \det(B_{j_1 j_2 \dots j_m}) &= \epsilon_{k_1 k_2 \dots k_m} \det(B_{k_1 k_2 \dots k_m}) \text{ no implied sum on } k'_i \text{'s.} \end{aligned}$$

Now from the definition of determinant for the m -by- m matrix AB we have

$$\begin{aligned}
\det(AB) &= \sum_{1 \leq l_1, \dots, l_m \leq m} \epsilon_{l_1 l_2 \dots l_m} \left(\sum_{k=1}^n a_{1k} b_{kl_1} \right) \dots \left(\sum_{k=1}^n a_{mk} b_{kl_m} \right) \\
&= \sum_{1 \leq k_1, \dots, k_m \leq n} (a_{1k_1} \dots a_{mk_m}) \sum_{1 \leq l_1, \dots, l_m \leq m} \epsilon_{l_1 l_2 \dots l_m} b_{k_1 l_1} \dots b_{k_m l_m} \\
&= \sum_{1 \leq k_1, \dots, k_m \leq n} (a_{1k_1} \dots a_{mk_m}) \det(B_{k_1 \dots k_m}) \\
&= \sum_{1 \leq k_1, \dots, k_m \leq n} (a_{1k_1} \dots a_{mk_m} \epsilon_{k_1 \dots k_m}) \det(B_{j_1 \dots j_m}) \\
&= \sum_{1 \leq k_1, \dots, k_m \leq n} (\epsilon_{k_1 \dots k_m} a_{1k_1} \dots a_{mk_m}) \det(B_{j_1 \dots j_m}) \\
&= \sum_{1 \leq j_1 \leq j_2 \leq \dots \leq j_m \leq n} \det(A_{j_1 \dots j_m}) \det(B_{j_1 \dots j_m})
\end{aligned}$$

If any two j 's are equal then $\det(A_{j_1 \dots j_m}) = 0$

In the case of A and $B = A^T$ we immediately have

$$\det(AA^T) = \sum_{1 \leq j_1 \leq j_2 \leq \dots \leq j_m \leq n} [\det(A_{j_1 \dots j_m})]^2$$

25 Resultant of Two Polynomials

Following the demonstration in [7], consider two polynomials

$$\begin{aligned}
f(x) &= \sum_{i=0}^n f_i x^{n-i} = f_0 x^n + f_1 x^{n-1} + f_2 x^{n-2} + \dots + f_{n-1} x + f_n \\
g(x) &= \sum_{j=0}^m g_j x^{m-j} = g_0 x^m + g_1 x^{m-1} + g_2 x^{m-2} + \dots + g_{m-1} x + g_m
\end{aligned}$$

Let us assume that $f(x)$ and $g(x)$ have a common factor $(x - \alpha)$. Then

$$\begin{aligned}
f(x) &= (x - \alpha)f_1(x) \quad \text{and} \\
g(x) &= (x - \alpha)g_1(x).
\end{aligned}$$

Then

$$f(x)g_1(x) = g(x)f_1(x) = \frac{f(x)g(x)}{x - \alpha}.$$

Assuming that

$$\begin{aligned}
f_1(x) &= - \sum_{i=0}^{n-1} z_i x^{n-1-i} = -(z_0 x^{n-1} + z_1 x^{n-2} + z_2 x^{n-3} + \dots + z_{n-2} x + z_{n-1}) \\
g_1(x) &= \sum_{j=0}^{m-1} y_j x^{m-1-j} = y_0 x^{m-1} + y_1 x^{m-2} + y_2 x^{m-3} + \dots + y_{m-2} x + y_{m-1}
\end{aligned}$$

By equating like powers of x , we can write out a linear system corresponding to the equation

$$f(x)g_1(x) - g(x)f_1(x) = 0$$

we find

$$\begin{aligned}
-g(x)f_1(x) = & g_0z_0x^{n+m-1} + g_0z_1x^{n+m-2} + g_0z_2x^{n+m-3} + \dots + g_0z_{n-2}x^{m+1} + g_0z_{n-1}x^m + \\
& g_1z_0x^{n+m-2} + g_1z_1x^{n+m-3} + \dots + g_1z_{n-3}x^{m+1} + g_1z_{n-2}x^m + g_1z_{n-1}x^{m-1} + \\
& g_2z_0x^{n+m-3} + g_2z_1x^{n+m-4} + \dots + g_2z_{n-3}x^m + g_2z_{n-2}x^{m-1} + g_2z_{n-1}x^{m-2} + \\
& \vdots + \\
& \dots + (g_{m-2}z_{n-1} + g_{m-1}z_{n-2} + g_mz_{n-3})x^2 + (g_{m-1}z_{n-1} + g_mz_{n-2})x + g_mz_{n-1}
\end{aligned}$$

$$\begin{aligned} g(x)g_1(x) = & f_0y_0x^{n+m-1} + f_0y_1x^{n+m-2} + f_0y_2x^{n+m-3} + \dots + f_0y_{m-2}x^{n+1} + f_0y_{n-1}x^n + \\ & f_1y_0x^{n+m-2} + f_1y_1x^{n+m-3} + \dots + f_1y_{n-3}x^{n+1} + f_1y_{m-2}x^n + f_1y_{n-1}x^{n-1} + \\ & f_2y_0x^{n+m-3} + f_2y_1x^{n+m-4} + \dots + f_2y_{n-3}x^n + f_2y_{m-2}x^{n-1} + f_2y_{n-1}x^{n-2} + \\ & \vdots + \\ & \dots + (f_{n-2}y_{m-1} + f_{n-1}y_{m-2} + f_ny_{m-3})x^2 + (f_{n-1}y_{m-1} + f_ny_{m-2})x + f_ny_{m-1} \end{aligned}$$

In order to satisfy $f(x)g_1(x) - g(x)f_1(x) = 0$, all the terms multiplying like powers of x each must vanish. This can be arranged as a linear system in the vector $[z_0, z_1, \dots, z_{n-1}, y_0, y_2, \dots, y_{m-1}]$. The coefficient matrix of this linear system is equal to the transpose of the Sylvester Resultant matrix which is defined as

$$R_{m,n}(g, f) = \underbrace{\left[\begin{array}{ccc} g_0 & \cdots & g_m \\ & \ddots & \cdots & \ddots \\ f_0 & \cdots & f_n & \cdots & g_m \\ & \ddots & \cdots & \ddots \\ & & f_0 & \cdots & f_n \end{array} \right]}_{m+n} \left. \begin{array}{l} \left. \vphantom{\begin{matrix} g_0 \\ \vdots \\ f_0 \end{matrix}} \right\} n \text{ lines} \\ \left. \vphantom{\begin{matrix} g_m \\ \vdots \\ f_n \end{matrix}} \right\} m \text{ lines} \end{array} \right\}$$

The corresponding linear system that sets all the coefficients of the same powers of x to 0 in $f(x)g_1(x) - g(x)f_1(x) = 0$ is

$$\begin{bmatrix} g_0 & \cdots & g_m & & \\ & \ddots & & \ddots & \\ & & g_0 & \cdots & g_m \\ f_0 & \cdots & f_n & & \\ & \ddots & & \ddots & \\ & & f_0 & \cdots & f_n \end{bmatrix}^T \begin{bmatrix} z_0 \\ \vdots \\ z_{n-1} \\ y_0 \\ \vdots \\ y_{m-1} \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

The criterion for a non-trivial solution to the above linear system is

$$\det \begin{bmatrix} g_0 & \cdots & g_m & & \\ & \ddots & \cdots & \ddots & \\ f_0 & \cdots & f_n & \cdots & g_m \\ & \ddots & \cdots & \ddots & \\ & & f_0 & \cdots & f_n \end{bmatrix} = 0$$

The rank of the matrix $R_{m,n}(g, f)$ also gives the multiplicity of the common root of $f(x)$ and $g(x)$. The leading term of this determinant is $g_0^n f_n^m$. The terms of in the expansion an $(n+m) \times (n+m)$ matrix M of the form $\sum_{\sigma} \epsilon_{r_1, r_2, r_3, \dots, r_{n+m}} m_{1r_1} m_{2r_2} m_{3r_3} \dots m_{(n+m)r_{n+m}}$ where the sum is over all permutations σ of the numbers $[1, 2, \dots, n+m]$ and $\text{sgn}(\sigma) = \epsilon_{r_1, r_2, r_3, \dots, r_{n+m}}$ is the sign of the permutation. In the case of the Sylvester Resultant the terms take the form

$$\text{sgn}(\sigma) g_0^{\nu_0} g_1^{\nu_1} \dots g_m^{\nu_m} f_0^{\mu_0} f_1^{\mu_1} \dots f_n^{\mu_n},$$

where

$$n = \nu_0 + \nu_1 + \nu_2 + \dots + \nu_m$$

$$m = \mu_0 + \mu_1 + \mu_2 + \dots + \mu_n$$

Factoring out $g_0^n f_0^m$ the generic term in the expansion of the determinant becomes

$$\text{sgn}(\sigma) g_0^n f_0^m \left(\frac{g_1}{g_0}\right)^{\nu_1} \left(\frac{g_2}{g_0}\right)^{\nu_2} \dots \left(\frac{g_m}{g_0}\right)^{\nu_m} \left(\frac{f_1}{f_0}\right)^{\mu_1} \left(\frac{f_2}{f_0}\right)^{\mu_2} \dots \left(\frac{f_n}{f_0}\right)^{\mu_n}$$

Hence the Sylvester Resultant has the form

$$R_{m,n}(g, f) = g_0^n f_0^m \phi(\alpha, \beta)$$

where ϕ is an integral function of the roots α_i and β_i . As the determinant vanishes for any common roots $\alpha_i = \beta_k$, the determinant is divisible by $(\alpha_i - \beta_k)$ and the determinant is divisible by

$$g_0^n f_0^m \prod_{i,k} (\alpha_i - \beta_k)$$

Theorem 25.1 $R_{m,n}(g, f)$ is a homogenous polynomial with integer coefficients g_i, f_j .

(i) $R_{m,n}(g, f)$ is homogenous of degree n in $g_m, \dots, g_0$ and degree m in $f_n, \dots, f_0$.

(ii) If g_i and f_i are regarded as having degree i . then $R_{m,n}(g, f)$ is homogenous of degree mn .

Proof It is obvious that $R_{m,n}(g, f)$ is a homogenous polynomial with integer coefficients in the coefficients g_i, f_j , of total degree $m+n$. Moreover, to replace g_i by tg_i and f_j by uf_j in the Sylvester matrix means that we multiply each of the first m rows by t and each of the last n rows by u , and thus the determinant $R_{m,n}(g, f)$ by $t^m u^n$, which shows part (i). It is here best to treat g_i, f_j, t and u as different indeterminants and do the calculations in $F(g_0, \dots, g_m, f_0, \dots, f_n, t, u)$. Similarly, (ii) follows because to replace each g_i by $t^i g_i$ and each f_j by $t^j f_j$ in $R_{m,n}(g, f)$ yields the same result as multiplying the i th row by t^{m+i} for $i = 1, 2, \dots, m$ and by t^i for $i = m+1, \dots, m+n$, and the j th column by t^{-j} . The overall effect is to multiply the determinant $R_{m,n}(g, f)$ by t^K where $K = mn + \sum_{i=1}^{m+n} i - \sum_{j=1}^{m+n} j = mn$. ■

26 Abel's Method of Rationalizing a Ratio

As an application of Sylvester's Resultant, let us apply the theorem to the case of two polynomials which are related to Abel's Proof of the insolubility of the quintic equation in terms of radicals. In particular let us examine Abel's method of rationalizing a polynomial function of radicals. Let

$$\begin{aligned} f(x) &= x^p - R \\ g(x) &= a_{p-1}x^{p-1} + a_{p-2}x^{p-2} + \dots + a_2x^2 + a_1x + a_0, \end{aligned}$$

where R is not a p th power of an element in the field of coefficients. $f(x)$ and $g(x)$ will have a common root if the Sylvester Resultant formed by these two functions vanishes. The roots of $f(x) = 0$ take the form $z_k = \omega^k R^{\frac{1}{p}}$ and we're assuming that $R^{\frac{1}{p}}$ is an irrationality in the domain containing the coefficients. Also with respect to Abel's rationalization method we need to show that

$$\prod_{k=0}^{p-1} g(z_k) = G(z^p) = G(R) \text{ where } G(R) \text{ is rational in the domain of the coefficients of } g(x). \quad (26)$$

Using $f(x)$ and $g(x)$ we construct the Sylvester Resultant

$$\det(M) = \prod_{k=0}^{p-1} g(z_k) = \det \begin{bmatrix} 1 & 0 & \dots & -R & & \\ 0 & 1 & \ddots & \dots & \ddots & \\ \vdots & & \ddots & \dots & \ddots & \\ 0 & \dots & & 1 & \dots & -R \\ a_{p-1} & & \dots & & a_0 & \\ & & \ddots & & \dots & \ddots \\ & & & a_{p-1} & \dots & a_0 \end{bmatrix}.$$

Since the roots of $f(x) = x^p - R$ are exactly $z_k = \omega^k R^{\frac{1}{p}}$, substituting them into the formula yields

$$\det(M) = \prod_{k=0}^{p-1} g(z_k).$$

Because computing the determinant $\det(M)$ only involves additions, subtractions, and multiplication of the matrix entries, the final polynomial expression outputted by the determinant can only contain powers of combinations of the terms in the matrix M . Therefore the resulting polynomial must be a function of $x^p = R$ and the coefficients and must therefore be a rational expression. The above is a demonstration of Abel's method of Rationalization.

27 Substitutions and Almost Symmetric Functions

Substitution refers to the set of permutations of n elements which form the Symmetric Group.

Theorem 27.1 *All the substitutions of $x_1, x_2, \dots, x_n$ which leave unaltered a rational function $\phi(x_1, x_2, \dots, x_n)$ form a group G .*

Let ϕ_s denote the function obtained by applying to ϕ the substitution s . If a and b are two substitutions which leave ϕ unaltered, then $\phi_a = \phi, \phi_b = \phi$. Then

$$(\phi_a)_b = (\phi)_b = \phi_b = \phi, \text{ or } \phi_{ab} = \phi$$

Hence the product of ab is one of the substitutions which leave ϕ unaltered and the set G has the group property.

The group G is called the *group of the function* ϕ , while ϕ is said to *belong to the group* G

Theorem 27.2 *Every substitution can be expressed as a product of transpositions in various ways.*

Theorem 27.3 *Of the various decompositions of a given substitution s into a product of transpositions, all contain an even number of transpositions (an even substitution), or all contain an odd number of transpositions (an odd substitution).*

Corollary 27.4 *The totality of even substitutions on n letters forms a group, called the Alternating Group on n letters.*

Consider n elements $x = \{x_1, x_2, x_3, \dots, x_n\}$. Consider the rational function

$$\phi(x) = \prod_{i < j} (x_i - x_j) = \begin{matrix} (x_1 - x_2)(x_1 - x_3)(x_1 - x_4) & \dots & (x_1 - x_n) \\ & (x_2 - x_3)(x_2 - x_4) & \dots & (x_2 - x_n) \\ & & (x_3 - x_4) & \dots & (x_3 - x_n) \\ & & & \dots & \\ & & & & (x_{n-1} - x_n) \end{matrix}$$

consisting of the product of all the difference of the elements.

$\phi(x)$ is related to the Vandemonde determinant

$$\phi(x) = V(x) = \begin{vmatrix} 1 & x_1 & x_1^2 & x_1^3 & \dots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & x_2^3 & \dots & x_2^{n-1} \\ 1 & x_3 & x_3^2 & x_3^3 & \dots & x_3^{n-1} \\ \vdots & & & & & \\ 1 & x_n & x_n^2 & x_n^3 & \dots & x_n^{n-1} \end{vmatrix}$$

The square of $V(x)$ is a well-known symmetric function, the discriminant Δ . Therefore $V(x)$ has two values equal numerically with opposite signs, viz. $\sqrt{\Delta}$ and $-\sqrt{\Delta}$. $V(x)$ is an example of an alternating function. Any transposition alters the sign of $V(x)$. For example, the transposition $(x_1 x_2)$ affects only the terms in the first and second lines of the product, and replaces them by

$$\begin{matrix} (x_2 - x_1)(x_2 - x_3)(x_2 - x_4) & \dots & (x_2 - x_n) \\ (x_1 - x_3)(x_1 - x_4) & \dots & (x_1 - x_n). \end{matrix}$$

Hence, if s is the product of an even number of transpositions, it leaves $v(x)$ unaltered; if s is the product of an odd number of transpositions, it changes $V(x)$ into $-V(x)$.

Conjugate values of Multiple-valued Function and Conjugate Groups

Theorem 27.5 (Lagrange) *The order of a subgroup is a divisor of the order of the group.*

Proof Consider a group G of N and a subgroup H composed of the substitutions

$$h_1 = I, h_2, h_3, \dots, h_P. \quad (27)$$

If G contains no further substitutions, then $N = P$ and the theorem is true. Now let G contain a substitution g_2 not in H . Then G also must contain

$$g_2, h_2g_2, h_3g_2, \dots, h_Pg_2. \quad (28)$$

This set are all distinct and all different from the substitutions (27), since $h_\alpha g_2 = h_\beta$ requires that $g_2 = h_\alpha^{-1}h_\beta$ = a substitution of H contrary to the assumption that g_2 is distinct from the elements of H . Hence the substitutions combined now give $2P$ distinct substitutions of G . If there are no other substitutions in G , then $N = 2P$ and the theorem is true. If there are any other distinct substitutions g_3 in G not in (27) or (28), then G contains

$$g_3, h_2g_3, h_3g_3, \dots, h_Pg_3. \quad (29)$$

Moreover, these substitutions are all different from (27) and (28), since $h_\alpha g_3 = h_\beta g_2$ requires that $g_3 = h_\alpha^{-1}h_\beta g_2$ and shall belong to (28), contrary to assumption. We now have $3P$ distinct substitutions of G . Either $N = 3P$ or else G contains a substitution g_4 not in the sets (27), (29), (29). In the latter case G contains the products

$$g_4, h_2g_4, h_3g_4, \dots, h_Pg_4, \quad (30)$$

all of which are distinct and all different from the sets (27), (29), (29), so that we have $4P$ distinct substitutions. Proceeding in this way, we finally reach a last set of P substitutions

$$g_\nu, h_2g_\nu, h_3g_\nu, \dots, h_Pg_\nu, \quad (31)$$

since the order of H is finite. Hence $N = \nu P$.

Definition The number ν is called the *index* of the subgroup H under G and is denoted by $\nu = [G : H]$.

Corollary 27.6 *The order of any group H of substitutions on n letters is a divisor of $n!$. H is a subgroup of the symmetric group G_n on n letters.*

Theorem 27.7 *The period of any substitution contained in a group G of order N is a divisor of N .*

Corollary 27.8 (Applied to the Symmetric Group) *The order of any substitution group on a set of n elements is an exact divisor of $N = n!$, the quotient being the number of distinct value of the corresponding multiple-valued function.*

Let us arrange the N substitutions of G in a rectangular array using the substitutions of a subgroup H in the first row:

$$\begin{array}{cccccc} h_1 = I & h_2 & h_3 & \dots & h_P \\ g_2 & h_2g_2 & h_3g_2 & \dots & h_Pg_2 \\ g_3 & h_2g_3 & h_3g_3 & \dots & h_Pg_3 \\ \vdots & \dots & \dots & \dots & \vdots \\ g_\nu & h_2g_\nu & h_3g_\nu & \dots & h_Pg_\nu \end{array} \quad (32)$$

The $g_1 = I, g_2, g_3, \dots, g_\nu$ multiply the elements of H on the right hand side forming right cosets. Similarly, a rectangular array for the substitutions could be formed by using left cosets.

Theorem 27.9 *If ψ is a rational function of $x_1, x_2, \dots, x_n$ belonging to a subgroup H of index ν , then ψ is ν -valued under the full set of substitutions of G .*

Proof Apply to ψ all the N substitutions of G arranged in a rectangular array as in (32). All the substitutions belonging to the same row give the same answer since

$$\psi_{h_i g_\alpha} = (\psi_{h_i})_{g_\alpha} = (\psi)_{g_\alpha} = \psi_{g_\alpha}.$$

Hence there result at most ν values. But, if

$$\psi_{g_\alpha} = \psi_{g_\beta}$$

then $\psi_{g_\alpha g_\beta^{-1}} = \psi$, so that $g_\alpha g_\beta^{-1} h_i \in H$, leaving ψ unaltered. Hence $g_\alpha = h_i g_\beta$, contrary to the assumption made in forming the rectangular array.

Regarding the orders of a group H and a larger group G containing H as a subgroup; that is to say, the order r of H is an exact divisor of the order G which contains H as a subgroup, the quotient m being the number of distinct values of a multiple-valued function unaltered by the substitutions of G which are obtained by the substitutions of H .

Definition The ν distinct functions $\psi, \psi_{g_2}, \psi_{g_3}, \dots, \psi_{g_\nu}$ are called **conjugate values** of ψ under G .

Theorem 27.10 *The ρ distinct values which a rational function $\psi(x_1, \dots, x_n)$ takes when operated on by all $n!$ substitutions are the roots of an equation of degree ρ whose coefficients are rational functions of the elementary symmetric functions.*

Formation of Functions of a Given Group

We define the Galois Function having $n!$ values for all the substitutions of the symmetric group

$$\psi_1 = \alpha_1 x_1 + \alpha_2 x_2 + \alpha_3 x_3 + \dots + \alpha_n x_n,$$

in which $\alpha_1, \alpha_2, \dots, \alpha_n$ are n distinct arbitrary constants. This function is called the Galois Function. If we use all the P substitutions of H to form the functions $\psi_1, \psi_2, \dots, \psi_P$ by the substitutions of H any symmetric function of these values will be unaltered by the substitutions of H . In particular $\phi_1 = (y - \psi_1)(y - \psi_2) \dots (y - \psi_P)$ will be unaltered by the substitutions of H , and altered to a different value by the substitutions not in H . The function ϕ_1 is not therefore unaltered by the substitutions of any wider group that contains H as a subgroup. Noting the forms of the expression of the sums of powers of the roots in terms of the coefficients, we see that instead of the coefficients of the powers of y in ϕ_1 we may take sums of powers of $\psi_1, \psi_2, \dots, \psi_P$ up to the P^{th} and deduce that one at least of such P sums of powers will provide a function unaltered by the substitutions of H and altered by any other substitution of the symmetric group.

Theorem 27.11 *Every integral symmetric function of the distinct values of any integral multiple-valued function of n elements is a symmetric function of the elements themselves.*

Proof Let $F(\phi_1, \phi_2, \dots, \phi_\rho)$ be any rational integral symmetric function of the ρ -values. Any substitution S whatever affecting the elements applied to these ρ -values either leaves any function unchanged or replaces it by one of the others. No two of the resulting values can be equal, for if $\phi_i S$ were equal to $\phi_j S$ it would follow, by applying the substitution S^{-1} , that $\phi_i = \phi_j$ which is contrary to the assumptions. Consequently the ρ values of ϕ are reproduced by S in some order or other. The symmetric function F therefore remains unchanged by any substitution, and is consequently a symmetric element of the elements themselves.

Corollary 27.12 *The ρ distinct values of an integral multiple-valued function are roots of an equation whose coefficients are integral symmetric functions of the elements themselves.*

Theorem 27.13 (Lagrange Theorem on functions belonging to the same group) *Two rational functions belonging to the same group are rationally expressed each in terms of the other.*

Proof Let ϕ_1 and ψ_1 be two functions belonging to the same group

$$H = \{I, S_2, S_3, \dots, S_r\},$$

of order r and degree n , each of these functions having ρ distinct values, where $r * \rho = N$. Any substitution not contained in H will convert ϕ_1 into another of its values, say ϕ_k and at the same time convert ψ_1 into ψ_k . By operating all possible substitutions, ρ pairs of values $\phi_1, \psi_1 : \phi_2, \psi_2 : \dots : \phi_\rho, \psi_\rho$ are obtained. Now, in the first place, the rational function

$$\sum \phi^i \psi^j = \phi_1^i \psi_1^j + \phi_2^i \psi_2^j + \dots + \phi_k^i \psi_k^j + \dots + \phi_\rho^i \psi_\rho^j \quad (33)$$

is clearly a symmetric function of the elements, for it appears by the same reasoning as before that any substitution acting on this function will merely permute the terms leaving the sum unaltered. Therefore $\sum \phi^i \psi^j$ is a symmetric function of the elements.

If we now take $j = 1$ and assign to i all the values $0, 1, 2, \dots, \rho - 1$ in succession, we obtain the following ρ equations linear in $\psi_1, \psi_2, \dots, \psi_\rho$:

$$\begin{array}{ccccccccc} \psi_1 & + & \psi_2 & + & \dots & + & \psi_\rho & = & T_0 \\ \phi_1 \psi_1 & + & \phi_2 \psi_2 & + & \dots & + & \phi_\rho \psi_\rho & = & T_1 \\ \phi_1^2 \psi_1 & + & \phi_2^2 \psi_2 & + & \dots & + & \phi_\rho^2 \psi_\rho & = & T_2 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & = & \dots \\ \phi_1^{\rho-1} \psi_1 & + & \phi_2^{\rho-1} \psi_2 & + & \dots & + & \phi_\rho^{\rho-1} \psi_\rho & = & T_{\rho-1} \end{array} \quad (34)$$

where $T_0, T_1, \dots, T_{\rho-1}$ are all symmetric functions of $x_1, x_2, \dots, x_n$.

We can write the above as a linear system

$$\begin{bmatrix} 1 & 1 & 1 & 1 & \dots & 1 \\ \phi_1 & \phi_2 & \phi_3 & \phi_4 & \dots & \phi_\rho \\ \phi_1^2 & \phi_2^2 & \phi_3^2 & \phi_4^2 & \dots & \phi_\rho^2 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \phi_1^{\rho-1} & \phi_2^{\rho-1} & \phi_3^{\rho-1} & \phi_4^{\rho-1} & \dots & \phi_\rho^{\rho-1} \end{bmatrix} \begin{bmatrix} \psi_1 \\ \psi_2 \\ \vdots \\ \psi_\rho \end{bmatrix} = \begin{bmatrix} T_0 \\ T_1 \\ \vdots \\ T_{\rho-1} \end{bmatrix} \quad (35)$$

Linear systems such as Eq.(35) can be solved using Cramer's rule. Alternatively, in order to understand better the relationship between ϕ_i and ψ_i let us consider a method based on transforming non-homogenous linear systems into sets of corresponding homogeneous linear systems in each of the unknown parameters. To this effect consider the linear system [4].

$$\begin{array}{ccccccc} x & + & y & + & z & = & T_0 \\ \alpha x & + & \beta y & + & \gamma z & = & T_1 \\ \alpha^2 x & + & \beta^2 y & + & \gamma^2 z & = & T_2 \end{array} \quad (36)$$

By adding a fourth equation to the above we can isolate one of the variables, say y , and determine the functional dependence of y on β and the symmetric functions of the coefficients. Adding $y = y$ to the above system and converting it into a system of homogenous equations by introducing a 4th unknown w we obtain the system

$$\begin{aligned}
0 + y + 0 &= y \\
x + y + z &= T_0 \\
\alpha x + \beta y + \gamma z &= T_1 \\
\alpha^2 x + \beta^2 y + \gamma^2 z &= T_2
\end{aligned} \tag{37}$$

The above system can be converted into a Homogeneous System by moving the final column to the left-hand side of the equal sign.

$$\begin{aligned}
0 + y + 0 - y &= 0 \\
x + y + z - T_0 &= 0 \\
\alpha x + \beta y + \gamma z - T_1 &= 0 \\
\alpha^2 x + \beta^2 y + \gamma^2 z - T_2 &= 0
\end{aligned} \tag{38}$$

Writing the above system using a matrix of coefficients and introducing the 4th undetermined variable we find

$$\begin{bmatrix} 0 & 1 & 0 & -y \\ 1 & 1 & 1 & -T_0 \\ \alpha & \beta & \gamma & -T_1 \\ \alpha^2 & \beta^2 & \gamma^2 & -T_2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \tag{39}$$

In order for this homogenous system to be consistent with Eq.(36) the determinant of the matrix of coefficients must vanish.

$$\begin{vmatrix} 0 & 1 & 0 & -y \\ 1 & 1 & 1 & -T_0 \\ \alpha & \beta & \gamma & -T_1 \\ \alpha^2 & \beta^2 & \gamma^2 & -T_2 \end{vmatrix} \tag{40}$$

Since the determinant vanishes we can factor out the -1 from the last column and write the consistency condition as

$$\begin{vmatrix} 0 & 1 & 0 & y \\ 1 & 1 & 1 & T_0 \\ \alpha & \beta & \gamma & T_1 \\ \alpha^2 & \beta^2 & \gamma^2 & T_2 \end{vmatrix} = 0. \tag{41}$$

Multiplying the above determinant by the non-singular determinant

$$\begin{vmatrix} 1 & 1 & 1 & 0 \\ \alpha & \beta & \gamma & 0 \\ \alpha^2 & \beta^2 & \gamma^2 & 0 \\ 0 & 0 & 0 & 1 \end{vmatrix} \tag{42}$$

we find

$$\begin{vmatrix} 0 & 1 & 0 & y \\ 1 & 1 & 1 & T_0 \\ \alpha & \beta & \gamma & T_1 \\ \alpha^2 & \beta^2 & \gamma^2 & T_2 \end{vmatrix} \begin{vmatrix} 1 & \alpha & \alpha^2 & 0 \\ 1 & \beta & \beta^2 & 0 \\ 1 & \gamma & \gamma^2 & 0 \\ 0 & 0 & 0 & 1 \end{vmatrix} = \begin{vmatrix} 1 & \beta & \beta^2 & y \\ s_0 & s_1 & s_2 & T_0 \\ s_1 & s_2 & s_3 & T_1 \\ s_2 & s_3 & s_4 & T_2 \end{vmatrix} = 0, \tag{43}$$

where

$$s_i = \alpha^i + \beta^i + \gamma^i$$

and the s_i are symmetric functions in α, β , and γ .

Since the s_i 's are sums of powers of α, β, γ and are symmetric functions, they can be expressed rationally in terms of the elementary symmetric functions. By expanding the final determinant in Eq. (43) along the top row and setting the expansion equal to zero to satisfy the consistency requirement we see that we have y expressed as a quadratic function of β along with the sums of powers of the three quantities α, β, γ . Summarizing the results obtained – we expand the above determinant along the top row and solve for y .

$$y = \frac{\begin{vmatrix} s_1 & s_2 & T_0 \\ s_2 & s_3 & T_1 \\ s_3 & s_4 & T_2 \end{vmatrix} - \beta \begin{vmatrix} s_0 & s_2 & T_0 \\ s_1 & s_3 & T_1 \\ s_2 & s_4 & T_2 \end{vmatrix} + \beta^2 \begin{vmatrix} s_0 & s_1 & T_0 \\ s_1 & s_2 & T_1 \\ s_2 & s_3 & T_2 \end{vmatrix}}{\begin{vmatrix} s_0 & s_1 & s_2 \\ s_1 & s_2 & s_3 \\ s_2 & s_3 & s_4 \end{vmatrix}}. \quad (44)$$

The right-hand-side depends quadratically on β and other expressions which are symmetric functions of α, β , and γ and hence demonstrate the claim made in Theorem (27.13).

The denominator of Eq. (44) is a function of the squares of the differences of α, β , and γ :

$$\Delta(\alpha, \beta, \gamma) = \begin{vmatrix} s_0 & s_1 & s_2 \\ s_1 & s_2 & s_3 \\ s_2 & s_3 & s_4 \end{vmatrix} = (\alpha - \beta)^2 (\alpha - \gamma)^2 (\beta - \gamma)^2 \quad (45)$$

Using this result we can write the solution for y as

$$\Delta(\alpha, \beta, \gamma)y = \begin{vmatrix} s_1 & s_2 & T_0 \\ s_2 & s_3 & T_1 \\ s_3 & s_4 & T_2 \end{vmatrix} - \beta \begin{vmatrix} s_0 & s_2 & T_0 \\ s_1 & s_3 & T_1 \\ s_2 & s_4 & T_2 \end{vmatrix} + \beta^2 \begin{vmatrix} s_0 & s_1 & T_0 \\ s_1 & s_2 & T_1 \\ s_2 & s_3 & T_2 \end{vmatrix}. \quad (46)$$

Returning to the original problem Eq. (34), we can similarly express the solution for ψ_j in terms of ϕ_j in the form of a determinant:

$$\begin{vmatrix} 1 & \phi_j & \phi_j^2 & \dots & \phi_j^{\rho-1} & \psi_j \\ s_0 & s_1 & s_2 & \dots & s_{\rho-1} & T_0 \\ s_1 & s_2 & s_3 & \dots & s_\rho & T_1 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ s_{\rho-1} & s_\rho & s_{\rho+2} & \dots & s_{2\rho-2} & T_{\rho-1} \end{vmatrix} = 0, \quad (47)$$

where $T_0, T_1, \dots, T_{\rho-1}$ are all symmetric functions of $x_1, x_2, \dots, x_n$ and $s_k = \phi_1^k + \phi_2^k + \phi_3^k + \dots + \phi_\rho^k$.

The solution of these equations takes the form

$$\Delta(\phi_1, \phi_2, \phi_3, \dots, \phi_\rho)\psi_j = A_0\phi_j^{\rho-1} + A_1\phi_j^{\rho-2} + \dots + A_{\rho-1}, \quad (48)$$

where Δ is the product of the square of the differences of all the ϕ 's (the square of the Vandermonde determinant) and $A_0, A_1, \dots, A_{\rho-1}$ are all symmetric in $x_1, x_2, \dots, x_n$.

Therefore ψ_j can be rationally expressed as a rational function of ϕ_j and the elementary symmetric functions of $x_1, x_2, \dots, x_n$ (the coefficients of the polynomial that has $x_1, x_2, \dots, x_n$ as roots).

Lagrange Interpolation Formula:

Given a set of $k + 1$ nodes $\{x_0, x_1, \dots, x_k\}$, which must all be distinct, $x_j \neq x_m$ for indices $j \neq m$, the *Lagrange basis* for polynomials of degree $\leq k$ for those nodes is the set of polynomials $\{l_0(x), l_1(x), \dots, l_k(x)\}$ each of degree k which take values $l_j(x_m) = 0$ if $m \neq j$ and $l_j(x_j) = 1$. Using the Kronecker delta this can be written $l_k(x_m) = \delta_{jm}$. Each basis polynomial can be explicitly described by the product:

$$\begin{aligned} l_j(x) &= \frac{(x - x_0)}{(x_j - x_0)} \frac{(x - x_1)}{(x_j - x_1)} \cdots \frac{(x - x_{j-1})}{(x_j - x_{j-1})} \frac{(x - x_{j+1})}{(x_j - x_{j+1})} \cdots \frac{(x - x_k)}{(x_j - x_k)} \\ &= \prod_{\substack{0 \leq m \leq k \\ m \neq j}} \frac{x - x_m}{x_j - x_m} \end{aligned}$$

Notice that the numerator $\prod_{m \neq j} (x - x_m)$ has k roots at the nodes $\{x_m\}_{m \neq k}$ while the denominator $\prod_{m \neq j} (x_j - x_m)$ scales the resulting polynomial so that $l_j(x_j) = 1$.

The Lagrange interpolating polynomial for those nodes through the corresponding values $\{y_0, y_1, \dots, y_k\}$ is the linear combination

$$L(x) = \sum_{j=0}^k y_j l_j(x),$$

Each basis polynomial has degree k , so the sum $L(x)$ has degree $\leq k$, and it interpolates the data because

$$L(x_m) = \sum_{j=0}^k y_j l_j(x_m) = \sum_{j=0}^k y_j \delta_{mj} = y_m,$$

Theorem 27.14 (Extension of Lagrange's Theorem to Non Symmetric Functions) *If two rational functions of any set of variables are such that one remains unchanged by all the substitutions of the group to which the other belongs, the first is expressible by means of the second in the form of an integral polynomial whose coefficients are rational symmetric functions of the variables.*

Another statement of Lagrange's Theorem:

If a rational function $\phi(x_1, x_2, \dots, x_n)$ remains unaltered by all the substitutions which leaves another rational function $\psi(x_1, x_2, \dots, x_n)$ unaltered, then ϕ is a rational function of ψ and $c_1, c_2, \dots, c_n$. Note that the group, G to which ϕ belongs contains the group, H to which ψ belongs, i.e. $H \subseteq G$. G is the wider group. Also ϕ is therefore the more symmetric function – a wider set of substitutions keep it unaltered.

Proof

$$\begin{array}{cccc|ccc} I & h_2 & \dots & h_P & \psi = \psi_1 & \phi = \phi_1 \\ g_2 & h_2 g_2 & \dots & h_P g_2 & \psi_{g_2} = \psi_2 & \phi_{g_2} = \phi_2 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ g_\nu & h_2 g_\nu & \dots & h_P g_\nu & \psi_{g_\nu} = \psi_\nu & \phi_{g_\nu} = \phi_\nu \end{array} \quad (49)$$

The $\psi_1, \psi_2, \dots, \psi_\nu$ are all distinct; but $\phi_1, \phi_2, \dots, \phi_\nu$ need not be distinct since ϕ belongs to a group G which may be bigger than H . Under any substitution s on $x_1, x_2, \dots, x_n$, the functions $\psi_1, \psi_2, \dots, \psi_\nu$ are merely permuted. Moreover, if s replaces ψ_i by ψ_j , it replaces ϕ_i by ϕ_j . Set (using the Lagrange Interpolation formula)

$$\begin{aligned} g(t) &= (t - \psi_1)(t - \psi_2) \dots (t - \psi_\nu), \\ \lambda(t) &= g(t) \left(\frac{\phi_1}{t - \psi_1} + \frac{\phi_2}{t - \psi_2} + \dots + \frac{\phi_\nu}{t - \psi_\nu} \right) \end{aligned}$$

so that $\lambda(t)$ is an integral function of degree $\nu - 1$ in t . Since $\lambda(t)$ remains unaltered under every substitution of s , its coefficients are rational symmetric functions of $x_1, x_2, \dots, x_n$ and hence are rational functions of the elementary symmetric functions.

Now we're assuming here that $x_1, x_2, \dots, x_n$ denote indeterminant quantities since $\psi_1, \psi_2, \dots, \psi_\nu$ are algebraically distinct, so that $g'(\psi)$ is not identically zero. In the case where special values are assigned to $x_1, x_2, \dots, x_n$ such that two or more of the functions $\psi_1, \psi_2, \dots, \psi_\nu$ become numerically equal, then $g'(\psi) = 0$, and ϕ is not a rational function of $\psi, c_1, \dots, c_n$. Assuming that we're working with indeterminant $x_1, x_2, \dots, x_n$, and taking $\psi_1 = \psi$ for t , we get

$$\lambda(\psi_1) = [(\psi_1 - \psi_2)(\psi_1 - \psi_3) \dots (\psi_1 - \psi_\nu)] \phi_1$$

and in general

$$\phi = \frac{\lambda(\psi)}{g'(\psi)}.$$

28 Cauchy's 1815 Theorem

Cauchy's 1815 theorem says that if a function of n variables takes more than two distinct values when you permute the variables, then it must take at least p distinct values, where p is the largest prime less than or equal to n . The number of distinct values always divides n factorial. Symmetric functions take one value, and the only way to get exactly two are by use of the alternating functions, like the Vandermonde product.

Theorem 28.1 (Cauchy's theorem:) *For a function of n variables, $f = f(x_1, \dots, x_n)$, the number of distinct values under all permutations of the variables is either 1, 2, or at least p , where p is the largest prime less than or equal to n .*

Proof Let f be a function of n (distinct) variables $f = f(x_1, \dots, x_n)$. Permuting the variables produces up to $n!$ possible inputs, but we assume f to take only R distinct values under these permutations. By Lagrange's theorem $n!$ is divisible R . Assume for contradiction purposes that $2 < R < p$. Consider a p -cycle σ (a cyclic permutation of order p on p of the variables, fixing the rest). The sequence of function values obtained by applying successive powers of σ to a fixed initial ntuple of variables:

$$f(x), f(\sigma(x)), f(\sigma^2(x)), \dots, f(\sigma^{p-1}(x)).$$

There are at most p distinct values in the above list (since σ^p is the identity). Let

$$v_k = f(\sigma^k(x)) \text{ for } k = 0, 1, \dots, p-1,$$

stand for these p values and let us apply the **Pigeonhole Principle** (i.e., if you have more pigeons than pigeonholes, at least one hole has more than one pigeon). These p values are drawn from the set of R assumed distinct possible values that f can ever take, and we assume $R < p$. These values satisfy the cyclic relation: $v_{k+1} = f(\sigma^{k+1}(x))$ is obtained from v_k by applying σ (mod p). By the pigeonhole principle at least two of these p values must be equal.

$$v_r = v_s \text{ for some } 0 \leq s < r \leq p-1.$$

Let $d = r - s$ (so $1 \leq d \leq p - 1$).

Now apply powers of σ repeatedly. Because σ shifts the indices cyclically (i.e., applying σ moves v_k to $v_{(k+1) \pmod p}$), the assumption $v_r = v_s$ means the sequence is periodic with a smaller step. The equality propagates under shifts: Apply σ any number of times to both sides of $v_r = v_s$. This gives: $v_{r+m} = v_{s+m \pmod p}$ for any integer m . In other words, shifting the indices by the same amount preserves the equality. The index differences are multiples of d : The indices where we know equalities hold are exactly the index positions that differ by multiples of $d \pmod p$. That is, for any integer m : $v_k = v_{(k+m-d) \pmod p}$, whenever the equality chain reaches there.

Primality of p makes d a generator: Crucially because p is prime and d is between 1 and $p - 1$, we have $\gcd(d, p) = 1$. Therefore, d has a multiplicative inverse modulo p , and the multiples

$$\{0 \cdot d, 1 \cdot d, 2 \cdot d, \dots, (p - 1) \cdot d\} \pmod p$$

hit every residue class modulo p exactly once and we cycle through all residues modulo p . Starting from any fixed index k , we can reach any other index l by adding a suitable multiple of d modulo p . Therefore $v_k = v_l$ for all k, l . In group theory terms: the subgroup generated by d modulo p is the entire cyclic group $\mathbb{Z}/p\mathbb{Z}$. Therefore entire orbit of $f(\sigma^k x)$ for $k = 0, 1, 2, \dots, p - 1$ collapses to a single value.

Conclusion:

The shifts by multiples of d generate the entire cyclic group of order p . It follows that all the v_k are equal (the orbit collapses to a single value). In particular, $v_0 = v_1$, i.e., $f(x) = f(\sigma(x))$. Since the initial tuple x was arbitrary, this holds for every starting assignment: σ leaves f invariant (does not change its value). By choosing different p -cycles (or by conjugacy/relabeling of variables), this applies to all p -cycles in S_n .

This implies that all p -cycles leave f invariant (fix every value). One can then construct a 3-cycle as a product (or commutator) of two such p -cycles. Thus, 3-cycles also fix values. A 3-cycle is a product of two transpositions. Analysis of transpositions then shows that either f is fully symmetric ($R = 1$) or it takes exactly two values (like the sign/alternating case, e.g., Vandermonde product). But we assumed $R > 2$, a contradiction. You can't get three or more without hitting at least p values. ■

Thus, no such R with $2 < R < p$ is possible. The pigeonhole step forces p -cycles (and then smaller cycles) to preserve values, collapsing the possibilities to $R = 1$ or 2. This is the heart of Cauchy's 1815 argument (sometimes phrased in terms of "forms" or "arrangement" of the variables). Modern presentations often frame it in terms of the action of the symmetric group S_n on the values, with stabilizers and orbits (see the **Orbit-Stabilizer Theorem**) but the core pigeonhole idea on powers of a cycle remains the same.

Why Primality of p Is Essential

If p were composite, say d shares a common factor with p , the multiples of $d \pmod p$ would only generate a proper subgroup, and you might only force equality within cosets rather than full invariance under σ . The primality ensures the generated subgroup is the whole group, forcing full invariance. This is the precise bridge from the pigeonhole collision to invariance under the full p -cycle. Once you have invariance under all p -cycles, the rest of the proof proceeds by constructing smaller cycles (e.g., 3-cycles) as products/commutators and analyzing transpositions.

29 Wantzel's proofs

Theorem 29.1 *Every two valued function of n variables is of the form $S_1 \pm S_2\sqrt{\Delta}$, where S_1 and S_2 are integral symmetric functions, and Δ is the discriminant.*

Proof A two-valued function must belong to the group of order $\frac{1}{2}N$. The only group of this order is the alternating group to which the function $\sqrt{\Delta}$ belongs. The theorem therefore follows as an immediate consequence of the fundamental theorem Thm (27.13).

An Second Proof

Let two values of the function be denoted by ϕ_1 and ϕ_2 and let G_1 and G_2 be the corresponding groups, each of order $\frac{1}{2}N$. In the first place, these two groups must be identical; for if any substitution S of G_1 were to change ϕ_2 to its second value ϕ_1 , then S^{-1} would change ϕ_1 into ϕ_2 ; but this is impossible, since S^{-1} and S each belong to G_1 . Every substitution therefore of G_1 must belong to G_2 and vice-versa.

To show now that these groups coincided with the alternating group, consider the function $\psi = \phi_1 - \phi_2$. Any substitution which belongs to the common group leaves this unaltered; any other will change ϕ_1 to ϕ_2 and ϕ_2 to ϕ_1 , and will therefore change the sign of ψ ; some transposition, $(x_\alpha x_\beta)$ for example, will have this effect, for no group can include all the transpositions without coinciding with the symmetric group. It is easily inferred that $\psi_1 - \psi_2$ is divisible by $x_\alpha - x_\beta$, and hence by the product of all the differences, since ψ^2 is symmetric.

The quotient of ψ by $\sqrt{\Delta}$ is symmetric. To prove this, let $(\sqrt{\Delta})^m$ be the highest power of $\sqrt{\Delta}$ which occurs in ψ . The quotient of ψ by $(\sqrt{\Delta})^m$ is symmetric, since, if not, it would be an alternating function, and would again contain $\sqrt{\Delta}$ as a factor, which is contrary to assumption. It follows immediately that m is an odd number and that the quotient of ψ by $\sqrt{\Delta}$ is symmetric. Writing therefore $\psi_1 - \psi_2 = A\sqrt{\Delta}$ and $\phi_1 + \phi_2 = B$, where A and B are both symmetric, we have

$$\phi_1 = S_1 + S_2\sqrt{\Delta} \text{ and } \phi_2 = S_1 - S_2\sqrt{\Delta},$$

where S_1 and S_2 are both symmetric functions of the variables $x_1, x_2, \dots, x_n$. The groups G_1 and G_2 obviously coincide with the group of $\sqrt{\Delta}$ – the Alternating Group.

Theorem 29.2 *Permutations are elements of the Symmetric Group and act on products of functions homomorphically because the action naturally distributes over multiplication. By definition, a permutation rearranges inputs, which respects function evaluation and standard pointwise multiplication. The following operations define the action of the Symmetric Group:*

$$\begin{aligned} \sigma \cdot (f + g) &= \sigma \cdot f + \sigma \cdot g; \\ \sigma \cdot (fg) &= (\sigma \cdot f)(\sigma \cdot g), \\ \tau \cdot (\sigma \cdot f) &= (\tau\sigma) \cdot f \end{aligned}$$

for $\sigma, \tau \in S_n$ and $f, g \in F[x_1, \dots, x_n]$. S_n is the set of permutations on n elements $[x_1, x_2, \dots, x_n]$.

Proof 1. Define the Permutation Action A permutation σ acts on a function f by precomposition:

$$(\sigma \cdot f)(x) = f(\sigma^{-1}(x))$$

2. Evaluate the Action on a Product of Functions We apply the permutation to the pointwise product of two functions, $(fg)(x) = f(x)g(x)$.

$$[\sigma \cdot (fg)](x) = (fg)(\sigma^{-1}(x)).$$

3. Unpack the Function Evaluation Evaluate the product at the permuted input $\sigma^{-1}(x)$:

$$[\sigma \cdot (fg)](x) = f(\sigma^{-1}(x)) \cdot g(\sigma^{-1}(x)).$$

4. Regroup using the Action Definition By the definition of the action in Step 1, we rewrite the scalar outputs:

$$[\sigma \cdot (fg)](x) = (\sigma \cdot f)(x) \cdot (\sigma \cdot g)(x).$$

5. Conclude for the Pointwise Product Since this holds for every element $x \in X$, the functions are equivalent:

$$\sigma \cdot (fg) = (\sigma \cdot f)(\sigma \cdot g).$$

Because this action distributes over the product of functions, it satisfies the required structural property (a homomorphism) for group actions on algebraic structures.

Theorem 29.3 *The alternating functions are the only unsymmetrical functions of n variables of which a power can be symmetric.*

Proof It is sufficient to prove this theorem for prime powers; for if there exists a function $F(x_1, x_2, \dots, x_n)$ such that F^{pq} is symmetric and p is a prime, then there is also a function $\phi = F^q$ such that ϕ^p is symmetric.

Therefore let

$$\phi^p = S, \text{ a symmetric function.}$$

Since the group of ϕ which is unsymmetric, cannot contain all the transpositions, let $\sigma = (x_\alpha x_\beta)$ be a transposition which converts ϕ into ϕ_j ; we have

$$\phi_j^p = \phi^p = S.$$

and therefore $\phi_j = \omega \phi$, where ω is a p th root of unity. Hence

$$\sigma \phi = \phi_j = \omega \phi,$$

and operating again with σ

$$\sigma^2 \phi = \omega \sigma \phi = \omega^2 \phi,$$

but $\sigma^2 = I$ and therefore $\omega^2 = 1$, and consequently $p = 2$.

Since ϕ^s is symmetric, ϕ is an alternating function. ■

Theorem 29.4 *For any number, n , of independent elements there is no multiple-valued function of which a power is two-valued when $n > 4$; and when $n = 3$, or $n = 4$, if there is any such power it is third power.*

Concentrating on prime numbers, and supposing that ϕ is a multiple-valued function whose p power is two-valued we have

$$\phi^p = S_1 + S_2 \sqrt{\Delta}. \tag{50}$$

The group of ϕ cannot contain all the circular substitutions of the third order, for if it did, then this group would coincide with the alternating group, and ϕ would be two-valued. Let $\sigma = (x_\alpha, x_\beta, x_\gamma)$ be such a substitutions not contained in the group of ϕ and suppose $\sigma \phi = \phi_j$. From Eq. (50), since $S_1 + S_2 \sqrt{\Delta}$ is unaltered by σ , we have

$$\phi^p = \phi_j^p,$$

hence $\phi_j = \omega\phi$, where ω is a p^{th} root of unity. Operating again twice in succession with σ we obtain

$$\begin{aligned}\sigma\phi &= \omega\phi, \\ \sigma^2\phi &= \omega\sigma\phi = \omega^2\phi, \\ \sigma^3\phi &= \omega^2\sigma\phi = \omega^3\phi,\end{aligned}$$

whence, since $\sigma^3 = 1$ (three cycle), we have $\omega^3 = 1$, and therefore $p = 3$.

Again, when the number of elements is greater than 4, there are circular substitutions of the fifth order, and those cannot all be contained in the group of ϕ (or else it will be two-valued and belong to the Alternating Group). Let τ be one of those not contained in this group and $\tau\phi = \phi_j$. We have, as before, from Eq. (50), by applying this substitution (which does not affect the right-hand side),

$$\phi^p = \phi_j^p = S_1 + S_2\sqrt{\Delta}.$$

Hence proceeding as before, we have $\tau\phi = \omega\phi$ and operating again on this and the succeeding equations with τ we easily find $\tau^5\phi = \omega^5\phi$; whence $\omega^5 = 1$, since $\tau^5 = 1$, and it is proved that $p = 5$. Now this results being incompatible with the previous value of p obtained, we infer that when the number of elements is greater than 4, it is impossible to find any multiple-valued function ϕ , a prime power of which will be two-valued.

There are actually, when n is not greater than 4, multiple-valued functions, a third power of which is two-valued.

30 The Commutator Subgroup (The Abelian Quotient)

Definition Let G be a group. An element $g \in G$ is called a commutator if $g = aba^{-1}b^{-1}$ for elements $a, b \in G$. The smallest subgroup that contains all commutators of G is called the commutator subgroup or derived subgroup of G , and is denoted by G' .

A highly practical relationship emerges with the commutator subgroup G' . The quotient G/G' forms the largest Abelian (commutative) group that can be created from G . If you “mod out” by the commutator subgroup, all order-dependent operations are eliminated.

Proposition 30.1 *Let G be a group with commutator subgroup G' .*

- (a) *The subgroup G' is normal in G , and the factor group G/G' is Abelian.*
- (b) *If N is any normal subgroup of G , then the factor group G/N is Abelian, if and only if $G' \subseteq N$.*

Proof (a) Let $x \in G'$ and let $g \in G$. Then $gxg^{-1}x^{-1} \in G'$, and since $x \in G'$, we must have $gxg^{-1} = gxg^{-1}x^{-1}x \in G'$.

The factor group G/G' must be Abelian since for any cosets aG', bG' , because G' is normal, we have $xG' = G'x$ for all $x \in G$.

$$aG'bG'a^{-1}G'b^{-1}G' = abG'G'a^{-1}b^{-1}G'G' = aba^{-1}b^{-1}G' = G'.$$

Proof (b) Let N be a normal subgroup of G . If $N \supseteq G'$, then G/N is a homomorphic image of G/G' and must be Abelian. Conversely, suppose that G/N is Abelian. Then $aNbN = bNaN$ for all $a, b \in G$, or simply $aba^{-1}b^{-1}N = N$, showing that every commutator of G belongs to N . This implies $G' \subseteq N$. ■

Theorem 30.2 *The Symmetric Group G on n letters is not solvable unless $n \leq 4$.*

Proof Let $G = S_0 \supseteq S_1 \supseteq S_2 \supseteq \dots \supseteq S_s$ be any chain of subgroups, each normal in the preceeding with cyclic quotient-group S_{k-1}/S_k ; we shall prove by induction on s that S_s must contain every 3-cycle (ijk) . This will imply that $S_s > I$, and so, that G cannot be solvable. Since $S_0 = G$ contains every 3-cycle, it is sufficient to show that if S_{k-1} contains every 3-cycle, then so does S_k . First note that if the permutations ϕ and ψ are both in S_{k-1} , then their so-called commutator $\gamma = \phi^{-1}\psi^{-1}\phi\psi$ is in S_s . To see this, consider their images ϕ' and ψ' in S_{k-1}/S_k . The latter, being cyclic, is commutative; hence

$$\gamma' = \phi'^{-1}\psi'^{-1}\phi'\psi' = \phi'^{-1}\psi'^{-1}\psi'\phi' = I \text{ in } S_{k-1}/S_k,$$

which implies that $\gamma \in S_k$. But in the special case when $\phi = (ijk)$ and $\psi = (jkm)$, where i, j, k are given and l, m are any two other letters (such letters exist unless $n \leq 4$), we have

$$\gamma = (jli)(mkj)(ilk)(jkm) = (ijk) \in S_k, \text{ for all } i, j, k.$$

This proves that S_k contains every 3-cycle, as desired.

The quotient chain doesn't terminate on the Identity, therefore the group G is not solvable unless $n \leq 4$. The 3-cycles generate the Alternating group which is simple and has no normal subgroups.

31 Hamilton

Sylvester's dialytic elimination method (and the resulting dialytic resolvent) can be used as an alternate approach to clear denominators and eliminate variables, yielding the same final polynomial equations as Abel's method of multiplying conjugate algebraic expressions to clear denominators of radical expressions and roots of unity. While Niels Henrik Abel's famous 1826 paper on the impossibility of solving the general quintic equation relied on multiplying by algebraic conjugates to clear denominators systematically, James Joseph Sylvester's later dialytic method (developed in the 1840s) reformulates the elimination problem into linear algebra. Instead of constructing field extensions and tracking field conjugates, Sylvester's approach creates a system of linear equations by treating powers of the variable as independent unknowns, solving it via a single determinant

1. Formulated the Algebraic Elimination Suppose Abel's method encounters an expression involving an algebraic denominator, such as $y = \frac{A(x)}{B(x)}$, where x is a root of a known polynomial $P(x) = 0$. Abel would clear the denominator by finding the algebraic conjugates of $B(x)$ relative to the roots of $P(x)$ and multiplying them.

To use Sylvester's resolvent instead, we rewrite the relationship as a system of two polynomial equations sharing a common root x : $P(x) = 0$ (The base polynomial equation) $R(x, y) = y \cdot B(x) - A(x) = 0$ (The relation defining y) Our goal is to eliminate x entirely to find a polynomial equation purely in terms of y .

2. Construct the Dialytic Matrix Sylvester's method eliminates x by generating a set of linearly independent equations. If $P(x)$ has degree n and $R(x, y)$ has degree m with respect to x , we multiply $P(x)$ by successive powers of x up to x^{m-1} , and multiply $R(x, y)$ by powers of x up to x^{n-1} . This creates a system of $n + m$ equations: $x^k \cdot P(x) = 0$ for $k = 0, 1, \dots, m - 1$ $x^j \cdot R(x, y) = 0$ for $j = 0, 1, \dots, n - 1$ We treat each power of x (from x^0 up to x^{n+m-1}) as a distinct, independent linear variable.

3. Compute the Sylvester Resolvent For a non-trivial solution to exist for this linear system, the determinant of the coefficients must equal zero. This determinant is the Sylvester Resultant (or Sylvester Resolvent), denoted as $\text{Res}(P, R, x)$. The matrix takes the form:

$$\text{Res}(P, R, x) = \det \begin{pmatrix} p_n & p_{n-1} & \dots & p_0 & 0 & \dots & 0 \\ 0 & p_n & p_{n-1} & \dots & p_0 & \dots & 0 \\ \vdots & & \ddots & & & \ddots & \vdots \\ r_m(y) & r_{m-1}(y) & \dots & r_0(y) & 0 & \dots & 0 \\ 0 & r_m(y) & r_{m-1}(y) & \dots & r_0(y) & \dots & 0 \\ \vdots & & \ddots & & & \ddots & \vdots \end{pmatrix} = 0$$

Evaluating this determinant yields a pure polynomial equation $Q(y) = 0$, completely clearing the denominator without ever needing to calculate explicit field conjugates or Galois orbits.

Conclusion Sylvester's dialytic resolvent acts as a perfect mathematical alternative to Abel's conjugate multiplication for clearing denominators. By framing the algebraic relation as a simultaneous polynomial system, Sylvester bypasses the need to explicitly compute field conjugates, instead achieving the exact same polynomial reduction through the evaluation of a single matrix determinant.

To see exactly how Sylvester's method serves as a direct alternative, we can look at a representative reduction from Niels Henrik Abel's work. In his 1826 impossibility proof, Abel builds algebraic functions of increasing "orders" by adjoining radicals. He routinely deals with expressions where a new variable y is a rational function of a root x , where x satisfies a lower-degree base polynomial equation. Let's isolate a core algebraic step typical of Abel's paper: reducing an expression where y is defined by a denominator containing a quadratic root x , where x satisfies the base cubic equation: $P(x) = x^3 + ax + b = 0$ Suppose the relationship defining the new variable y is given by the rational function: $y = \frac{1}{x^2+x}$

1. Formulate the Two Polynomials To clear the denominator using Sylvester's method instead of multiplying by algebraic conjugates, we first transform the rational function into a second polynomial equation, $R(x, y) = 0$, grouped by powers of x : $y(x^2 + x) - 1 = 0 \implies R(x, y) = yx^2 + yx - 1 = 0$ Now we have a system of two polynomial equations sharing the common root x : $P(x) = 1x^3 + 0x^2 + ax + b = 0$ (Degree $n = 3$) $R(x, y) = yx^2 + yx - 1 = 0$ (Degree $m = 2$ with respect to x)

2. Set Up the Rows of the Sylvester Matrix The Sylvester matrix will have a total dimension of $(n+m) \times (n+m)$, which means a 5×5 matrix. We shift the coefficients of each polynomial across the columns: Rows 1 & 2 (from $P(x)$): Multiply $P(x)$ by x^1 and x^0 . Rows 3, 4, & 5 (from $R(x, y)$): Multiply $R(x, y)$ by x^2 , x^1 , and x^0 . The columns correspond to the descending independent powers of x : $(x^4 \ x^3 \ x^2 \ x^1 \ x^0)$.

3. The Explicit Sylvester Matrix S
$$S = \begin{pmatrix} 1 & 0 & a & b & 0 \\ 0 & 1 & 0 & a & b \\ y & y & -1 & 0 & 0 \\ 0 & y & y & -1 & 0 \\ 0 & 0 & y & y & -1 \end{pmatrix}$$

4. Clearing the Denominator To completely eliminate x and clear the denominator, you take the determinant of this matrix and set it to zero ($\det(S) = 0$). Evaluating this 5×5 determinant yields a pure polynomial equation in terms of y , a , and b . This polynomial gives you the exact same result Abel would get by finding the other conjugate roots of the cubic equation and computing: $(y(x_1^2 + x_1) - 1)(y(x_2^2 + x_2) - 1)(y(x_3^2 + x_3) - 1) = 0$

After clearing the denominators to ensure all algebraic expressions are expressed as polynomials in the radicals, Niels Henrik Abel's 1826 proof proceeds through a structured combination of field extensions, permutation group properties, and a final proof by contradiction. Abel did not use modern Galois group vocabulary, but his strategy directly mapped out what we now call the Galois theory of fields.

1. The Structure of an Algebraic Solution Abel begins by formalizing what a general "solution by radicals" must look like. If a general quintic equation is solvable, its roots must be expressible by a finite chain of algebraic operations. He structures the general form of a root as a polynomial in a radical: $x = q_0 + q_1 R^{1/m} + q_2 R^{2/m} + \dots + q_{m-1} R^{(m-1)/m}$. Here, m is a prime number, and the quantities q_i and R are either rational functions of the original coefficients, or they belong to a lower "order" field extension previously built up by adjoining other radicals.

2. Formulating Abel's Theorem on Radical Functions The core pivot of the proof relies on a fundamental theorem Abel establishes about the algebraic dependencies of these radicals. He proves that if a root of a general quintic can be written in the form above, then the radical itself, $R^{1/m}$, must be expressible as a rational function of the roots of the quintic and the coefficients. By applying permutations to the roots of the general quintic, Abel studies how the value of $R^{1/m}$ behaves. Because the coefficients of the general quintic are independent, independent variables, permuting the roots causes $R^{1/m}$ to cycle through its m distinct algebraic values (the m -th roots of unity multiplied by the radical).

3. Restricting the Prime Index m Abel analyzes what values the prime degree m of the outermost radical can take. A function of 5 independent variables can only take a specific number of distinct values when its variables are permuted. By invoking Cauchy's earlier work on the number of values a symmetric or partially symmetric function can take, Abel shows that the number of values a rational function of 5 variables can take under permutation must be a divisor of $5! = 120$. He narrows down the possible values of the prime index m . The only primes that can mathematically divide these algebraic combinations for 5 variables are $m = 2$ and $m = 5$.

4. Bounding the Value Multiplicity (The Cauchy Constraint) Abel then looks at a critical theorem proved by Augustin-Louis Cauchy: a non-symmetric function of 5 variables cannot take fewer than 5 distinct values unless it takes exactly 2 values (which corresponds to the alternating group A_5 , generated by the square root of the discriminant). This creates a rigid algebraic trap: If $m = 5$, the radical expression must take exactly 5 distinct values under root permutations. Abel shows that the only way a function of 5 variables can take exactly 5 values under permutation is if it behaves like a specific linear combination of the roots (similar to a Lagrange resolvent).

5. The Final Contradiction Abel explicitly sets up the equations for these 5-valued functions. He assumes a root looks like: $x = q_0 + q_1 R^{1/5} + q_2 R^{2/5} + q_3 R^{3/5} + q_4 R^{4/5}$. By examining the algebraic behavior of the individual terms under 3-cycle and 5-cycle permutations, he derives a system of equations for the components q_i . He proves that for the general quintic, it is impossible for these equations to hold simultaneously. The permutations of the 5 independent roots require the individual components to vanish or collide in a way that contradicts the assumption that x is a root of a general equation with independent coefficients. Because the assumption that a general root can be expressed as a chain of radicals leads directly to an algebraic impossibility, Abel concludes that the general quintic equation cannot be solved by radicals.

Summary of Abel's Path After Clearing Denominators

Defines Root Anatomy: Expresses the hypothetical root as a polynomial of a prime-degree radical $R^{1/m}$.

Inverts the Radical: Proves that $R^{1/m}$ must be a rational function of the quintic's roots.

Applies Permutations: Analyzes how $R^{1/m}$ changes values when the roots are permuted.

Invokes Cauchy's Bound: Restricts the possible values of m to 2 or 5 based on the combinatorics of 5 variables.

Forces Contradiction: Demonstrates that the algebraic conditions required to form a 5-valued function cannot be met by the coefficients of a general quintic.

To understand how Augustin-Louis Cauchy's work on permutations constrained Niels Henrik Abel, we have to look at the combinatorics of switching variables.

When Abel proved that the outermost radical of a hypothetical solution, $v = R^{1/m}$, must be a rational function of the five roots $(x_1, x_2, x_3, x_4, x_5)$, he created a bridge between algebra and combinatorics. Because the coefficients of a general quintic are completely independent, permuting the roots creates a set of distinct values for that function. Cauchy's theorem placed a strict mathematical ceiling on exactly how many values such a function could take, narrowing Abel's options down to a dead end.

1. The Core of Cauchy's Theorem (1815) In 1815, Cauchy published a groundbreaking theorem concerning symmetric and non-symmetric functions. He proved a rigid restriction for functions of n independent variables

Theorem 31.1 (Cauchy's Number of Values Theorem:) *If a non-symmetric function of n independent variables changes its value when the variables are permuted, the number of distinct values it can take (its index) must be at least the largest prime factor of n . Furthermore, for $n = 5$, a function cannot have 3 or 4 distinct values. It can only have 1 (fully symmetric), 2, 5, or more (up to $5! = 120$) distinct values.*

32 Some Results in the Theory of Numbers

Definition If a and b are two integers such that $a - b$ is divisible by n , we say that a is congruent to b modulo n , and write $a \equiv b \pmod{n}$, or merely $a \equiv b(n)$.

Theorem 32.1 *Lagrange's theorem.*

The order of any subgroup of a finite group is a factor of the order of the group.

The order of a finite group G is a multiple of the order of every one of its subgroups.

Suppose H is a subgroup of G , where the order of G is n and that of H is m .

The whole group G may be decomposed into cosets gH relative to H . The number of elements in each coset must be m , since the coset gH is formed by taking all the elements of H and pre-multiplying them by g , and all the elements so formed are different by the Cancellation Law. But the cosets are completely disjoint. Hence if there are r cosets we must have that $n = mr$, and so m is a factor of n .

Each element a of G generates a cyclic subgroup, whose order is simple the order of a . Therefore we have:

Corollary 32.2 *(Corollary 1): Every element of a finite group G has as order a divisor of the order of G .*

Corollary 32.3 *(Corollary 2): Every group G of prime order p is cyclic.*

Proof For the cyclic subgroup A generated by any element $a \neq e$ in such a group has an order $n > 1$ dividing p . But this implies $n = p$, and so $G = A$ is cyclic. ■

Corollary 32.4 *(Corollary 3): The only abstract groups of order four are the cyclic group of that order and the four-group (Viergruppe).*

Proof If a group G has order 4, contains an element of order 4, it is cyclic. Otherwise, by Corollary 1, all elements of G except e must have order 2. Call them a, b, c . By the cancellation law, ab cannot be either $ac = a, eb = b$, of $aa = e$; hence $ab = e$. By symmetry, $ac = ca = b, bc = cb = a, ba = c$. But these, together with $a^2 = b^2 = c^2 = e$, and $ex = xe = x$ for all x , give the multiplication table of the four-group. ■

Theorem 32.5 (*Fermat's Little Theorem*): If p is prime, $a^p \equiv a(p)$, and if a is not a multiple of p , then $a^{p-1} \equiv 1(p)$

Proof (1) The numbers $a, 2a, 3a, \dots, (p-1)a$ are all different modulo p if a is not a multiple of p , and so they must be $1, 2, \dots, (p-1)$ modulo p in some order. Thus

$$a \cdot 2a \cdot 3a \dots (p-1)a \equiv 1 \cdot 2 \cdot 3 \dots (p-1).$$

Hence $a^{(p-1)} \equiv 1(p)$. Thus $a^p \equiv a(p)$ if a is not a multiple of p , and this latter result is true also if a is a multiple of p , since then both sides are 0. Hence it is true for all a . ■

Proof (2) The multiplication group mod p (excluding zero) has $(p-1)$ elements. The order of any element a of this group is then a divisor of $(p-1)$, by Corollary 1, so that $a^{(p-1)} \equiv 1(p)$ whenever $a \not\equiv 0(p)$. If we multiply by a or both sides, we obtain the desired congruence, except for the case $a \equiv 0(p)$, for which the conclusion is trivially true. ■

Proof (3) For a fixed prime p , let $P(n)$ be the proposition that $n^p \equiv n(p)$. Then $P(0)$ and $P(1)$ are obvious. In the binomial expansion for $(n+1)^p$, every coefficient except the first and the last is divisible by p , hence $(n+1)^p \equiv n^p + 1(p)$, whence $P(n)$ implies $(n+1)^p \equiv n+1(p)$, which is the proposition $P(n+1)$. ■

Theorem 32.6 (*Wilson's Theorem*): Let $p > 2$ be a prime number. The field $\mathbb{Z}_p$ is the splitting field of $x^p - x$.

Proof Since $\mathbb{Z}_p$ is the splitting field of $x^p - x$, we have

$$x^p - x = x(x-1)(x-2) \dots (x-(p-1)).$$

Thus

$$x^{(p-1)} - 1 = (x-1)(x-2) \dots (x-(p-1)),$$

and substituting $x = 0$ gives

$$-1 = (-1)(-2) \dots (-(p-1)).$$

There are an even number of factors on the right hand side, so the minus signs cancel out, leaving

$$(p-1)! \equiv -1 \pmod{p},$$

which is Wilson's Theorem.

33 Chinese Remainder Theorem

Consider a sequence of congruence equations:

$$\begin{aligned} x &\equiv a_1 \pmod{n_1} \\ &\vdots \\ x &\equiv a_k \pmod{n_k}, \end{aligned}$$

where the n_i are coprime in pairs. Let $N = n_1 n_2 \dots n_k$ and define $N_i = N/n_i$ be the product of all moduli but one. As the n_i are pairwise coprime, N_i and n_i are coprime. Therefore, by the Bezout identity, there exist integers M_i and m_i such that

$$M_i N_i + m_i n_i = 1.$$

Therefore a solution to the system of congruences is

$$x = \sum_{i=1}^k a_i M_i N_i.$$

Definition We restate the result as: if the n_i are pairwise coprime, the map

$$x \pmod{N} \rightarrow (x \pmod{n_1}, \dots, x \pmod{n_k})$$

defines a ring homomorphism

$$\mathbb{Z}/N\mathbb{Z} \cong \mathbb{Z}/n_1\mathbb{Z} \times \dots \times \mathbb{Z}/n_k\mathbb{Z}$$

between the ring of integers modulo N and the direct product of rings of integers modulo the n_i . This means that for doing a sequence of arithmetic operations in $\mathbb{Z}/N\mathbb{Z}$, one may do the same computation independently in each $\mathbb{Z}/n_i\mathbb{Z}$ and then get the result by applying the isomorphism (from the right to the left).

$$R/(\cap A_i) \cong R/(A_1) \times \dots \times R/(A_k)$$

We have a surjective map and know the kernel is the intersection of all the A_i 's. If you take a map and quotient by the kernel, then that is isomorphic to the image – which is the whole set (everything) because the map is surjective. Therefore the LHS is isomorphic to the RHS.

34 Finite Fields

Theorem 34.1 (*Characteristic of a Field*): In a field F , all non-zero elements have the same additive order. This is called the characteristic of F and is said to be zero if all non-zero elements have infinite order.

Proof Suppose that any given non-zero element y has finite order m . Then if x is any other non-zero element, $(mx)y = m(xy) = x(my) = 0$. But $y \neq 0$ and so $mx = 0$. This is a step that could not be performed in the case of rings, which may have zero divisors. Also, suppose $rx = 0$ for some $r < m$. Then $x(ry) = (rx)y = 0$ as before and since $x \neq 0$, $ry = 0$, which is impossible since $r < m$ and m is the order of y . Hence x also has order m .

Theorem 34.2 If F has a finite characteristic, this must be a prime.

Proof Suppose F has as characteristic a composite number hk . If a is any non-zero element of F , a^2 is non-zero and so has order hk , so that $hka^2 = 0$, i.e., $(ha)(ka) = 0$ and so either ha or ka is 0, which is impossible, since both h and k are less than hk and a must have order hk .

Theorem 34.3 If F has characteristic p , it contains a subfield isomorphic to the field of residues modulo p .

Proof Consider the unity 1 and denote 1+1 by 2, 1+1+1 by 3, etc. Then 1 has order p , and so $p \cdot 1 = 0$ (i.e., the sum of the p 1's is the zero element). Now consider the subset of elements $0, 1, 2, \dots, (p-1)$. The sum and difference of any two of these is in the set as are 0 and 1. Furthermore the product of any two is in the set, since the product is the ordinary product of residues modulo p , by the Distributive Law. For example,

$$2 \cdot 3 = (1+1)(1+1+1) = 1+1+1+1+1+1 = 6.$$

Since sum and product behave as they do for residues modulo p the subset is isomorphic to the set of residues modulo p under residue sum and product and so, since p is prime, inverses exist in the subset and the subset is isomorphic to the field of residues modulo p .

Theorem 34.4 *Let F be a field of characteristic 0, it contains a subfield isomorphic to the field of rational numbers.*

Proof The unity 1 has infinite order and so the elements $0, 1, 2, \dots$ are all distinct. Consider the subset of all elements m/n where m and n are co-prime, n is non-zero and m/n means $m \cdot n^{-1}$. It is easily seen that this subset contains the sum, difference, product and quotient of any two of its elements (except dividing by 0, of course) and also contains 0 and 1, and so is a subfield, and is trivially isomorphic to the field of rationals.

Definition A finite field is a field which has a finite number of elements.

Theorem 34.5 *Any finite field has a characteristic p , for p a finite prime (i.e., it cannot have characteristic 0).*

For by Theorem 34.4 a field with characteristic 0 contains an infinite subfield and so must itself be infinite.

Theorem 34.6 *Let F be a finite field of characteristic p . The F has p^n elements, for some positive integer n .*

Proof Recall that if F has characteristic p , then the ring homomorphism $\phi : \mathbb{Z} \rightarrow F$ defined by $\phi(n) = n \cdot 1$ for all $n \in \mathbb{Z}$ has kernel $p\mathbb{Z}$, and thus the image of ϕ is a subfield K of F isomorphic to $\mathbb{Z}_p$. Since F is finite, it must certainly have finite dimension as a vector space over K , say $[F : K] = n$. If $v_1, v_2, \dots, v_n$ is a basis for F over K , then each element of F has the form $a_1 v_1 + a_2 v_2 + \dots + a_n v_n$ for elements $a_1, a_2, \dots, a_n \in K$. Thus to define an elements of F there are n coefficients a_i , and for each coefficient there are p choices, since K has only p elements. Therefore the total number of ways in which an element in F can be defined is p^n . ■

Theorem 34.7 *Let F be a finite field with p^n elements. Then F is the splitting field of the polynomial $x^{p^n} - x$ over the prime subfield of F .*

Proof Let F be a finite field of characteristic p . Then as in Theorem 34.6 the field F is an extension of degree n of its prime subfield K , which is isomorphic to $\mathbb{Z}_p$. Since F has p^n elements, the order of the multiplicative group $F^\times$ of non-zero elements of F is $p^n - 1$. Therefore $x^{p^n-1} \equiv 1$ for all $0 \neq x \in F$, and so $x^{p^n} = x$ for all $x \in F$. The polynomial $f(x) = x^{p^n} - x$ has at most p^n roots, and so its roots must be precisely the elements of F . Thus F is the splitting field of $f(x)$ over K . ■

35 Linear Spaces

A linear space $\mathbb{L}_p$ of dimension p is the totality of points of $\mathbb{R}_n$ into which a fixed point P is carried by the vectors of a p -dimensional vector space L of $\mathbb{R}_n$. Suppose there are given $p + 1$ points

$$P_k = (x_k^1, x_k^2, \dots, x_k^n), \quad k = 0, 1, \dots, p$$

which do not lie in any linear space of dimension $< p$. Let us consider the vectors $\mathbf{a}_k = \overrightarrow{P_0 P_k}, k = 1, 2, \dots, p$. These p vectors are linearly independent. For if they were linearly dependent, then the vector space spanned by them would have dimension $< p$. By applying these vectors to this vector space to P_0 , we obtain a linear space which on the one hand has dimension $< p$, and on the other contains the points $P_0, P_1, \dots, P_p$, which contradicts our assumption.

All points of a linear space are equivalent to one another because any point of it can be used to obtain the entire linear space.

36 Linear Equations

Let a system of m linear equations in n unknowns $x^1, x^2, \dots, x^n$ be given in the following form:

$$\begin{array}{cccc} a_{11}x^1 & + a_{12}x^2 & + \dots & + a_{1n}x^n = b_1 \\ a_{21}x^1 & + a_{22}x^2 & + \dots & + a_{2n}x^n = b_2 \\ \dots & \dots & \dots & \dots \\ a_{i1}x^1 & + a_{i2}x^2 & + \dots & + a_{in}x^n = b_i \\ \dots & \dots & \dots & \dots \\ a_{m1}x^1 & + a_{m2}x^2 & + \dots & + a_{mn}x^n = b_m \end{array} \quad (51)$$

which can be written in the following form based on the rules of matrix multiplication

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{i1} & a_{i2} & \dots & a_{in} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x^1 \\ x^2 \\ \dots \\ x^i \\ \dots \\ x^n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \dots \\ b_i \\ \dots \\ b_m \end{bmatrix} \quad (52)$$

To every column of the matrix of coefficients we can associate a vector

$$\mathbf{a}_k = \{a_{1k}, a_{2k}, \dots, a_{mk}\}, \quad k = 1, 2, \dots, n.$$

The original system of equations can be expressed as a single vector equation

$$\mathbf{a}_1x^1 + \mathbf{a}_2x^2 + \dots + \mathbf{a}_nx^n = \mathbf{b}, \quad (53)$$

where $\mathbf{b} = \{b_1, b_2, \dots, b_m\}$.

Consider the vector space $\mathbb{L}$ spanned by the vectors $\mathbf{a}_k, k = 1, \dots, n$ in $\mathbb{R}_m$. If Eq.(53) is solvable for $x^1, x^2, \dots, x^n$, then $\mathbf{b}$ must be a linear combination of the $\mathbf{a}_k$ and therefore belongs to $\mathbb{L}$. Conversely if $\mathbf{b}$ belongs to $\mathbb{L}$ then Eq.(53) is solvable.

Consider the vector space $\mathbb{L}'$ spanned by the vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n, \mathbf{b}$. A vector $\mathbf{c}$ in $\mathbb{L}'$ is of the form

$$\mathbf{c} = \lambda_1\mathbf{a}_1 + \lambda_2\mathbf{a}_2 + \dots + \lambda_n\mathbf{a}_n + \lambda\mathbf{b}.$$

If $\mathbf{b}$ belongs to $\mathbb{L}$, then Eq. (53) is satisfied by certain numbers $x^1, x^2, \dots, x^n$ which can be substituted for $\mathbf{b}$ and so $\mathbf{c}$ is a linear combination of the $\mathbf{a}_k, k = 1, \dots, n$ and therefore belongs to $\mathbb{L}$. In this case $\mathbb{L}' \subseteq \mathbb{L}$ and also $\mathbb{L} \subseteq \mathbb{L}'$ and therefore $\mathbb{L}'$ and $\mathbb{L}$ are identical.

Theorem 36.1 *The equation (53) is solvable for the x^i , if and only if, the maximal number of linearly independent vectors among $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$ is the same as the maximal number of linearly independent vectors among $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n, \mathbf{b}$.*

Definition The **rank** of a matrix is equal to the maximal number of independent column vectors in the matrix.

Consider the following matrix

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \dots & \dots & \dots & \dots & \dots \\ a_{i1} & a_{i2} & \dots & a_{in} & b_i \\ \dots & \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{bmatrix}$$

obtained by appending the vector $\{b_1, b_2, \dots, b_m\}$ to the final column. This matrix is called the **augmented** matrix of the system of equations (51).

Theorem 36.2 Eqs. (51) are solvable for the x^i if and only if the rank of the coefficient matrix of the system (51) is equal to the rank of the augmented matrix

The following system of equations is called a **homogenous in the x^i**

$$\begin{aligned} a_{11}x^1 + a_{12}x^2 + \dots + a_{1n}x^n &= 0 \\ a_{21}x^1 + a_{22}x^2 + \dots + a_{2n}x^n &= 0 \\ \dots & \dots \dots \dots \dots \dots \dots \\ a_{i1}x^1 + a_{i2}x^2 + \dots + a_{in}x^n &= 0 \\ \dots & \dots \dots \dots \dots \dots \dots \\ a_{m1}x^1 + a_{m2}x^2 + \dots + a_{mn}x^n &= 0 \end{aligned} \tag{54}$$

Any set of numbers $x^1, x^2, \dots, x^n$ which satisfy Eq. (54) are now taken to be the components of an n -dimensional vector $\mathbf{r} = \{x^1, x^2, \dots, x^n\}$. We call such a vector a *vector solution* of the system (54). The totality of vector solutions of the system of equations (54) forms a vector space.

Theorem 36.3 If the coefficient matrix of the system of equations (54) has rank r , then the set of vector solutions of this system is an $(n - r)$ -dimensional vector space.

Theorem 36.4 The system of equations (54) has a non-trivial solution, i.e. a solution $x^1, x^2, \dots, x^n$ such that not all the x^i vanish, if and only if, the rank of the matrix of (54) is less than n .

Theorem 36.5 If the number of equations in the system of homogenous equations (54) is less than the number of unknowns, then the system must have non-trivial solutions.

Theorem 36.6 All the solutions of a non-homogenous system of equations (51) are of the form $\mathbf{z} = \mathbf{r} + \eta$, where $\mathbf{r}$ is a fixed solution of the non-homogenous system (51) and η runs through all solutions of the corresponding homogenous system (54).

Let s be the maximal number of linearly independent rows of the matrix row vectors. Let us assume that the rows of the homogenous system are ordered so that the first s rows are independent to start with. This involves no loss of generality since it does not affect that rank. The i -th equation of this system is, by definition of the scalar product of two vectors, equivalent to the vector equation

$$\mathbf{a}_i \cdot \mathbf{r} = 0,$$

where we set $\mathbf{r} = \{x^1, x^2, \dots, x^n\}$.

Every system of n numbers $x^1, x^2, \dots, x^n$ satisfying the first s equation of (the re-ordered) (54), satisfies all of the equations of (54). Because the first s -rows are linearly independent, all the rows satisfy a relation of the form

$$\mathbf{a}_k = \lambda_1^k \mathbf{a}_1 + \lambda_2^k \mathbf{a}_2 + \dots + \lambda_s^k \mathbf{a}_s, \text{ for } 1 \leq k \leq m.$$

If we apply the distributive law on the right, we obtain

$$\mathbf{a}_k \cdot \mathbf{r} = \lambda_1^k(\mathbf{a}_1 \cdot \mathbf{r}) + \lambda_2^k(\mathbf{a}_2 \cdot \mathbf{r}) + \dots + \lambda_s^k(\mathbf{a}_s \cdot \mathbf{r}).$$

Since $\mathbf{a}_i \cdot \mathbf{r} = 0$ for $i = 1, 2, \dots, s$ the right hand side of this last equation becomes 0, so that

$$\mathbf{a}_k \cdot \mathbf{r} = 0, \quad k = 1, 2, \dots, m,$$

and in particular for $k = s + 1, s + 2, \dots, m$ as was to be proved.

The vector space of all the solutions of the first s equations of (54) is thus identical with the space of solutions of the entire system (54). Its dimension is this $n - r$ since r is the rank of the matrix of coefficients in (52). It thus follows that the rank of the matrix of the first s equations, i.e. of the matrix which consists only of the first s rows of (54) is equal to $n - (n - r) = r$. But the column vectors of this matrix are vectors with s components. Thus the maximal number of linearly independent column vectors of this matrix is $\leq s$ since by previous theorem any $s + 1$ vectors of s -dimensional vector space are linearly dependent.

Therefore the maximal number of linearly independent column vectors of a matrix is at most as large as the maximal number of linearly independent row vectors.

Now consider the transpose of the matrix of coefficients of (54). The maximal number of independent column vectors of this matrix is s , and the maximal number of row vectors is r . Applying the previous result we also obtain $s \leq r$. Hence we have $r \leq s$ and $s \leq r$ and therefore $s = r$. Thus we have proved

Theorem 36.7 *The maximal number of linearly independent column vectors of a matrix is equal to the maximal number of linearly independent row vectors of that matrix.*

Theorem 36.8 *With any given vector space L of $\mathbb{R}_n$, we can always associate a system of homogenous linear equations in n unknowns such that all the vectors of L , and no others, are vector solutions of this system.*

Let $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ be a basis of L , where we set $\mathbf{a}_i = \{\mathbf{a}_{i1}, \mathbf{a}_{i2}, \dots, \mathbf{a}_{in}\}$. If in addition we set

$$\mathbf{r} = \{x^1, x^2, \dots, x^n\}$$

then the equations

$$\mathbf{a}_i \cdot \mathbf{r} = 0, \quad i = 1, 2, \dots, p, \tag{55}$$

form a system of homogenous linear equations in the unknowns $x^1, x^2, \dots, x^n$ whose matrix is

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{i1} & a_{i2} & \dots & a_{in} \\ \dots & \dots & \dots & \dots \\ a_{p1} & a_{p2} & \dots & a_{pn} \end{bmatrix}$$

Let p be the dimension of L . The rank of this system must be $n - p$. The totality of vectors which are orthogonal to L is a vector space L' of dimension $n - p$. It consists precisely of all the vector solutions of (55).

We now seek a those vectors which are othogonal to the vector space L' just found, We chose a basis $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_{n-p}$ of L' . The $\mathbf{b}_i$ as vectors of L' are orthogonal to L , and this in particular to $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$. Thus, for $i = 1, 2, \dots, p$, we have

$$\mathbf{b}_1 \cdot \mathbf{a}_i = 0, \quad \mathbf{b}_2 \cdot \mathbf{a}_2 = 0, \quad \dots, \quad \mathbf{b}_{n-p} \cdot \mathbf{a}_i = 0.$$

Now just as for L , the totality of all vectors orthogonal to L' consists of all the vector solutions of the equations

$$\mathbf{b}_1 \cdot \mathbf{r} = 0, \quad \mathbf{b}_2 \cdot \mathbf{r} = 0, \quad \dots, \quad \mathbf{b}_{n-p} \cdot \mathbf{r} = 0. \quad (56)$$

Thus the $\mathbf{a}_i, i = 1, 2, \dots, p$ are orthogonal to L' . But the vectors which are orthogonal to L' form a vector space L'' of dimension $n - (n - p) = p$. Since the vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$ are in L'' and are linearly independent, then form a basis of L'' . Therefore L'' is identical with L . Thus the vectors of L are precisely the vector solutions of equations (56), i.e. of a certain system of $n - p$ linear homogenous equations.

Theorem 36.9 *The point solutions of a solvable system of equations of the type (51) form a linear space. This space has dimensions $n - rm$ where r is the rank of the coefficient matrix of the system. Conversely, every linear space may be represented as the totality of all point solutions of some suitable system of linear equations.*

37 Some Vector Theorems (3-dimensions)

$$\mathbf{B} \times \mathbf{C} = \det \begin{bmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ B_x & B_y & B_z \\ C_x & C_y & C_z \end{bmatrix} = \mathbf{i}(B_y C_z - C_y B_z) + \mathbf{j}(B_z C_x - B_x C_z) + \mathbf{k}(B_x C_y - B_y C_x)$$

Theorem 37.1 (BAC-CAB)

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{c}(\mathbf{a} \cdot \mathbf{b})$$

$$(\mathbf{a} \times \mathbf{b}) \times \mathbf{c} = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{a}(\mathbf{b} \cdot \mathbf{c})$$

Proof Given the expression for $\mathbf{B} \times \mathbf{C}$ we have

$$\begin{aligned} \mathbf{a} \times (\mathbf{b} \times \mathbf{c}) &= \det \begin{bmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_x & a_y & a_z \\ (b_y c_z - b_z c_y) & (b_z c_x - b_x c_z) & (b_x c_y - b_y c_x) \end{bmatrix} \\ &= \mathbf{i}(a_y(b_x c_y - b_y c_x) - a_z(b_z c_x - b_x c_z)) + \\ &\quad \mathbf{j}(a_z(b_y c_z - c_y b_z) - a_x(b_x c_y - b_y c_x)) + \\ &\quad \mathbf{k}(a_x(b_z c_x - c_z b_x) - a_y(b_y c_z - b_z c_y)) \end{aligned}$$

Prove component-by-component[8]. The $\mathbf{i}^{th}$ component can be written as

$$b_x(a_y c_y + a_z c_z) - c_x(a_y b_y + a_z b_z) = b_x(a_x c_x + a_y c_y + a_z c_z) - c_x(a_x b_x + a_y b_y + a_z b_z) = b_x(\mathbf{a} \cdot \mathbf{c}) - c_x(\mathbf{a} \cdot \mathbf{b}).$$

The $\mathbf{j}^{th}$ component can be written as

$$b_y(a_z c_z + a_x c_x) - c_y(a_z b_z + a_x b_x) = b_y(a_x c_x + a_y c_y + a_z c_z) - c_y(a_x b_x + a_y b_y + a_z b_z) = b_y(\mathbf{a} \cdot \mathbf{c}) - c_y(\mathbf{a} \cdot \mathbf{b}).$$

The k^{th} component can be written as

$$b_z(a_x c_x + a_y c_y) - c_z(a_x b_x + a_y b_y) = b_z(a_x c_x + a_y c_y + a_z c_z) - c_z(a_x b_x + a_y b_y + a_z b_z) = b_z(\mathbf{a} \cdot \mathbf{c}) - c_z(\mathbf{a} \cdot \mathbf{b}).$$

Putting back the respective unit vectors and adding we find the vector sum to be equal to

$$\mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{c}(\mathbf{a} \cdot \mathbf{b}).$$

This demonstrates the proof that

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{c}(\mathbf{a} \cdot \mathbf{b}).$$

The second part of the theorem is proven in a similar manner:

$$(\mathbf{a} \times \mathbf{b}) \times \mathbf{c} = -\mathbf{c} \times (\mathbf{a} \times \mathbf{b}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{a}(\mathbf{b} \cdot \mathbf{c}),$$

using the first half of the proof along with the substitutions:

$$\mathbf{a} \rightarrow -\mathbf{c} \quad \mathbf{b} \rightarrow \mathbf{a} \quad \mathbf{c} \rightarrow \mathbf{b}$$

viz.

$$(\mathbf{a} \times \mathbf{b}) \times \mathbf{c} = -\mathbf{c} \times (\mathbf{a} \times \mathbf{b}) \rightarrow \mathbf{a}(-\mathbf{c} \cdot \mathbf{b}) - \mathbf{b}(-\mathbf{c} \cdot \mathbf{a}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{a}(\mathbf{b} \cdot \mathbf{c}).$$

■

Theorem 37.2

$$(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = (\mathbf{a} \cdot \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) - (\mathbf{a} \cdot \mathbf{d})(\mathbf{b} \cdot \mathbf{c})$$

Proof

$$\mathbf{x} \cdot (\mathbf{c} \times \mathbf{d}) = (\mathbf{x} \times \mathbf{c}) \cdot \mathbf{d}$$

Let $\mathbf{x} = \mathbf{a} \times \mathbf{b}$. Then

$$\begin{aligned} (\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) &= \{(\mathbf{a} \times \mathbf{b}) \times \mathbf{c}\} \cdot \mathbf{d} \\ &= \{\mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{a}(\mathbf{b} \cdot \mathbf{c})\} \cdot \mathbf{d} \\ &= (\mathbf{a} \cdot \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) - (\mathbf{a} \cdot \mathbf{d})(\mathbf{b} \cdot \mathbf{c}) \\ &= \det \begin{bmatrix} \mathbf{a} \cdot \mathbf{c} & \mathbf{a} \cdot \mathbf{d} \\ \mathbf{b} \cdot \mathbf{c} & \mathbf{b} \cdot \mathbf{d} \end{bmatrix} \end{aligned}$$

Theorem 37.3

$$(\mathbf{a} \times \mathbf{b}) \times (\mathbf{c} \times \mathbf{d}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c} \times \mathbf{d}) - \mathbf{a}(\mathbf{b} \cdot \mathbf{c} \times \mathbf{d}) = \mathbf{c}(\mathbf{a} \cdot \mathbf{b} \times \mathbf{d}) - \mathbf{d}(\mathbf{a} \cdot \mathbf{b} \times \mathbf{c})$$

Proof

$$\mathbf{x} \times (\mathbf{c} \times \mathbf{d}) = \mathbf{c}(\mathbf{x} \cdot \mathbf{d}) - \mathbf{d}(\mathbf{x} \cdot \mathbf{c})$$

Let $\mathbf{x} = \mathbf{a} \times \mathbf{b}$. Then

$$\begin{aligned} \text{I: } (\mathbf{a} \times \mathbf{b}) \times (\mathbf{c} \times \mathbf{d}) &= \mathbf{c}(\mathbf{a} \times \mathbf{b} \cdot \mathbf{d}) - \mathbf{d}(\mathbf{a} \times \mathbf{b} \cdot \mathbf{c}) \\ &= \mathbf{c}(\mathbf{a} \cdot \mathbf{b} \times \mathbf{d}) - \mathbf{d}(\mathbf{a} \cdot \mathbf{b} \times \mathbf{c}) \quad \blacksquare \end{aligned}$$

Also

$$(\mathbf{a} \times \mathbf{b}) \times \mathbf{y} = \mathbf{b}(\mathbf{a} \cdot \mathbf{y}) - \mathbf{a}(\mathbf{b} \cdot \mathbf{y})$$

Let $\mathbf{y} = \mathbf{c} \times \mathbf{d}$. Then

$$\text{II: } (\mathbf{a} \times \mathbf{b}) \times (\mathbf{c} \times \mathbf{d}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c} \times \mathbf{d}) - \mathbf{a}(\mathbf{b} \cdot \mathbf{c} \times \mathbf{d}) \quad \blacksquare$$

Theorem 37.4 (Compound Vector Scalar Product)

$$(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{b} \times \mathbf{c}) \times (\mathbf{c} \times \mathbf{a}) = (\mathbf{a} \cdot \mathbf{b} \times \mathbf{c})^2$$

Proof

$$\mathbf{x} \times (\mathbf{c} \times \mathbf{a}) = \mathbf{c}(\mathbf{x} \cdot \mathbf{a}) - \mathbf{a}(\mathbf{x} \cdot \mathbf{c})$$

Let $\mathbf{x} = \mathbf{b} \times \mathbf{c}$, then

$$\begin{aligned} (\mathbf{b} \times \mathbf{c}) \times (\mathbf{c} \times \mathbf{a}) &= \mathbf{c}(\mathbf{b} \times \mathbf{c} \cdot \mathbf{a}) - \mathbf{a}(\mathbf{b} \times \mathbf{c} \cdot \mathbf{c}) \\ &= \mathbf{c}(\mathbf{a} \cdot \mathbf{b} \times \mathbf{c}) - \mathbf{a}(\mathbf{b} \cdot \mathbf{c} \times \mathbf{c}) \\ &= \mathbf{c}(\mathbf{a} \cdot \mathbf{b} \times \mathbf{c}). \end{aligned}$$

Therefore

$$\begin{aligned} (\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{b} \times \mathbf{c}) \times (\mathbf{c} \times \mathbf{a}) &= (\mathbf{a} \times \mathbf{b}) \cdot \mathbf{c}(\mathbf{a} \cdot \mathbf{b} \times \mathbf{c}) \\ &= (\mathbf{a} \times \mathbf{b} \cdot \mathbf{c})(\mathbf{a} \cdot \mathbf{b} \times \mathbf{c}) \\ &= (\mathbf{a} \cdot \mathbf{b} \times \mathbf{c})^2 \quad \blacksquare \end{aligned}$$

Definition The operator ∇ is given by

$$\nabla = \mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z}.$$

Definition The Divergence of a vector field is given by

$$\nabla \cdot \mathbf{a} = \frac{\partial}{\partial x} a_x + \frac{\partial}{\partial y} a_y + \frac{\partial}{\partial z} a_z.$$

Definition The Laplacian (Divergence of the Gradient) operator is given by

$$\nabla^2 = \nabla \cdot \nabla = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}.$$

Definition The Curl of a vector field is given by

$$\nabla \times \mathbf{a} = \det \begin{bmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ a_x & a_y & a_z \end{bmatrix}$$

Theorem 37.5

$$\nabla \cdot (\mathbf{a} \times \mathbf{b}) = \mathbf{b} \cdot (\nabla \times \mathbf{a}) - \mathbf{a} \cdot (\nabla \times \mathbf{b}).$$

Proof

$$\nabla \cdot (\mathbf{a} \times \mathbf{b}) = \det \begin{bmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ a_x & a_y & a_z \\ b_x & b_y & b_z \end{bmatrix}$$

$$\begin{aligned}
&= [\partial_x(a_y b_z - a_z b_y) - \partial_y(a_x b_z - a_z b_x) + \partial_z(a_x b_y - a_y b_x)] \\
&= b_z \partial_x a_y - b_y \partial_x a_z - b_z \partial_y a_x + b_x \partial_y a_z + b_y \partial_z a_x - b_x \partial_z a_y \\
&+ a_y \partial_x b_z - a_z \partial_x b_y - a_x \partial_y b_z + a_z \partial_y b_x + a_x \partial_z b_y - a_y \partial_z b_x \\
&= [b_x(\partial_y a_z - \partial_z a_y) - b_y(\partial_x a_z - \partial_z a_x) + b_z(\partial_x a_y - \partial_y a_x)] \\
&- [a_x(\partial_y b_z - \partial_z b_y) - a_y(\partial_x b_z - \partial_z b_x) + a_z(\partial_x b_y - \partial_y b_x)] \\
&= \mathbf{b} \cdot (\nabla \times \mathbf{a}) - \mathbf{a} \cdot (\nabla \times \mathbf{b}) \quad \blacksquare
\end{aligned}$$

Theorem 37.6

$$\nabla \times (\mathbf{a} \times \mathbf{b}) = (\mathbf{b} \cdot \nabla) \mathbf{a} - (\mathbf{a} \cdot \nabla) \mathbf{b} + (\nabla \cdot \mathbf{b}) \mathbf{a} - (\nabla \cdot \mathbf{a}) \mathbf{b}.$$

Theorem 37.7

$$\nabla \times (\nabla \times \mathbf{a}) = \nabla(\nabla \cdot \mathbf{a}) - (\nabla \cdot \nabla) \mathbf{a} = \nabla(\nabla \cdot \mathbf{a}) - \nabla^2 \mathbf{a}.$$

38 Reciprocal Vector Space (3-dimensions)

Take three non-coplanar vectors $\mathbf{a}, \mathbf{b}, \mathbf{c}$ which form a right-handed system of vectors and form the vector scalar product whose magnitude is the volume, V , of the parallelepiped with edges given by $\mathbf{a}, \mathbf{b}, \mathbf{c}$

$$V = \mathbf{a} \cdot \mathbf{b} \times \mathbf{c}.$$

The set of vectors $\mathbf{a}', \mathbf{b}', \mathbf{c}'$ which are reciprocal to $\mathbf{a}, \mathbf{b}, \mathbf{c}$ is given by

$$\begin{aligned}
\mathbf{a}' &= \frac{\mathbf{b} \times \mathbf{c}}{\mathbf{a} \cdot \mathbf{b} \times \mathbf{c}} \\
\mathbf{b}' &= \frac{\mathbf{c} \times \mathbf{a}}{\mathbf{a} \cdot \mathbf{b} \times \mathbf{c}} \\
\mathbf{c}' &= \frac{\mathbf{a} \times \mathbf{b}}{\mathbf{a} \cdot \mathbf{b} \times \mathbf{c}}
\end{aligned}$$

The reciprocal basis $\mathbf{a}', \mathbf{b}', \mathbf{c}'$ satisfy the following identities

$$\begin{aligned}
\mathbf{a} \cdot \mathbf{a}' &= 1, & \mathbf{b} \cdot \mathbf{a}' &= 0, & \mathbf{c} \cdot \mathbf{a}' &= 0 \\
\mathbf{b} \cdot \mathbf{b}' &= 1, & \mathbf{a} \cdot \mathbf{b}' &= 0, & \mathbf{c} \cdot \mathbf{b}' &= 0 \\
\mathbf{c} \cdot \mathbf{c}' &= 1, & \mathbf{a} \cdot \mathbf{c}' &= 0, & \mathbf{b} \cdot \mathbf{c}' &= 0
\end{aligned}$$

Theorem 38.1 (Reciprocal Basis has Reciprocal Volume = 1/V)

$$\mathbf{a}' \cdot \mathbf{b}' \times \mathbf{c}' = \frac{1}{V}$$

Proof

$$\begin{aligned} \mathbf{a}' &= \frac{\mathbf{b} \times \mathbf{c}}{V} \\ \mathbf{b}' &= \frac{\mathbf{c} \times \mathbf{a}}{V} \\ \mathbf{c}' &= \frac{\mathbf{a} \times \mathbf{b}}{V} \end{aligned}$$

$$\begin{aligned} \mathbf{a}' \cdot \mathbf{b}' \times \mathbf{c}' &= \frac{(\mathbf{b} \times \mathbf{c}) \cdot (\mathbf{c} \times \mathbf{a}) \times (\mathbf{a} \times \mathbf{b})}{V^3} = \frac{(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{b} \times \mathbf{c}) \times (\mathbf{c} \times \mathbf{a})}{V^3} \\ &= \frac{(\mathbf{a} \cdot \mathbf{b} \times \mathbf{c})^2}{V^3} = \frac{V^2}{V^3} = \frac{1}{V} \quad \blacksquare \end{aligned}$$

39 Vandermonde Determinant

The Vandermonde Determinant for n variables $x = [x_1, x_2, \dots, x_n]$ is given by

$$V_n(x) = \begin{bmatrix} 1 & x_1 & x_1^2 & x_1^3 & \dots & x_1^n \\ 1 & x_2 & x_2^2 & x_2^3 & \dots & x_2^n \\ 1 & x_3 & x_3^2 & x_3^3 & \dots & x_3^n \\ \vdots & & & & & \\ 1 & x_n & x_n^2 & x_n^3 & \dots & x_n^n \end{bmatrix}$$

Except for the first column, subtract x_1 times the previous column to the rest of the columns. Since a multiple of any column added to any other column does not change the determinant, we can write

$$\det V_n(x) = \det \begin{bmatrix} 1 & 0 & 0 & 0 & \dots & 0 \\ 1 & x_2 - x_1 & x_2(x_2 - x_1) & x_2^2(x_2 - x_1) & \dots & x_2^{n-1}(x_2 - x_1) \\ 1 & x_3 - x_1 & x_3(x_3 - x_1) & x_3^2(x_3 - x_1) & \dots & x_3^{n-1}(x_3 - x_1) \\ \vdots & & & & & \\ 1 & x_n - x_1 & x_n(x_n - x_1) & x_n^2(x_n - x_1) & \dots & x_n^{n-1}(x_n - x_1) \end{bmatrix}$$

Expanding the determinant using the elements in the first row we have

$$\det V_n(x) = \det \begin{bmatrix} x_2 - x_1 & x_2(x_2 - x_1) & x_2^2(x_2 - x_1) & \dots & x_2^{n-1}(x_2 - x_1) \\ x_3 - x_1 & x_3(x_3 - x_1) & x_3^2(x_3 - x_1) & \dots & x_3^{n-1}(x_3 - x_1) \\ \vdots & & & & \\ x_n - x_1 & x_n(x_n - x_1) & x_n^2(x_n - x_1) & \dots & x_n^{n-1}(x_n - x_1) \end{bmatrix}$$

Factoring out $x_2 - x_1$ from the first row, $x_3 - x_1$ from the second row, and so on until the last row we have

$$\det V_n(x_1, x_2, \dots, x_n) = \left[\prod_{i=2}^n (x_i - x_1) \right] \det \begin{bmatrix} 1 & x_2 & x_2^2 & \dots & x_2^{n-1} \\ 1 & x_3 & x_3^2 & \dots & x_3^{n-1} \\ \vdots & & & & \\ 1 & x_n & x_n^2 & \dots & x_n^{n-1} \end{bmatrix} = \prod_{i=2}^n (x_i - x_1) \det V_{n-1}(x_2, x_3, \dots, x_n)$$

Proceeding in a similar way by subtracting x_2 times the previous column except for the first column we can continue

$$\det V_n(x_1, x_2, \dots, x_n) = \left[\prod_{i=2}^n (x_i - x_1) \right] \left[\prod_{i=3}^n (x_i - x_2) \right] \det V_{n-2}(x_3, x_4, \dots, x_n)$$

To arrive at

$$\det V_n(x_1, x_2, \dots, x_n) = \prod_{\substack{1 \leq j \\ i < j}}^n (x_j - x_i) = \prod_{1 \leq i < j \leq n} (x_j - x_i)$$

40 Cayley-Hamilton Theorem

The characteristic equation for eigenvalues $0 = \det |A - \lambda I|$ can be expanded as a polynomial in λ

$$\phi(\lambda) = \det |A - \lambda I| = a_n \lambda^n + a_{n-1} \lambda^{n-1} + \dots + a_1 \lambda + a_0, \quad a_n = (-1)^n.$$

And the characteristic equation can be written in the form $\phi(\lambda) = 0$.

If in any polynomial $f(\lambda) = \sum c_i \lambda^i$ we substitute the matrix A for λ and multiply the constant term c_0 by I , we obtain a matrix denoted by $f(A)$.

Theorem 40.1 (*Cayley-Hamilton*) *Any square matrix satisfies its characteristic equation.*

Proof Let $\phi(\lambda)$ be the characteristic determinant of A . We want to show that $\phi(A) = 0$. Since the elements of $A - \lambda I$ are linear functions of λ , and the elements of its adjoint [the transpose of the matrix of signed minors of $A - \lambda I$ - denoted by $\text{Adj}-(A - \lambda I)$] are $(n-1)$ -rowed determinants, they are polynomials in λ of degree $\leq n-1$. Let C stand for the $\text{Adj}-(A - \lambda I)$. If the element in the i -th row and j -th column of C is $\sum_k c_{ijk} \lambda^k$, then

$$C = \sum_{k=0}^{n-1} C_k \lambda^k, \quad C_k = (c_{ijk}) \quad (i, j = 1, 2, \dots, n)$$

Using the relationship between the adjoint and the determinant, $\phi(\lambda) = \det |A - \lambda I|$, we can write

$$(A - \lambda I)C = \phi(\lambda)I.$$

This expression can be expanded as

$$A \sum_{k=0}^{n-1} C_k \lambda^k - \lambda \sum_{k=0}^{n-1} C_k \lambda^k = \sum_{k=0}^n a_k \lambda^k I$$

Equating the terms free of λ and the coefficients of $\lambda, \lambda^1, \dots, \lambda^{n-1}, \lambda^n$ we get

$$\begin{aligned} AC_0 &= a_0 I, \\ AC_1 - C_0 &= a_1 I, \\ AC_2 - C_1 &= a_2 I, \\ &\dots \dots \\ AC_{n-1} - C_{n-2} &= a_{n-1} I, \\ -C_{n-1} &= a_n I. \end{aligned}$$

Multiply these equations on the left by $I, A, A^2, \dots, A^{n-1}, A^n$ respectively and add; we get

$$0 = a_0 I + a_1 A + a_2 A^2 + \dots + a_{n-1} A^{n-1} + a_n A^n = \phi(A),$$

and we see that the matrix A satisfies its own characteristic equation. ■

41 Sylvesters Criterion for Positive Definite Hermitian Matrices

Theorem 41.1 (*Sylvester's Criterion*) *An $n \times n$ Hermitian matrix M is positive-definite if and only if all the leading principal minors are positive.*

Proof Let M be an $n \times n$ Hermitian positive definite matrix. Then $0 \leq x^\dagger M x$ if x is not the zero vector. Let M_k be the k -th leading principal minor (the first k -rows and k -columns in the upper left hand corner of M). Select a non-zero vector x as follows

$$x = \begin{bmatrix} x_1 \\ \vdots \\ x_k \\ 0 \\ \vdots \\ 0 \end{bmatrix} = \begin{bmatrix} \vec{x} \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

Then we have that $0 \leq x^\dagger M x = \vec{x}^\dagger M_k \vec{x}$. Therefore, since $\vec{x} \neq \vec{0}$ is otherwise arbitrary, then all the eigenvalues of M_k must be positive and hence $\det |M_k| > 0$

To prove the reverse implication, we use induction. The general form of an $(n+1) \times (n+1)$ Hermitian matrix is

$$M_{n+1} = \begin{bmatrix} M_n & \vec{v} \\ \vec{v}^\dagger & d \end{bmatrix}$$

Denote the $(n+1)$ dimensional vector x as

$$x = \begin{bmatrix} \vec{x} \\ x_{n+1} \end{bmatrix}$$

Then

$$x^\dagger M_{n+1} x = \vec{x}^\dagger M_n \vec{x} + x_{n+1} \vec{x}^\dagger \vec{v} + \bar{x}_{n+1} \vec{v}^\dagger \vec{x} + d |x_{n+1}|^2$$

By completing the square, this last expression is equal to

$$\begin{aligned} & (\vec{x}^\dagger + \vec{v}^\dagger M_n^{-1} \bar{x}_{n+1}) M_n (\vec{x} + x_{n+1} M_n^{-1} \vec{v}) - |x_{n+1}|^2 \vec{v}^\dagger M_n^{-1} \vec{v} + d |x_{n+1}|^2 \\ & = (\vec{x} + \vec{c})^\dagger M_n (\vec{x} + \vec{c}) + |x_{n+1}|^2 (d - \vec{v}^\dagger M_n^{-1} \vec{v}), \end{aligned}$$

where $\vec{c} = x_{n+1} M_n^{-1} \vec{v}$. M_n^{-1} exists because the eigenvalues of M_n are all positive. The first term is positive by the induction hypothesis. Using this useful determinant expansion

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \det |A| \det |D - C A^{-1} B|$$

We have that

$$\det |M_{n+1}| = \det |M_n| (d - \vec{v}^\dagger M_n^{-1} \vec{v}) > 0,$$

which implies $(d - \vec{v}^\dagger M_n^{-1} \vec{v}) > 0$. Hence $x^\dagger M_{n+1} x > 0$ (is also positive definite). ■

42 Notes on Riemannian Hypersurfaces

Let us summarize some facts about Riemannian Geometry:

Tangent Vectors $\mathbf{r}_j = \partial \mathbf{r} / \partial x^j$, $\mathbf{a} = a^i \mathbf{r}_i$, second derivatives: $\mathbf{r}_{ij} = \frac{\partial^2 \mathbf{r}}{\partial x^i \partial x^j}$

Metric: $g_{ij} = \mathbf{r}_i \cdot \mathbf{r}_j = g_{ji}$ (symmetric).

Dot product: $\mathbf{a} \cdot \mathbf{b} = g_{ij} a^i b^j$.

Consider the determinant

$$g = \det |g_{ij}| = \begin{vmatrix} g_{11} & g_{12} & \cdots & g_{1n} \\ g_{21} & g_{22} & \cdots & g_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ g_{i1} & g_{i2} & \cdots & g_{in} \\ \cdots & \cdots & \cdots & \cdots \\ g_{n1} & g_{n2} & \cdots & g_{nn} \end{vmatrix} \quad (57)$$

We suppose, here and throughout, that g is not zero. Let Δ^{ij} be the cofactor of g_{ij} in this determinant, so that

$$g_{mr} \Delta^{ms} = g_{rm} \Delta^{sm} = \delta_r^s g,$$

which follows from the ordinary rules for developing a determinant.

Let us construct new quantities g^{ij} [10]. The values of the components of g^{ij} are equal to the cofactor of the g_{ij} , divided by the full determinant g ,

$$g^{kl} = g^{-1} \Delta^{kl}. \quad (58)$$

Alternatively the cofactor of g_{ij} can be expressed in terms of g^{ij} as follows

$$\Delta^{ij} = g g^{ij}. \quad (59)$$

g^{ij} satisfies the following equations:

$$g_{ik} g^{kl} = \delta_i^l \text{ inverse} \quad (60)$$

Now g_{ij} is symmetric and we can also show that g^{ij} is symmetric. Multiply Eq. (60) by $g_{ls} g^{ir}$. The left hand side becomes

$$g_{ik} g^{kl} g_{ls} g^{ir} = g_{ki} g^{ir} g^{kl} g_{sl} = \delta_k^r g^{kl} g_{sl} = g_{sl} g^{rl},$$

while the right hand side becomes

$$\delta_i^l g_{ls} g^{ir} = g_{is} g^{ir} = g_{si} g^{ir} = \delta_s^r.$$

Therefore we obtain

$$g_{sl} g^{rl} = \delta_s^r \text{ inverse} \quad (61)$$

Comparing Eqs (60) and (61), we find that

$$g^{kl} = g^{lk}$$

Since g_{ij} is symmetric, it is obvious that g^{ij} should be symmetric also.

Let us examine the implications of the above applied to surfaces embedded in $\mathbb{R}^n$. A *curve* is defined as the totality of points given by the equations

$$x^r = f^r(u) \quad (r = 1, 2, \dots, n). \quad (62)$$

Here u is a parameter and $f^r(u)$ are n functions.

Next consider the totality of points given by

$$x^r = f^r(t^1, t^2, \dots, t^m) \quad (r = 1, 2, \dots, n), \quad (63)$$

where the t 's are parameters and $m < n$. This set of points forms V_m , a subspace of V_n . According to the Implicit Function Theorem it is possible to eliminate the parameters from (63). As long as the system is differentiable enough and the the Jacobian of the first m equations does not vanish, we can find m functions $g^i(x^1, \dots, x^m)$ such that

$$t^i = g^i(x^1, \dots, x^m) \text{ for } i = 1, \dots, m \quad (64)$$

which implies that the remaining functions $x^r, r = m + 1, \dots, n$ can be written in terms of the first m coordinates $x^1, \dots, x^m$ as follows:

$$x^r = f^r(x^1, \dots, x^m) \text{ for } r = m + 1, \dots, n \quad (65)$$

In the case of $m = n - 1$ there is only one such relation and so, elimination gives just one equation

$$\phi(x^1, x^2, \dots, x^n) = x^n - f^n(x^1, \dots, x^m) = 0. \quad (66)$$

In this case, V_{n-1} divides portions of V_n into two parts for which respectively ϕ is positive and negative. V_{n-1} is commonly referred to as a *hypersurface* in V_n .

Let V_n be a region of $\mathbb{R}^n$, bounded by a smooth (i.e. sufficiently differentiable) surface B_{n-1} , which is an $n - 1$ dimensional closed manifold. We take B_{n-1} as an example of a hypersurface V_{n-1} . We has suppose that B_{n-1} is parameterized by $n - 1$ independent parameters $t^\alpha (\alpha = 1, 2, \dots, n - 1)$. Let the hypersurface B_{n-1} be defined implicitly by the equation (66).

On the surface of B_{n-1} , $x^i = x^i(t^\alpha)$, and therefore

$$\frac{\partial \phi}{\partial x^j} \frac{\partial x^j}{\partial t^\alpha} = 0. \quad (67)$$

Equation (67) may be regarded as $n - 1$ linear conditions upon the n quantities $\partial \phi / \partial x^j$. Since the matrix

$$\left[\frac{\partial x^j}{\partial t^\alpha} \right]$$

is assumed to have its full rank value $n - 1$ on B_{n-1} , the mutual ratios of the partials $\partial \phi / \partial x^j$ are fully determined by (67).

For future reference let us define a set of determinants with respect to the change of variables between the x^i and the t^α

$$D_k = (-1)^{k-1} \det \left[\frac{\partial(x^1, \dots, x^{k-1}, x^{k+1}, \dots, x^n)}{\partial(t^1, \dots, t^{n-1})} \right], \quad k = 1, 2, \dots, n \quad (68)$$

The n determinants D_k satisfy the $n - 1$ linear conditions

$$D_k \frac{\partial x^k}{\partial t^\alpha} = 0, \quad (69)$$

which can be established by noting that the left hand side can be written as an n -by- n determinant which has two rows the same and therefore vanishes. In view of (67, conditions (69) imply that the $\partial\phi/\partial x^k$ and D_k are proportional

$$\frac{\partial\phi}{\partial x^k} = \alpha D_k,$$

for some scalar factor of proportionality α . Hence the $\partial\phi/\partial x^k$ determine the unit normal to B_{n-1} , namely

$$n_k = \frac{1}{\sqrt{(\nabla\phi)^2}} \frac{\partial\phi}{\partial x^k} \quad (70)$$

$$n_k = \frac{D_k}{\sqrt{g^{ij} D_i D_j}} = \frac{D_k}{D}, \quad (71)$$

where

$$D^2 = g^{ij} D_i D_j = g^{ij} \frac{\partial\phi}{\partial x^i} \frac{\partial\phi}{\partial x^j} \alpha^{-2}.$$

43 Hypersurfaces and Concepts of Area

Consider an $n - 1$ dimensional hypersurface, B_{n-1} , embedded in a Riemannian manifold M_n (which we assume to be sufficiently differentiable). Let $x^1, \dots, x^n$ be local coordinates in M_n and suppose that B_{n-1} is represented locally in the form

$$x^j = x^j(t^1, \dots, t^{n-1}), \quad j = 1, \dots, n$$

The metric tensor of M_n is denoted by g_{ij} . and the metric tensor on B_{n-1} by $\bar{g}_{\alpha\beta}$. These two tensors are related by

$$\bar{g}_{\alpha\beta} = g_{ij} \frac{\partial x^i}{\partial t^\alpha} \frac{\partial x^j}{\partial t^\beta} = \frac{\partial x^i}{\partial t^\alpha} g_{ij} \frac{\partial x^j}{\partial t^\beta} \quad (72)$$

The element of area (or volume) in M_n is

$$dV = \sqrt{g} dx^1 \dots dx^n, \quad (73)$$

where g is the determinant of the metric tensor in M_n . Similarly the element of area (volume) in the hypersurface B_{n-1} is given by

$$dS = \sqrt{\bar{g}} dt^1 \dots dt^{n-1}. \quad (74)$$

We will use (72) to define the element of area in B_{n-1} based on expansion of the determinant of $\bar{g}$ in terms of products of the various minors in a manner similar to the Cauchy-Binet formula or the Laplace expansion of a determinant. Let S_n stand for the set of permutations of the numbers $\{1, 2, \dots, n\}$ and let P represent one of the members of S_n .

$\bar{g}_{\alpha\beta}$ is an $(n-1)$ -by- $(n-1)$ matrix and we can consider its determinant.

$$\bar{g} = \det |\bar{g}| = \sum_{P \in S_{n-1}} \epsilon_{p_1 p_2 \dots p_{n-1}} \bar{g}_{1p_1} \bar{g}_{2p_2} \dots \bar{g}_{n-1p_{n-1}} \quad (75)$$

We can think of the right hand side of (72) as representing the matrix product of three matrices.

$$\begin{aligned} J &= \left[\frac{\partial x^i}{\partial t^\alpha} \right] \text{ is } n\text{-by-}(n-1), \\ G &= [g] \text{ is } n\text{-by-}n \end{aligned}$$

and (72) can be written in the form:

$$\begin{aligned} \bar{g}_{\alpha\beta} &= \sum_{i,j=1}^n \left[\frac{\partial x^i}{\partial t^\alpha} \right]_{\alpha i}^T [g]_{ij} \left[\frac{\partial x^j}{\partial t^\beta} \right]_{j\beta} = \frac{\partial x^i}{\partial t^\alpha} g_{ij} \frac{\partial x^j}{\partial t^\beta} \text{ summation convention on } i \text{ and } j \\ [\bar{g}_{\alpha\beta}] &= J^T G J \end{aligned}$$

Inserting this computation into (75) we have $\bar{g}_{\alpha\beta}$ is an $(n-1)$ -by- $(n-1)$ matrix and we can consider its determinant. Let $\bar{g} = \det |\bar{g}_{\alpha\beta}|$.

$$\begin{aligned} \bar{g} &= \sum_{P \in S_{n-1}} \epsilon_{p_1 p_2 \dots p_{n-1}} \sum_{\substack{i_1, \dots, i_{n-1}=1 \\ k_1, \dots, k_{n-1}=1}}^n (J_{1i_1}^T G_{i_1 k_1} J_{k_1 p_1}) (J_{2i_2}^T G_{i_2 k_2} J_{k_2 p_2}) \dots (J_{n-1 i_{n-1}}^T G_{i_{n-1} k_{n-1}} J_{k_{n-1} p_{n-1}}) \\ &= \sum_{P \in S_{n-1}} \epsilon_{p_1 p_2 \dots p_{n-1}} \sum_{\substack{i_1, \dots, i_{n-1}=1 \\ k_1, \dots, k_{n-1}=1}}^n (J_{1i_1}^T \dots J_{n-1 i_{n-1}}^T) (G_{i_1 k_1} \dots G_{i_{n-1} k_{n-1}}) (J_{k_1 p_1} \dots J_{k_{n-1} p_{n-1}}) \\ &= \sum_{\substack{i_1, \dots, i_{n-1}=1 \\ k_1, \dots, k_{n-1}=1}}^n (J_{1i_1}^T \dots J_{n-1 i_{n-1}}^T) (G_{i_1 k_1} \dots G_{i_{n-1} k_{n-1}}) \sum_{P \in S_{n-1}} \epsilon_{p_1 p_2 \dots p_{n-1}} (J_{k_1 p_1} \dots J_{k_{n-1} p_{n-1}}) \end{aligned}$$

Examining the last term in the sum on the right hand side we find a sum over the $n-1$ set $\{p_1, p_2, \dots, p_{n-1}\}$ for a particular choice of index values: $\{k_1, \dots, k_{n-1}\}$

$$\sum_{P \in S_{n-1}} \epsilon_{p_1 p_2 \dots p_{n-1}} (J_{k_1 p_1} \dots J_{k_{n-1} p_{n-1}}) = \epsilon_{k_1 k_2 \dots k_{n-1}} |J_k| = \epsilon_{k_1 k_2 \dots k_{n-1}} (-1)^{1+k} D_k \quad (76)$$

where the index k is not included in $\{k_1, \dots, k_{n-1}\}$ and final result involves the respective determinant D_k .

At this point notice that the sum over the $n-1$ variables $k_1, k_2, \dots, k_{n-1}$ involves summing over all values from 1 to n for each k_i , but that these values must be distinct. If $k_i = k_j$ for some pair i, j , then the sum is zero because of the presence of the permutation symbol $\epsilon_{k_1 k_2 \dots k_{n-1}}$. This means that there are $n-1$ distinct k values chosen from the full set of n values, therefore one of the k_i values is to be omitted in this sum. Let k stand for this missing value and J_k stand for the matrix J with the k -th row deleted. The set of k_i values in any set is also propagated on to the G_{ik} matrices and the corresponding k -th column of G will not contribute this particular set of choices. There will be n such cases and therefore we insert a sum over $k = 1, 2, \dots, n$ to account for each case in the following. The sum over the $i_1, \dots, i_{n-1}$ indices follows that same way.

$$\begin{aligned}
\bar{g} &= \sum_{k=1}^n \sum_{\substack{i_1, \dots, i_{n-1}=1 \\ k_1, \dots, k_{n-1}=1, \neq k}}^n (J_{1i_1}^T \dots J_{n-1i_{n-1}}^T) (G_{i_1 k_1} \dots G_{i_{n-1} k_{n-1}}) \epsilon_{k_1 k_2 \dots k_{n-1}} (-1)^{1+k} D_k \\
&= \sum_{i,k=1}^n \sum_{\substack{i_1, \dots, i_{n-1}=1, \neq i}}^n (J_{1i_1}^T \dots J_{n-1i_{n-1}}^T) \epsilon_{i_1 i_2 \dots i_{n-1}} (-1)^{i+k} \text{cofactor}(g_{ik}) (-1)^{1+k} D_k \\
&= \sum_{i,k=1}^n \sum_{\substack{i_1, \dots, i_{n-1}=1, \neq i}}^n \epsilon_{i_1 i_2 \dots i_{n-1}} (J_{1i_1}^T \dots J_{n-1i_{n-1}}^T) (-1)^{i+k} \Delta^{ik} (-1)^{1+k} D_k \\
&= \sum_{i,k=1}^n (-1)^{1+i} D_i (-1)^{i+k} \Delta^{ik} (-1)^{1+k} D_k \\
&= \sum_{i,k=1}^n D_i g^{ik} D_k = g \sum_{i=1}^n D^i D_i = g D^2,
\end{aligned}$$

where $g = |g_{ij}|$ as before and the sum is over the n values of i and k which correspond to the rows or columns which are crossed out to form the n minors each of dimension $(n-1)$ -by- $(n-1)$. Hence we find that

$$\bar{g} = g D^2.$$

We finally derive the relationship between the element of surface are on the hyperplane H and the metric of the n -dimensional Riemann M_n space that H is embedded inside.

$$dS = \sqrt{\bar{g}} dt^1 \dots dt^{n-1} = \sqrt{g D^2} dt^1 \dots dt^{n-1} \quad (77)$$

If the $n-1$ dimensional hypersurface H which is embedded in Cartesian $\mathbb{R}^n$ is given in parametric form as follows:

$$\mathbf{r}(x) = \{x, f(x)\}$$

Then $g_{ij} = I_n$ (n -by- n identity matrix) and J is an n -by- $n-1$ matrix:

$$J = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \\ \frac{\partial f}{\partial x^1} & \frac{\partial f}{\partial x^2} & \dots & \frac{\partial f}{\partial x^{n-1}} \end{bmatrix} \quad (78)$$

$$J^T = \begin{bmatrix} 1 & 0 & \dots & 0 & \frac{\partial f}{\partial x^1} \\ 0 & 1 & \dots & 0 & \frac{\partial f}{\partial x^2} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 & \frac{\partial f}{\partial x^n} \end{bmatrix} \quad (79)$$

Working out the minors, we find $D_n = 1$ (cross out the last row of J), and crossing out any of the other rows $i < n$ inserts a 0 in the i -th diagonal and shifts all the other rows $i+1, \dots, n$ up one row. The resulting determinant is equal to $D_k = (-1)^{1+k} \frac{\partial f}{\partial x_k}$

Therefore we obtain

$$\bar{g} = \sum_{k=1}^n D_k = \left(\frac{\partial f}{\partial x^1}\right)^2 + \left(\frac{\partial f}{\partial x^2}\right)^2 + \dots + \left(\frac{\partial f}{\partial x^n}\right)^2 + 1 \quad (80)$$

And the resulting magnitude, given that $g_{ij} = I_n$ is $\sqrt{D^2} = \sqrt{1 + |\nabla f|^2}$ and

$$dS = \sqrt{1 + |\nabla f|^2} dt^1 \dots dt^{n-1}. \quad (81)$$

44 Application to the Green Theorem

In Riemannian Geometry the divergence of a vector $\mathbf{b}$ is given by

$$\nabla \cdot \mathbf{b} = \frac{1}{\sqrt{g}} \frac{\partial}{\partial x^i} (\sqrt{g} b^i) \text{ summation convention}$$

Let us form the integral of the divergence of $\mathbf{b}$ over a region R in V_n bounded by the closed surface B_{n-1}

$$\int_R \nabla \cdot \mathbf{b} dV = \int_R \frac{1}{\sqrt{g}} \frac{\partial}{\partial x^i} (\sqrt{g} b^i) dV = \int_R \frac{1}{\sqrt{g}} \frac{\partial}{\partial x^i} (\sqrt{g} b^i) \sqrt{g} dx^1 \dots dx^n \quad (82)$$

Now let us integrate the differentiated terms over the range of the appropriate variable for each one. We obtain an integral over the boundary surface B_{n-1} : for the k term this is (k not summed)

$$\int_B \sqrt{g} b^k dx^1 \dots dx^{k-1} dx^{k+1} \dots dx^n = \int_B \sqrt{g} b^k D_k dt^1 \dots dt^{n-1}, \quad (83)$$

where D_k is given as in (68) represents the transformation of the integration variables from the x^i to the coordinates as given on B_{n-1} .

From the above set of equations we see that

$$\begin{aligned} \int_{R_n} \nabla \cdot \mathbf{b} dV_n &= \int_{B_{n-1}} b^k D_k \sqrt{g} dt^1 \dots dt^{n-1} \\ &= \int_{B_{n-1}} b^k n_k \sqrt{g} D dt^1 \dots dt^{n-1} \\ &= \int_{B_{n-1}} \mathbf{b} \cdot \mathbf{n} dS, \end{aligned}$$

where dS is given by (77).

Theorem 44.1 (Divergence Theorem)

$$\iiint_V A^k_{,k} dV = \iint_S A^k \nu_k dS$$

45 Extension and Stokes' Theorem

In a space of n dimensions, the **Generalized Kronecker delta** $\delta^{k_1 k_2 \dots k_m}_{s_1 s_2 \dots s_m}$ for any positive integer m is defined as

$$\delta^{k_1 k_2 \dots k_m}_{s_1 s_2 \dots s_m} = +1 \text{ if } k_1 k_2 \dots k_m \text{ are distinct integers selected from the range } 1, 2, \dots, n, \\ \text{and if } s_1, s_2, \dots, s_m \text{ is an } \textit{even} \text{ permutation of } k_1 k_2 \dots k_m.$$

$$\delta^{k_1 k_2 \dots k_m}_{s_1 s_2 \dots s_m} = -1 \text{ if } k_1 k_2 \dots k_m \text{ are distinct integers selected from the range } 1, 2, \dots, n, \\ \text{and if } s_1, s_2, \dots, s_m \text{ is an } \textit{odd} \text{ permutation of } k_1 k_2 \dots k_m.$$

$\delta_{s_1 s_2 \dots s_m}^{k_1 k_2 \dots k_m} = 0$ if any two $k_1 k_2 \dots k_m$ are, equal, or if any two $s_1, s_2, \dots, s_m$ are equal, or if the set of numbers $\{k_1, k_2, \dots, k_m\}$ differs, apart from order, from the set $\{s_1, s_2, \dots, s_m\}$.

In a space of n dimensions, in which no metric is assigned, consider a subspace V_m of m dimensions $m \leq n$ defined by the parametric equations

$$x^k = x^k(y^1, y^2, \dots, y^m), \quad k = 1, 2, \dots, n.$$

In this V_m consider a certain region of R_m . Let us divide R_m into cells by m families of surfaces

$$f^\alpha(y) = c^\alpha, \quad \alpha = 1, 2, \dots, m.$$

where c^α are constants, each taking on a number of discrete values and so forming a family of surfaces.

It is easy to attach a meaning to the statement that a point P of V_m lies in a certain cell, because P may be said to lie between two surfaces of the same family, say family α , if the expression $f^\alpha(y) - c^\alpha$ changes sign when we change c^α from the value belonging to one of these surfaces to the value belonging to the other. Here y refers to the parameter values at the point P .

Now pick one of the cells and let A be the point corresponding to one of the corners in the cell. Through A there passes one surface of each family. Let the corresponding constants be $c^1, c^2, \dots, c^m$. Passing along the edge for which c^1 alone changes, we arrive at another corner B_1 for which the constants have values $c^1 + \Delta c^1, c^2, \dots, c^m$. In passing from A to B_1 the parameters change from y^α at A to $y^\alpha + \Delta_1 y^\alpha$ at B_1 and the coordinates from x^k to $x^k + \Delta_1 x^k$. Similarly the parameter values and the coordinates of all corners $B_1, B_2, \dots, B_m$ of the cell reached by traveling from A along an edge of the cell can be written as

$$\begin{aligned} y^\alpha + \Delta_\beta y^\alpha, \\ x^\alpha + \Delta_\beta x^\alpha. \end{aligned}$$

and we form the determinants

$$\Delta^{k_1 \dots k_m} = \begin{vmatrix} \Delta_1 x^{k_1} & \Delta_1 x^{k_2} & \dots & \Delta_1 x^{k_m} \\ \Delta_2 x^{k_1} & \Delta_2 x^{k_2} & \dots & \Delta_2 x^{k_m} \\ \dots & \dots & \dots & \dots \\ \Delta_m x^{k_1} & \Delta_m x^{k_2} & \dots & \Delta_m x^{k_m} \end{vmatrix} \quad (84)$$

Introducing the generalized Kronecker delta $\delta_{s_1 s_2 \dots s_m}^{k_1 k_2 \dots k_m}$, these determinants can be written compactly using the Summation Convention

$$\Delta^{k_1 \dots k_m} = \delta_{s_1 s_2 \dots s_m}^{k_1 k_2 \dots k_m} \Delta_1 x^{s_1} \Delta_2 x^{s_2} \dots \Delta_m x^{s_m} \quad (85)$$

Now if the edges had been taken in a different order, then the various determinants might have opposite signs, but apart from this the determinants are independent of the order taken. Also if we have used a different corner to start from the determinants formed similarly would have different values. But since we are going to be working with infinitesimal, the differences will be of higher order than the determinants in (85). Using differential displacements, $d_\beta x^i$ (where β indexes the β -th edge of the cell) we have

$$d\tau_m^{k_1 \dots k_m} = \delta_{s_1 s_2 \dots s_m}^{k_1 k_2 \dots k_m} d_1 x^{s_1} d_2 x^{s_2} \dots d_m x^{s_m} \quad (86)$$

The set of quantities $d\tau_m^{k_1 \dots k_m}$ is called the *extension* of the infinitesimal m -cell. No metrical concepts have entered

into the definition of extension. Extension is the nearest concept closest to the idea of elementary volume one can define in a non-metrical space.

The edges of the infinitesimal m -cell can be written as

$$d_\beta x^k = \frac{\partial x^k}{\partial y^\alpha} d_\beta y^\alpha \quad (87)$$

Theorem 45.1 (Stoke's Theorem)

$$\oint_C A_p \frac{dx^p}{ds} ds = - \iint_S \epsilon^{pqr} A_{q,r} \nu_p dS,$$

where $\frac{dx^p}{ds}$ is the unit tangent vector to the closed curve C and ν_p is the positive unit normal to the surface S which has C as its boundary.

46 Applications In Mechanics

Definition Lie Algebras. Here $F = \mathbb{R}$ or $\mathbb{C}$. A Lie Algebra of F is a pair $(\mathfrak{g}, [\cdot, \cdot])$, where $\mathfrak{g}$ is a vector space over F and

$$[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$$

is an F -bilinear map satisfying the following properties

$$\begin{aligned} X, Y] &= -[Y, X] \\ [X, [Y, Z]] + [Z, [X, Y]] + [Y, [Z, X]] &= 0, \end{aligned}$$

The later is the Jacobi identity.

Definition Structure Constants: Given the elements X_a, X_b and X_c in a Lie Algebra we define the Structure Constraints c_{ab}^c to be the coefficients in the expansion of the Lie Bracket:

$$[X_a, X_b] = \sum_c c_{ab}^c X_c.$$

References

- [1] F.M. Hall, “Introduction to Abstract Algebra (Vol 2)”, Cambridge University Press (1969).
- [2] Garrett Birkoff and Saunders MacLane, “A Survey of Modern Algebra”, The MacMillan Company, New York (1957)..
- [3] O. Schreier and F. Sperner, “Introduction to Modern Algebra and Matrix Theory”, Second Edition, Chelsea Publishing (1959).
- [4] William Snow Burnside and Arthur William Panton, “The Theory of Equations with an Introduction to the Theory of Binary Algebraic Forms, Volume 2”, Dover Publications (1960).
- [5] Jim Brown, “Abel and the Insolvability of the Quintic”, Mathematics Department The Ohio State University (2013).
- [6] Peter Pesic, “Abel’s Proof”, The MIT Press, Cambridge, Massachusetts (2003).
- [7] Edgar Dehn, “Algebraic Equations : An Introduction to the Theories of Lagrange and Galois”, Dover Publications (1960).
- [8] Murray R. Spiegel, “Vector Analysis and an Introduction to Tensor Analysis”, Schaum’s Outline Series, McGraw-Hill Book Company (1960).
- [9] M.H. Protter and C.B. Morrey, “A First Course in Real Analysis”, Springer-Verlag (1977).
- [10] Peter Gabriel Bergmann, “Introduction to the Theory of Relativity”, Dover (1976).