

MATH681  
Mathematical Logic

Pingbang Hu

March 21, 2023

## Abstract

This is a graduate-level mathematical logic course taught by [Matthew Harrison-Trainor](#), aiming to obtain insights into all other branches of mathematics, such as algebraic geometry, analysis, etc. Specifically, we will cover model theory beyond the basic foundational ideas of logic.

While there are no required textbooks, some books do cover part of the material in the class. For example, Marker's *Model Theory: An Introduction* [[Mar02](#)], Hodges's *A Shorter Model Theory* [[HH97](#)], and Hinman's *Fundamentals of Mathematical Logic* [[Hin05](#)].



# Contents

<b>1</b>	<b>Language, Logic, and Structures</b>	<b>2</b>
1.1	Languages and Structures . . . . .	3
1.2	Embeddings and Isomorphisms . . . . .	4
1.3	Terms . . . . .	5
1.4	Formulas . . . . .	6
1.5	Truths . . . . .	7
<b>2</b>	<b>Soundness, Completeness, and Compactness</b>	<b>12</b>
2.1	Theories . . . . .	12
2.2	Elementary Embeddings . . . . .	13
2.3	Definable Sets . . . . .	14
2.4	Proofs . . . . .	15
2.5	Soundness Theorem . . . . .	18
2.6	Completeness and Compactness Theorems . . . . .	21
<b>3</b>	<b>The Beginning of Model Theory</b>	<b>29</b>
3.1	Complete Theories . . . . .	29
3.2	A Detour to Algebraically Closed Fields . . . . .	29
3.3	The ACF Theory . . . . .	36
3.4	Up and Down . . . . .	38
3.5	Back and Forth . . . . .	43
<b>4</b>	<b>Quantifier Elimination and Algebraic Applications</b>	<b>47</b>
4.1	Quantifier Elimination . . . . .	47
4.2	Ultrapowers . . . . .	56

# Chapter 1

## Language, Logic, and Structures

### Lecture 1: Introduction to Mathematical Logic

The goal of mathematical logic is to obtain insights into other areas of mathematics – algebra, analysis, combinatorics, and so on, by formalizing the **process** of mathematics. 5 Jan. 14:30

**Remark.** More concretely, there are different branches:

- (a) Model Theory: Study subsets of an object defined by a **formula** (i.e., first-order logic).
- (b) Computability Theory / Recursion Theory: Formalizing what it means to have an algorithm and studying relative computability.
- (c) Set Theory: Study the structure of the mathematical universe.
- (d) Proof Theory: Study the syntactic nature of **proofs**.

In this class, we study model theory in nature; specifically, we will cover

- basic definitions of logic:
  - What is a **formula**?
  - What does it mean for a **formula** to be **true**?
  - What is a **proof**?
- **Soundness** & **completeness theorems**:
  - Anything **provable** is **true**.
  - Anything **true** is **provable**.
- Compactness theorem:
  - Non-standard objects exist.
- Using compactness theorem for applications:
  - **Chevalley's theorem**.

The main theme of this course will be *syntax* v.s. *semantics*:

Syntax	v.s.	Semantics
<b>proofs</b>		<b>truth</b>
form of a <b>formula</b>		mathematical <b>structures</b>
number and type of quantifiers		<b>isomorphisms, embeddings</b>

And this is what this chapter aiming to address. We will turn to other topics based on these.

## 1.1 Languages and Structures

Let's start with the fundamental object, [language](#).

**Definition 1.1.1 (Language).** A *language*  $\mathcal{L}$  consists of:

- a set  $\mathcal{F}$  of function symbols  $f$  with arities  $n_f$ ;
- a set  $\mathcal{R}$  of relation symbols  $R$  with arities  $n_R$ ;
- a set  $\mathcal{C}$  of constant symbols  $c$ .

A [language](#) is also sometimes called a *signature*, in which case we use  $\sigma$  rather than  $\mathcal{L}$ .

**Note.** A constant is the same as a 0-ary function.

**Remark.** Any or all sets in [Definition 1.1.1](#) might be empty.

**Example (Graph).** The [language](#) of graphs,  $\mathcal{L}_{\text{graph}} = \{E\}$  where  $E$  is a binary (2-ary) relation symbol.

**Example (Ring).** The [language](#) of rings,  $\mathcal{L}_{\text{ring}} = \{0, 1, +, \cdot, -\}$ , where  $0, 1$  are constants,  $+, \cdot$  are binary functions, and  $-$  is a unary function.

**Example (Ordered ring).** The [language](#) of ordered rings,  $\mathcal{L}_{\text{ord}} = \mathcal{L}_{\text{ring}} \cup \{\leq\}$  where  $\leq$  is the binary relation for an ordered ring.

Then, given a [language](#), we can now interpret it in the following way.

**Definition 1.1.2 (Structure).** Given a [language](#)  $\mathcal{L}$ , an  $\mathcal{L}$ -*structure*  $\mathcal{M}$  consists of:

- a non-empty set  $M$  called the *universe*, *domain*, or *underlying set* of  $\mathcal{M}$ ;
- for each function symbol  $f \in \mathcal{F}$ , a function  $f^{\mathcal{M}}: M^{n_f} \rightarrow M$ ;
- for each relation symbol  $R \in \mathcal{R}$ , a relation  $R^{\mathcal{M}} \subseteq M^{n_R}$ ;
- for each constant symbol  $c \in \mathcal{C}$ , an element  $c^{\mathcal{M}} \in M$ .

**Notation (Interpretation).** The *interpretation* of symbols  $f, R, c$  in  $\mathcal{M}$  is  $f^{\mathcal{M}}, R^{\mathcal{M}}, c^{\mathcal{M}}$ , respectively.

Basically, a [structure](#) gives meaning to the symbols from the [language](#), and we often write

$$\mathcal{M} = (M, f^{\mathcal{M}}, \dots, R^{\mathcal{M}}, \dots, c^{\mathcal{M}}, \dots) = (M, f^{\mathcal{M}}, R^{\mathcal{M}}, c^{\mathcal{M}}: f \in \mathcal{F}, R \in \mathcal{R}, c \in \mathcal{C}).$$

**Notation.** We usually use  $\mathcal{M}, \mathcal{N}, \dots, \mathcal{A}, \mathcal{B}, \dots$  to refer to [structures](#), and  $M, N, \dots, A, B, \dots$  for the domains.<sup>a</sup>

<sup>a</sup>Some people use  $|\mathcal{M}|$  for the domain of  $\mathcal{M}$ .

It's time to look at some examples.

**Example.** The rationals  $\mathbb{Q}$  and integers  $\mathbb{Z}$  are both  $\mathcal{L}_{\text{ring}}$ -structures.

**Proof.** Clearly, the domain is the set of rationals, and naively, we let  $+^{\mathbb{Q}} = +$  in  $\mathbb{Q}$ ,  $0^{\mathbb{Q}} = 0$  in  $\mathbb{Q}$ ,  $1^{\mathbb{Q}} = 1$  in  $\mathbb{Q}$ , etc. In this way,  $\mathbb{Q} = (\mathbb{Q}, 0, 1, +, \cdot, -)$  is an  $\mathcal{L}_{\text{ring}}$ -structure. Similarly,  $\mathbb{Z} = (\mathbb{Z}, 0, 1, +, \cdot, -)$  is as well. \*

While the [language](#) we have seen are all intuitively correct with their name, e.g.,  $\mathcal{L}_{\text{ring}}$ ,  $\mathcal{L}_{\text{ord}}$ , and  $\mathcal{L}_{\text{graph}}$ , they are really just the high-level abstraction of the objects in the subscript.

**Example.** Nothing forces an  $\mathcal{L}_{\text{ring}}$ -structure to be a ring.

**Proof.** Since an  $\mathcal{L}_{\text{ring}}$ -structure is just any [structure](#) with two binary functions, a unary function, and two constants interpreting the symbols of the [language](#); hence we can define an  $\mathcal{L}_{\text{ring}}$ -structure  $\mathcal{M}$  as

- $\mathcal{M} = \{0, 5, 11\}$ ;
- $0^{\mathcal{M}} = 5$ ;
- $1^{\mathcal{M}} = 11$ ;
- $+^{\mathcal{M}}$  is the constant function 0;
- $\cdot^{\mathcal{M}}$  is the function 5;
- $-^{\mathcal{M}}$  is the identity.

This is clearly not a ring since it fails nearly every axiom of a ring. \*

**Note.** Later, we will talk about theories that let us restrict to [structures](#) we want.

## 1.2 Embeddings and Isomorphisms

We can now consider the relation between [structures](#).

**Definition 1.2.1 (Embedding).** Let  $\mathcal{M}$  and  $\mathcal{N}$  be  $\mathcal{L}$ -structures. A map  $\eta: \mathcal{M} \rightarrow \mathcal{N}$  is an  $\mathcal{L}$ -embedding if it is one-to-one and preserves the [interpretation](#) of all symbols of  $\mathcal{L}$ :

- (a) for each function symbol  $f \in \mathcal{F}$  of arity  $n_f$ , and  $a_1, \dots, a_{n_f} \in M$ ,

$$\eta(f^{\mathcal{M}}(a_1, \dots, a_{n_f})) = f^{\mathcal{N}}(\eta(a_1), \dots, \eta(a_{n_f}));$$

- (b) for each relation symbol  $R \in \mathcal{R}$  of arity  $n_R$ , and  $a_1, \dots, a_{n_R} \in M$ ,

$$(a_1, \dots, a_{n_R}) \in R^{\mathcal{M}} \Leftrightarrow (\eta(a_1), \dots, \eta(a_{n_R})) \in R^{\mathcal{N}};$$

- (c) for each constant symbol  $c \in \mathcal{C}$ ,  $c^{\mathcal{M}} = c^{\mathcal{N}}$ .

From the definition, an  $\mathcal{L}$ -embedding is an injection, and naturally, we have the following.

**Definition 1.2.2 (Isomorphism).** An  $\mathcal{L}$ -isomorphism is a bijective  $\mathcal{L}$ -embedding.

**Definition 1.2.3 (Automorphism).** An  $\mathcal{L}$ -automorphism of  $\mathcal{M}$  is an  $\mathcal{L}$ -isomorphism from  $\mathcal{M}$  to  $\mathcal{M}$ .

**Definition.** Let  $\mathcal{M}$  and  $\mathcal{N}$  be  $\mathcal{L}$ -structures. Suppose  $M \subseteq N$  and the inclusion map  $\iota: \mathcal{M} \hookrightarrow \mathcal{N}$  is an  $\mathcal{L}$ -embedding.

**Definition 1.2.4 (Substructure).**  $\mathcal{M}$  is a *substructure* of  $\mathcal{N}$ .

**Definition 1.2.5 (Extension).**  $\mathcal{N}$  is an *extension* of  $\mathcal{M}$ .

**Example.** Ring embeddings are  $\mathcal{L}_{\text{ring}}$ -embeddings.

This generalizes the notions of embedding and isomorphism for many mathematical structures.

**Remark.** Asking that  $\eta$  be injective is the same as (b) in Definition 1.2.1 for the relation  $=$  since

$$a = b \in \mathcal{M} \Leftrightarrow \eta(a) = \eta(b) \in \mathcal{N}.$$

The notion of substructure is language sensitive. For groups, there are two possible languages:

- (a)  $\mathcal{L}_1 = \{e, \cdot\}$ ;
- (b)  $\mathcal{L}_2 = \{e, \cdot, {}^{-1}\}$ , i.e., with the unary inverse operation.

While both seem valid at the first glance, we should use the second one.

To see why, if we use  $\mathcal{L}_2$ , the substructure of a group is the same thing as a subgroup. But if we use  $\mathcal{L}_1$ , then  $(\mathbb{N}, +, 0)$  is a substructure of  $(\mathbb{Z}, +, 0)$ , while  $\mathbb{N}$  is not a group for sure.<sup>1</sup>

Similarly, we include  $-$  in  $\mathcal{L}_{\text{ring}}$  for a similar reason as in the previous example.

**Example.** An  $\mathcal{L}_{\text{ring}}$ -substructure of a field will be a subring, not a subfield. If we want subfields, use  $\mathcal{L}_{\text{ring}} \cup \{{}^{-1}\}$ .<sup>a</sup>

<sup>a</sup>We can set  $0^{-1} = 0$ , but never use this.

## Lecture 2: Formulas and First-Order Logic

We start by asking that given a function symbol  $f$  of arity  $n$ , could we replace  $f$  with an  $(n+1)$ -ary  $R$  relation to represent its graph? 10 Jan. 14:30

**Example.** Let  $\mathcal{L}$  be a language with only relation symbols. Let  $\mathcal{A}$  be an  $\mathcal{L}$ -structure. For any  $B \subseteq A$ , there is a substructure  $\mathcal{B}$  of  $\mathcal{A}$  with domain  $B$ .

**Proof.** For each relation symbol  $R$ , letting  $R^{\mathcal{B}} = R^{\mathcal{A}} \cap B^{n_R}$  will make  $\mathcal{B}$  a substructure of  $\mathcal{A}$ .  $\circledast$

The above is not true for function symbols though.

**Example.** If  $G = (\mathbb{Z}, 0, +)$ , then  $\mathbb{N}$  is not the domain of a subgroup. So if we took  $\mathcal{L} = \{0, +, {}^{-1}\}$ , where  $0$  is the unary relation,  $+$  is the ternary relation, and  ${}^{-1}$  is the binary relation, an  $\mathcal{L}$ -substructure of a group might not be a subgroup.

### 1.3 Terms

Intuitive, an  $\mathcal{L}$ -formula is an expression built using the symbols in a language  $\mathcal{L}$ ,  $=$ , the logical connectives  $\wedge, \vee, \neg$ , and variable symbols  $v_1, v_2, \dots, x, y, z$ , and also quantifiers  $\exists$  and  $\forall$ .

**Definition 1.3.1 (Term).** Given a language  $\mathcal{L}$ , the set of  $\mathcal{L}$ -terms are defined inductively by:

- (a) each constant symbol is a term;
- (b) each variable symbol  $v_1, \dots$  is a term;
- (c) if  $f$  is a function symbol, and  $t_1, \dots, t_{n_f}$  are terms, then  $f(t_1, \dots, t_{n_f})$  is a term.

If  $\mathcal{M}$  is an  $\mathcal{L}$ -structure, and  $t$  is a term involving only variables among  $v_1, \dots, v_n$ , then  $t$  has an interpretation  $t^{\mathcal{M}}: M^n \rightarrow M$  as a function as follows. On input  $a_1, \dots, a_n \in M$ ,

- (a) if  $t$  is a constant  $c$ ,  $t^{\mathcal{M}}(a_1, \dots, a_n) = c^{\mathcal{M}}$ .

<sup>1</sup>Simply observe that both  $(\mathbb{N}, 0, +), (\mathbb{Z}, 0, +)$  are  $\mathcal{L}_1$ -structures.

- (b) if  $t$  is a variable  $v_i$ ,  $t^{\mathcal{M}}(a_1, \dots, a_n) = v_i$ ;  
 (c) if  $t$  is  $f(s_1, \dots, s_k)$ , then  $t^{\mathcal{M}}(a_1, \dots, a_n) = f^{\mathcal{M}}(s_1^{\mathcal{M}}(a_1, \dots, a_n), \dots, s_k^{\mathcal{M}}(a_1, \dots, a_n))$ .

**Intuition.** We are basically substituting for variables and evaluating the expression.

**Example.** In  $(\mathbb{R}, 0, 1, +, \cdot, -)$ , a **term** is essentially just a polynomial with integer coefficients, assuming we interpret them in a ring. Technically, a **term** looks like

$$\cdot(+ (1, 1), + (x, y)),$$

but we will write **terms** the natural way, i.e.,

$$(1 + 1)(x + y).$$

Also, we will use  $\underline{n}$  or  $n$  to represent the **term**  $\underline{n} = \underbrace{1 + 1 + \dots + 1}_{n \text{ times}}$ . So we could write the above **term** as  $2 \cdot (x + y)$ .

## 1.4 Formulas

A **term** is just a building block of **formulas**, as we now see.

**Definition 1.4.1 (Formula).** The set of  $\mathcal{L}$ -formulas is defined inductively:

- (a) If  $s, t$  are **terms**, then  $s = t$  is a *formula*.
- (b) If  $R$  is a relation symbol of arity  $n_R$  and  $s_1, \dots, s_{n_R}$  are **terms**, then  $R(s_1, \dots, s_{n_R})$  is a *formula*.
- (c) If  $f$  is a **formula**, then  $\neg f$  is a *formula*.
- (d) If  $\varphi$  and  $\psi$  are **formulas**, then  $\varphi \wedge \psi$  and  $\varphi \vee \psi$  are *formulas*.
- (e) If  $\varphi$  is a **formula** and  $v_i$  are variables, then  $\exists v_i \varphi$  and  $\forall v_i \varphi$  are *formulas*.

**Notation (Atomic).** Definition 1.4.1 (a) and (b) are called *atomic*.

**Notation (Quantifier-free).** Definition 1.4.1 (a), (b), (c), and (d) are called *quantifier-free*.

This logic is called *first-order logic* (FO logic), since the quantifiers range over elements of the **structures**, but not over, e.g., subsets.

**Example.** We can say that an element  $x$  of a ring has a square root by  $\exists y \ y^2 = x$ .

**Example.** A group is torsion of order 2 can be said by  $\forall x \ x \cdot x = e$ .

**Example.** We can write down all the field/group/... axioms as **formulas**.

### 1.4.1 Bounded and Free Variables

Notice that for the first example, the **formula**  $\exists y \ y^2 = x$  only has meaning if we assign what  $x$  is. In this case, we say that  $y$  is *bound* by  $\exists y$ . But this is local:



**Example.** Consider

$$y = 1 \wedge \exists y \, y^2 = x,$$

while the first appearance of  $y$  is free, the second appearance of  $y$  is bound by (in the scope of)  $\exists y$ .

While our definitions work perfectly fine with the above example, but sometimes we don't want this to happen. In such a case, we simply replace the bound instances of  $y$  with a new variable  $z$ . This idea of variables being free or bound is defined formally as follows.

**Definition 1.4.2 (Free variable).** The *free variables*  $\text{FV}(\varphi)$  of a *formula*  $\varphi$  are defined inductively:

- (a)  $\text{FV}(s = t)$  is the set of variables showing up in  $s$  or  $t$ .
- (b)  $\text{FV}(R(s_1, \dots, s_{n_R}))$  is the set of variables showing up in  $s_1, \dots, s_{n_R}$ .
- (c)  $\text{FV}(\neg\varphi) = \text{FV}(\varphi)$ .
- (d)  $\text{FV}(\varphi \wedge \psi) = \text{FV}(\varphi \vee \psi) = \text{FV}(\varphi) \cup \text{FV}(\psi)$ .
- (e)  $\text{FV}(\exists x \, \varphi) = \text{FV}(\forall x \, \varphi) = \text{FV}(\varphi) \setminus \{x\}$ .

**Example.**  $\text{FV}(\exists y \, y^2 = x) = \{x\}$ .

**Example.**  $\text{FV}(\forall x \, x \cdot x = e) = \emptyset$ .

**Definition 1.4.3 (Sentence).** A *formula*  $\varphi$  is called a *sentence* if it has no *free variables*.

**Notation.** If  $\varphi$  is a *formula* with *free variables* among  $x_1, \dots, x_n$  we often write  $\varphi(x_1, \dots, x_n)$ .

**Remark.** So given  $\varphi(x_1, \dots, x_n)$ , we know that  $\varphi$  has no other *free variables* than  $x_1, \dots, x_n$ .

**Example.** It's valid to write  $\varphi(x, y, z) := x = y$ .

## 1.5 Truths

Finally, we define the notion of *truth*.

**Definition 1.5.1 (Truth).** Given an  $\mathcal{L}$ -structure  $\mathcal{M}$ , let  $\varphi(x_1, \dots, x_n)$  be an  $\mathcal{L}$ -formula and let  $a_1, \dots, a_n \in M$ . Then we say  $\varphi$  is *true* of  $\bar{a}$  in  $\mathcal{M}$ ,<sup>a</sup> denoted as  $\mathcal{M} \models \varphi(\bar{a})$ , as follows:

- (a) If  $\varphi$  is  $s = t$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if  $s^{\mathcal{M}}(\bar{a}) = t^{\mathcal{M}}(\bar{a})$ .
- (b) If  $\varphi$  is  $R(t_1, \dots, t_{n_R})$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if  $(t_1^{\mathcal{M}}(\bar{a}), \dots, t_{n_R}^{\mathcal{M}}(\bar{a})) \in R^{\mathcal{M}}$ .
- (c) If  $\varphi$  is  $\neg\psi$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if  $\mathcal{M} \not\models \psi(\bar{a})$ .
- (d) If  $\varphi$  is  $\psi_1 \wedge \psi_2$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if  $\mathcal{M} \models \psi_1(\bar{a})$  and  $\mathcal{M} \models \psi_2(\bar{a})$ .
- (e) If  $\varphi$  is  $\psi_1 \vee \psi_2$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if  $\mathcal{M} \models \psi_1(\bar{a})$  or  $\mathcal{M} \models \psi_2(\bar{a})$ .
- (f) If  $\varphi$  is  $\exists y \, \psi(\bar{x}, y)$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if there's  $b \in M$  such that  $\mathcal{M} \models \psi(\bar{a}, b)$ .
- (g) If  $\varphi$  is  $\forall y \, \psi(\bar{x}, y)$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if for all  $b \in M$  such that  $\mathcal{M} \models \psi(\bar{a}, b)$ .

<sup>a</sup>Or  $\mathcal{M}$  satisfies  $\varphi(\bar{a})$ .

**Remark.** Every formula is true, or its negation is.

## Lecture 3: Logical Consequence and Equivalence

### 1.5.1 Implications

12 Jan. 14:30

**Notation** (Material implication). The *material implication*  $\varphi \rightarrow \psi$  between two formulas  $\varphi, \psi$  is an abbreviation of  $\neg\varphi \vee \psi$ .

**Notation.** We use  $\varphi \leftrightarrow \psi$  as an abbreviation of  $((\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi))$ .

Essentially,  $\rightarrow$  and  $\leftrightarrow$  is different from  $\Rightarrow$  and  $\Leftrightarrow$ , where the former are only shown in formula. Now, consider the language of graphs  $\mathcal{L}_{\text{graph}} = \{E\}$ , let's see some examples.

**Example.** An undirected graph can be written as

$$\forall x \forall y (xEy \rightarrow yEx).$$

**Example.** A vertex has at least three neighbors can be written as

$$\varphi(x) := \exists u \exists v \exists w (xEu \wedge xEv \wedge xEw \wedge u \neq v \wedge v \neq w \wedge u \neq w)$$

in non-reflexive graphs.

**Example.** For a vertex has exactly three neighbors,

$$\psi(x) := \exists u \exists v \exists w \forall y (xEu \wedge xEv \wedge xEw \wedge u \neq v \wedge v \neq w \wedge u \neq w \wedge (y = u \vee y = v \vee y = w \vee \neg yEx)).$$

**Problem.** Can we say that  $x$  has an even number of neighbors?

**Answer.** We can't. Some things are not expressible in FO logic. ⊗

**Example.** For a vertex  $x$  has a path of length 4 to  $y$ ,

$$\Theta(x, y) := \exists u \exists v \exists w (xEu \wedge uEv \wedge vEw \wedge wEy).$$

We can also express that there is a path of length at most 4.

**Problem.** Can we say that there is a path from  $x$  to  $y$ ?

**Answer.** We still can't! Not in FO logic (using compactness theorem). ⊗

**Remark.** When we prove results by induction on formulas, we only need to prove for  $\neg, \wedge, \exists$ , instead of for both  $\wedge, \vee$ , and both  $\exists$  and  $\forall$ .

**Proof.** Since we can view  $\varphi \vee \psi$  as an abbreviation for  $\neg(\neg\varphi \wedge \neg\psi)$  and  $\forall x \varphi$  as an abbreviation for  $\neg(\exists x \neg\varphi)$ . ⊗

**Remark (Sheffer stroke).** In fact, we can get  $\wedge, \vee, \neg$  from one logical connective, e.g., the *sheffer stroke*  $\uparrow$ , which is defined as

$$\varphi \uparrow \psi := \neg(\varphi \wedge \psi),$$

and we can use  $\uparrow$  to define  $\neg, \vee, \wedge$ .

**Notation.** Let  $\Phi$  be a (possibly infinite) set of **sentences**, we write  $\mathcal{M} \models \Phi$  if  $\mathcal{M} \models \varphi$  for all  $\varphi \in \Phi$ .

### 1.5.2 Logical Consequences and Equivalent

**Definition 1.5.2** (Logical consequence). Let  $\Phi$  be a set of **sentences**, and  $\varphi$  be a **sentence**. We say that  $\varphi$  is a *logical consequence* of  $\Phi$ , written  $\Phi \models \varphi$ , if  $\mathcal{M} \models \varphi$  whenever  $\mathcal{M} \models \Phi$ .

If  $\Phi = \emptyset$  is the empty set, **Definition 1.5.2** is written as  $\models \varphi$ , i.e.,  $\varphi$  is **true** in all  $\mathcal{L}$ -structures.<sup>2</sup>

**Definition 1.5.3** (Equivalent). Given two **formulas**  $\varphi, \psi$ ,  $\varphi(\bar{x})$  and  $\psi(\bar{x})$  are *equivalent* if

$$\models \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x})).$$

**Problem.** Two **sentences**  $\varphi$  and  $\psi$  are **equivalent** if and only if  $\varphi \models \psi$  and  $\psi \models \varphi$ .

DIY

**As previously seen.**  $\mathcal{A}$  is a **substructure** of  $\mathcal{B}$ , or  $\mathcal{A} \subseteq \mathcal{B}$ , means that  $A \subseteq B$  and  $\text{id}: A \hookrightarrow B$  is an  $\mathcal{L}$ -embedding.

**Proposition 1.5.1.** Suppose that  $\mathcal{A}$  is a **substructure** of  $\mathcal{B}$ , and  $\varphi(\bar{x})$  is a **quantifier-free formula**. Let  $\bar{a} \in \mathcal{A}$ ,<sup>a</sup> then  $\mathcal{A} \models \varphi(\bar{a})$  if and only if  $\mathcal{B} \models \varphi(\bar{a})$ .

<sup>a</sup>Formally, we need to write  $\mathcal{A}$  to be the Cartesian product with a fixed length.

**Proof.** We start with **terms** by proving that if  $t$  is a **term** and  $\bar{b} \in \mathcal{A}$ , then  $t^{\mathcal{A}}(\bar{b}) = t^{\mathcal{B}}(\bar{b})$ . The proof is induction on **terms**.

- (a) If  $t$  is a constant symbol  $c$ , then  $t^{\mathcal{A}}(\bar{b}) = c^{\mathcal{A}} = c^{\mathcal{B}} = t^{\mathcal{B}}(\bar{b})$ .
- (b) If  $t$  is a variable  $x_i$ , then  $t^{\mathcal{A}}(\bar{b}) = b_i = t^{\mathcal{B}}(\bar{b})$ .
- (c) If  $t$  is a function symbol  $f(s_1, \dots, s_n)$  where  $s_i$  are **terms**, then  $t^{\mathcal{A}}(\bar{b}) = f^{\mathcal{A}}(s_1^{\mathcal{A}}(\bar{b}), \dots, s_n^{\mathcal{A}}(\bar{b}))$ .  
By the induction hypothesis,  $s_i^{\mathcal{A}}(\bar{b}) = s_i^{\mathcal{B}}(\bar{b}) \in \mathcal{A}$ , and hence

$$t^{\mathcal{B}}(\bar{b}) = f^{\mathcal{B}}(s_1^{\mathcal{B}}(\bar{b}), \dots, s_n^{\mathcal{B}}(\bar{b})) = f^{\mathcal{A}}(s_1^{\mathcal{A}}(\bar{b}), \dots, s_n^{\mathcal{A}}(\bar{b})) = t^{\mathcal{A}}(\bar{b}),$$

i.e.,  $f^{\mathcal{B}} \upharpoonright_{\mathcal{A}} = f^{\mathcal{A}}$ , so  $t^{\mathcal{A}}(\bar{b}) = t^{\mathcal{B}}(\bar{b})$ .

Now we turn to **formulas**, and prove that for  $\varphi$  **quantifier-free**, then  $\mathcal{A} \models \varphi(\bar{a}) \Leftrightarrow \mathcal{B} \models \varphi(\bar{a})$  for  $\bar{a} \in \mathcal{A}$ . The proof is, again, induction on **formulas**.<sup>a</sup>

- (a) If  $\varphi$  is  $s = t$ , then  $s^{\mathcal{A}}(\bar{a}) = s^{\mathcal{B}}(\bar{a})$  and  $t^{\mathcal{A}}(\bar{a}) = t^{\mathcal{B}}(\bar{a})$ , so

$$\mathcal{A} \models \varphi(\bar{a}) \Leftrightarrow s^{\mathcal{A}}(\bar{a}) = t^{\mathcal{A}}(\bar{a}) \Leftrightarrow s^{\mathcal{B}}(\bar{a}) = t^{\mathcal{B}}(\bar{a}) \Leftrightarrow \mathcal{B} \models \varphi(\bar{a}).$$

- (b) If  $\varphi$  is  $R(s_1, \dots, s_n)$ , then

$$\mathcal{A} \models \varphi(\bar{a}) \Leftrightarrow (s_1^{\mathcal{A}}(\bar{a}), \dots, s_n^{\mathcal{A}}(\bar{a})) \in R^{\mathcal{A}} \Leftrightarrow (s_1^{\mathcal{B}}(\bar{a}), \dots, s_n^{\mathcal{B}}(\bar{a})) \in R^{\mathcal{B}} \Leftrightarrow \mathcal{B} \models \varphi(\bar{a}).$$

- (c) If  $\varphi$  is  $\neg\psi$ ,

$$\mathcal{A} \models \varphi(\bar{a}) \Leftrightarrow \mathcal{A} \not\models \psi(\bar{a}) \Leftrightarrow \mathcal{B} \not\models \psi(\bar{a}) \Leftrightarrow \mathcal{B} \models \varphi(\bar{a}),$$

where we use the induction hypothesis in the second  $\Leftrightarrow$ .

<sup>2</sup>Recall that we always have a **language**  $\mathcal{L}$  implicitly.

(d) If  $\varphi$  is  $\psi_1 \vee \psi_2$ ,

$$\mathcal{A} \models \varphi(\bar{a}) \Leftrightarrow \mathcal{A} \models \psi_1(\bar{a}) \text{ or } \mathcal{A} \models \psi_2(\bar{a}) \Leftrightarrow \mathcal{B} \models \psi_1(\bar{a}) \text{ or } \mathcal{B} \models \psi_2(\bar{a}) \Leftrightarrow \mathcal{B} \models \varphi(\bar{a}),$$

where we use the induction hypothesis in the second  $\Leftrightarrow$ .

<sup>a</sup>Recall that we only need to show one of  $\vee$  or  $\wedge$ , and here we pick  $\vee$  and treat  $\wedge$  as an abbreviation. ■

**As previously seen** (Characteristic). Given a field  $K$ , the *characteristic*  $p$  of  $K$  is the number of 1 you need to add 1 in order to get 0, i.e.,  $\underbrace{1 + 1 + \dots + 1}_p = 0$ .

**Example.** Let  $L$  be a subfield of  $K$ , for each  $p > 0$ ,  $\varphi_p := \underbrace{1 + 1 + \dots + 1}_p = 0$ , which says the characteristic  $p$ .  $\varphi_p$  is **quantifier-free**, so

$$L \models \varphi_p \Leftrightarrow K \models \varphi_p.$$

**Example.** Consider  $\mathbb{Z} = (\mathbb{Z}, 0, 1, +, -, \cdot)$ , and let  $\varphi(x) := \neg \exists y \ y + y = x$ . We see that  $\mathbb{Z} \models \varphi(1)$  but  $\mathbb{Q} \models \neg \varphi(1)$ .

**Proposition 1.5.2.** Suppose that  $\mathcal{A}$  is a **substructure** of  $\mathcal{B}$ , and  $\varphi(\bar{x}, y_1, \dots, y_n)$  is a **quantifier-free formula**. Let  $\bar{a} \in \mathcal{A}$ , then

- (a) if  $\mathcal{A} \models \exists y_1 \dots \exists y_n \ \varphi(\bar{a}, y_1, \dots, y_n)$ , then  $\mathcal{B} \models \exists y_1 \dots \exists y_n \ \varphi(\bar{a}, y_1, \dots, y_n)$ ;
- (b) if  $\mathcal{B} \models \forall y_1 \dots \forall y_n \ \varphi(\bar{a}, y_1, \dots, y_n)$ , then  $\mathcal{A} \models \forall y_1 \dots \forall y_n \ \varphi(\bar{a}, y_1, \dots, y_n)$ .

**Proof.** Suppose that  $\mathcal{A} \models \exists y_1 \dots \exists y_n \ \varphi(\bar{a}, y_1, \dots, y_n)$ , so there are  $b_1, \dots, b_n \in \mathcal{A}$  such that  $\mathcal{A} \models \varphi(\bar{a}, b_1, \dots, b_n)$ . Since  $\varphi$  is **quantifier-free**, so  $\mathcal{B} \models \varphi(\bar{a}, b_1, \dots, b_n)$  from **Proposition 1.5.1**, and hence  $\mathcal{B} \models \exists y_1 \dots \exists y_n \ \varphi(\bar{a}, y_1, \dots, y_n)$ .

On the other hand, it's easy to see that (b) is implied by (a). ■

**Notation** (Existential). In **Proposition 1.5.2**, **formulas** as in (a) are called *existential* ( $\exists_1$  or  $\exists$ ) *formulas*.

**Notation** (Universal). In **Proposition 1.5.2**, **formulas** as in (b) are called *universal* ( $\forall_1$  or  $\forall$ ) *formulas*.

**Example.** Recall  $\mathcal{L}_1 = \{e, \cdot\}$ ,  $\mathcal{L}_2 = \{e, \cdot, {}^{-1}\}$ .

- Associativity:  $\forall x \forall y \forall z \ (xy)z = x(yz)$ .
- Identity:  $\forall x \ ex = xe$ .

These are  $\forall$ -**formulas** in either **language**.

- Inverses in  $\mathcal{L}_1$ :  $\forall x \exists y \ xy = yx = e$ , which is **not** an  $\forall$ -**formula**.
- Inverses in  $\mathcal{L}_2$ :  $\forall x \ xx^{-1} = x^{-1}x = e$ , which is an  $\forall$ -**formula**.

Hence, group axioms in  $\mathcal{L}_1$  are not **universal**, but in  $\mathcal{L}_2$  they are.

The above discrepancy is the reason why  $\mathcal{L}_2$  is better than  $\mathcal{L}_1$ , i.e.,  $\mathcal{L}_1$ -**substructure** might not be a group.

**Problem.** Show that  $\forall x \exists y \, xy = yx = e$  in the above example is not **equivalent** to an  $\forall$ -**formula**.

## Lecture 4: Theories and Axioms

17 Jan. 14:30

**Example.** Let  $\mathcal{L}_1 = \{E\}$ , where  $E$  is a binary relation representing edge relation; and  $\mathcal{L}_2 = \{V, E, I\}$ , where  $V, E$  are unary relations and  $I$  is a binary relation representing incidence such that  $I(v, e)$  for  $v \in V, e \in E$  means that  $v$  is a vertex on edge  $e$ . Then,

- Let  $G$  be a graph, viewed as an  $\mathcal{L}_1$ -**structure**. A **substructure** of  $G$  is an induced subgraph  $H \subseteq G$  such that any edge in  $G$  between two vertices of  $H$  is in  $H$ .
- If we view  $G$  as an  $\mathcal{L}_2$ -**substructure**, a **substructure** is a subgraph  $H$  such that  $H$  has some vertices and edges from  $G$ .<sup>a</sup>

<sup>a</sup>But there might be edges in  $H$  with no vertices, which can be fixed by having two functions  $I_1(e) = v, I_2(e) = w$  when  $e: v \rightarrow w$ .

**Remark.** The difference is that for  $\mathcal{L}_1$ , having an edge is **quantifier-free**, while in  $\mathcal{L}_2$  is **existential**. To elaborate a bit further, for  $\mathcal{L}_2$ ,  $vEw$  is **quantifier-free**, while in  $\mathcal{L}_2$ ,

$$\exists (v \in V \wedge w \in V \wedge e \in E \wedge I(v, e) \wedge I(w, e))$$

is not **quantifier-free**.

## Chapter 2

# Soundness, Completeness, and Compactness

In this chapter, we're going to formalize [proofs](#), including what do we mean by “having a proof” of a statement, and study properties of which.

### 2.1 Theories

Let's start by the notion of [theory](#).

**Definition 2.1.1** (Theory). An  $\mathcal{L}$ -theory is a set of  [\$\mathcal{L}\$ -sentences](#).

**Definition 2.1.2** (Model).  $\mathcal{M}$  is a *model* of a [theory](#)  $T$ , written as  $\mathcal{M} \models T$ , if  $\mathcal{M} \models \varphi$  for all  $\varphi \in T$ .

**Note.** Not every [theory](#) has a [model](#), e.g.,  $\{\exists x \, x \neq x\}$ .

The above note motivates the following.

**Definition 2.1.3** (Satisfiable). A [theory](#) is *satisfiable* if it has a [model](#).

**Definition 2.1.4** (Elementary class). A class  $\mathcal{K}$  of  [\$\mathcal{L}\$ -structures](#)  $\mathcal{M}$  is called an *elementary class* if there is an  [\$\mathcal{L}\$ -theory](#)  $T$  such that

$$\mathcal{K} = \{\mathcal{M} \mid \mathcal{M} \models T\}.$$

One way to get an [elementary class](#) is to take an  [\$\mathcal{L}\$ -structure](#)  $\mathcal{M}$  and take the [full theory](#).

**Definition 2.1.5** (Full theory). The *full theory*  $\text{Th}(\mathcal{M})$  of an  [\$\mathcal{L}\$ -structure](#)  $\mathcal{M}$  is defined as  $\text{Th}(\mathcal{M}) = \{\varphi \mid \mathcal{M} \models \varphi\}$ .

From the definition,  $\mathcal{M} \models \text{Th}(\mathcal{M})$ , and  $\text{Th}(\mathcal{M})$  characterizes the [structures](#) satisfying the same [sentences](#) as  $\mathcal{M}$ .

**Definition 2.1.6** (Complete). A [theory](#)  $T$  is *complete* if for any [sentence](#)  $\varphi$ , either  $\varphi \in T$  or  $\neg\varphi \in T$ .

**Remark.**  $\text{Th}(\mathcal{M})$  is [complete](#).

**Definition 2.1.7** (Elementarily equivalent).  $\mathcal{M}$  and  $\mathcal{N}$  are *elementarily equivalent*  $\mathcal{M} \equiv \mathcal{N}$  if for all [sentences](#)  $\varphi$ ,

$$\mathcal{M} \models \varphi \Leftrightarrow \mathcal{N} \models \varphi.$$

**Remark (Non-standard model of arithmetic).** There are  $\mathcal{N} \models \text{Th}(\mathbb{N})$ , but  $\mathcal{N}$  is not isomorphic to  $\mathbb{N}$ .  $\mathcal{N}$  is called a *non-standard model of arithmetic*, and  $\mathcal{N}$  might have *infinite element* larger than all of  $\mathbb{N}$ . Here,  $\mathbb{N} = (\mathbb{N}, 0, 1, +, \cdot, -)$

**Example.**  $\mathbb{Z} \oplus \mathbb{Z} \not\cong \mathbb{Z}$  as groups.

The other way to define a **theory** is to write down axioms.

**Example (Infinite set).** Let  $\mathcal{L} = \emptyset$ , and let  $T$  consist of

$$\varphi_n := \exists x_1 \dots \exists x_n \bigwedge_{i \neq j} x_i \neq x_j.$$

**Example (Linear order).** Let  $\mathcal{L} = \{\leq\}$ , and let  $T$  consist of the axioms of linear orders, e.g.,

$$\forall x \forall y (x \leq y \wedge y \leq x \rightarrow x = y).$$

There are other interesting **theories** of linear orders, e.g., dense ones.

**Example (Dense linear order).** Consider

$$\forall x \forall y (x < y \rightarrow \exists z x < z < y),$$

where we use  $a < b$  as shorthand of saying  $a \leq b \wedge a \neq b$ .

**Example (Group).** In  $\mathcal{L}_{\text{group}} = \{e, \cdot, {}^{-1}\}$ , let  $T$  be the group axioms.

Other **theories** of groups include Abelson group, divisible, etc.

**Definition 2.1.8 (Finitely axiomatizable).** A **theory** is *finitely axiomatizable* if it has a finite set of axioms.

Given a **theory**, consider  $T^{\models} = \{\varphi \mid T \models \varphi\}$ ,<sup>1</sup> so  $\mathcal{M} \models T$  if and only if  $\mathcal{M} \models T^{\models}$ . Often we think of  $T$  and  $T^{\models}$  as the same. A **theory**  $T$  is **finitely axiomatizable** if there is a finite  $\Phi$  such that  $T^{\models} = \Phi^{\models}$ .

## 2.2 Elementary Embeddings

Let's now consider the following notion.

**Definition 2.2.1 (Elementary embedding).** Let  $\mathcal{M}$  and  $\mathcal{N}$  be  $\mathcal{L}$ -structures, and  $f: \mathcal{M} \rightarrow \mathcal{N}$  an  $\mathcal{L}$ -embedding. Then  $f$  is an *elementary embedding* if for any **formula**  $\varphi(\bar{x})$  and  $\bar{a} \in \mathcal{M}$ ,

$$\mathcal{M} \models \varphi(\bar{a}) \Leftrightarrow \mathcal{N} \models \varphi(f(\bar{a})).$$

**Definition 2.2.2 (Elementary substructure).** If  $f: \mathcal{M} \hookrightarrow \mathcal{N}$  is a **elementary embedding** where  $\mathcal{M}$  is a **substructure** of  $\mathcal{N}$ , then  $\mathcal{M}$  is an *elementary substructure* of  $\mathcal{N}$ , written as  $\mathcal{M} \preceq \mathcal{N}$ .

**Example.** As groups,  $\mathbb{Z} \hookrightarrow \mathbb{Q}$  is not **elementary**. In fact,  $\mathbb{Z} \not\equiv \mathbb{Q}$ . Whereas, if  $f: \mathcal{M} \hookrightarrow \mathcal{N}$  is an **elementary embedding**,  $\mathcal{M} \equiv \mathcal{N}$ .<sup>a</sup>

<sup>a</sup>And also much more is true.

<sup>1</sup>Recall **Definition 1.5.2**.

**Proposition 2.2.1.** Every **isomorphism** is an **elementary embedding**.

**Proof.** Let  $f: \mathcal{M} \rightarrow \mathcal{N}$  be an **isomorphism**. We will argue by induction on **formulas**  $\varphi$ , that for all  $\bar{a} \in M$ ,

$$\mathcal{M} \models \varphi(\bar{a}) \Leftrightarrow \mathcal{N} \models \varphi(f(\bar{a})).$$

Firstly, observe that all cases except quantifiers are the same as **Proposition 1.5.1**. For quantifiers, suppose that  $\varphi(\bar{x})$  is  $\exists y \psi(\bar{x}, y)$  and  $\mathcal{M} \models \varphi(\bar{a})$ . This means that there is  $b \in M$  such that  $\mathcal{M} \models \psi(\bar{a}, b)$ . By the induction hypothesis,  $\mathcal{N} \models \psi(f(\bar{a}), f(b))$ , so  $\mathcal{N} \models \varphi(f(\bar{a}))$ .

Now suppose  $\mathcal{N} \models \varphi(f(\bar{a}))$ , then there is  $c \in N$  such that  $\mathcal{N} \models \psi(f(\bar{a}), c)$ . Since  $f$  is an **isomorphism**, so there is a  $b \in M$  such that  $f(b) = c$ . By the induction hypothesis,  $\mathcal{M} \models \psi(\bar{a}, b)$ , so  $\mathcal{M} \models \varphi(\bar{a})$ . ■

**Corollary 2.2.1.** If  $\mathcal{M} \cong \mathcal{N}$ , then  $\mathcal{M} \equiv \mathcal{N}$ .

## 2.3 Definable Sets

Consider the following.

**Definition 2.3.1** (Definable). Let  $\mathcal{M}$  be an  **$\mathcal{L}$ -structure**, then  $X \subseteq M^n$  is *definable* if there is a **formula**  $\varphi(x_1, \dots, x_n, \bar{y})$  and  $\bar{b} \in M$  such that

$$X = \{\bar{a} \in M^n \mid \mathcal{M} \models \varphi(\bar{a}, \bar{b})\}.$$

**Notation** (Define). We say that  $\varphi(\bar{x}, \bar{b})$  *defines*  $X$  over  $\bar{b}$ , written as  $X = \varphi(\mathcal{M}, \bar{b})$ .

**Notation** (Parameter). The tuple  $\bar{b}$  is called the *parameters* when  $X$  is **definable** over  $\bar{b}$ .

**Remark.** Sometimes  $X$  is **definable** without **parameters**, or **definable** over  $\emptyset$ .

**Example.** Take  $\mathbb{R} = (\mathbb{R}, 0, 1, +, \cdot, -)$  in  $\mathcal{L}_{\text{ring}}$ , then  $\leq = \{(a, b) : a \leq b\}$  is **definable**.

**Example.** Let  $\mathbb{Z} = (\mathbb{Z}, +, -, \cdot, 0, 1)$ , then  $\mathbb{N}$  is  **$\emptyset$ -definable** in  $\mathbb{Z}$  by<sup>a</sup>

$$\mathbb{N} = \{z \in \mathbb{Z} : \exists u, v, x, y \ u^2 + v^2 + x^2 + y^2 = z\}.$$

<sup>a</sup>From the **Langrange's four-square theorem**, which says that every natural number is the sum of four squares.

**Example.**  $\mathbb{Z}$  is  **$\emptyset$ -definable** in  $\mathbb{Q} = (\mathbb{Q}, +, -, \cdot, 0, 1)$ . This is a result of Julia Robinson [Rob49], and the formulation is very complicated.

**Problem.** How does one show that a set is not **definable**? For example,  $\mathbb{R}$  is not **definable** in  $\mathbb{C} = (\mathbb{C}, 0, 1, +, \cdot, -)$ .

## Lecture 5: Hilbert-Style Deductive System

We start by asking whether  $\mathbb{R}$  is **definable** in  $\mathbb{C} = (\mathbb{C}, 0, 1, +, \cdot, -)$ ?

19 Jan. 14:30

**Proposition 2.3.1.** Let  $\mathcal{M}$  be an  **$\mathcal{L}$ -structure**, and let  $X \subseteq M^n$  be a set which is **definable** over  $\bar{a}$ .



Then any **automorphism** of  $\mathcal{M}$  that fixes  $\bar{a}$  pointwise<sup>a</sup> fixes  $X$  setwise.<sup>b</sup>

<sup>a</sup>If  $\bar{a} = (a_1, \dots, a_m)$ , then  $f(a_i) = a_i$ .

<sup>b</sup>If  $b \in X$ , then  $f(b) \in X$ .

**Proof.** Let  $f$  be an **automorphism** of  $\mathcal{M}$  fixing  $\bar{a}$  pointwise, and  $X = \{\bar{b} \in M^n : \mathcal{M} \models \varphi(\bar{b}, \bar{a})\}$ . Fix  $\bar{b}$ , and suppose  $\bar{b} \in X$ , so  $\mathcal{M} \models \varphi(\bar{b}, \bar{a})$ . Because  $f$  is an **elementary embedding** from **Proposition 2.2.1**,

$$\mathcal{M} \models \varphi(f(\bar{b}), f(\bar{a})) \Rightarrow \mathcal{M} \models \varphi(f(\bar{b}), \bar{a}),$$

hence  $f(\bar{b}) \in X$ . Similarly, if  $\bar{b} \notin X$ ,  $\mathcal{M} \models \neg\varphi(\bar{b}, \bar{a}) \Rightarrow \mathcal{M} \models \neg\varphi(f(\bar{b}), \bar{a})$ , so  $f(\bar{b}) \notin X$ . ■

**Remark.** If  $X$  is  **$\emptyset$ -definable**, it is fixed setwise by any **automorphism**.

**Example.**  $\mathbb{N}$  is fixed setwise by any **automorphism** of the ring  $\mathbb{Z}$ . In fact, the only **automorphism** of  $\mathbb{Z}$  is the identity.

**Example.**  $\mathbb{N}$  is not  **$\emptyset$ -definable** in  $\mathbb{Z} = (\mathbb{Z}, 0, +)$ .

**Proof.** Consider an **automorphism**  $f(x) = -x$  of the group  $\mathbb{Z}$ , which does not fix  $\mathbb{N}$  setwise. ⊗

**Problem.** Is  $\mathbb{N}$  **definable** in  $\mathbb{Z} = (\mathbb{Z}, 0, +)$  over some parameters  $\bar{a}$ ?

**Answer.** For example, if  $\bar{a} = (1)$ , then  $f$  does not fix 1. In fact, any **automorphism** fixing 1 also fixes all of  $\mathbb{Z}$ , but  $\mathbb{N}$  is not **definable** in  $(\mathbb{Z}, 0, +)$ . To prove this we need **compactness**. ⊗

**As previously seen.** Given a field  $F$ , then  $F(a) \cong F(b)$  if  $a$  and  $b$  have the same minimal polynomial over  $F$  or if both do not satisfy any polynomial over  $F$ .

**Example.**  $\mathbb{Q}(\pi) \cong \mathbb{Q}(e)$  because  $\pi$  and  $e$  are both transcendental.

We now return to the big question: is  $\mathbb{R}$  **definable** in  $\mathbb{C} = (\mathbb{C}, 0, 1, +, \cdot, -)$ ? If  $f: \mathbb{Q}(a) \rightarrow \mathbb{Q}(b)$  such that  $a \mapsto b$ , then there is an **automorphism**  $\hat{f}: \mathbb{C} \rightarrow \mathbb{C}$  such that  $a \mapsto b$ , i.e.,  $\hat{f}$  extends  $f$ . In other words, we need to find such an  $f$  with  $a \in \mathbb{R}$  and  $b \notin \mathbb{R}$ .

**Example.**  $a = \pi$ ,  $b = i\pi$  are both transcendental.

**Example.**  $a$  is a real  $\sqrt[4]{2}$ ,  $b$  is a complex  $\sqrt[4]{2}$ .

The above two examples show that  $\mathbb{R}$  is not  **$\emptyset$ -definable** in  $\mathbb{C}$ . In fact,  $\mathbb{R}$  is not **definable** over any  $\bar{a}$  because there are elements of  $\mathbb{R}$  and  $\mathbb{C} \setminus \mathbb{R}$  transcendental over any  $\bar{a}$ .

**Intuition.** There are so many  $a, b$  such that given any  $\bar{a}$ , we can still find a pair that works.

## 2.4 Proofs

There are all sorts of different proof systems, and the one we use is the so-called Hilbert-style deductive system. Before that, we first see some common notions.

**Notation** (Schema). A *schema* is written in symbols for **formulas**, variables, etc.

**Example.**  $\varphi \rightarrow (\psi \rightarrow \varphi)$  is a **schema**, i.e., an infinite set with all possible choices of  $\varphi$  and  $\psi$ .

Specifically, every **logical axiom** is written in **schema**, meaning that any instance of a symbol for a **formula**, e.g.,  $\varphi$ , can be replaced by any **formula**.

**Definition 2.4.1 (Generalization).** A **formula**  $\varphi$  is a *generalization* of a **formula**  $\psi$  if  $\varphi$  is  $\forall x_1 \dots \forall x_n \psi$  where  $x_1, \dots, x_n$  are variables.

**Notation (Hypothesis).** *Hypotheses* are **formulas** that we may assume in a **proof**.

**Definition 2.4.2 (Proof).** A *proof* is a sequence of **formulas**  $\{\varphi_i\}_{i=1}^n$  such that  $\varphi_n$  is the conclusion, and each **formula** is either an **axiom** or is obtained from the previous **formulas** by a **rule of inference**.

Moreover, for a **proof** based on a set of **hypotheses**  $\Gamma$ , then in addition to a **logical axiom**, we can assert a **formula**  $\varphi \in \Gamma$ . If we prove  $\psi$  using  $\Gamma$  as **hypotheses**, we write  $\Gamma \vdash \psi$ .

**Definition 2.4.3 (Valid).** If we **prove**  $\psi$  without **hypotheses**, we write  $\vdash \psi$  and say  $\psi$  is *valid*.

**Definition 2.4.4 (Logical axioms).** The *logical axioms* are the following **formulas** written in **schema**, as well as all of their **generalizations**:

**Definition 2.4.5 (Propositional axioms).** The *propositional axioms* are

- (A1)  $\varphi \rightarrow (\psi \rightarrow \varphi)$ .
- (A2)  $(\varphi \rightarrow (\psi \rightarrow \theta)) \rightarrow ((\varphi \rightarrow \psi) \rightarrow (\varphi \rightarrow \theta))$ .
- (A3)  $(\neg \varphi \rightarrow \neg \psi) \rightarrow ((\neg \varphi \rightarrow \psi) \rightarrow \varphi)$ .

(A4)  $\forall x \varphi(x, \dots) \rightarrow \varphi(t, \dots)$  where  $t$  is any **term**.

(A5)  $[\forall x (\varphi \rightarrow \psi)] \rightarrow [(\forall x \varphi) \rightarrow (\forall x \psi)]$ .

(A6)  $\varphi \rightarrow \forall x \varphi$ , where  $x$  is not **free** in  $\varphi$ .

**Definition 2.4.6 (Axioms for equality).** The *axioms for equality* is

- (A7) for any **terms**  $t, u, v, \dots$ , function symbols  $f$ , and relation symbols  $R$ ,
  - (a)  $t = t$ .
  - (b)  $t = u \rightarrow u = t$ .
  - (c)  $(t = u \wedge u = v) \rightarrow (t = v)$ .
  - (d)  $(u_1 = t_1 \wedge \dots \wedge u_{n_f} = t_{n_f}) \rightarrow f(u_1, \dots, u_{n_f}) = f(t_1, \dots, t_{n_f})$ .
  - (e)  $(u_1 = t_1 \wedge \dots \wedge u_{n_R} = t_{n_R}) \rightarrow (R(u_1, \dots, u_{n_R}) \leftrightarrow R(t_1, \dots, t_{n_R}))$ .

**Definition 2.4.7 (Rule of inference).** From  $\varphi$  and  $\varphi \rightarrow \psi$ , deduces  $\psi$ .<sup>a</sup>

<sup>a</sup>This is called **modus ponens**.

These **formulas** might have **free variables**.

**Example.** A **proof** from calculus of a limit, e.g.,  $\forall \epsilon \exists \delta \dots$ . And we start by stating

1. let  $\epsilon > 0$ ,
2. choose  $\delta = \epsilon$ ,

$$\vdots$$

$$n. |f(x) - f(y)| < \epsilon.$$

We should interpret **free variables** as anything.

**As previously seen** (Propositional logic).  $(p \wedge q) \vee (r \wedge \neg q)$ .

**Remark.** We can check whether the **propositional axioms** are **true** with a truth table.

**Definition 2.4.8 (Propositional tautology).** A *propositional tautology* is a boolean combination  $\vee, \wedge, \neg$  of **formulas**  $\varphi_1, \dots, \varphi_n$  which is **true** via a truth table assigning true or false to each of  $\varphi_1, \dots, \varphi_n$ .

So instead of using **propositional axioms**, we could instead allow as **logical axioms** any **propositional tautology**. To prove **completeness**, we will need 5 **propositional tautologies**. We will **prove** some of these, but take others on faith.

**Remark.** **Propositional axioms** are enough to **prove** all **propositional tautologies**.

**Notation.** We write  $\Gamma \vdash_{\mathcal{L}} \varphi$  if there is a **proof** of  $\varphi$  from  $\Gamma$  in the **language**  $\mathcal{L}$ .

**Note.** Passing to a larger **language** will not let you **prove** more, so we can just write  $\vdash$ .

## Lecture 6: Soundness Theorem

To see why **propositional axioms** are enough to **prove** all **propositional tautologies**, we see one example. 24 Jan. 14:30

**Problem.** **Prove**  $\varphi \rightarrow \varphi$  using **propositional axioms**.

**Answer.** We see that

1.  $\varphi \rightarrow ((\psi \rightarrow \varphi) \rightarrow \varphi)$  from (A1), where  $\psi$  is any **formula** (possibly  $\psi = \varphi$ ).
2.  $[\varphi \rightarrow ((\psi \rightarrow \varphi) \rightarrow \varphi)] \rightarrow [(\varphi \rightarrow (\psi \rightarrow \varphi)) \rightarrow (\varphi \rightarrow \varphi)]$  from (A2).
3.  $(\varphi \rightarrow (\psi \rightarrow \varphi)) \rightarrow (\varphi \rightarrow \varphi)$  from (MP) and the two above.
4.  $\varphi \rightarrow (\psi \rightarrow \varphi)$  from (A1).
5.  $\varphi \rightarrow \varphi$  from (MP) and the two above.

⊛

In general, we can **prove**

- |   |  |
|---|--|
| (a) $\varphi \rightarrow \varphi$ ;         | (d) $(\varphi \rightarrow \psi) \rightarrow ((\neg\varphi \rightarrow \psi) \rightarrow \psi)$ ; |
| (b) $\varphi \rightarrow \neg\neg\varphi$ ; |  |
| (c) $\neg\neg\varphi \rightarrow \varphi$ ; | (e) $\varphi \rightarrow (\psi \rightarrow (\varphi \rightarrow \psi))$ ,                        |
| and so on.                                  |  |

**Note.** As we said, we may replace **propositional axioms** by every **propositional tautologies**.

Some **proof** system also have a second rule about universal quantifiers, but in our system, we have built this into the axioms. We can prove, as a theorem, what the other proof systems take as a rule.

**Theorem 2.4.1.** If  $\Gamma \vdash \varphi$ , and  $x$  does not occur **freely** in  $\Gamma$ , then  $\Gamma \vdash \forall x \varphi$ .

**Proof.** Fix  $\Gamma$  and  $x$ , we use *induction on proofs*. Consider the set  $\{\varphi \mid \Gamma \vdash \forall x \varphi\}$ , we will show that this set contains all the **logical axioms**, **formulas** from  $\Gamma$ , and is closed under **modus ponens**.<sup>a</sup>

- (a) If  $\varphi$  is a **logical axiom**, so is its **generalization**  $\forall x \varphi$ , so  $\Gamma \vdash \forall x \varphi$ .
- (b) If  $\varphi \in \Gamma$ , then  $x$  is not **free** in  $\varphi$ , so from (A6),  $\varphi \rightarrow \forall x \varphi$ , and from (MP),  $\forall x \varphi$ . The above are based on  $\Gamma$ , hence  $\Gamma \vdash \forall x \varphi$ .
- (c) Suppose  $\Gamma \vdash \forall x \varphi$  and  $\Gamma \vdash \forall x (\varphi \rightarrow \psi)$ , we want to show that  $\Gamma \vdash \forall x \psi$ .
  1. By (A5),  $\forall x (\varphi \rightarrow \psi) \rightarrow (\forall x \varphi \rightarrow \forall x \psi)$ ,  $\Gamma$  **proves** this.
  2. By (MP),  $\Gamma \vdash \forall x \varphi \rightarrow \forall x \psi$ .
  3. By (MP) again,  $\Gamma \vdash \forall x \psi$ .

■

<sup>a</sup>Thus, if  $\Gamma \vdash \theta$ , then  $\theta \in \{\varphi \mid \Gamma \vdash \forall x \varphi\}$ .

**Corollary 2.4.1.** If  $\vdash \varphi$ , then  $\vdash \forall x \varphi$ . So the **generalization** of anything **valid** is also **valid**.

We now ask a critical question: is our **proof** system a good one?

## 2.5 Soundness Theorem

The first thing we should check is whether our **proofs** are **sound**.

**Definition 2.5.1 (Sound).** A **proof** system is *sound* if any **provable sentence**  $\varphi$  is **true**.

The idea is that if an  **$\mathcal{L}$ -sentence**  $\varphi$  is **provable**, then it is **true** in all  **$\mathcal{L}$ -structures**, i.e., every thing we **prove** should be **true**, in other words, we can't **prove** wrong things.

**Lemma 2.5.1 (Soundness).** If  $\Gamma$  is a set of  **$\mathcal{L}$ -sentences** and  $\varphi$  is a **sentence**, and  $\Gamma \vdash_{\mathcal{L}} \varphi$ , then  $\Gamma \models \varphi$ .

**Proof.** Suppose that  $\Gamma \vdash \varphi$ , let  $\psi_1, \psi_2, \dots, \psi_n = \varphi$  be such a **proof**.<sup>a</sup> Let  $\bar{x} = (x_1, \dots, x_m)$  be the **free variable** that appears in the  $\psi_i$ . Let  $\mathcal{M}$  be an  **$\mathcal{L}$ -structure**,  $\mathcal{M} \models \Gamma$ . To show  $\mathcal{M} \models \varphi$ , we show that by induction on  $i$ , for all  $\bar{a} \in M^m$ ,  $\mathcal{M} \models \psi_i(\bar{a})$ . For  $\psi_i$ , we have three cases.

- (a) If  $\psi_i \in \Gamma$ , then  $\mathcal{M} \models \Gamma$  so  $\mathcal{M} \models \psi_i$ .
- (b) If  $\psi_i$  is a (**generalization** of) a **logical axiom**, then we can check that  $\mathcal{M} \models \psi_i(\bar{a})$ . For example, if  $\psi_i$  is (A1),  $\theta \rightarrow (\gamma \rightarrow \theta)$ , it's easy to check that

$$\mathcal{M} \models \theta(\bar{a}) \rightarrow (\gamma(\bar{a}) \rightarrow \theta(\bar{a})).$$

- (c) If there are  $j, k < i$  such that  $\psi_k$  is  $\psi_j \rightarrow \psi_i$ , from inductive hypothesis, for all  $\bar{a}$ ,  $\mathcal{M} \models \psi_j(\bar{a}), \mathcal{M} \models \psi_k(\bar{a})$ , then  $\mathcal{M} \models \psi_j(\bar{a}) \rightarrow \psi_i(\bar{a})$ . Checking our definition of **truth**, we get  $\mathcal{M} \models \psi_i(\bar{a})$ .

■

<sup>a</sup>Some  $\psi_i$  might be **formulas**, but  $\varphi$  should be a **sentence**.

There are remarks to make about some obvious properties of  $\vdash_{\mathcal{L}}$ .

**Remark.** If  $\varphi \in \Gamma$ , then  $\Gamma \vdash \varphi$ .

**Remark.** If  $\Delta \subseteq \Gamma$ , and  $\Delta \vdash \varphi$ , then  $\Gamma \vdash \varphi$ .

**Remark.** If  $\Gamma \vdash_{\mathcal{L}} \varphi$ , and  $\mathcal{L}^+ \supseteq \mathcal{L}$ , then  $\Gamma \vdash_{\mathcal{L}^+} \varphi$ .

**Remark.** If  $\Gamma \vdash \varphi$ , then there is a finite  $\Delta \subseteq \Gamma$  such that  $\Delta \vdash \varphi$ .

We can prove the following.

**Theorem 2.5.1 (Deduction theorem).** For any set of formulas  $\Gamma$ , formulas  $\theta$  and  $\psi$ ,

$$\Gamma \cup \{\theta\} \vdash \psi \Leftrightarrow \Gamma \vdash \theta \rightarrow \psi.$$

**Proof.** The backward direction is easier. Suppose  $\Gamma \vdash \theta \rightarrow \psi$ , then  $\Gamma \cup \{\theta\} \vdash \psi$  since we can have a proof like:

1.  $\theta$
- $\vdots$  (the proof of  $\Gamma \vdash \theta \rightarrow \psi$ )
- $n$ .  $\theta \rightarrow \psi$
- $n + 1$ .  $\psi$ .

Now, suppose that  $\Gamma \cup \{\theta\} \vdash \psi$ , then there is a proof  $\psi_1, \dots, \psi_n = \psi$  from  $\Gamma \cup \{\theta\}$ . We argue inductively that  $\Gamma \vdash \theta \rightarrow \psi_i$ . For  $i$ , we have three cases.

- (a) If  $\psi_i \in \Gamma$  or it is a logical axiom. By (A1),  $\psi_i \rightarrow (\theta \rightarrow \psi_i)$ , so  $\Gamma \vdash \theta \rightarrow \psi_i$ .
- (b) If  $\psi_i = \theta$ . Then  $\Gamma \vdash \theta \rightarrow \theta$  by (A1) and (A2) from here, hence  $\Gamma \vdash \theta \rightarrow \psi_i$ .
- (c) If  $\psi_i$  follows from  $\psi_j, \psi_k = \psi_j \rightarrow \psi_i$ , using (MP) with  $j, k < i$ .
  1. From the induction hypothesis,  $\Gamma \vdash \theta \rightarrow \psi_j$  and  $\Gamma \vdash \theta \rightarrow (\psi_j \rightarrow \psi_i)$ .
  2. By (A2),  $\Gamma \vdash [\theta \rightarrow (\psi_j \rightarrow \psi_i)] \rightarrow [(\theta \rightarrow \psi_j) \rightarrow (\theta \rightarrow \psi_i)]$ .
  3. By (MP),  $\Gamma \vdash (\theta \rightarrow \psi_j) \rightarrow (\theta \rightarrow \psi_i)$ .
  4. By (MP),  $\Gamma \vdash \theta \rightarrow \psi_i$ .

■

## Lecture 7: Soundness, Completeness, and Compactness

**Proposition 2.5.1 (Contraposition).** If  $\Gamma \cup \{\varphi\} \vdash \neg\psi$ , then  $\Gamma \cup \{\psi\} \vdash \neg\varphi$ .

**Proof.** Suppose  $\Gamma \cup \{\varphi\} \vdash \neg\psi$ , by the deduction theorem says that  $\Gamma \vdash \varphi \rightarrow \neg\psi$ . From (A1), (A2), and (A3), we can prove  $(\varphi \rightarrow \neg\psi) \rightarrow (\psi \rightarrow \neg\varphi)$ . By (MP),  $\Gamma \vdash \psi \rightarrow \neg\varphi$ , then from the deduction theorem,  $\Gamma \cup \{\psi\} \vdash \neg\varphi$ . ■

Now we introduce an important notion.

**Definition 2.5.2 (Consistent).** A theory  $T$  is *consistent* if for all  $\varphi$ , it is not the case that  $T \vdash \varphi$  and  $T \vdash \neg\varphi$ .

**Definition 2.5.3 (Inconsistent).** If a theory  $T$  is not consistent, then it's *inconsistent*.

We could make the same definition for a set of formulas.

26 Jan. 14:30

**Proposition 2.5.2** (Proof by contradiction). If  $\Gamma \cup \{\varphi\}$  is **inconsistent**, then  $\Gamma \vdash \neg\varphi$ .

**Proof.** There is  $\psi$  such that  $\Gamma \cup \{\varphi\} \vdash \psi$  and  $\Gamma \cup \{\varphi\} \vdash \neg\psi$ , so  $\Gamma \vdash \varphi \rightarrow \psi$  and  $\Gamma \vdash \varphi \rightarrow \neg\psi$  by the **deduction theorem**. Using (A1), (A2), and (A3), we can prove that

$$(\varphi \rightarrow \psi) \rightarrow ((\varphi \rightarrow \neg\psi) \rightarrow \neg\varphi).$$

By (MP),  $\Gamma \vdash (\varphi \rightarrow \neg\psi) \rightarrow \neg\varphi$ , and by (MP) again, we have  $\Gamma \vdash \neg\varphi$ . ■

**Proposition 2.5.3.** If a **theory**  $T$  is **consistent**, and  $\varphi$  is a **sentence**, then either  $T \cup \{\varphi\}$  or  $T \cup \{\neg\varphi\}$  is **consistent**.

**Proof.** If they were both **inconsistent**,  $T \vdash \neg\varphi$  and  $T \vdash \neg\neg\varphi$ , so  $T$  would be **inconsistent**  $\nmid$  ■

**Note.** The above is also true for **formulas**.

**Remark.** If  $T$  is **inconsistent**, then  $T \vdash \varphi$  for any  $\varphi$ .

**Proof.** If  $T$  is **inconsistent**, then  $T \cup \{\neg\varphi\}$  is **inconsistent** for all  $\varphi$ . Hence, from **proof by contradiction**,  $T \vdash \neg\neg\varphi$  for all  $\varphi$ , which is just  $T \vdash \varphi$ . ⊛

**Definition 2.5.4** (Maximal). A **theory**  $T$  is *maximal* if it is **consistent** and for all **sentences**  $\varphi$ , either  $\varphi \in T$  or  $\neg\varphi \in T$ .

In particular, if  $T \vdash \varphi$ , then  $\varphi \in T$ .

**Intuition.** Basically, a **maximal consistent theory** has opinion on everything.

Now, we want to see that given a **consistent theory**, whether we can extend it to a **maximal** one. To do this, we need the following.

**Definition.** Let  $(P, \leq)$  be a **partially ordered set**.

**Definition 2.5.5** (Chain). A *chain* is a set  $C \subseteq P$  such that for every  $p, q \in C$ , either  $p \leq q$  or  $q \leq p$ .

**Definition 2.5.6** (Upper bound). If  $X \subseteq P$  is a set, an *upper bound* for  $X$  is an element  $p \in P$  such that  $p \geq q$  for all  $q \in X$ .

**Definition 2.5.7** (Maximal). An element  $p \in P$  is *maximal* if there is no  $q \in P$  with  $q > p$ .

**Note.** Note that a **maximal** element might not be greater than everything, there is just nothing greater than it.

**Theorem 2.5.2** (Zorn's lemma). Let  $(P, \leq)$  be a **partially ordered set**. If every non-empty **chain** in  $P$  has an **upper bound**, then  $P$  has a **maximal** element.

**Theorem 2.5.3.** Any **consistent theory**  $T$  can be extended to a **maximal consistent theory**  $T' \supseteq T$ .

**Proof.** We first consider the case that  $T$  is countable by considering  $\mathcal{L}$  is countable since if  $\mathcal{L}$  is countable, then there are only countable many **formulas** since there are only countable many **formulas** of each length.

**Claim.** The result holds for  $\mathcal{L}$  being countable.

**Proof.** Firstly, list out all sentences  $\varphi_1, \varphi_2, \dots$ , start with  $T_0 = T$ . Given  $T_i$  consistent, one of  $T_i \cup \{\varphi_i\}$  or  $T_i \cup \{\neg\varphi_i\}$  is consistent from Proposition 2.5.3. Let  $T_{i+1}$  be one of these that is consistent. Let  $T^* = \bigcup_i T_i$ , which is maximal, and we now show that  $T^*$  is consistent.

Suppose not, then  $T^* \vdash \theta$  and  $T^* \vdash \neg\theta$  for some  $\theta$ . In this case, there is some  $T_i$  such that  $T_i \vdash \theta$  and  $T_i \vdash \neg\theta$  because proofs are finite, with  $T_i$  being consistent, a contradiction  $\nmid \quad \circledast$

**Claim.** The result holds for arbitrary  $\mathcal{L}$ .

**Proof.** For arbitrary  $\mathcal{L}$ , let  $(P, \leq)$  be the set of consistent theories extending  $T_i$  ordered by inclusion. Let  $C$  be a non-empty chain, and let  $T^* = \bigcup_{T' \in C} T' \supseteq T$ .

We see that  $T^*$  is consistent because if  $T^* \vdash \theta$  and  $T^* \vdash \neg\theta$ , there are finitely many formulas used in those proofs, from, say,  $T_1, \dots, T_n \in C$ . Because  $C$  is a chain, by reordering, we may assume that  $T_1 \subseteq \dots \subseteq T_n$ . So  $T_n \vdash \theta$  and  $T_n \vdash \neg\theta$ , contradicting the consistency of  $T_n$ , so  $T^*$  is consistent, i.e.,  $T^* \in P$ . Furthermore,  $T^*$  is an upper bound on  $C$ ,<sup>a</sup> so  $(P, \leq)$  has a maximal consistent theory  $T^* \supseteq T$  from Zorn's lemma.

If  $T^*$  is not maximal, then there is  $\varphi$  where  $\varphi \notin T^*$ ,  $\neg\varphi \notin T^*$ . From Proposition 2.5.3, one of  $T^* \cup \{\varphi\}$  or  $T^* \cup \{\neg\varphi\}$  is consistent, hence in  $P$ , contradicting to  $T^*$  being maximal  $\nmid \quad \circledast$

<sup>a</sup>Note that  $C$  is arbitrary.

■

**Remark.** We can do that same proof for any  $\mathcal{L}$  using transfinite recursion for the uncountable case.

Motivated by Lemma 2.5.1 and Theorem 2.5.3, we close this section with the following.

**Theorem 2.5.4 (Soundness).** Let  $T$  be a theory and  $\varphi$  be a sentence.

- (a) If  $T \vdash \varphi$ , then  $T \models \varphi$ .
- (b) If  $T$  is satisfiable, then it is consistent.

**Proof.** (a) is exactly Theorem 2.5.4. For (b), let  $\mathcal{M} \models T$ , suppose that  $T$  was inconsistent, then  $T \vdash \varphi$  and  $T \vdash \neg\varphi$  for some  $\varphi$ . By (a),  $T \models \varphi$  and  $T \models \neg\varphi$ , so  $\mathcal{M} \models \varphi$  and  $\mathcal{M} \models \neg\varphi$ . But  $\mathcal{M} \models \neg\varphi$  means  $\mathcal{M} \not\models \varphi$ , so this is a contradiction, hence  $T$  is consistent. ■

## 2.6 Completeness and Compactness Theorems

After knowing our proof system is sound, we now ask the converse: is our proof system complete?

**Definition 2.6.1 (Complete).** A proof system is complete if any true sentence  $\varphi$  is provable.

And indeed, this is the case.

**Theorem 2.6.1 (Completeness).** Let  $T$  be a theory and  $\varphi$  be a sentence.

- (a) If  $T \models \varphi$ , then  $T \vdash \varphi$ .
- (b) If  $T$  is consistent, then it is satisfiable.

(b) implies (a) is easy to see. Suppose that  $T \models \varphi$ , so  $T \cup \{\neg\varphi\}$  is not satisfiable. By (b),  $T \cup \{\neg\varphi\}$  is inconsistent. By proof by contradiction,  $T \vdash \varphi$ . One important consequence of the completeness theorem is the compactness theorem.

**Theorem 2.6.2 (Compactness).** Let  $T$  be a **theory** and  $\varphi$  be a **sentence**.

- (a) If  $T \models \varphi$ , then there is a finite  $T_0 \subseteq T$  such that  $T_0 \models \varphi$ .
- (b)  $T$  is **satisfiable** if and only if every finite subset of  $T$  is **satisfiable**.

**Proof.** Consider the following.

- (a\*) If  $T \vdash \varphi$ , then there is a finite  $T_0 \subseteq T$  such that  $T_0 \vdash \varphi$ .
- (b\*) If  $T$  is **consistent** if and only if every finite subset of  $T$  is **consistent**.

We see that (a\*) and (b\*) are true because **proofs** are finite, and **soundness** and **completeness** translate directly between (a) and (a\*) (and (b) and (b\*)). ■

**Remark.** The **compactness theorem** does have something to do with topological compactness; consider the topological space of **complete satisfiable theories**, with the basic open sets being the sets

$$U_\varphi := \{T : T \models \varphi\},$$

then this topological space is compact.

Let's see one cool example using **compactness**.

**Example (Construction of non-standard model of arithmetic).** Let  $\mathcal{L} = \{0, 1, +, \cdot, -, <\}$ , and  $\mathcal{L}^* = \mathcal{L} \cup \{c\}$ , where  $c$  is a new constant symbol. Let

$$T = \text{Th}_{\mathcal{L}}(\mathbb{N}) \cup \{c > \underline{n} \mid n \in \mathbb{N}\},$$

then  $T$  is finitely **satisfiable**.

**Proof.** Given  $T_0 \subseteq T$  finite,  $T_0 \subseteq \text{Th}_{\mathcal{L}}(\mathbb{N}) \cup \{c > \underline{n}_1, \dots, c > \underline{n}_\ell\}$ , and may assume they are equal and show that  $T_0$  is **satisfiable**. Let  $\mathcal{N}$  be the  $\mathcal{L} \cup \{c\}$ -**structure** which is the **expansion** of the  $\mathcal{L}$ -**structure**  $\mathbb{N}$ , with

$$c^{\mathcal{N}} = 1 + \max(n_1, \dots, n_\ell),$$

then  $\mathcal{N} \models T_0$ , and  $T_0$  is **satisfiable**. By **compactness**,  $T$  is **satisfiable**, say  $\mathcal{A} \models T$ . Then  $\mathcal{A} \equiv \mathbb{N}$  and  $\mathcal{A}$  contains an element  $c^{\mathcal{A}}$  bigger than  $1, 1 + 1, 1 + 1 + 1, \dots$ , but  $\mathcal{A} \not\equiv \mathbb{N}$ , so  $\mathcal{A}$  is a **non-standard model of arithmetic**. ⊛

We now start a long journey toward proving **completeness theorem**, specifically (b).

## Lecture 8: Henkin Constants

### 2.6.1 Henkin Construction

31 Jan. 14:30

To prove **Theorem 2.6.1 (b)**, we need an additional definition and a technical lemma due to Henkin.

**Definition 2.6.2 (Henkin constant).** An  $\mathcal{L}^*$ -**theory**  $T^*$  has *Henkin constants* if for each **formula**  $\varphi(x)$  with one **free variable**, there is a constant symbol  $c \in \mathcal{L}^*$  such that

$$(\exists x \varphi(x)) \rightarrow \varphi(c) \text{ is in } T^*.$$

We see that the above is equivalent to

$$(\neg \forall x \varphi(x)) \rightarrow \neg \varphi(c) \text{ is in } T^*,$$

and we will use this version ( $\forall$ ) and view  $\exists$  being a shorthand for  $\neg \forall \neg$ ; also, we will use  $\rightarrow$  and  $\neg$  as primitive, and  $\wedge, \vee$  are shorthand.



**Lemma 2.6.1.** If  $\Gamma \vdash \varphi(c)$ , and  $c$  does not occur in  $\Gamma$  or in  $\varphi(x)$ , then there is a variable  $y$  not appearing in  $\varphi(x)$ , such that  $\Gamma \vdash \forall y \varphi(y)$ . Moreover, there is a **proof** of  $\forall y \varphi(y)$  in which  $c$  does not appear.

**Proof.** Let  $\alpha_1(c), \dots, \alpha_n(c) = \varphi(c)$  be a **proof** of  $\varphi(c)$  from  $\Gamma$ , and let  $y$  be a variable not appearing in this **proof**. We claim that  $\alpha_1(y), \dots, \alpha_n(y) = \varphi(y)$  is still a valid **proof** of  $\varphi(y)$ . There are three cases to consider (for each  $i = 1, \dots, n$ ):

- (a) If  $\alpha_i(c)$  is in  $\Gamma$ , then  $c$  does not actually occur in  $\alpha_i(c)$  because it does not appear in  $\Gamma$ . So  $\alpha_i(y)$  is the same as  $\alpha_i(c)$ , hence in  $\Gamma$ .
- (b) If  $\alpha_i(c)$  is a **logical axiom**, then  $\alpha_i(y)$  is a **logical axiom** as well. For most of these it is easy to check, but for (A6), i.e.,  $\varphi \rightarrow \forall x \varphi$  if  $x$  is not **free** in  $\varphi$ , there is a little more. But  $y$  did not appear in  $\alpha_i(c)$ , so  $y \neq x$ , and substituting  $y$  for  $c$  will not stop  $x$  from being not **free**.
- (c) If  $\alpha_i(c)$  follows by (MP) from  $\alpha_j(c)$  and  $\alpha_k(c) = \alpha_j(c) \rightarrow \alpha_i(c)$  for  $j, k < i$ , then  $\alpha_i(y)$  follows by (MP) from  $\alpha_j(y)$  and  $\alpha_k(y) = \alpha_j(y) \rightarrow \alpha_i(y)$ .

So  $\Gamma \vdash \varphi(y)$  and the **proof** does not involve  $c$ . Let  $\Phi \subseteq \Gamma$  be the subset of  $\Gamma$  that was used in the **proof**, so  $y$  does not appear in  $\Phi$ , hence  $\Phi \vdash \varphi(y)$  and  $\Phi \vdash \forall y \varphi(y)$ , so  $\Gamma \vdash \forall y \varphi(y)$ . ■

So Lemma 2.6.1 implies that we have  $\Gamma \vdash \varphi(y)$  and the **proof** does not involve  $c$ . And sometimes, we want to be able to choose the variable  $y$  from above.

**Corollary 2.6.1.** If  $\Gamma \vdash \varphi(c)$ , and  $c$  does not occur in  $\Gamma$  or in  $\varphi(x)$ , then  $\Gamma \vdash \forall x \varphi(x)$ . Moreover, there is a **proof** of  $\forall x \varphi(x)$  not involving  $c$ .<sup>a</sup>

<sup>a</sup>Here,  $x$  is any variable that does not appear in  $\varphi(c)$ .

**Proof.** We know that for some  $y$ ,  $\Gamma \vdash \forall y \varphi(y)$ , (A4) says  $\forall y \varphi(y) \rightarrow \varphi(x)$ . So  $\forall y \varphi(y) \vdash \varphi(x)$  since  $x$  does not appear in  $\forall y \varphi(y)$ ,  $\forall y \varphi(y) \vdash \forall x \varphi(x)$ . ■

**Note.**  $x$  might appear in  $\Gamma$ .

**Theorem 2.6.3.** Let  $T$  be a **consistent  $\mathcal{L}$ -theory**. There is a **language**  $\mathcal{L}^* \supseteq \mathcal{L}$  and  $T^* \supseteq T$  a **consistent  $\mathcal{L}^*$ -theory** such that  $T^*$  has **Henkin constants**. We can choose  $\mathcal{L}^*$  such that  $|\mathcal{L}^*| = |\mathcal{L}| + \aleph_0$ , and all new symbols in  $\mathcal{L}^*$  are constants.

**Proof.** Let  $\mathcal{L}_0 = \mathcal{L}$  and  $T_0 = T$ . Let  $\mathcal{L}_1$  be the **expansion** of  $\mathcal{L}_0$  by adding a new constant symbol  $c_\varphi$  for each  **$\mathcal{L}_0$ -formula**  $\varphi$  w.r.t. the **Henkin** construction. First, we show that after this procedure,  $T_0$  is still a **consistent  $\mathcal{L}_1$ -theory**.

**Remark.** Technically,  $\vdash$  is really  $\vdash_{\mathcal{L}}$ , so this is a key step for seeing that it does not matter.

**Claim.**  $T_0$  is still a **consistent  $\mathcal{L}_1$ -theory** after the **expansion** of  $\mathcal{L}_0$ .

**Proof.** If not, there is a **proof** of a **contradiction** from  $T_0$ , and which uses only finitely many of the new constants symbols. By Corollary 2.6.1, we can replace these constants one-by-one by variables, e.g., if the original **contradiction** was  $\varphi(c_1, \dots, c_n)$  and  $\neg \varphi(c_1, \dots, c_n)$ , then  $T_0$  proves  $\forall x_1, \dots, \forall x_n \varphi(x_1, \dots, x_n)$  and  $\forall x_1, \dots, \forall x_n \neg \varphi(x_1, \dots, x_n)$ . Moreover, these **proofs** take place in  $\mathcal{L}_0$ , so by (A4),  $T_0 \vdash_{\mathcal{L}_0} \varphi(x_1, \dots, x_n)$ , and  $T_0 \vdash_{\mathcal{L}_0} \neg \varphi(x_1, \dots, x_n) \not\vdash$  ⊗

To construct  $T_1$  w.r.t. the **Henkin** construction, it's natural to consider the following: if  $\varphi$  is of the form  $\neg \forall x \psi(x)$ , then let

$$\theta_\varphi := (\neg \forall x \psi(x)) \rightarrow \neg \psi(c_\varphi), \text{ i.e., } (\exists x \neg \psi(x)) \rightarrow \neg \psi(c_\varphi),$$

otherwise, let  $\theta_\varphi := \forall x (x = x)$  (trivially **true**). Let  $\Theta = \{\theta_\varphi \mid \varphi \text{ an } \mathcal{L}_0\text{-formula}\}$ , and we let that  $T_1 = T_0 \cup \Theta$ . We claim that  $T_1$  is still **consistent**.

**Claim.**  $T_1 = T_0 \cup \Theta$  is a consistent  $\mathcal{L}_1$ -language after the expansion of  $\mathcal{L}_0$ .

**Proof.** If not, then there are  $\varphi_1, \dots, \varphi_{m+1}$  such that  $T_0 \cup \{\theta_{\varphi_1}, \dots, \theta_{\varphi_m}, \theta_{\varphi_{m+1}}\}$  is inconsistent. Taking  $m$  to be as small as possible,  $T_0 \cup \{\theta_{\varphi_i}\}_{i=1}^m$  is consistent, so  $T_0 \cup \{\theta_{\varphi_i}\}_{i=1}^m \vdash \neg\theta_{\varphi_{m+1}}$  with  $\varphi_{m+1}$  being of the form  $\neg\forall x \psi(x)$ ,  $\theta_{\varphi_{m+1}}$  is  $\neg\forall x \psi(x) \rightarrow \neg\psi(c_\varphi)$ . By (A1), (A2), (A3),

$$T_0 \cup \{\theta_{\varphi_1}, \dots, \theta_{\varphi_m}\} \vdash \neg\forall x \psi(x) \text{ and } T_0 \cup \{\theta_{\varphi_1}, \dots, \theta_{\varphi_m}\} \vdash \psi(c_{\varphi_{m+1}}).$$

Since  $c_{\varphi_{m+1}}$  does not appear in  $T_0 \cup \{\theta_{\varphi_i}\}_{i=1}^m$ , so  $T_0 \cup \{\theta_{\varphi_i}\}_{i=1}^m \vdash \forall x \psi(x)$ , i.e.,  $T_0 \cup \{\theta_{\varphi_i}\}_{i=1}^m$  is inconsistent, contradicting to the fact that  $m$  is the smallest choice  $\nless^a$   $\textcircled{*}$

<sup>a</sup>If  $m = 0$ , then we violate the consistency of  $T_0$ .

It might be that  $T_1$  does not have Henkin constants since there are new  $\mathcal{L}_1$ -formulas which are not  $\mathcal{L}_0$ -formulas. But we know that  $T_1$  does have Henkin constants for  $\mathcal{L}_0$ -formulas, hence we can repeat that process and keep fixing things. In general, given  $T_i$  and  $\mathcal{L}_i$ , define a  $T_{i+1}$  and  $\mathcal{L}_{i+1}$  in the above way. Since each  $T_i$  is consistent, so  $T^* = \bigcup T_i$  is an  $\mathcal{L}^* = \bigcup \mathcal{L}_i$ -theory. Note that  $T^*$  is consistent as a nested union of consistent theories, and  $T^*$  has Henkin constants because every  $\mathcal{L}^*$ -formula  $\varphi$  is an  $\mathcal{L}_i$ -formula for some  $i$ , and  $\theta_\varphi \in T_{i+1} \subseteq T^*$ .

**Intuition.** This is like “chasing its own tail,” which basically fixes new errors introduced every time and then takes the union in the end.

Finally, we want to show that  $|\mathcal{L}^*| = |\mathcal{L}| + \aleph_0$ . Given  $\mathcal{L}_i$ , we define  $\mathcal{L}_{i+1}$  to be  $\mathcal{L}_i$  plus new constants  $c_\varphi$  for  $\varphi$  on  $\mathcal{L}_i$ -formula. Then, we have

$$|\mathcal{L}_{i+1}| \leq |\mathcal{L}_i| + \underbrace{|\mathcal{L}_i|}_{\# \text{ of } \mathcal{L}_i\text{-formulas}} + \aleph_0 = |\mathcal{L}_i| + \aleph_0.$$

So for all  $i$ ,  $|\mathcal{L}_i| \leq |\mathcal{L}| + \aleph_0$ , and  $\mathcal{L}^* = \bigcup_i \mathcal{L}_i$  is a countable union, so  $|\mathcal{L}^*| \leq |\mathcal{L}| + \aleph_0$ , and in fact,  $|\mathcal{L}^*| = |\mathcal{L}| + \aleph_0$ .  $\blacksquare$

## 2.6.2 Proof of Completeness Theorem

After proving Theorem 2.6.3, we see that to prove Theorem 2.6.1 (b), we can proceed by:

1. extend  $T^*$  to a maximal theory  $T^{**}$ ;<sup>2</sup>
2. turn  $T^{**}$  into a model. The elements of the model are constant symbols from  $\mathcal{L}^*$ , modulo the equivalence relation  $c \sim d$  if  $c = d$  is in  $T^{**}$ , i.e.,  $T^{**} \vdash c = d$ .

Thankfully, the first step is easy from Theorem 2.5.3, so we just need to show the second step, and we're done.

## Lecture 9: Proving the Completeness Theorem

To finish the proof of Theorem 2.6.1 (b), we follow the plan mentioned last lecture, and prove the following. 2 Feb. 14:30

**Theorem 2.6.4.** If  $T$  is a maximal consistent  $\mathcal{L}$ -theory with Henkin constants, then  $T$  has a model.

**Proof.** The model we build is called a “canonical model.” Let  $\mathcal{C}$  be the set of constants in  $\mathcal{L}$ , and define an equivalence relation  $\sim$  on  $\mathcal{C}$  by  $c \sim d$  if and only if  $c = d$  is in  $T$ .

**Claim.** The relation  $\sim$  on  $\mathcal{C}$  defined by  $c \sim d \Leftrightarrow c = d \in T$  is an equivalence relation.

<sup>2</sup>Which still has Henkin constants.

**Proof.** We check the axioms for being an equivalence relation.

- (a)  $c \sim c$  because  $c = c$  is in  $T$  by (A7) (a).<sup>a</sup>
- (b) If  $c \sim d$ , then  $c = d$  is in  $T$  so  $d = c$  is in  $T$  by (A7) (b), i.e.,  $d \sim c$ .
- (c) If  $c \sim d$  and  $d \sim e$ , then  $c = d$  and  $d = e \in T \Rightarrow c = e \in T$  by (A7) (c), so  $c \sim e$ .

⊗

<sup>a</sup>Otherwise,  $c \neq c$  is in  $T$  from the maximality, so  $T \vdash c \neq c$  with  $T \vdash c = c$ , so  $T$  would be inconsistent.

Let  $[c]$  be the equivalence class of  $c$ . Define an  $\mathcal{L}$ -structure  $\mathcal{M}$  with domain  $M = \mathcal{C} / \sim = \{[c] \mid c \in \mathcal{C}\}$ , with functions, relations, and constants defined as follows:

- (a)  $c^{\mathcal{M}} = [c]$ .
- (b)  $R^{\mathcal{M}}([c_1], \dots, [c_n])$  **true** if  $R(c_1, \dots, c_n)$  is in  $T$ . This is well-defined by (A7) (e).
- (c)  $f^{\mathcal{M}}([c_1], \dots, [c_n]) = [d]$  if  $f(c_1, \dots, c_n) = d$  is in  $T$ . Such a  $d$  exists because  $\exists x f(c_1, \dots, c_n) = x$ , i.e.,  $\neg \forall x f(c_1, \dots, c_n) \neq x$ , is in  $T$ .<sup>b</sup> If this is in  $T$ , then there is a Henkin constant  $d$  with  $f(c_1, \dots, c_n) = d$  in  $T$ . To show that this is well-defined, from (A7) (d), i.e.,

$$(t_1 = u_1 \wedge \dots \wedge t_n = u_n) \rightarrow f(t_1, \dots, t_n) = f(u_1, \dots, u_n).$$

So if  $[c_1] = [d_1], \dots, [c_n] = [d_n]$ , then  $c_1 = d_1, \dots, c_n = d_n$  are in  $T$ . So  $f(c_1, \dots, c_n) = f(d_1, \dots, d_n)$  is in  $T$  by (A7) (d). If  $a$  and  $b$  are constants such that  $f(c_1, \dots, c_n) = a$  and  $f(d_1, \dots, d_n) = b$  are in  $T$ , so  $a = b$  is in  $T$  by (A7) (c), i.e., the transitivity of  $=$ .

Now we need to show that  $\mathcal{M} \models T$ , i.e., we claim that

$$\mathcal{M} \models \varphi([c_1], \dots, [c_n]) \Leftrightarrow \varphi(c_1, \dots, c_n) \text{ is in } T.$$

We prove this by induction on terms and then formulas.

1. **Terms:** Show that  $t^{\mathcal{M}}([c_1], \dots, [c_n]) = [d]$  if and only if  $t(c_1, \dots, c_n) = d$  is in  $T$ .

- (a) If  $t$  is a constant  $e$ ,  $t^{\mathcal{M}}([c_1], \dots, [c_n]) = e^{\mathcal{M}} = [e]$ , and

$$[e] = t^{\mathcal{M}}([c_1], \dots, [c_n]) = [d] \Leftrightarrow [e] = [d] \Leftrightarrow e = d \text{ is in } T.$$

- (b) If  $t$  is  $x_i$ ,  $t^{\mathcal{M}}([c_1], \dots, [c_n]) = [c_i]$ . This is equal to  $[d]$  if and only if  $c_i = d$  is in  $T$ .

- (c) Suppose that  $t(x_1, \dots, x_n) = f(s_1(x_1, \dots, x_n), \dots, s_m(x_1, \dots, x_n))$ . Let

$$[d_i] = s_i^{\mathcal{M}}([c_1], \dots, [c_n]),$$

by the inductive hypothesis,  $d_i = s_i(c_1, \dots, c_n)$  is in  $T$ . Let  $[e] = f^{\mathcal{M}}([d_1], \dots, [d_m]) = t^{\mathcal{M}}([c_1], \dots, [c_n])$ . By the definition of  $f$ ,  $e = f(d_1, \dots, d_m)$  is in  $T$ . By (A7) (d),

$$e = f(s_1(c_1, \dots, c_n), \dots, s_m(c_1, \dots, c_n))$$

is in  $T$ . This is the direction ( $\Rightarrow$ ).

Now suppose that  $t(c_1, \dots, c_n) = e'$  is in  $T$ . We want to show that  $[e] = [e']$ , i.e.,  $e = e'$  is in  $T$ . Since  $e = t(c_1, \dots, c_n)$  is in  $T$ , and  $e' = t(c_1, \dots, c_n)$  is in  $T$ . By (A7) (c),  $e = e'$  is in  $T$ , so  $[e'] = [e] = t^{\mathcal{M}}([c_1], \dots, [c_n])$ . This is ( $\Leftarrow$ ).

2. **Formulas:** Show that  $\mathcal{M} \models \varphi([c_1], \dots, [c_n])$  if and only if  $\varphi(c_1, \dots, c_n)$  is in  $T$ .<sup>c</sup>

- (a) If  $\varphi$  is  $s(x_1, \dots, x_n) = t(x_1, \dots, x_n)$ :

( $\Rightarrow$ ) If  $\mathcal{M} \models s([c_1], \dots, [c_n]) = t([c_1], \dots, [c_n])$ ,

$$s^{\mathcal{M}}([c_1], \dots, [c_n]) = t^{\mathcal{M}}([c_1], \dots, [c_n]).$$

Let  $[d]$  be this element equal to the above, so  $d = s(c_1, \dots, c_n)$  and  $d = t(c_1, \dots, c_n)$  are in  $T$  so  $\underbrace{s(c_1, \dots, c_n) = t(c_1, \dots, c_n)}_{\varphi(c_1, \dots, c_n)}$  is in  $T$  by (A7) (c).

( $\Leftarrow$ ) If  $s(c_1, \dots, c_n) = t(c_1, \dots, c_n)$  is in  $T$ , let

$$[d] = s^{\mathcal{M}}([c_1], \dots, [c_n]) \text{ and } [e] = t^{\mathcal{M}}([c_1], \dots, [c_n]),$$

so  $d = s(c_1, \dots, c_n)$  and  $e = t(c_1, \dots, c_n)$  are in  $t$ , so  $d = e$  is in  $t$ , and  $[e] = [d]$ .

(b) If  $\varphi$  is  $R(t_1(\bar{x}), \dots, t_m(\bar{x}))$ : Let  $[d_i] = t_i^{\mathcal{M}}([c_1], \dots, [c_n])$ ,

$$\begin{array}{ccc} R^{\mathcal{M}}([d_1], \dots, [d_m]) \text{ is true} & \Longleftrightarrow & R(d_1, \dots, d_m) \text{ is in } T \\ \Updownarrow & & \Updownarrow \\ R^{\mathcal{M}}(t_1^{\mathcal{M}}[\bar{c}], \dots, t_m^{\mathcal{M}}[\bar{c}]) \text{ is true} & & R(t_1(\bar{c}), \dots, t_m(\bar{c})) \text{ is in } T \\ \Updownarrow & & \\ \mathcal{M} \models \varphi([c_1], \dots, [c_n]) & & \end{array}$$

(c) If  $\varphi$  is  $\neg\psi$ : Then

$$\mathcal{M} \models \varphi(\bar{c}) \Leftrightarrow \mathcal{M} \not\models \psi(\bar{c}) \Leftrightarrow \psi(\bar{c}) \text{ is not in } T \Leftrightarrow \varphi(\bar{c}) \text{ is in } T$$

where the last  $\Leftrightarrow$  follows from the fact that  $T$  is maximal and consistent.

(d) If  $\varphi$  is  $\psi \rightarrow \theta$ :

- If  $\psi(\bar{c}) \rightarrow \theta(\bar{c})$  is in  $T$ : then if  $\psi(\bar{c})$  is in  $T$ , then  $\theta(\bar{c})$  is in  $T$  by (MP). then by the induction hypotheses, if  $\mathcal{M} \models \psi(\bar{c})$ , then  $\mathcal{M} \models \theta(\bar{c})$ .
- If  $\mathcal{M} \models \psi(\bar{c}) \rightarrow \theta(\bar{c})$ : then either  $\mathcal{M} \models \theta(\bar{c})$  or  $\mathcal{M} \models \neg\psi(\bar{c})$ . So either
  - i.  $\theta(\bar{c})$  is in  $T$ : by (A1),  $\theta(\bar{c}) \rightarrow (\psi(\bar{c}) \rightarrow \theta(\bar{c}))$ , so  $\psi(\bar{c}) \rightarrow \theta(\bar{c})$  is in  $T$ .
  - ii.  $\neg\psi(\bar{c})$  is in  $T$ :  $T \cup \{\psi(\bar{c})\}$  is now inconsistent, so  $T \cup \{\psi(\bar{c})\} \vdash \theta(\bar{c})$ . From the deductive theorem,  $T \vdash \psi(\bar{c}) \rightarrow \theta(\bar{c})$ . Because  $T$  is maximal and consistent,  $\psi(\bar{c}) \rightarrow \theta(\bar{c})$  is in  $T$ .

<sup>b</sup>Otherwise,  $\forall x f(c_1, \dots, c_n) \neq x$  is in  $T$ . By (A4),  $f(c_1, \dots, c_n) \neq f(c_1, \dots, c_n)$  is in  $T$ , contradicts to (A7) (a).

<sup>c</sup>In particular, for a sentence  $\varphi$ ,  $\mathcal{M} \models \varphi \Leftrightarrow \varphi$  is in  $T$ , and so  $\mathcal{M} \models T$ .

## Lecture 10: Introduction to Model Theory

Let's start by finishing the proof of Theorem 2.6.4.

7 Feb. 14:30

**Proof of Theorem 2.6.4 (Continued).** There's one final case left:

- (e) If  $\varphi$  is  $\forall x \psi(x, \bar{y})$ : Because  $T$  has Henkin constants, there is  $d$  such that  $\neg\forall x \psi(x, \bar{c}) \rightarrow \neg\psi(d, \bar{c})$  is in  $T$ .
- If  $\varphi(c_1, \dots, c_n)$  is not in  $T$ , i.e.,  $\forall x \psi(x, \bar{c})$  is in  $T$ , then since  $T$  is maximal,  $\neg\forall x \psi(x, \bar{c})$  is in  $T$ . So by (MP),  $\neg\psi(d, \bar{c})$  is in  $T$ . So,  $\mathcal{M} \models \neg\psi([d], [\bar{c}])$  by induction hypotheses, hence  $\mathcal{M} \models \neg\forall x \psi(x, [\bar{c}])$ , i.e.,  $\mathcal{M} \not\models \varphi([\bar{c}])$ .
  - If  $\mathcal{M} \models \varphi([\bar{c}])$ , then  $\mathcal{M} \models \forall x \psi(x, [\bar{c}])$ , so there is  $[e]$  such that  $\mathcal{M} \models \neg\psi([e], [\bar{c}])$ . Hence,  $\neg\psi(e, \bar{c})$  is in  $T$ . Suppose for a contradiction that  $\varphi(\bar{c})$ , i.e.,  $\forall x \psi(x, \bar{c})$  is in  $T$ , by (A4),  $\forall x \psi(x, \bar{c}) \rightarrow \psi(e, \bar{c})$ , so  $\psi(e, \bar{c})$  is in  $T$  by maximality and by consistency. But then  $T$  is inconsistent, a contradiction  $\nmid$  Hence  $\varphi(\bar{c})$  is not in  $T$ .

Thus,  $\mathcal{M} \models T$ , so  $T$  is satisfiable, proving the theorem. ■

**Remark.** We see that when proving the above, when we talk about  $\mathcal{M}$ , the witness comes for free, while for  $T$ , we need **Henkin constants** for getting a witness.

Now, we can complete the proof of **completeness theorem** by putting everything together.

**Claim.** The **completeness theorem (b)** holds.

**Proof.** We see that

1. **Theorem 2.6.3:** There is a **consistent**  $T^* \supseteq T$  and  $\mathcal{L}^*$ -**theory** (with  $\mathcal{L}^* \supseteq \mathcal{L}$ ) and  $T^*$  has **Henkin constants**.
2. **Theorem 2.5.3:** There is a **maximal consistent**  $\mathcal{L}^*$ -**theory**  $T^{**} \supseteq T^*$ , where  $T^{**}$  still has **Henkin constants**.
3. **Theorem 2.6.4:**  $T^{**}$  has a **model**  $\mathcal{M}^*$  an  $\mathcal{L}^*$ -**structure**. Let  $\mathcal{M}$  be the **reduct** of  $\mathcal{M}^*$  to an  $\mathcal{L}$ -**structure**.

Hence,  $\mathcal{M} \models T$ . ⊛

**As previously seen** (Problem set 1). Let  $\mathcal{L}^* \supseteq \mathcal{L}$ . If  $\mathcal{M}^*$  is an  $\mathcal{L}^*$ -**structure**, then by ignoring the **interpretation** of the symbols in  $\mathcal{L}^* - \mathcal{L}$ , we get an  $\mathcal{L}$ -**structure**  $\mathcal{M}$ .

**Notation** (Reduct).  $\mathcal{M}$  is a *reduct* of  $\mathcal{M}^*$ .

**Notation** (Expansion).  $\mathcal{M}^*$  is an *expansion* of  $\mathcal{M}$ .

**Remark.** We see that  $\vdash$  and  $\models$  are the same.

### 2.6.3 Consequences of Completeness Theorem

#### Size of Models

Now, let's step back and look at the proof of the **completeness theorem**, and ask the following.

**Problem.** When we did the **Henkin** construction of  $\mathcal{M}^* \models T^{**}$ , how big was  $M$ ?

This can be answered by the following.

**Theorem 2.6.5.** If  $T$  is a **satisfiable**  $\mathcal{L}$ -**theory**, then it has a **model** of size at most  $|\mathcal{L}| + \aleph_0$ .

**Proof.** Since  $|M| \leq |\mathcal{L}^*|$  since  $\mathcal{M} = \mathcal{C} / \sim$ , and in step one,  $|\mathcal{L}^*| \leq |\mathcal{L}| + \aleph_0$ , so  $|M| \leq |\mathcal{L}| + \aleph_0$ . ■

**Example.** Let  $\mathcal{L} = \{f\}$ ,  $T$  says that  $f$  is injective but not surjective.

**Example.** Let  $\mathcal{L} = \{\leq\}$ ,  $T$  says that  $\leq$  is a linear order with no greatest element.

**Example.** Let  $\mathcal{L} = \emptyset$ ,  $T$  says that there are at least  $n$  elements for each  $n$ .

#### Single Stroke and Double Stroke Style Deduction

As previously seen,  $\vdash$  and  $\models$  are actually  $\vdash_{\mathcal{L}}^a$  and  $\models_{\mathcal{L}}^b$

<sup>a</sup>Proofs can only use  $\mathcal{L}$ -formulas.

<sup>b</sup>Only looking at  $\mathcal{L}$ .

**Remark.** Suppose  $\mathcal{L} \supseteq \mathcal{L}_0$ , and  $\Gamma$  a set of  $\mathcal{L}_0$ -sentences,  $\varphi$  on  $\mathcal{L}_0$ -sentence.

- (a)  $\Gamma \models_{\mathcal{L}_0} \varphi \Leftrightarrow \Gamma \models_{\mathcal{L}_1} \varphi$ .
- (b)  $\Gamma \vdash_{\mathcal{L}_0} \varphi \Leftrightarrow \Gamma \vdash_{\mathcal{L}_1} \varphi$ .

**Proof.** (a) and (b) are equivalent by the **completeness theorem**, and we prove (a).

Suppose  $\Gamma \models_{\mathcal{L}_0} \varphi$ . Suppose  $\mathcal{M}_1$  is an  $\mathcal{L}_1$ -structure such that  $\mathcal{M}_1 \models \Gamma$ . Let  $\mathcal{M}_0$  be the **reduct** of  $\mathcal{M}_1$  to  $\mathcal{L}_0$  and  $\mathcal{M}_0 \models \Gamma$ , so  $\mathcal{M}_0 \models \varphi$ , then  $\mathcal{M}_1 \models \varphi$ , thus  $\Gamma \models_{\mathcal{L}_1} \varphi$ .

Now, suppose  $\Gamma \models_{\mathcal{L}_1} \varphi$ . Suppose  $\mathcal{M}_0$  is an  $\mathcal{L}_0$ -structure with  $\mathcal{M}_0 \models \Gamma$ . Expand  $\mathcal{M}_0$  to an  $\mathcal{L}_1$ -structure  $\mathcal{M}_1$  in any way.  $\mathcal{M}_1 \models \Gamma$ , so  $\mathcal{M}_1 \models \varphi$ . Thus,  $\mathcal{M}_0 \models \varphi$ , so  $\Gamma \models_{\mathcal{L}_0} \varphi$ . \*

What is important about the **proof system**?

- (1) **Soundness** and **completeness**,  $\vdash \Leftrightarrow \models$ .
- (2) **Proofs** are finite, and use only finitely many hypotheses  $\Rightarrow$  **compactness**.

## Computational Properties

Consider the following.

**Definition 2.6.3 (Computably enumerable).** A set is *computably enumerable* or *computable listable* if there is a program that lists out its elements.

If  $\mathcal{L}$  is finite, or computable (complete list of symbols and their arities), we have the following.

- (a) We can compute with **formulas**.
- (b) Given a **formula**, it's computable to check whether it's a **logical axiom**.
- (c) It's computable to check whether a **proof** is valid.
- (d) If  $\Gamma$  is a **computably enumerable** set of **sentences**,  $\{\varphi: \Gamma \vdash \varphi\}$  is also **computably enumerable**.<sup>3</sup>
- (e) There is no program that given  $\varphi$  can decide whether  $\vdash \varphi$  at least for  $\mathcal{L} = \{E\}$ ,  $E$  binary.

<sup>3</sup>We can list out all the valid **proofs** from  $\Gamma$  of any  $\varphi$ .

# Chapter 3

## The Beginning of Model Theory

We now discuss various properties of the [models](#) of some [theories](#) in our interest. In particular, we care about the size of the [models](#), and how different [models](#) with different size relate to each other.

### 3.1 Complete Theories

Let's start with a proposition.

**Proposition 3.1.1.** Let  $T$  be an  $\mathcal{L}$ -theory with an infinite [model](#), and let  $\kappa$  be an infinite cardinal with  $\kappa \geq |\mathcal{L}|$ . Then  $T$  has a [model](#) of cardinality  $\kappa$ .

**Proof.** Let  $\mathcal{C}$  be a set of  $\kappa$ -many new constants, and let  $\mathcal{L}^* = \mathcal{L} \cup \mathcal{C}$ . Let

$$T^* = T \cup \{c \neq d \mid c, d \in \mathcal{C} \text{ distinct}\}.$$

If  $\mathcal{M} \models T^*$ , then  $|M| \geq \kappa$ ; also, if  $T^*$  is [satisfiable](#), it has a [model](#) of size at most  $|\mathcal{L}^*| = \kappa$  since

$$\kappa = |\mathcal{C}| \leq |\mathcal{L}^*| \leq |\mathcal{C}| + |\mathcal{L}| \leq \kappa + \kappa = \kappa$$

from [Theorem 2.6.5](#). Hence, if  $T^*$  is [satisfiable](#), it has a [model](#)  $\mathcal{M}$  with  $|M| = \kappa$ .

**Claim.**  $T^*$  is [satisfiable](#).

**Proof.** It's enough to show that every finite  $\Gamma \subseteq T^*$  is [satisfiable](#) from the [compactness theorem](#). Let  $\mathcal{M}$  be infinite, and  $\Gamma \subseteq T^*$  finite, then we can write

$$\Gamma \subseteq T \cup \{c_i \neq c_j \mid i, j = 1, \dots, n, i \neq j\}$$

for  $c_1, \dots, c_n \in \mathcal{C}$  since only finitely many  $c_i$  are involved. Without loss of generality,  $\Gamma = T \cup \{c_i \neq c_j \mid i, j = 1, \dots, n, i \neq j\}$ . Pick  $a_1, \dots, a_n \in M$ , distinct, we then turn  $\mathcal{M}$  into an  $\mathcal{L}^*$ -structure  $\mathcal{M}^*$  with  $c_i^{\mathcal{M}^*} = a_i$ ,<sup>a</sup> resulting in  $\mathcal{M}^* \models \Gamma$ . ⊕

<sup>a</sup>And each other  $d \in \mathcal{C}$  with  $d^{\mathcal{M}^*} = a_1$ .

■

## Lecture 11: Algebraically Closed Fields

### 3.2 A Detour to Algebraically Closed Fields

9 Feb. 14:30

[Algebraically closed fields](#) are a great example of a *tame* theory (as opposed to e.g.,  $\mathbb{N}$ , which are not *tame*). We detour to discuss some important and related definitions for the future discussion.

### 3.2.1 Rings

All rings  $R$  we refer to will be commutative.

**Definition 3.2.1 (Ideal).** Let  $R$  be a ring. An *ideal*  $I$  of  $R$  is a set  $I \subseteq R$  such that

- (a)  $0 \in I$ ;
- (b) if  $a, b \in I$ , then  $a + b \in I$ ;
- (c) if  $a \in I$  and  $r \in R$ ,  $ra \in I$ .

**Intuition.** An *ideal* is trying to act as a set of “zeros” (in order to be further mod out).

**Definition 3.2.2 (Proper).** An *ideal* is *proper* if  $1 \notin I$ , equivalently,  $I \neq R$ .

**Definition.** Let  $I$  be a *proper ideal*.

**Definition 3.2.3 (Radical).**  $I$  is *radical* if  $a^n \in I$ , then  $a \in I$ .

**Definition 3.2.4 (Prime).**  $I$  is *prime* if  $ab \in I$ , then  $a \in I$  or  $b \in I$ .

**Definition 3.2.5 (Maximal).**  $I$  is *maximal* if there is no *proper ideal*  $J \supsetneq I$ .

**Remark.** *Maximal*  $\supseteq$  *Prime*  $\supseteq$  *Radical*.

**Definition 3.2.6 (Polynomial ring).** Let  $R$  be a ring. Then  $R[x_1, \dots, x_n]$  is the *polynomial ring* with coefficients in  $R$  on indeterminates  $x_1, \dots, x_n$ .

**Example.** Let  $K$  be a field,  $S \subseteq K^n$ , and  $I \subseteq K[x_1, \dots, x_n]$  defined as

$$I = \{f(\bar{x}) \mid f(\bar{s}) = 0 \text{ for all } \bar{s} \in S\}.$$

Then  $I$  is a *radical ideal*.

**Definition 3.2.7 (Ideal generation).** Let  $R$  be a ring. The *ideal*  $I$  *generated* by the set  $\{x_1, \dots, x_n \in R\}$ , denoted as  $I = (x_1, \dots, x_n)$ , is given by

$$I = \{r_1x_1 + \dots + r_nx_n \mid r_i \in R\}.$$

**Intuition.** The *ideal generated* by  $\{x_i\}$  is the “smallest” *ideal* containing all  $x_i$ ’s.

**Definition 3.2.8 (Principal ideal).** An *ideal* is a *principal ideal* if it’s *generated* by a single element.

**Definition 3.2.9 (Principal ideal ring).** A ring  $R$  is a *principal ideal ring* if all its *ideals* are *principal*.

**As previously seen** (Zero divisor). If  $a, b \neq 0$ , but  $ab = 0$ , then  $a$  and  $b$  are *zero divisors* of the ring  $R$ .



**Definition 3.2.10** (Integral domain). A nontrivial ring with no **zero divisors** is called an *integral domain*.<sup>a</sup>

<sup>a</sup>Some authors will just call *domain*.

**Definition 3.2.11** (Principal ideal domain). An **integral domain** where all **ideals** are **principal** is called a *principal ideal domain* or *PID*.

**Theorem 3.2.1.**  $K[x]$  is a **PID**, i.e., every **ideal** is **generated** by one element as  $I = (f(x)) = \{g(x)f(x) \mid g(x) \in K[x]\}$ .<sup>a</sup>

<sup>a</sup>It's clear that  $K[x]$  is an **integral domain**.

**Proof.** We can let  $g$  be the polynomial of the least degree in  $I$ . Then for any other  $h \in I$ , by long division,  $h = gs + r$ , with  $\deg(r) < \deg(g)$ . But then  $r = h - gs \in I$ , so if  $r$  has lower degree than  $g$ ,  $r = 0$ , hence  $h = gs \in (g)$ . ■

If it's too much to ask for an **ideal generated** by a single element, then we might as well consider the finite case.

**Definition 3.2.12** (Noetherian ring). A ring  $R$  is *Noetherian* if every **ideal**  $I$  of  $R$  is finitely generated.

**Remark.** Equivalently, there is no infinite proper ascending chain of **ideals**.

**Theorem 3.2.2** (Hilbert basis theorem). If  $R$  is a **Noetherian ring**, then  $R[x]$  is also **Noetherian**. In particular,  $K[x_1, \dots, x_n]$  is **Noetherian** and so every **ideal** in  $K[x_1, \dots, x_n]$  is finitely generated.

**As previously seen** (Ring homomorphism). Let  $R, S$  be rings. A *ring homomorphism*  $\varphi: R \rightarrow S$  is a map satisfies

- (a)  $\varphi(x +_R y) = \varphi(x) +_S \varphi(y)$  for  $x, y \in R$ ;
- (b)  $\varphi(x \times_R y) = \varphi(x) \times_S \varphi(y)$  for  $x, y \in R$ ;
- (c)  $\varphi(1_R) = 1_S$ .

**Theorem 3.2.3.** If  $\alpha: R \rightarrow S$  is a **ring homomorphism**, then  $\ker \alpha$  is an **ideal** of  $R$ , and the induced map  $\bar{\alpha}: R / \ker \alpha \rightarrow S$  is injective.

**Theorem 3.2.4.** Let  $R$  be a ring, and  $I$  an **ideal** of  $R$ .<sup>a</sup>

- (a)  $R / I$  is an **integral domain** if and only if  $I$  is a **prime**.
- (b)  $R / I$  is a field if and only if  $I$  is **maximal**.

<sup>a</sup>Then  $\pi: R \rightarrow R / I$  is a **ring homomorphism** with kernel  $I$ .

### 3.2.2 Field Extensions

Now, we can talk about **field extension**.

**Definition 3.2.13** (Field extension). If  $K \subseteq L$  is a subfield of  $L$ , we call  $L / K$  a *field extension*.

Given a **field extension**  $L / K$ , then we have that  $L$  is a  $K$ -vector space, which suggests the following natural notion.

**Definition 3.2.14 (Degree).** The *degree*  $[L: K]$  of  $L / K$  is the dimension of the  $K$ -vector space  $L$ .

**Notation** (Finite extension). If  $[L: K]$  is finite, then we say  $L / K$  is a *finite extension*.

**Example.**  $\mathbb{C}$  is a [field extension](#) over  $\mathbb{R}$  with  $[\mathbb{C}: \mathbb{R}] = 2$ .

**Proof.** Since  $\mathbb{C}$  is an  $\mathbb{R}$ -vector space with basis  $\{1, i\}$ . \*

**Example.**  $\mathbb{Q}(\sqrt{2})$  is a [field extension](#) over  $\mathbb{Q}$  with  $[\mathbb{Q}(\sqrt{2}): \mathbb{Q}] = 2$ .

**Proof.** Since  $\mathbb{Q}(\sqrt{2})$  is a  $\mathbb{Q}$ -vector space with basis  $\{1, \sqrt{2}\}$ . \*

The following is the powerful way to calculate the [degree](#) of a [field extension](#) if it can be constructed by a “tower” of [field extensions](#).

**Theorem 3.2.5.** If  $M / L$  and  $L / K$  are [field extensions](#), then  $[M: K] = [M: L][L: K]$ .

### 3.2.3 Algebraically Closed Fields

We care about [field extensions](#)  $L / K$  that are [algebraic](#). This start from defining what does it mean by a single element  $a \in L$  is [algebraic](#) over  $K$ .

**Definition.** Let  $L / K$  be a [field extension](#), and  $a \in L$ .

**Definition 3.2.15 (Algebraic).** If there is a non-zero  $f(x) \in K[x]$  such that  $f(a) = 0$ , then  $a$  is *algebraic* over  $K$ .

**Definition 3.2.16 (Transcendental).** If  $a$  is not [algebraic](#), then it is *transcendental* over  $K$ .

**Definition 3.2.17 (Minimal polynomial).** If  $a$  is [algebraic](#) over  $K$ , there is a non-zero, monic<sup>a</sup>  $f(x) \in K[x]$  of least degree such that  $f(a) = 0$  which we call the *minimal polynomial* of  $a$  over  $K$ .

<sup>a</sup>This is a common practice.

**Intuition.** An [algebraic](#) number  $a$  is the root of some polynomials  $f$  in this [polynomial ring](#), and we can find the [minimal](#) such  $f$ .

**As previously seen** (Irreducible). A non-zero non-unit of an [integral domain](#)  $R$  is *irreducible* if it cannot be written as the product of two non-units.

**Note.** A [minimal polynomial](#) is [irreducible](#).

**Remark.** If  $f(x)$  is a [minimal polynomial](#), then  $(f(x)) = \{g(x) \in K[x] \mid g(a) = 0\}$ .

**Example.** Consider a [field extension](#)  $\mathbb{R} / \mathbb{Q}$  with  $a = \sqrt{2} \in \mathbb{R}$ . Then the [minimal polynomial](#) is  $f(x) = x^2 - 2$ .

**Theorem 3.2.6.** Let  $L / K$  be a [field extension](#) and  $a \in L$ , then  $a$  is [algebraic](#) over  $K$  if and only

if  $n = [K(a) : K] < \infty$ . Furthermore, if  $a$  is **algebraic** over  $K$ , then  $n$  is the degree of the **minimal polynomial** of  $a$ , and  $1, a, \dots, a^{n-1}$  is a basis for  $K(a)$  as a  $K$ -vector space.

**Proof idea.** Think about  $f(a) = a^n + r_{n-1}a^{n-1} + \dots + r_1a + r_01 = 0$ . ■

The following example illustrates how can we combine **Theorem 3.2.5** and **Theorem 3.2.6**,

**Example.** Let  $f(x) = x^2 - 2$ ,  $\mathbb{Q}(\sqrt{2}) = \{a1 + b\sqrt{2} \mid a, b \in \mathbb{Q}\}$ .



**Theorem 3.2.7.** Let  $L/K$  be a **field extension**,  $a \in L$ , and  $f(x) \in K[x]$  be the **minimal polynomial** of  $a$  over  $K$ .

(a)  $K[x]/(f(x)) \cong K(a)$ .<sup>a</sup>

(b) If  $b \in L$  has the same **minimal polynomial** as  $a$ , then  $K(a) \cong K[x]/(f(x)) \cong K(b)$ .

<sup>a</sup>Let  $x \in K[x]$ , then  $\bar{x} = x + (f(x)) \in K[x]/(f(x))$ , i.e.,  $\bar{x}$  is a root of  $f$ , hence the isomorphism is given by  $\bar{x} \mapsto a$ .

**Example.** Let  $a = \sqrt{2}, b = -\sqrt{2}$ , and  $f(x) = x^2 - 2$  with  $K = \mathbb{Q}$ . Then

$$\begin{aligned} \mathbb{Q}(\sqrt{2}) &\cong \mathbb{Q}[x]/(x^2 - 2) \cong \mathbb{Q}(-\sqrt{2}); \\ a + b\sqrt{2} &\mapsto [a + bx] \mapsto a - b\sqrt{2}. \end{aligned}$$

Then, it's now natural to talk about a **algebraic extension**.

**Definition 3.2.18 (Algebraic extension).** Let  $L/K$  be a **field extension**. Then  $L$  is an *algebraic extension* of  $K$  if all  $a \in L$  are **algebraic** over  $K$ .

If  $a$  is **algebraic** over  $K$ , then  $K(a)/K$  is **algebraic**: If  $b \in K(a)$ , then  $K(b) \subseteq K(a)$ , so  $[K(b) : K] \leq [K(a) : K] < \infty$ , so  $b$  is **algebraic** over  $K$ .

**Theorem 3.2.8.** If  $M/L$  and  $L/K$  are **algebraic extensions**, then  $M/K$  is an **algebraic extension**.

**Proof.** Let  $a \in M$ , and let  $b_1, \dots, b_n \in L$  be the coefficients of the **minimal polynomial** of  $a$  over  $L$ . Then  $b_1, \dots, b_n$  are **algebraic** over  $K$ . Since

$$\begin{aligned} [K(a) : K] &\leq [K(a, b_1, \dots, b_n) : K] \\ &= [K(a, b_1, \dots, b_n) : K(b_1, \dots, b_n)] \cdot [K(b_1, \dots, b_n) : K(b_2, \dots, b_n)] \cdots [K(b_n) : K]. \end{aligned}$$

Since each of these is a finite **extension**, so  $[K(a) : K] < \infty$ . ■

**Definition 3.2.19 (Algebraically closed).** A field  $L$  is *algebraically closed* if any non-constant  $f(x) \in L[x]$  has a root in  $L$ .

**Definition 3.2.20 (Algebraic closure).** If  $L/K$ , then  $L$  is an *algebraic closure* of  $K$  if  $L$  is **algebraically closed** and an **algebraic extension** of  $K$ .

**Remark.** Over an algebraically closed field  $K$ , any polynomial  $f(x) \in K[x]$  factors completely into  $f(x) = (x - a_1) \cdots (x - a_n)$  for  $n = \deg f$ .

**Example.**  $\mathbb{C}$  is algebraically closed, while  $\mathbb{R}$  is not.

**Example.**  $\mathbb{C}$  is the algebraic closure of  $\mathbb{R}$ , and  $[\mathbb{C} : \mathbb{R}] = 2$ .

**Example.**  $\mathbb{Q}^{\text{alg}} = \{a \in \mathbb{C} \mid a \text{ is algebraic over } \mathbb{Q}\}$  is the algebraic closure of  $\mathbb{Q}$ .

If  $L$  is algebraically closed, any  $f(x) \in L[x]$  factors completely as  $f(x) = (x - a_1) \cdots (x - a_n)$  and  $a_1, \dots, a_n$  are the only roots of  $f$ .

**Theorem 3.2.9.** Every field  $K$  has an algebraic closure. If  $L/K$  and  $M/K$  are algebraic closures over  $K$ , then  $L \cong_K M$ .<sup>a</sup>

<sup>a</sup>There exists  $\alpha: L \rightarrow M$  such that  $\alpha(a) = a$  for  $a \in K$ .

**Proof.** First, we show the existence. Let  $f_1, f_2, \dots$  be (non-constant) polynomials over  $K$ . Start with  $K = K_0$ , let  $g_1(x)$  be an irreducible factor of  $f_1(x)$  and consider

$$K_1 := K_0[x] / (g_1(x)).$$

Since  $g_1$  is irreducible,  $(g_1(x))$  is maximal, so  $K_1$  is a field with a root of  $f_1$ . Now, we build

$$K = K_0 \subseteq K_1 \subseteq K_2 \subseteq \dots \subseteq K^* = \bigcup_i K_i$$

in the same way such that  $K_i$  contains a root of  $f_i(x)$ . Since any  $f(x) \in K$  has a root in  $K^*$ , so  $K^*/K$  is algebraic. Now, we do the same construction for  $K^*$  to get

$$K \subseteq K^* \subseteq K^{**} \subseteq K^{***} \subseteq \dots \subseteq L = \bigcup K^{*...},$$

then  $L$  is algebraically closed since any non-constant polynomial with coefficients in  $L$  actually has coefficients in one of the  $K^{*...}$ , so it has a root in the next field. Now we prove the uniqueness.

**Lemma 3.2.1.** An algebraically closed field  $L$  has no proper algebraic extensions  $M$ .

**Proof.** If  $a \in M$  is algebraic over  $L$  for some  $M$ , the minimal polynomial  $f(x)$  of  $a$  factors completely (irreducible), so  $f(x) = x - r$  for  $r \in L$  with  $f(a) = 0$ , i.e.,  $a = r$ , so  $M = L$ . ■

**Lemma 3.2.2.** Let  $L/K$  algebraic,  $M/K$  algebraically closed. Then there is an embedding  $\alpha: L \rightarrow M$  fixing  $K$ .

**Proof.** Consider the case that  $L = K(a)$ <sup>a</sup> with  $a$  algebraic over  $K$ , and let  $f(x)$  be the minimal polynomial of  $a$  over  $K$ . Then there is a root  $b \in M$  of  $f$  with  $K(a) \cong K[x]/(f) \cong K(b) \subseteq L$  from Theorem 3.2.7. Let this isomorphism be our  $\alpha$ . ■

<sup>a</sup>Once this is done, repeat iteratively and get the general case by using Zorn's lemma or transfinite induction.

Hence, if  $L/K$  and  $M/K$  are algebraic closures over  $K$ , there is an embedding  $\alpha: L \rightarrow M$  over  $K$ .

Finally, since  $M/\alpha(L)$  is an algebraic extension, and  $\alpha(L) \cong L$  is algebraically closed, by Lemma 3.2.1,  $M = \alpha(L)$ , so  $\alpha$  is an isomorphism  $L \rightarrow M$  over  $K$ . ■

## Lecture 12: The ACF Theory and Categorical

**Definition 3.2.21 (Characteristic).** A field  $F$  has finite *characteristic*  $p > 0$  if  $\underbrace{1 + \dots + 1}_{p \text{ times}} = 0$ .

**Remark.**  $p$  is always prime, otherwise,  $F$  has *characteristic*  $p = 0$ , i.e.,  $1 + \dots + 1 \neq 0$ , always.

The following notion comes up naturally.

**Definition 3.2.22 (Prime field).** The *prime field*  $\mathbb{F}_p$  in *characteristic*  $p$  such that  $\mathbb{F}_p = \mathbb{Q}$  if  $p = 0$ ,  $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$  if  $p > 0$ .

**Definition 3.2.23 (Transcendence basis).** Let  $L/K$  be a *field extension*. A set  $S \subseteq L$  is called a *transcendence basis* of  $L/K$  if  $S$  is algebraically independent<sup>a</sup> and  $L$  is an *algebraic extension* of  $K(S)$ , i.e.,  $S$  is maximal.

<sup>a</sup>No  $a_1, \dots, a_n \in S$  have non-zero polynomial  $f(x_1, \dots, x_n) \in K[\bar{x}]$  with  $f(a_1, \dots, a_n) = 0$ .

**Remark.** Every *field extension* has a *transcendence basis* (and all *transcendence basis* have the same size).

**Proof.** On a combinatorial level, this is exactly the same as the proof that any two bases for a vector space have the same cardinality. \*

**Example.** Let  $K(t_1, \dots, t_n)$  be the fraction field of  $K[x_1, \dots, x_n]$ , then  $\{t_1, \dots, t_n\}$  is a *transcendence basis* for  $K(t_1, \dots, t_n)$  over  $K$ .

**Definition 3.2.24 (Transcendence degree).** The *transcendence degree* of  $L$  over  $K$  is the cardinality of any *transcendence basis*.

If we do not specify  $K$ , then  $K$  is the *prime field*  $K = \mathbb{F}_p$ .

**Theorem 3.2.10.** Any two *algebraically closed* fields of the same *characteristic*  $p$  and *transcendence degree* are isomorphic.

**Proof.** Let  $L, K$  be those fields, with *transcendence basis*  $S, T$  over  $\mathbb{F}_p$  with  $|S| = |T|$ .  $L$  is the *algebraic closure* of  $\mathbb{F}_p(S)$  and  $K$  is the *algebraic closure* of  $\mathbb{F}_p(T)$ . There is a bijection  $f: S \rightarrow T$ , and then  $f$  extends to  $\bar{f}: \mathbb{F}_p(S) \rightarrow \mathbb{F}_p(T)$  such that

$$\bar{f}\left(\frac{\sum_{\alpha} r_{\alpha} \bar{x}^{\alpha}}{\sum_{\alpha} s_{\alpha} \bar{x}^{\alpha}}\right) = \frac{\sum_{\alpha} r_{\alpha} f(\bar{x})^{\alpha}}{\sum_{\alpha} s_{\alpha} f(\bar{x})^{\alpha}},$$

where  $r_{\alpha}, s_{\alpha} \in \mathbb{F}_p$  and  $\bar{x}^{\alpha}$  is some monomial from  $S$ , e.g.,  $x_1^2 x_2$  for  $x_1, x_2 \in S$ .<sup>a</sup>

$\mathbb{F}_p(S)$  and  $\mathbb{F}_p(T)$  are the same (up to isomorphism), but the *algebraic closures* are unique from [Theorem 3.2.9](#), so  $K \cong L$  via an isomorphism extending  $\bar{f}$ . ■

<sup>a</sup> $\alpha$  can be thought as a tuple, in the case of  $x_1^2 x_2$ ,  $\alpha = (2, 1)$ .

The above proof actually shows more.

**Corollary 3.2.1.** If  $L/K$  and  $M/K$  are *field extensions* with *transcendence bases*  $S$  and  $T$ , and  $\alpha: S \rightarrow T$  is a bijection, then  $\alpha$  extends to an isomorphism  $L \cong_K M$ .

If we apply this inside a single *algebraically closed* field, we have the following.

**Theorem 3.2.11.** Let  $K$  be the *algebraic closure* of  $k$ , and  $L, M$  be subfields of  $K$  which *extend*  $k$ . Suppose that  $\alpha: M \rightarrow L$  is an isomorphism fixing  $k$ , then  $\alpha$  extends to an automorphism of  $K$ .

### 3.3 The ACF Theory

Finally, we are ready to introduce the [theory](#) we're going to study, which is called ACF. It turns out that the [models](#) of which are exactly the [algebraically closed](#) fields with nice properties we're going to discuss.

**Definition 3.3.1 (ACF).** ACF is the [theory](#) of [algebraically closed](#) fields consists of field axioms and [formulas](#) that for every  $n \geq 1$ ,

$$\forall a_0 \dots \forall a_n (a_n \neq 0 \rightarrow \exists b \ a_n b^n + a_{n-1} b^{n-1} + \dots + a_0 = 0).$$

**Remark.** The [models](#) of [ACF](#) are exactly the [algebraically closed](#) fields with the [language](#)  $\mathcal{L} = \mathcal{L}_{\text{ring}} = \{0, 1, +, -, \cdot\}$ .

**Notation** ( $\text{ACF}_p$ ). For a prime  $p > 0$ , let  $\text{ACF}_p := \text{ACF} \cup \{\underbrace{1 + \dots + 1}_p = 0\}$ .

**Notation** ( $\text{ACF}_0$ ). Let  $\text{ACF}_0 := \text{ACF} \cup \{\underbrace{1 + \dots + 1}_n \neq 0 \mid n \in \mathbb{N}\}$ .

**Definition 3.3.2 (Categorical).** Let  $\kappa$  be an infinite cardinal and  $T$  be an [L-theory](#). We say  $T$  is  $\kappa$ -categorical if for any  $\mathcal{M}, \mathcal{N} \models T$  of size  $\kappa$ , we have  $\mathcal{M} \cong \mathcal{N}$ .

**Definition 3.3.3 (Countably categorical).** If  $\kappa$  is countable, then  $T$  is *countably categorical*.

**Definition 3.3.4 (Uncountably categorical).** If  $\kappa$  is uncountable, then  $T$  is *uncountably categorical*.

We see that for being [uncountably categorical](#), we only need one uncountable  $\kappa$ .

**Example.**  $(\mathbb{Q}, \leq)$  is [countably categorical](#).

**Lemma 3.3.1.** If  $K$  has [transcendence degree](#)  $\lambda$ , then  $|K| = \lambda + \aleph_0$ .

**Proof.** Let  $K$  be [algebraic](#) over  $\mathbb{F}_p(S)$ , where  $S$  is a [transcendence basis](#) of size  $\lambda$ . By counting,  $|\mathbb{F}_p(S)| = \lambda + \aleph_0$ , so  $|\mathbb{F}_p(S)[x]| = \lambda + \aleph_0$ . But since each element of  $K$  satisfies some polynomials, and each polynomial has finitely many roots in  $K$ , so  $|K| = \lambda + \aleph_0$ . ■

**Theorem 3.3.1.** For each  $p$ ,  $\text{ACF}_p$  is  $\kappa$ -categorical for every uncountable  $\kappa$ .

**Proof.** Let  $L, K$  be a [model](#) of  $\text{ACF}_p$  for size  $\kappa$ . From [Theorem 3.2.10](#), if  $L, K$  have [transcendence degree](#)  $\kappa$ , then they are isomorphic. With the application of [Lemma 3.3.1](#), we're done. ■

**Example.**  $\mathbb{Q}^{\text{alg}}$ , the [algebraic closure](#) of  $\mathbb{Q}$ , has size  $\aleph_0$  with [transcendence degree](#) 0.

**Example.**  $\mathbb{Q}(t)^{\text{alg}}$ , the [algebraic closure](#) of  $\mathbb{Q}(t) \cong \mathbb{Q}(\pi)$ , has size  $\aleph_0$  with [transcendence degree](#) 1.

The above implies  $\mathbb{Q}^{\text{alg}} \not\cong \mathbb{Q}(t)^{\text{alg}}$ , which lead to the following.

**Remark.**  $\text{ACF}_0$  is not [countably categorical](#). The same with  $\text{ACF}_p$  for  $p > 0$ .

**Proof.** Notice that

$$\mathbb{Q}(t)^{\text{alg}} = \{z \in \mathbb{C} \mid z \text{ is algebraic over } \mathbb{Q}(\pi)\},$$

which is countable. With  $\mathbb{Q}^{\text{alg}} \not\cong \mathbb{Q}(t)^{\text{alg}}$ , we have the result.  $\circledast$

**Note.** ACF is not uncountably categorical.

**Theorem 3.3.2 (Vaught's test).** Let  $T$  be a satisfiable  $\mathcal{L}$ -theory with no finite models. If  $T$  is  $\kappa$ -categorical for some infinite  $\kappa \geq |\mathcal{L}|$ , then  $T$  is complete.

**Proof.** Suppose  $T$  is not complete, then we can pick  $\varphi$  with  $T \not\models \varphi$  and  $T \not\models \neg\varphi$ , i.e.,  $T \cup \{\varphi\}$  and  $T \cup \{\neg\varphi\}$  are satisfiable. By a consequence of the proof of completeness theorem (with a compactness argument),

- $T \cup \{\varphi\}$  has a model  $\mathcal{M}$  of size  $\kappa$ , and
- $T \cup \{\neg\varphi\}$  has a model  $\mathcal{N}$  of size  $\kappa$ .

But  $T$  is  $\kappa$ -categorical, so  $\mathcal{M} \cong \mathcal{N}$ , which is a contradiction  $\blacksquare$

**Corollary 3.3.1.**  $\text{ACF}_p$  is complete for each  $p$ .

**Proof.** Follows immediately by Theorem 3.3.1 and Vaught's test.  $\blacksquare$

The axioms for  $\text{ACF}_p$  completely determines all first-order facts about algebraically closed fields of characteristic  $p$ .

**Remark.**  $\{\varphi \mid \text{ACF} \models \varphi\}$  and  $\{\varphi \mid \text{ACF}_p \models \varphi\}$  can be listed computably.

**Proof.** Since the axioms for ACF or  $\text{ACF}_p$  can be listed computably.  $\circledast$

**Definition 3.3.5 (Decidable).** A theory  $T$  is *decidable* if there is a program that given  $\varphi$ , it determines whether  $T \models \varphi$  or  $T \not\models \varphi$ .

**Corollary 3.3.2.**  $\text{ACF}_p$  is decidable for each  $p$ .

**Proof.** Given  $\varphi$ , either  $\text{ACF}_p \models \varphi$  or  $\text{ACF}_p \models \neg\varphi$  since  $\text{ACF}_p$  is complete. By looking for a proof of  $\varphi$  and a proof of  $\neg\varphi$ , eventually we will find one, telling us whether  $\text{ACF}_p \models \varphi$ .  $\blacksquare$

**Corollary 3.3.3.** ACF is decidable.

**Proof.** Given  $\varphi$ , the algorithm simultaneously

- looks for a proof of  $\text{ACF} \vdash \varphi$ , and
- looks for  $p$  such that  $\text{ACF}_p \vdash \neg\varphi$  (hence  $\text{ACF} \not\models \varphi$ ).<sup>a</sup>

If  $\text{ACF} \models \varphi$ , then we will halt at the first case. Now suppose  $\text{ACF} \not\models \varphi$ , i.e., there is  $\mathcal{M} \models \text{ACF}$  such that  $\mathcal{M} \models \neg\varphi$ , and also, there is  $p$  such that  $\mathcal{M} \models \text{ACF}_p$ . Since  $\text{ACF}_p$  is complete,  $\text{ACF}_p \models \neg\varphi$ , so the search of the second case will halt, hence the whole search will eventually halt.  $\blacksquare$

<sup>a</sup>It might not be true that  $\text{ACF} \models \neg\varphi$ , we don't know.

## Lecture 13: Upward Löwenheim-Skolem Theorem

Another consequence of completeness is that since  $\mathbb{C} \models \text{ACF}_0$ , if  $K$  is any algebraically closed field of characteristic 0,  $\mathbb{C} \equiv K$ , i.e., we have the following. 16 Feb. 14:30

**Remark.** The sentences true of  $\mathbb{C}$  are exactly the same as the sentences true of any algebraically closed field.

Essentially, the idea is that if one proves an algebraic statement about the complex numbers by analytic techniques of complex analysis, then there will be a proof of the same algebraic statement using purely algebraic tools, which works in any algebraically closed field of characteristic 0. The compactness theorem also gives connections to fields of finite characteristic.

**Theorem 3.3.3 (Leftschetz principle).** Let  $\mathcal{L}$  be the language of rings. For an  $\mathcal{L}$ -sentence  $\varphi$ , the following are equivalent:

- (i)  $\varphi$  is true in  $\mathbb{C}$ ;
- (ii)  $\varphi$  is true in every algebraically closed field of characteristic 0;
- (iii)  $\varphi$  is true in some algebraically closed fields of characteristic 0;
- (iv) there is a number  $n$  such that  $\varphi$  is true in all algebraically closed fields of characteristic  $p > n$ ;
- (v) for each number  $n$ ,  $\varphi$  is true in all algebraically closed fields of characteristic  $p > n$ .

**Proof.** Let  $K \models \text{ACF}_0$ , then since it's complete, we know that  $K \models \varphi \Leftrightarrow \text{ACF}_0 \models \varphi$ , which proves the first three. Others are left as homework. ■

We can use the Leftschetz principle to prove the following.

**Theorem 3.3.4 (Ax-Grothendieck theorem).** Let  $f: \mathbb{C}^n \rightarrow \mathbb{C}^n$  be a polynomial map.<sup>a</sup> If  $f$  is injective, then it's surjective. More generally, this is true for any  $K \models \text{ACF}_p$  for any  $p$ .

<sup>a</sup>I.e.,  $f(\bar{x}) = (f_1(\bar{x}), \dots, f_n(\bar{x}))$  where  $f_1, \dots, f_n$  are polynomials.

**Proof.** The claim can be expressed by the sentences, so by Leftschetz principle, it's enough to prove that if for  $K = \overline{\mathbb{F}_p}$ , for each  $p > 0$ .

Let  $f: \overline{\mathbb{F}_p}^n \rightarrow \overline{\mathbb{F}_p}^n$  be an injective polynomial map and  $\bar{y} \in \overline{\mathbb{F}_p}^n$ . Then there is a finite subfield  $L \subseteq \overline{\mathbb{F}_p}$  which contains  $\bar{y}$  and the coefficients of  $f$ . Then,  $f$  restricts to an injective function  $L^n \rightarrow L^n$ , which is surjective because  $L^n$  is finite, so  $\exists \bar{x} \in L^n$  such that  $f(\bar{x}) = \bar{y}$ . ■

## 3.4 Up and Down

After introducing categorical, and seeing that ACF as an example, we now study something else: when does a model exists w.r.t. some original model, and what size is it?

### 3.4.1 Diagrams

One way to capture the structure in terms of theory is using the so-called “diagrams”.

**Definition.** Let  $\mathcal{M}$  be an  $\mathcal{L}$ -structure. Let  $\mathcal{L}_{\mathcal{M}} \supseteq \mathcal{L}$  be the expanded language with a new constant symbol  $\underline{a}$  for each  $a \in M$ .

**Definition 3.4.1 (Atomic diagram).** The atomic diagram of  $\mathcal{M}$  is the  $\mathcal{L}_{\mathcal{M}}$ -theory

$$\text{Diag}(\mathcal{M}) := \{\varphi(\underline{a}_1, \dots, \underline{a}_n) \mid \mathcal{M} \models \varphi(m_1, \dots, m_n) \text{ and } \varphi \text{ is atomic or negated of atomic}\}.$$

**Definition 3.4.2 (Elementary diagram).** The elementary diagram of  $\mathcal{M}$  is the  $\mathcal{L}_{\mathcal{M}}$ -theory

$$\text{Diag}_{\text{el}}(\mathcal{M}) := \{\varphi(\underline{a}_1, \dots, \underline{a}_n) \mid \mathcal{M} \models \varphi(m_1, \dots, m_n) \text{ and } \varphi \text{ an } \mathcal{L}\text{-formula}\}.$$



**Intuition.** Basically both  $\text{Diag}(\mathcal{M})$  and  $\text{Diag}_{\text{el}}(\mathcal{M})$  contain the information about the **structure** but in the form of a **theory**.

**Notation.** There's a canonical way of **expanding**  $\mathcal{M}$  to an  $\mathcal{L}_{\mathcal{M}}$ -**structure** with  $\underline{a}^{\mathcal{M}} := a$ , i.e., we write  $a$  for both the symbol and the element.

**Lemma 3.4.1.** Let  $\mathcal{N}$  be an  $\mathcal{L}_{\mathcal{M}}$ -**structure**.

- (a) If  $\mathcal{N} \models \text{Diag}(\mathcal{M})$  then, viewing  $\mathcal{N}$  as an  $\mathcal{L}$ -**structure**, there is an **embedding**  $f: \mathcal{M} \rightarrow \mathcal{N}$ .
- (b) If  $\mathcal{N} \models \text{Diag}_{\text{el}}(\mathcal{M})$ , then there is an **elementary  $\mathcal{L}$ -embedding** of  $\mathcal{M}$  into  $\mathcal{N}$ .

**Proof.** Take  $f(a) = \underline{a}^{\mathcal{N}}$ , then  $\mathcal{N} \models \text{Diag}(\mathcal{M})$  means exactly that  $f$  is an **embedding**, and  $\mathcal{N} \models \text{Diag}_{\text{el}}(\mathcal{M})$  means that  $f$  is an **elementary embedding**. ■

### 3.4.2 Upward Löwenheim-Skolem theorem

**Theorem 3.4.1** (Upward Löwenheim-Skolem theorem). Let  $\mathcal{M}$  be an infinite  $\mathcal{L}$ -**structure** and let  $\kappa$  be an infinite cardinal  $\kappa \geq |\mathcal{M}| + |\mathcal{L}|$ . Then there is an  $\mathcal{L}$ -**structure**  $\mathcal{N}$  of cardinality  $\kappa$  such that  $j: \mathcal{M} \rightarrow \mathcal{N}$  is **elementary**.

**Proof.**  $\text{Diag}_{\text{el}}(\mathcal{M})$  is **satisfiable** since  $\mathcal{M} \models \text{Diag}_{\text{el}}(\mathcal{M})$ , by **Proposition 3.1.1** it has a **model**  $\mathcal{N}$  of cardinality  $\kappa \geq |\mathcal{L}_{\mathcal{M}}| = |\mathcal{M}| + |\mathcal{L}|$ , so an **elementary embedding** exists  $\mathcal{M} \rightarrow \mathcal{N}$  by **Lemma 3.4.1**. ■

**Intuition.** The **upward Löwenheim-Skolem theorem** says that every **structure** is an **elementary sub-structure** of many much bigger **structures**.

**As previously seen.** Our very first application of the **compactness theorem**: in **the construction of the non-standard model of arithmetic**, we built  $\mathcal{N} \models \text{Th}(\mathbb{N})$  not **isomorphic** to  $\mathbb{N}$ , which is exactly like this. Every element of  $\mathbb{N}$  can already be expressed as a **term** of the form  $1 + \dots + 1$  without adding any new constants.

### 3.4.3 Downward Löwenheim-Skolem theorem

A “downward” version of the **upward Löwenheim-Skolem theorem** also exists, which says that big **models** contain smaller **elementary substructures**. This will take some more work to prove, so we first need a test for this.

**Proposition 3.4.1** (Tarski-Vaught test). Let  $\mathcal{M}$  be a **substructure** of  $\mathcal{N}$ . Then  $\mathcal{M}$  is an **elementary substructure** of  $\mathcal{N}$  if and only if for any **formula**  $\varphi(x, \bar{y})$  and  $\bar{a} \in M^n$ , if there is  $b \in N$  such that  $\mathcal{N} \models \varphi(b, \bar{a})$ , then there is  $c \in M$  such that  $\mathcal{N} \models \varphi(c, \bar{a})$ .

**Proof.** The forward direction follows from the fact that  $\mathcal{M}$  is an **elementary substructure**, so the **truth** of  $\exists x \varphi(x, \bar{y})$  is proved since  $\mathcal{M} \models \exists x \varphi(x, \bar{a})$  if and only if  $\mathcal{N} \models \exists x \varphi(x, \bar{a})$ .

For the backward direction, suppose the condition holds. We show that  $\mathcal{M} \models \varphi(\bar{a}) \Leftrightarrow \mathcal{N} \models \varphi(\bar{a})$  by induction on  $\varphi$ . Since  $\mathcal{M}$  is a **substructure** of  $\mathcal{N}$ , this is true for all **quantifier-free formulas**, and in particular for the **atomic formulas**.

Suppose that the claim is true for  $\psi$ , then

$$\mathcal{M} \models \neg\psi(\bar{a}) \Leftrightarrow \mathcal{M} \not\models \psi(\bar{a}) \Leftrightarrow \mathcal{N} \not\models \psi(\bar{a}) \Leftrightarrow \mathcal{N} \models \neg\psi(\bar{a}),$$

so the claim is also true for  $\neg\psi$ . Similarly, suppose the claim holds for  $\varphi, \psi$ . Then,

$$\mathcal{M} \models (\varphi \wedge \psi)(\bar{a}) \Leftrightarrow \mathcal{M} \models \varphi(\bar{a}) \text{ and } \mathcal{M} \models \psi(\bar{a}) \Leftrightarrow \mathcal{N} \models \varphi(\bar{a}) \text{ and } \mathcal{N} \models \psi(\bar{a}) \Leftrightarrow \mathcal{N} \models (\varphi \wedge \psi)(\bar{a}).$$

Finally, suppose the claim holds for  $\varphi(x, \bar{y})$ , then

$$\mathcal{M} \models \exists x \varphi(x, \bar{a}) \Rightarrow \exists b \in M \mathcal{M} \models \varphi(b, \bar{a}) \Rightarrow \exists b \in M \mathcal{N} \models \varphi(b, \bar{a}) \Rightarrow \mathcal{N} \models \exists x \varphi(x, \bar{a}),$$

by induction hypotheses. Conversely,  $\mathcal{N} \models \exists x \varphi(x, \bar{a})$ , then  $\exists b \in N$  such that  $\mathcal{N} \models \varphi(b, \bar{a})$  by the condition from the statement, so  $\exists c \in M$  such that  $\mathcal{N} \models \varphi(c, \bar{a})$ . By the induction hypotheses, we further have  $\mathcal{M} \models \varphi(c, \bar{a})$ , hence  $\mathcal{M} \models \exists x \varphi(x, \bar{a})$ . ■

**Example.** The ring  $\mathbb{Z}$  is a [substructure](#) of  $\mathbb{Q}$ , but  $\mathbb{Q} \models \exists x (x + x = 1)$  while  $\mathbb{Z} \not\models \exists x (x + x = 1)$ .

## Lecture 14: Downward Löwenheim-Skolem theorem

We will also need to introduce the [Skolemizations](#) of [theories](#).

21 Feb. 14:30

**Definition 3.4.3** (Built-in Skolem function). We say an  $\mathcal{L}$ -theory  $T$  has *built-in Skolem functions* if for all  $\mathcal{L}$ -formulas  $\varphi(x, y_1, \dots, y_n)$ , there is a function symbol  $f$  such that

$$T \models \forall \bar{y} (\exists x \varphi(x, \bar{y}) \rightarrow \varphi(f(\bar{y}), \bar{y})).$$

**Intuition.** This is like a parametrized version of [Henkin constants](#).

**Lemma 3.4.2.** Let  $T$  be an  $\mathcal{L}$ -theory, then there are  $\mathcal{L}^* \supseteq \mathcal{L}$  and  $T^* \supseteq T$  an  $\mathcal{L}^*$ -theory such that  $T^*$  has [built-in Skolem functions](#). Moreover, if  $\mathcal{M} \models T$ , then we can [expand](#)  $\mathcal{M}$  to  $\mathcal{M}^* \models T^*$ . Finally, we can choose  $\mathcal{L}^*$  such that  $|\mathcal{L}^*| = |\mathcal{L}| + \aleph_0$ .

**Proof.** Start with  $\mathcal{L}_0 = \mathcal{L}$  and  $T_0 = T$ , we build  $\mathcal{L}_0 \subseteq \mathcal{L}_1 \subseteq \dots$  and  $T_0 \subseteq T_1 \subseteq \dots$  and let  $\mathcal{L}^* = \bigcup_i \mathcal{L}_i$  and  $T^* = \bigcup_i T_i$ . Given  $\mathcal{L}_i$  and  $T_i$ , define

$$\mathcal{L}_{i+1} = \mathcal{L}_i \cup \{f_\varphi \mid \varphi(x, \bar{y}) \text{ is an } \mathcal{L}_i\text{-formula}\}$$

where the arity of  $f_\varphi$  is the same as  $\bar{y}$ , and

$$T_{i+1} = T_i \cup \{\forall \bar{y} (\exists x \varphi(x, \bar{y}) \rightarrow \varphi(f_\varphi(\bar{y}), \bar{y}))\}.$$

We now argue that if  $\mathcal{M}_i \models T_i$ , we can [expand](#) it to a [model](#)  $\mathcal{M}_{i+1}$  of  $T_{i+1}$ . Pick  $c \in M_i$  a “default value.” Given  $\varphi$  and  $\bar{a}$ , define  $f_\varphi^{\mathcal{M}_{i+1}}(\bar{a})$  to be some  $b$  with  $\mathcal{M}_i \models \varphi(b, \bar{a})$  if such a  $b$  exists, or  $c$  otherwise.<sup>a</sup> Then,  $\mathcal{M}_{i+1} \models T_{i+1}$ . From this construction, we see that  $T^*$  has [built-in Skolem functions](#) since any  $\mathcal{L}^*$ -formula  $\varphi$  is in some  $\mathcal{L}_i$  and has a [Skolem function](#) in  $\mathcal{L}_{i+1}$ .

Now, suppose  $\mathcal{M} \models T$ , i.e.,  $\mathcal{M} = \mathcal{M}_0 \models T_0$ . From above,  $\mathcal{M}_0$  has an [expansion](#)  $\mathcal{M}_1 \models T_1$ , which has an [expansion](#)  $\mathcal{M}_2 \models T_2$ , etc. By [expanding](#)  $\mathcal{M}$  iteratively, we get a [model](#)  $\mathcal{M}^*$  of  $T^*$ .

Finally, at each step, we add one symbol for each  $\mathcal{L}_i$ -formula to  $\mathcal{L}_i$ , hence by counting,  $|\mathcal{L}_{i+1}| = |\mathcal{L}_i| + \aleph_0$ , hence  $|\mathcal{L}^*| = |\mathcal{L}| + \aleph_0$ . ■

<sup>a</sup>If such  $b$  doesn't exist, then the left-hand side is false, so we don't really care about the right-hand side.

**Note.** We see that this is a similar argument to when we added [Henkin constants](#), though it's simpler now because we can work semantically.

**Notation** (Skolemization). We call  $T^*$  in [Lemma 3.4.2](#) a *Skolemization* of  $T$ .

**Theorem 3.4.2** (Downward Löwenheim-Skolem theorem). Let  $\mathcal{M}$  be an  $\mathcal{L}$ -structure and  $X \subseteq M$ . Then there is an [elementary substructure](#)  $\mathcal{N}$  of  $\mathcal{M}$  with  $X \subseteq \mathcal{N}$  and  $|N| \leq |X| + |\mathcal{L}| + \aleph_0$ .

**Proof.** By expanding the language, we get  $\mathcal{M}^*$  and  $\mathcal{L}^*$ -structure with  $\text{Th}(\mathcal{M}^*)$  has built-in Skolem functions (where  $T = \text{Th}(\mathcal{M})$  in Lemma 3.4.2). Hence, by replacing  $\mathcal{M}$  by  $\mathcal{M}^*$ , etc., we may assume that we already had built-in Skolem functions.

Start with  $X_0 = X \cup \{c^{\mathcal{M}} \mid c \text{ a constant symbol}\}$ . Given  $X_i$ , define  $X_{i+1}$  as

$$X_{i+1} = X_i \cup \{f^{\mathcal{M}}(\bar{a}) \mid f \in \mathcal{L} \text{ a function symbol, } \bar{a} \in X_i^{n_f}\}.$$

Let  $N = \bigcup_i X_i$ , and let  $\mathcal{N}$  be the substructure of  $\mathcal{M}$  with domain  $N$ . This can be done by:

- for each function symbol  $f$ , let  $f^{\mathcal{N}}$  be the restriction of  $f^{\mathcal{M}}$  to  $N$ ;<sup>a</sup>
- for each relation symbol  $R$ , let  $R^{\mathcal{N}}$  be the restriction of  $R^{\mathcal{M}}$  to  $N$ ;
- for each constant symbol  $c$ , there is a Skolem function  $f$  such that  $f(x) = c^{\mathcal{M}} \in N$  for all  $x \in M$ , so let  $c^{\mathcal{N}} = c^{\mathcal{M}}$ .

Now, to show that  $\mathcal{N}$  is an elementary substructure of  $\mathcal{M}$ , we use the Tarski-Vaught test. Suppose that we have an  $\mathcal{L}$ -formula  $\varphi(x, \bar{y})$ ,  $\bar{a} \in N$ ,  $b \in M$  such that  $\mathcal{M} \models \varphi(b, \bar{a})$ , and we must replace  $b$  by  $c \in N$ . Since  $\mathcal{M} \models \exists x \varphi(x, \bar{a})$ , so  $\mathcal{M} \models \varphi(f_{\varphi}(\bar{a}), \bar{a})$ . But since  $\bar{a} \in N$ , so  $f_{\varphi}(\bar{a}) \in N$ , so the Tarski-Vaught test says  $\mathcal{N}$  is an elementary substructure of  $\mathcal{M}$ .

Finally, since  $|X_0| \leq |X| + |\mathcal{L}| + \aleph_0$ , with  $N$  being a countable union,  $|N| \leq |X| + |\mathcal{L}| + \aleph_0$ . ■

<sup>a</sup>From the definition of  $N$ , it is closed under the applications of functions  $f$ , so the restriction takes values in  $N$ .

**Notation** (Generated substructure).  $\mathcal{N}$  in the downward Löwenheim-Skolem theorem is called the substructure generated by  $X$ .

**Example** (Countable real closed field). Consider  $\mathbb{R} = (\mathbb{R}, 0, 1, +, -, \cdot, \leq)$ . Let  $X \subseteq \mathbb{R}$  be countable, e.g.,  $X = \emptyset$  or  $X = \{\pi, e\}$ . Then there is  $X \subseteq \mathcal{R} \preceq \mathbb{R}$  such that  $\mathcal{R}$  is countable. In particular,  $\text{Th}(\mathcal{R}) = \text{Th}(\mathbb{R})$  and  $\mathcal{R}$  is a countable real closed field, i.e.,

- $-1$  is not a sum of squares;
- for all  $a$ , there is  $b$  such that  $a = b^2$  or  $a = -b^2$ ;
- every odd degree polynomial has a root.

There is a whole theory of real closed fields just like for algebraically closed fields.

**Intuition.** Countable real closed fields are as algebraically closed as they can be while still being orderable.

**Example** (Skolem's paradox). Let  $\mathcal{L} = \{\in\}$  be the language of set theory, where  $\in$  a binary relation symbol. Let  $T = \text{ZFC}$ , and suppose ZFC is satisfiable,<sup>a</sup> i.e., there is a model  $\mathcal{M}$  such that  $\mathcal{M} \models T$ .

Then, there is a countable  $\mathcal{N} \preceq \mathcal{M}$ , in particular, there is a countable model of ZFC. We can then write down the sentence

$$\varphi := \text{“there is no bijection between } \mathbb{R}^{\mathcal{N}} \text{ and } \mathbb{N}^{\mathcal{N}}\text{”}$$

such that  $\mathcal{N} \models \varphi$ , in  $\mathcal{L}$ . Observe that  $\mathcal{N}$  thinks that it contains an uncountable set  $\mathbb{R}^{\mathcal{N}}$ , but  $\{a \in N \mid \mathcal{N} \models a \in \mathbb{R}^{\mathcal{N}}\} \subseteq N$  is countable! This is called *Skolem's paradox*.

<sup>a</sup>From Gödel's incompleteness theorem, in ZFC, one can't prove that ZFC is consistent.

We finish this section with two useful facts.

**Definition 3.4.4** (Universally axiomatizable). Let  $T$  be an  $\mathcal{L}$ -theory, then  $T$  is *universally axiomatizable*.

zable if there is a set  $\Gamma$  of **universal sentences** such that  $T \models \Gamma$  and  $\Gamma \models T$ .<sup>a</sup>

<sup>a</sup>I.e.,  $\mathcal{M} \models T$  if and only if  $\mathcal{M} \models \Gamma$ .

**Theorem 3.4.3.** Let  $T$  be an  **$\mathcal{L}$ -theory**.  $T$  is **universally axiomatized** if and only if whenever  $\mathcal{N} \models T$  and  $\mathcal{M} \subseteq \mathcal{N}$ , then  $\mathcal{M} \models T$ .

**Proof.** The forward direction is easy: suppose that  $T$  is **universally axiomatized** by  $\Gamma$ . If  $\mathcal{M} \subseteq \mathcal{N}$  and  $\mathcal{N} \models T$ , then  $\mathcal{N} \models \Gamma$ , and since  $\Gamma$  consists only of **universal formulas**,  $\mathcal{M} \models \Gamma$ , so  $\mathcal{M} \models T$ .

Now, to prove the backward direction, suppose that if  $\mathcal{N} \models T$ ,  $\mathcal{M} \subseteq \mathcal{N}$ , then  $\mathcal{M} \models T$ . Define

$$\Gamma = \{\varphi \text{ universal} \mid T \models \varphi\},$$

then  $T \models \Gamma$ . Now, we need to show that  $\Gamma \models T$ . We may assume that  $T$  is **satisfiable**<sup>a</sup> and let  $\mathcal{M} \models \Gamma$ , so we now want to prove that  $\mathcal{M} \models T$ . This can be done by finding  $\mathcal{N} \supseteq \mathcal{M}$  and  $\mathcal{N} \models T$ , then from our assumption we have  $\mathcal{M} \models T$ . We build such an  $\mathcal{N}$  by showing that  $\text{Diag}(\mathcal{M}) \cup T$  is **satisfiable**, and take the corresponding **model** to be  $\mathcal{N}$ . Then, trivially we have  $\mathcal{N} \models T$ .

This can be done by **compactness theorem**. Let  $\Delta \subseteq \text{Diag}(\mathcal{M}) \cup T$  be finite, then there is a finite set of **atomic** or negated **atomic formulas**  $\varphi_1, \dots, \varphi_\ell$  and  $m_1, \dots, m_k \in M$  such that

$$\Delta \subseteq \{\varphi_1(\bar{m}), \dots, \varphi_\ell(\bar{m})\} \cup T,$$

assume that they are actually equal. To show that  $\Delta$  is **satisfiable**, it is enough to show<sup>b</sup> that

$$\{\exists x_1 \dots \exists x_k (\varphi_1(\bar{x}) \wedge \dots \wedge \varphi_\ell(\bar{x}))\} \cup T$$

is **satisfiable**. If not, then since  $T$  is **satisfiable**,

$$T \models \forall x_1 \dots \forall x_k \neg(\varphi_1(\bar{x}) \wedge \dots \wedge \varphi_\ell(\bar{x})).$$

Since this is **universal**, hence in  $\Gamma$ , so it is **true** in  $\mathcal{M}$ . But it's also not **true** in  $\mathcal{M}$  since  $\mathcal{M} \models \varphi_1(\bar{m}) \wedge \dots \wedge \varphi_\ell(\bar{m})$ , a contradiction  $\nmid$  Hence,  $\Delta$  is **satisfiable**, so any finite subset is **satisfiable**, by **compactness theorem**, we're done. ■

<sup>a</sup>Since otherwise  $\Gamma \ni \forall x x \neq x$ , and hence  $\Gamma \models T$  trivially.

<sup>b</sup>Since the constant symbols  $m_1, \dots, m_k$  do not appear in  $T$ , we can interpret them as the witness to the  $\exists$ 's.

## Lecture 15: The Random Graph Theory

**Proposition 3.4.2.** Suppose  $\mathcal{M}_1 \preceq \mathcal{M}_2 \preceq \dots$ , and let  $\mathcal{M} = \bigcup_i \mathcal{M}_i$ . Then  $\mathcal{M}_i \preceq \mathcal{M}$  for all  $i$ .<sup>a</sup>

<sup>a</sup>Countability is not necessary.

**Proof.** By induction on **formulas**, for all  $i$  and  $\bar{a} \in M_i$ , we show that  $\mathcal{M}_i \models \varphi(\bar{a})$  if and only if  $\mathcal{M} \models \varphi(\bar{a})$ .

- (a) For  $\varphi$  is **atomic**, this is true since  $\varphi$  is **quantifier-free** and  $\mathcal{M}_i \subseteq \mathcal{M}$ .
- (b) For  $\neg, \vee, \wedge$ , exactly the same as the **Tarski-Vaught test**.
- (c) If  $\varphi$  is  $\exists y \psi(\bar{x}, y)$ :
  - If  $\mathcal{M}_i \models \exists y \psi(\bar{a}, y)$ , there is  $b \in M_i$  such that  $\mathcal{M}_i \models \psi(\bar{a}, b)$ . Then by the induction hypothesis,  $\mathcal{M} \models \psi(\bar{a}, b)$ , so  $\mathcal{M} \models \exists y \psi(\bar{a}, y)$ .
  - If  $\mathcal{M} \models \exists y \psi(\bar{a}, y)$ , then there is  $b \in M$  such that  $\mathcal{M} \models \psi(\bar{a}, b)$ . Since  $M = \bigcup_j \mathcal{M}_j$ , there is  $j \geq i$  such that  $b \in M_j$ . By the induction hypothesis,  $\mathcal{M}_j \models \psi(\bar{a}, b)$ , so  $\mathcal{M}_j \models \exists y \psi(\bar{a}, y)$ . Finally, since  $\mathcal{M}_i \preceq \mathcal{M}_j$ , so  $\mathcal{M}_i \models \exists y \psi(\bar{a}, y)$ .

23 Feb. 14:30

### 3.5 Back and Forth

We have examples of [uncountably categorical theories](#), but no examples of [countably categorical theories](#).

#### 3.5.1 Dense Linear Order Theory

The simplest example of a [countably categorical theory](#) is the [theory](#) of “linear orders (without endpoints),” denoted as [DLO](#).

**Definition 3.5.1 (DLO).** Let  $\mathcal{L} = \{\leq\}$ . The [theory](#) of *dense linear orders (without endpoints)*, denoted as DLO, has the axioms:

- (a)  $\leq$  is a linear order;
- (b)  $\forall x \forall y (x < y \rightarrow \exists z x < z < y)$  (the density axiom);
- (c)  $\forall x \exists y \exists z (y < x < z)$  (the no-endpoints axiom).

**Example.**  $(\mathbb{Q}, \leq)$  and  $(\mathbb{R}, \leq)$  are both DLO’s.

To create a new [dense linear orders](#), given  $\mathcal{M}_1, \mathcal{M}_2$  two DLO’s, define  $\mathcal{M}_1 + \mathcal{M}_2$  with domain  $M \sqcup N$  and has each element of  $M$  less than each element of  $N$ , and within  $M$  and  $N$ , the orderings are the same as in  $\mathcal{M}$  and  $\mathcal{N}$ . This is also a DLO.

**Example.**  $\mathbb{Q} + \mathbb{Q}$  and  $\mathbb{R} + \mathbb{R}$  are both DLO’s.

**Example.**  $\mathbb{R} + \mathbb{R} \not\cong \mathbb{R}$ .

**Proof.** Since  $\mathbb{R}$  has the least upper bound property while  $\mathbb{R} + \mathbb{R}$  does not (there is no least upper bound for the first copy). \*

**Example.**  $\mathbb{Q} + \mathbb{Q} \cong \mathbb{Q}$ .

**Proof.** For example, take some irrational, e.g.,  $\pi$ . Then  $\mathbb{Q} = \{x \mid x < \pi\} \cup \{x \mid x > \pi\}$ , and we observe that we have

$$\{x \mid x < \pi\} \cong \mathbb{Q} \cong \{x \mid x > \pi\},$$

and hence piecing them together we have  $\mathbb{Q} + \mathbb{Q} \cong \mathbb{Q}$ . \*

**Example.**  $\mathbb{Q} + \mathbb{R} \not\cong \mathbb{R}$ , so DLO is not  $|\mathbb{R}| = 2^{\aleph_0}$ -categorical. In fact, not  $\kappa$ -categorical for any  $\kappa \geq 2^{\aleph_0}$ .

We now show that DLO is actually [countably categorical](#).

**Theorem 3.5.1.** The [theory](#) DLO is [countably categorical](#) and hence [complete](#).

**Proof.** Let  $(A, \leq)$  and  $(B, \leq)$  be two countable DLO’s, and let  $a_1, a_2, \dots$  and  $b_1, b_2, \dots$  be a listing of  $A$  and  $B$ , respectively. We build an [isomorphism](#)  $f: A \rightarrow B$  stage-by-stage: at stage  $i$ , we have

- finite sets  $A_i \subseteq A$  and  $B_i \subseteq B$ , and
- a bijection  $f_i: A_i \rightarrow B_i$  called a *partial embedding*:<sup>a</sup> if  $a < a' \in A_i$ , then  $f_i(a) < f_i(a')$ .

In this way,  $f_i \subseteq f_{i+1}$ ,  $A_i \subseteq A_{i+1}$ , and  $B_i \subseteq B_{i+1}$ , and we need to make sure that

- $\bigcup_i A_i = A$ , i.e., each element of  $A$  is in the domain of  $f$  (ensured by odd stages);
- $\bigcup_i B_i = B$ , i.e., each element of  $B$  is in the range of  $f$  (ensured by even stages),

so  $f = \bigcup_i f_i$  is a bijection from  $A \rightarrow B$ . Then since each  $f_i$  is a partial **embedding**, so  $f$  is an  **$\mathcal{L}$ -embedding**, hence an **isomorphism**. This will prove that DLO is **countably categorical**.

The construction works as follows.

- Stage 0:  $A_0 = \emptyset$ ,  $B_0 = \emptyset$ ,  $f_0 = \emptyset$ .
- Stage  $i + 1 = 2k + 1$ : the goal is to make sure  $a_k \in A_{i+1} = \text{dom}(f_{i+1})$ :
  - if  $a_k \in A_i$  already, then do nothing, i.e.,  $A_{i+1} = A_i$ ,  $B_{i+1} = B_i$ ,  $f_{i+1} = f_i$ ;
  - otherwise,  $a_k \notin A_i$ , define  $f_{i+1} \supseteq f_i$  by adding  $a_k$  to  $A_{i+1} = A_i \cup \{a_k\}$ , and for elements  $a \in A_i$ ,  $f_{i+1}(a) = f_i(a)$ . Now, we have three possibilities:
    - \*  $a_k$  is less than all of  $A_i$ : choose  $b \in B$  less than all of  $B_i$ ;<sup>b</sup>
    - \*  $a_k$  is greater than all of  $A_i$ : similar to above;
    - \* there are  $a$  and  $a'$  in  $A_i$  such that  $a < a_k < a'$  with no other elements of  $A_i$  between  $a$  and  $a'$  since  $A_i$  is finite: pick  $b$  with  $f_i(a) < b < f_i(a')$ .<sup>c</sup>

In all cases, we can choose  $b$  and let  $B_{i+1} = B_i \cup \{b\}$  with  $f_{i+1}(a_k) = b$ .
- Stage  $i + 1 = 2k + 2$ : the goal is to make sure  $b_k \in B_{i+1} = \text{Im}(f_{i+1})$ : this is exactly the same, but in the other direction (e.g., working with  $f_i^{-1}$  rather than  $f_i$ ).

Now everything is checked, so DLO is **countably categorical** (hence **complete** by **Vaught test**). ■

<sup>a</sup>If  $f_i$  maps  $\bar{a}$  to  $\bar{b}$ , then  $\bar{a}$  satisfies the same **atomic** and negated **atomic formulas** in  $(A, \leq)$  that  $\bar{b}$  does in  $(B, \leq)$ .

<sup>b</sup>Such  $b$  exists since  $B_i$  is finite, and  $(B, \leq)$  has no left endpoint.

<sup>c</sup>Such  $b$  exists since  $B_i$  is finite and  $(B, \leq)$  is dense.

**Note (Back-and-forth).** We see that the above is the so-called *back-and-forth* argument.

**Corollary 3.5.1.**  $\mathbb{Q} + \mathbb{R} \equiv \mathbb{R}$ .

**Proof.** Since  $\text{Th}(\mathbb{Q} + \mathbb{R}) = \text{Th}(\mathbb{R}) = \{\varphi \mid \text{DLO} \models \varphi\}$ . ■

**Definition 3.5.2 (Complete).** A linear order is *complete* if every subset bounded above has a least upper bound.

**Corollary 3.5.2.** There is no first order **sentence**  $\varphi$  such that  $\mathcal{M} \models \varphi$  if and only if  $\mathcal{M}$  is a **complete** linear order.

### 3.5.2 Random Graph Theory

Another example of a **countably categorical theory** is the **theory of random graph**.

**Definition 3.5.3 (Random graph).** A *random graph* we will consider is constructed as follows. Firstly, fix countably infinitely many vertices  $v_1, v_2, \dots$ , and fix  $p$  such that  $0 < p < 1$ . For each pair of vertices, “flip a coin”: with probability  $p$ , put an edge; with  $1 - p$ , no edge.

Now, the question is, what graph do we get? It turns out to be interesting enough, so we will look into it:

**Remark.** With probability 1, we get the same graph up to isomorphism, no matter what  $p$  is.

Then, consider the following **theory**.


**Definition 3.5.4 (Random graph theory).** Let  $\mathcal{L} = \{E\}$ , where  $E$  is a binary relation. The *random graph theory*  $T$  has axioms:

- (a)  $\forall x \neg xEx$  and  $\forall x \forall y (xEy \rightarrow yEx)$  (irreflexive, undirected);
- (b)  $\exists x \exists y x \neq y$ ;
- (c) for each  $n$ , define  $\psi_n$  as

$$\psi_n := \forall x_1 \dots \forall x_n \forall y_1 \dots \forall y_n \left[ \bigwedge_{i=1}^n \bigwedge_{j=1}^n x_i \neq y_j \rightarrow \exists z \left( \bigwedge_{i=1}^n x_i E z \wedge \neg y_i E z \wedge z \neq x_i \wedge z \neq y_i \right) \right]$$

**Intuition (Extension axiom).** Think of  $\psi_n$  as an *extension axiom*: the property that for any finite disjoint sets  $X$  and  $Y$ , there is a vertex with an edge to each  $x \in X$  and no edge to each  $y \in Y$ . This axiom happens with probability  $p^{|X|} \cdot (1-p)^{|Y|}$  for a given  $z$  for some  $X, Y$ .

We see that a **model** of  $T$  is a graph with at least two elements with the **extension property**.



**Note.** In  $\psi_n$ 's, we allow there to be repetitions among the  $x$ 's and the  $y$ 's. In particular, if  $m \leq n$ , then  $\psi_n \models \psi_m$ .

## Lecture 16: Quantifier Elimination

**Remark.**  $T$  is actually **countably categorical**.

7 Mar. 14:30

**Proof.** We will show this on homework. But this is basically **Theorem 3.5.1**. ⊛

Now we show that it is also **satisfiable**. Fix  $N, p \in (0, 1)$ , and vertices  $\{1, \dots, N\}$ . Generate a **random graph**  $G_N$  on vertices  $\{1, \dots, N\}$  by, for each  $i, j \leq N, i \neq j$ , putting an edge between  $i$  and  $j$  with probability  $p$  and no edge with probability  $1 - p$ . Let  $\mathcal{G}_N$  be the class of all such graphs, and let  $p_N(\varphi)$  be the probability that a **randomly generated graph** on  $N$  vertices satisfies  $\varphi$ .

**Example.** If  $p = 1/2$ , all graphs in  $\mathcal{G}_N$  are equally likely.

Observe the following.

**Proposition 3.5.1.** For every  $n$ , then  $\lim_{N \rightarrow \infty} p_N(\psi_n) = 1$ .

**Proof.** Fix  $n$ , and let  $G$  be a **random graph** in  $\mathcal{G}_N, N > 2n$ . Fix  $x_1, \dots, x_n$ , and  $y_1, \dots, y_n$ , and  $z$  in  $G$ , all distinct. Consider the probability  $q$  that

$$\neg \left( \bigwedge_{i=1}^n E(x_i, z) \wedge \neg E(y_i, z) \right),$$

which is just  $1 - p^n(1-p)^n := q$  as we already know. The probability that

$$G \models \underbrace{\neg \exists z}_{\forall z \neg} \left( \bigwedge_{i=1}^n E(x_i, z) \wedge \neg E(y_i, z) \right)$$

is  $q^{N-2n}$ , where  $N - 2n$  is the number of possible  $z$ 's.<sup>a</sup> Let  $M$  be the number of different choices

of  $x_i$ 's and  $y_i$ 's, then we have

$$p_N(\neg\psi_n) \leq M \cdot q^{N-2n} \leq N^{2n} q^{N-2n}$$

by the union bound.<sup>b</sup> Since  $0 < q < 1$ , so  $p_N(\neg\psi_n) \rightarrow 0$ , i.e.,  $p_N(\psi_n) \rightarrow 1$  as  $N \rightarrow \infty$ . ■

<sup>a</sup>Notice that we already assume that  $x_i \neq z \wedge y_i \neq z$ .

<sup>b</sup>This bound is not allowing the  $x$ 's and  $y$ 's to have repetitions. But this doesn't change anything.

Then, we have the following.

**Theorem 3.5.2.**  $T$  is **satisfiable**.

**Proof.** From Proposition 3.5.1,  $\lim_{N \rightarrow \infty} p_N(\psi_n) = 1$ . In particular, for each  $n$ , there is  $N$  such that  $p_N(\psi_n) > 0$ , i.e., there is at least one  $G \in \mathcal{G}_N$  such that  $G \models \psi_n$ , hence  $G \models \psi_m$  for  $m \leq n$ .<sup>a</sup> This means that for every finite  $T^* \subseteq T$  is **satisfiable**, so  $T$  is **satisfiable** by the **compactness theorem**. ■

<sup>a</sup>Remember that for  $n > m$ ,  $\models \psi_n \rightarrow \psi_m$ .

Moreover, we have the following.

**Theorem 3.5.3** (Zero-one law for graphs). For any  $\mathcal{L}$ -sentence  $\varphi$ , we either have  $\lim_{N \rightarrow \infty} p_N(\varphi) = 0$  or  $\lim_{N \rightarrow \infty} p_N(\varphi) = 1$ . Moreover,  $T$  axiomatizes  $\{\varphi : \lim_{N \rightarrow \infty} p_N(\varphi) = 1\}$ , the “almost sure theory for graphs”, which is **decidable** and **complete**.

**Proof.** Since  $T$  is **countably categorical**, and has only infinite models, so it is **complete** by the **Vaught's test**. If  $T \models \varphi$ , there is some  $n$  such that  $\{\psi_n, (a), (b)\} \models \varphi$ . Since  $p_N(\psi_n) \leq p_N(\varphi)$ , so  $\lim_{N \rightarrow \infty} p_N(\varphi) = 1$  and the left-hand side goes to 1 as well. On the other hand, if  $T \not\models \varphi$ ,  $T \models \neg\varphi$ , so  $\lim_{N \rightarrow \infty} p_N(\neg\varphi) = 1$ , i.e.,  $\lim_{N \rightarrow \infty} p_N(\varphi) = 0$  from the same argument. ■

Before ending this section, we note that there are other things we may explore.

**Remark.** There are also Fraïssé constructions, and Ryll-Nardzewski theorem, etc.



## Chapter 4

# Quantifier Elimination and Algebraic Applications

### 4.1 Quantifier Elimination

Let's start with a definition.

**Definition 4.1.1** (Quantifier elimination). A theory  $T$  has (admits) *quantifier elimination* if for every formula  $\varphi(\bar{x})$  (with  $\bar{x}$  containing at least one variable), there is a quantifier-free  $\psi(\bar{x})$  such that

$$T \models \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x})).$$

The easiest example will be that DLO admits quantifier elimination. To prove this, we need some preliminaries.

**Note.** DLO has no constant symbols, so it has no quantifier-free sentences. For example,

$$\forall x \exists y y < x \leftrightarrow x = x,$$

where the right-hand side has a free variable. Another solution would be to allow  $\top$  and  $\perp$  for the true and false sentences.

**Lemma 4.1.1.** Let  $(A, \leq)$  and  $(B, \leq)$  be countable DLOs. Let  $a_1, \dots, a_n \in A$  have  $a_1 < a_2 < \dots < a_n$  and  $b_1, \dots, b_n \in B$  have  $b_1 < \dots < b_n$ . Then there is an isomorphism  $f: A \rightarrow B$  which maps  $a_i \mapsto b_i$ . Hence,

$$A \models \varphi(\bar{a}) \Leftrightarrow B \models \varphi(\bar{b}).$$

**Proof.** This is the same as the back-and-forth argument, but now we start with  $\bar{a} \mapsto \bar{b}$ . ■

**Theorem 4.1.1.** DLO admits quantifier elimination.

**Proof.** Fix  $\varphi(\bar{x})$ . If  $\varphi$  is actually a sentence then we're done since either

- DLO  $\models \varphi$ , so DLO  $\models \forall x (\varphi \leftrightarrow x = x)$ , or
- DLO  $\models \neg \varphi$ , so DLO  $\models \forall x (\varphi \leftrightarrow x \neq x)$ .

Now suppose  $\varphi(\bar{x})$  has at least one free variable,  $\bar{x} = (x_1, \dots, x_n)$ . Since DLO is complete, it's enough to find  $\psi(\bar{x})$  with  $\mathbb{Q} \models \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x}))$ . Let  $\sigma$  be a map from pairs  $i, j$  to  $\{1, 2, 3\}$ . Define<sup>a</sup>

$$\theta_\sigma(x_1, \dots, x_n) := \bigwedge_{\sigma(i,j)=1} x_i < x_j \wedge \bigwedge_{\sigma(i,j)=2} x_i = x_j \wedge \bigwedge_{\sigma(i,j)=3} x_j < x_i.$$

If  $\mathbb{Q} \models \theta_\sigma(\bar{a}) \wedge \theta_\sigma(\bar{b})$ , then  $\bar{a}$  and  $\bar{b}$  satisfy the same formulas by Lemma 4.1.1, so<sup>b</sup>

$$\Sigma := \{\sigma \mid \mathbb{Q} \models \exists \bar{x} (\theta_\sigma(\bar{x}) \wedge \varphi(\bar{x}))\} = \{\sigma \mid \mathbb{Q} \models \forall \bar{x} (\theta_\sigma(\bar{x}) \rightarrow \varphi(\bar{x}))\}.$$

If  $\Sigma = \emptyset$ , then  $\varphi(x) \leftrightarrow x \neq x$ ; if  $\Sigma \neq \emptyset$ , let  $\psi(\bar{x}) = \bigvee_{\sigma \in \Sigma} \theta_\sigma(\bar{x})$ , then  $\mathbb{Q} \models \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x}))$ . ■

<sup>a</sup>It's clear that some  $\theta_\sigma$  might be inconsistent.

<sup>b</sup>The second equality follows from the fact that if one such  $\bar{x}$  exists, all such  $\bar{x}$  work.

## Lecture 17: Quantifier Elimination

**Example.**  $\varphi(x, y) := \exists u (u > x \wedge u < y)$ . This is equivalent, in DLO, to  $\psi(x, y) := x < y$ .

9 Mar. 14:30

**Example.** In Problem Set 2, we have looked at  $\mathcal{L} = \emptyset$ . There, we showed that if  $A \subseteq B$  and both infinite, then  $A \preceq B$ . The same idea show that  $T = \{\exists x_1 \dots \exists x_n \bigwedge_{i \neq j} x_i \neq x_j \mid n \in \mathbb{N}\}$  admits quantifier elimination.

**Proposition 4.1.1.** Let  $T$  be a theory that admits quantifier elimination. If  $\mathcal{A}, \mathcal{B} \models T$  and  $\mathcal{A} \subseteq \mathcal{B}$ , then  $\mathcal{A} \preceq \mathcal{B}$ .


**Proof.** Given  $\phi(\bar{x})$  and  $\bar{a} \in A$ , then  $\phi(x)$  is equivalent to a quantifier-free  $\psi(\bar{x})$  (modulo  $T$ ). Then,

$$\mathcal{A} \models \phi(\bar{a}) \Leftrightarrow \mathcal{A} \models \psi(\bar{a}) \Leftrightarrow \mathcal{B} \models \psi(\bar{a}) \Leftrightarrow \mathcal{B} \models \phi(\bar{a}).$$

Next, we want to show that ACF admits quantifier elimination. First, we need a test for quantifier elimination.

**Theorem 4.1.2.** Let  $\mathcal{L}$  include at least one constant symbol  $c$ . Let  $T$  be an  $\mathcal{L}$ -theory, and  $\phi(\bar{x})$  an  $\mathcal{L}$ -formula. Then the following are equivalent.

- (a) There is a quantifier-free  $\psi(\bar{x})$  such that  $T \models \forall \bar{x} (\phi(\bar{x}) \leftrightarrow \psi(\bar{x}))$ .
- (b) If  $\mathcal{M}, \mathcal{N} \models T$ , and  $\mathcal{A}$  is a common substructure, then for all  $\bar{a} \in A$ ,  $\mathcal{M} \models \phi(\bar{a})$  if and only if  $\mathcal{N} \models \phi(\bar{a})$ .



**Proof.** (a) implies (b) is easy, since we have

$$\mathcal{M} \models \phi(\bar{a}) \Leftrightarrow \mathcal{M} \models \psi(\bar{a}) \Leftrightarrow \mathcal{A} \models \psi(\bar{a}) \Leftrightarrow \mathcal{N} \models \psi(\bar{a}) \Leftrightarrow \mathcal{N} \models \phi(\bar{a}).$$

To show (b) implies (a), we see that first, there are two easy cases:

- $T \models \forall \bar{x} \phi(\bar{x})$ : take  $\psi$  to be  $c = c$ ;
- $T \models \forall \bar{x} \neg \phi(\bar{x})$ : take  $\psi$  to be  $c \neq c$ .

Now, suppose we are not in the above two cases. Let  $\Gamma(\bar{x})$  be the set of all quantifier-free formulas  $\psi(\bar{x})$  such that  $T \models \forall \bar{x} (\phi(\bar{x}) \rightarrow \psi(\bar{x}))$ , and let  $\bar{d} = (d_1, \dots, d_n)$  be new constant symbols.

**Claim.** It's enough to show  $T \cup \Gamma(\bar{d}) \models \phi(\bar{d})$ .

**Proof.** If we can do this, then by **compactness**, there are  $\psi_1(\bar{x}), \dots, \psi_m(\bar{x})$  such that

$$T \models [\psi_1(\bar{d}) \wedge \dots \wedge \psi_m(\bar{d})] \rightarrow \phi(\bar{d})$$

By the choice of  $\Gamma$ , the  $\leftarrow$  direction also holds, hence

$$T \models [\psi_1(\bar{d}) \wedge \dots \wedge \psi_m(\bar{d})] \leftrightarrow \phi(\bar{d}),$$

i.e.,  $T \models \forall \bar{x} (\psi(\bar{x}) \leftrightarrow \phi(\bar{d}))$  where  $\psi(x) := \psi_1(\bar{x}) \wedge \dots \wedge \psi_m(\bar{x})$ . ⊗

**Claim.**  $T \cup \Gamma(\bar{d}) \models \phi(\bar{d})$ .

**Proof.** Suppose not. Then  $T \cup \Gamma(\bar{d}) \cup \{\neg\phi(\bar{d})\}$  is **satisfiable**.<sup>a</sup> Let  $\mathcal{M}$  be a **model** of  $T \cup \Gamma(\bar{d}) \cup \{\neg\phi(\bar{d})\}$ , and let  $\mathcal{A} \subseteq \mathcal{M}$  be the “submodel”<sup>b</sup> generated by  $\bar{d}$ . Notice that since  $c^{\mathcal{A}} \in \mathcal{A}$ , so  $\mathcal{A} \neq \emptyset$ . Every element of  $\mathcal{A}$  is  $t^{\mathcal{A}}(\bar{d}) = t^{\mathcal{M}}(\bar{d})$  for some **term**  $t$ . Also,  $\mathcal{A} \models \Gamma(\bar{d})$  because  $\mathcal{M} \models \Gamma(\bar{d})$  and  $\Gamma(\bar{d})$  is **quantifier-free**. We now show that  $T \cup \text{Diag}(\mathcal{A}) \cup \{\phi(\bar{d})\}$  is **satisfiable**, and take  $\mathcal{N}$  to be a **model**. If not, some finite subset is not **satisfiable**. There are **quantifier-free formulas**  $\gamma_1(\bar{d}), \dots, \gamma_\ell(\bar{d})$  in  $\text{Diag}(\mathcal{A})$  such that<sup>c</sup>

$$T \models [\gamma_1(\bar{d}) \wedge \dots \wedge \gamma_\ell(\bar{d})] \rightarrow \neg\phi(\bar{d}).$$

Equivalently,  $T \models \phi(\bar{d}) \rightarrow \neg[\gamma_1(\bar{d}) \wedge \dots \wedge \gamma_\ell(\bar{d})]$ , so  $T \models \forall \bar{x} (\phi(\bar{x}) \rightarrow [\neg\gamma_1(\bar{x}) \vee \dots \vee \neg\gamma_\ell(\bar{x})])$ . This implies that  $\neg\gamma_1(\bar{x}) \vee \dots \vee \neg\gamma_\ell(\bar{x})$  is in  $\Gamma(\bar{x})$ , hence  $\mathcal{A} \models \neg\gamma_1(\bar{d}) \vee \dots \vee \neg\gamma_\ell(\bar{d})$ . But  $\mathcal{A} \models \gamma_1(\bar{d}) \wedge \dots \wedge \gamma_\ell(\bar{d})$ , contradicts to the fact that  $\gamma_1(\bar{d}), \dots, \gamma_\ell(\bar{d})$  is in  $\text{Diag}(\mathcal{A})$ .

Hence, there is  $\mathcal{N} \models T \cup \text{Diag}(\mathcal{A}) \cup \{\phi(\bar{d})\}$ . This contradicts to (b):

$$\begin{array}{ccc} \mathcal{M} \models \neg\phi(\bar{d}) \text{ and } \mathcal{M} \models T & & \mathcal{N} \models \neg\phi(\bar{d}) \text{ and } \mathcal{N} \models T \\ & \nwarrow \subseteq \quad \nearrow & \\ & \mathcal{A} \ni \bar{d} & \end{array}$$

Hence,  $T \cup \Gamma(\bar{d}) \models \phi(\bar{d})$ . ⊗

<sup>a</sup>Notice that  $T \cup \Gamma(\bar{d})$  needs to be **satisfiable**, which is true from our assumption.

<sup>b</sup> $\mathcal{A}$  is the smallest **model** containing  $\bar{d}$ , and  $\bar{a}$  might not be a **model** of  $T$ .

<sup>c</sup>We can take  $\gamma_i(\bar{d})$  to be just about  $\bar{d}$  but not  $\gamma_i(\bar{a})$  for  $\bar{a} \in \mathcal{A}$  is because each  $\bar{a} \in \mathcal{A}$  is  $t^{\mathcal{A}}(\bar{d})$ , so we can replace  $\bar{a}$  by this **term**.

We now show that we can eliminate one  $\exists$  at a time. ■

**Lemma 4.1.2.** Let  $T$  be an  **$\mathcal{L}$ -theory**. Suppose that for every **quantifier-free formula**  $\gamma(\bar{x}, y)$ , there is a **quantifier-free formula**  $\psi(\bar{x})$  such that

$$T \models \forall \bar{x} (\exists y \gamma(\bar{x}, y) \leftrightarrow \psi(\bar{x})),$$

then  $T$  has **quantifier elimination**.

**Proof.** By induction on **formulas** with the hypotheses of **Theorem 4.1.2** for the  $\exists$  quantifier case. ■

By putting together the last two results, i.e., **Lemma 4.1.2** and **Theorem 4.1.2**, we get the following test for **quantifier elimination**. Compare this, for example, to **Vaught's test** for **elementary-substructure**.

**Corollary 4.1.1.** Let  $T$  be an  **$\mathcal{L}$ -theory** with at least one constant. Suppose that for all **quantifier-free**  $\psi(\bar{x}, y)$ , if  $\mathcal{M}, \mathcal{N} \models T$ , and  $\mathcal{A} \subseteq \mathcal{M}$  and  $\mathcal{A} \subseteq \mathcal{N}$ ,  $\bar{a} \in \mathcal{A}$  and  $b \in M^{\mathcal{A}}$  is such that  $\mathcal{M} \models \psi(\bar{a}, b)$ , then

there is  $c \in N$  such that  $\mathcal{N} \models \psi(\bar{a}, c)$ . Then  $T$  admits **quantifier elimination**.



<sup>a</sup>This is (b) for  $\exists y \psi(\bar{x}, y)$ .

#### 4.1.1 Quantifier Elimination for ACF

Now, we start with our favorite ACF, and show that it admits **quantifier elimination**.

**Theorem 4.1.3.** ACF admits **quantifier elimination**.

**Proof.** We use **Corollary 4.1.1**. Suppose that  $\mathcal{M}, \mathcal{N} \models \text{ACF}$ , and  $\mathcal{A} \subseteq \mathcal{M}, \mathcal{N}$  an **integral domain**. Let  $\psi(\bar{x}, y)$  be a **quantifier-free formula** and  $\bar{a} \in \mathcal{A}$ . Suppose there is  $b \in \mathcal{M}$  such that  $\mathcal{M} \models \psi(\bar{a}, b)$ .

Firstly, we replace  $\mathcal{N}$  by the **algebraic closure** of  $\mathcal{A}$ , then we can also assume that  $\mathcal{N} \subseteq \mathcal{M}$  since any **algebraic closure** of  $\mathcal{A}$  embeds in any **algebraically closed** field containing  $\mathcal{A}$ .



**Problem.** What does  $\psi$  look like?

**Answer.** Since it's **quantifier-free**, we may assume that it is  $\theta_1(\bar{x}, y) \vee \theta_2(\bar{x}, y) \vee \dots$ , where each  $\theta_i$  is a conjunction of **atomic** and negated **atomic formula**. We can replace  $\psi$  by whichever  $\theta_i$  is satisfied by  $b$ . ⊗

Now, since in the **language** of rings, an **atomic formula**  $\gamma(\bar{x}, y)$  is equivalent to a **formula** of the form  $p(\bar{x}, y) = 0$ , where  $p \in \mathbb{Z}[\bar{X}, Y]$ , so  $\psi(\bar{x}, y)$  is equal to

$$p_1(\bar{x}, y) = 0 \wedge \dots \wedge p_k(\bar{x}, y) = 0 \wedge q_1(\bar{x}, y) \neq 0 \wedge q_\ell(\bar{x}, y) \neq 0$$

for  $p_i, q_i \in \mathbb{Z}[\bar{X}, Y]$ . Then  $\psi(\bar{a}, y)$  says  $p_1(\bar{a}, y) = 0 \wedge \dots$  where  $p_i(\bar{a}, y)$  are now in  $\mathcal{A}[y]$ , i.e., polynomial in  $y$  with coefficient in  $\mathcal{A}$ . If any  $p_i(\bar{a}, y)$  is non-trivial, then  $b$  is a solution, so  $b \in \mathcal{N}$  since  $\mathcal{N}$  is **algebraically closed** (and we just take  $c = b$  for applying **Corollary 4.1.1**). Otherwise, assume  $p_i(\bar{a}, y)$  is trivial, so  $\psi(\bar{a}, y)$  is just

$$q_i(\bar{a}, y) \neq 0 \wedge \dots \wedge q_\ell(\bar{a}, y) \neq 0.$$

Since  $b$  satisfies this, each  $q_i(\bar{a}, y)$  is non-trivial, so  $q_i(\bar{a}, y) = 0$  has only finitely many solutions. But  $\mathcal{N}$  is infinite, so there is a  $c \in \mathcal{N}$  that is not a solution to any  $q_i(\bar{a}, y) = 0$ , so  $\mathcal{N} \models \psi(\bar{a}, c)$ . ■

## Lecture 18: Quantifier Elimination for Algebraically Closed Fields

#### 4.1.2 Definable and Constructible Sets

14 Mar. 14:30

**Problem 4.1.1.** What do we get from **quantifier elimination**?

**Answer.** Understand the **definable** sets, they are defined by **quantifier-free formulas**. ⊗

**Remark.** We see that

$$\varphi(\bar{x}) := \exists y p_n(\bar{x})y^n + \dots + p_1(\bar{x})y + p_0(\bar{x}) = 0$$

is equivalent to

$$\psi(\bar{x}) := p_n(\bar{x}) \neq 0 \vee \dots \vee p_1(\bar{x}) \neq 0 \vee p_0(\bar{x}) = 0.$$

**Definition 4.1.2 (Cofinite).** A *cofinite* subset of a set  $X$  is a subset  $A$  such that  $|A^c| < \infty$ .

**Proposition 4.1.2.** The *definable*<sup>a</sup> subsets of an *algebraically closed* field  $K$  are exactly the finite and *cofinite* sets.

<sup>a</sup>Using parameters.

**Proof.** Let  $\{a_1, \dots, a_n\}$  be a finite set. This is *definable* by  $x = a_1 \vee \dots \vee x = a_n$  using  $a_1, \dots, a_n$  as parameters; on the other hand, the complement of  $\{a_1, \dots, a_n\}$  is also *definable* by  $x \neq a_1 \wedge \dots \wedge x \neq a_n$ .<sup>a</sup>

Let  $X = \{x \in K \mid K \models \varphi(x, \bar{a})\}$  be a *definable* subset of  $K$ . By *quantifier elimination*, we may assume that  $\varphi$  is *quantifier-free*, so  $\varphi$  is a boolean combination of *atomic* and negated *atomic formulas*. Notice that an *atomic formula* is of the form

$$p_n(\bar{a})x^n + \dots + p_1(\bar{a})x + p_0(\bar{a}) = 0.$$

Hence, this *atomic formula defines* either a finite set or all of  $K$ . A negated *atomic formula defines* a *cofinite* set or  $\emptyset$ . Boolean combinations of finite and *cofinite* sets are finite or *cofinite*, so  $X$  is finite or *cofinite*. ■

<sup>a</sup>We did not use anything about fields here.

**Remark.** Proposition 4.1.2 is not true for  $X \subseteq K^2$ .

**Proof.**  $X = \{(x, y) \mid x^2 + y^2 + 1\}$ . ⊗

**Definition 4.1.3 (Strongly minimal).** A *theory*  $T$  is *strongly minimal* if for any  $\mathcal{M} \models T$ , and  $X \subseteq M$  *definable*,  $X$  is either finite or *cofinite*.

**Definition 4.1.4 (Algebraic).** Let  $K$  be a field, and  $X \subseteq K^n$ . We say that  $X$  is *algebraic* if there is a set  $S$  of polynomials over  $K$  such that  $X$  is the zero set of  $S$ .

**As previously seen.** Recall Definition 3.2.15 and compared it to the above.

**Example.**  $X = \{(x, y) \mid x^2 + y^2 = 1\}$  is *algebraic* since  $S = \{x^2 + y^2 - 1\}$ .

The complement of an *algebraic set* is usually not *algebraic*.

**Definition 4.1.5 (Constructible).** The *constructible* sets are the boolean combinations of *algebraic sets*.

**Remark.** The *constructible* sets are exactly the *definable* sets in  $K \models \text{ACF}$ .

**Proof.** The *definable* sets, by *quantifier elimination*, boolean combinations of sets *defined* by *atomic formulas*, which are *algebraic*. So *definable* implies *constructible*.

On the other hand, it is enough to see that *algebraic set* are *definable*. The issue is that  $S \subseteq K[\bar{X}]$  might be infinite. Let  $I$  be the *ideal generated* by  $S$ . Then the set  $X \subseteq K^n$  of common zeros of  $S$  is also the set of common zeros of  $I$ .<sup>a</sup> By the *Hilbert's basis theorem*,  $I = (f_1, \dots, f_m)$  is finitely *generated*. Hence,

$$X = \{\bar{a} \in K^n \mid f_1(\bar{a}) = 0 \wedge \dots \wedge f_m(\bar{a}) = 0\},$$

hence it's **definable**. ⊛

<sup>a</sup>Each  $f \in I$  is  $f = r_1 g_1 + \dots + r_n g_n$ , where  $g_1, \dots, g_n \in S$ .

**Theorem 4.1.4** (Chevalley's theorem). Let  $K$  be an **algebraically closed** field. Let  $X \subseteq K^n$  be **constructible**. Let  $p: K^n \rightarrow K^m$  be a polynomial map, i.e.,  $p(\bar{x}) = (q_1(\bar{x}), \dots, q_m(\bar{x}))$  for polynomials  $q_i$ . Then,  $p(X)$ , the image of  $X$  under  $p$ , is also **constructible**.

**Proof.** Since we know that **constructible** is the same as **definable**, so consider

$$p(X) = \{\bar{y} \mid \exists \bar{x} (\bar{x} \in X \wedge p(\bar{x}) = \bar{y})\}$$

where for  $\bar{x} \in X$ , there is a **formula** expressing this, and for  $p(\bar{x}) = \bar{y}$ ,  $y_1 = q_1(\bar{x}) \wedge \dots \wedge y_m = q_m(\bar{x})$ . Hence,  $p(X)$  is **definable** (hence **constructible**), since  $X$  was. ■

**Example.**  $p(x_1, x_2, x_3) = (x_1, x_3)$ .

**Theorem 4.1.5** (Weak Hilbert's Nullstellensatz). Let  $K$  be **algebraically closed**, and  $f_1, \dots, f_n \in K[\bar{x}]$ . Then there is  $\bar{a} \in K^m$  such that  $f_1(\bar{a}) = \dots = f_n(\bar{a}) = 0$  if and only if  $1 \notin (f_1, \dots, f_n)$ , i.e., there are no  $r_1, \dots, r_n \in K[\bar{x}]$  such that  $1 = r_1 f_1 + \dots + r_n f_n$ .

**Proof.** If  $1 \in (f_1, \dots, f_n)$ , there are  $r_1, \dots, r_n \in K[\bar{x}]$  such that  $1 = r_1 f_1 + \dots + r_n f_n$ . If  $\bar{a}$  was a common zero of  $f_1, \dots, f_n$ , then

$$1 = r_1(\bar{a})f_1(\bar{a}) + \dots + r_n(\bar{a})f_n(\bar{a}) = 0,$$

so no such  $\bar{a}$  exists.

Now, suppose  $1 \notin (f_1, \dots, f_n)$ , so  $(f_1, \dots, f_n) \neq K[\bar{x}]$ . Let  $I$  be a **maximal ideal** containing  $f_1, \dots, f_n$ , and let  $L = K[\bar{x}] / I$ , which is a **field extension** of  $K$ ,  $K \hookrightarrow L$ . There is  $\bar{a} \in L^m$  which is a common root of  $f_1, \dots, f_n$ , namely  $a_i = x_i + I$ . Let  $M$  be the **algebraic closure** of  $L$ , with  $\bar{a} \in M^n$ . By **quantifier elimination**,  $K \succeq M$ .  $M \models \exists \bar{y} f_1(\bar{y}) = 0 \wedge \dots \wedge f_n(\bar{y}) = 0$ , which is a **formula** about elements of  $K$  (the coefficients). Because  $K \succeq M$ ,  $K \models \exists \bar{y} f_1(\bar{y}) = 0 \wedge \dots \wedge f_n(\bar{y}) = 0$ , which says that  $f_1, \dots, f_n$  have a common zero of  $K$ . ■

**Note.** We leave the full version in the note, which relates to algebraic geometry (if you care).

**Remark.** The **weak Hilbert's Nullstellensatz** says that whether  $1 \in (f_1, \dots, f_n)$  is the only barrier for  $f_1, \dots, f_n$  having a common zero.

### 4.1.3 Algebraic Closure

**As previously seen** (Definable closure). The *definable closure*  $\text{dcl}(A)$  of  $A$  is the set of all  $b \in M$  which are **definable** over  $A$ .

On the homework, we looked at  $\text{dcl}$ , which is not really useful. In **strongly minimal theories**, we look at  $\text{acl}$  instead, which is very important.

**Definition 4.1.6** ((Model theoretical) algebraic closure). Let  $\mathcal{M}$  be a **structure** and  $A \subseteq M$ . Then the (*model-theoretic*) *algebraic closure*  $\text{acl}(A)$  of  $A$  is the set of all  $a \in M$  such that there are  $\bar{b} \in A$  and a **formula**  $\varphi(x, \bar{b})$  such that  $\mathcal{M} \models \varphi(a, \bar{b})$  and there are only finitely many other  $a'$  with  $\mathcal{M} \models \varphi(a', \bar{b})$ .

**Note.** There are some properties that  $\text{acl}$  always satisfies.

- $\text{dcl}(A) \subseteq \text{acl}(A)$ .

- $A \subseteq \text{acl}(A)$ .
- If  $A \subseteq B$ , then  $\text{acl}(A) \subseteq \text{acl}(B)$ .
- $\text{acl}(A) = \text{acl}(\text{acl}(A))$ .<sup>a</sup>
- If  $a \in \text{acl}(A)$ , then there is a finite  $F \subseteq A$  such that  $a \in \text{acl}(F)$ .

<sup>a</sup>This is a good exercise!

On homework 5, we will show that in **models** of a **strongly minimal theory**,  $\text{acl}$  also satisfies the “exchange property”:

**Remark (Exchange property).** If  $a \in \text{acl}(X \cup \{b\})$  and  $a \notin \text{acl}(X)$ , then  $b \in \text{acl}(X \cup \{a\})$ . This makes  $\text{acl}$  a **pregeometry** or **matroid**.

In **algebraically closed** fields,  $a \in \text{acl}(X)$  just means that  $a$  is **algebraic** over (the field generated by)  $X$ , i.e., there is  $p(y) \in F_X[\bar{x}]$  such that  $p(a) = 0$ . In vector spaces,  $a \in \text{acl}(X)$  if and only if  $a \in \text{span}(X)$ .

**Intuition.** This makes **model** has notions like dimensions, independence, etc.

## Lecture 19: acl in Algebraically Closed Fields

**Theorem 4.1.6.** Let  $K \models \text{ACF}$ , and  $A \subseteq K$ . The  $\text{acl}(A)$  is the set of all elements  $b \in K$  which are **algebraic** over  $A$ , i.e., that satisfy a non-zero polynomial with coefficients in the ring generated by  $A$ .

**Proof.** If  $p(x)$  is a non-zero polynomial over  $A$ , with  $p(b) = 0$ , then  $\{c \mid p(c) = 0\}$ , so  $p(x) = 0$  witnesses that  $b \in \text{acl}(A)$ .

Suppose that  $b \in \text{acl}(A)$ , as witnessed by  $\varphi(x, \bar{a})$  for  $\bar{a} \in A$ . We may assume that  $\varphi$  is **quantifier-free**. Write

$$\varphi(x, \bar{a}) = \psi_1(x, \bar{a}) \vee \dots \vee \psi_n(x, \bar{a}),$$

where each  $\psi_i$  is a conjunction of **atomic** or negated **atomic formulas**. We may replace  $\varphi$  with some  $\psi_i$ , choosing one that  $b$  satisfies. Then,

$$\varphi(x, \bar{a}) = p_1(x, \bar{a}) \wedge \dots \wedge p_k(x, \bar{a}) = 0 \wedge q_1(x, \bar{a}) \neq 0 \wedge \dots \wedge q_\ell(x, \bar{a}) \neq 0.$$

In order for this to have finitely many solutions, we must have  $k \geq 1$  and some  $p_i(x, \bar{a})$  non-zero polynomial in  $x$ . But then  $b$  satisfies a polynomial over  $A$ . ■

16 Mar. 14:30

### 4.1.4 Types

Now, unless we specify, we will now work in a **model** of a **strongly minimal theory**.

**Definition 4.1.7 (Independent).** A set  $X$  is *independent* if for all  $a \in X$ ,  $a \notin \text{acl}(X - \{a\})$ .

**Remark.** In homework, we will show that each set  $Y$  contains a set  $X \subseteq Y$  which is maximal **independent**.

**Notation (Basis).**  $X$  is called a *basis* of  $Y$  in the above.

**Remark.** If  $X_1, X_2$  are two **bases** for  $Y$ , then  $|X_1| = |X_2|$ .

**Notation (Dimension).** The *dimension*  $\dim Y$  of  $Y$  is this cardinality of its **basis**.

**Definition 4.1.8 (Type).** Let  $\mathcal{M}$  be an  $\mathcal{L}$ -structure,  $\bar{c} \in M$ ,  $A \subseteq M$ . The *type* of  $\bar{c}$  over  $A$  (in  $\mathcal{M}$ ) is

$$\text{tp}^{\mathcal{M}}(\bar{c}/A) = \{\varphi(\bar{x}, \bar{a}) \mid \mathcal{M} \models \varphi(\bar{c}, \bar{a}) \text{ and } \bar{a} \in A\}.$$

**Notation.** Where we omit  $/A$ , we just mean  $/\emptyset$ .

**Lemma 4.1.3.** Let  $T$  be a **complete strongly minimal theory**, and  $\mathcal{M}, \mathcal{N} \models T$ . Let  $\bar{a} \in M$  and  $\bar{b} \in N$  be **independent** tuples of the same size. Then  $\text{tp}^{\mathcal{M}}(\bar{a}) = \text{tp}^{\mathcal{N}}(\bar{b})$ , i.e.,  $\mathcal{M} \models \varphi(\bar{a}) \Leftrightarrow \mathcal{N} \models \varphi(\bar{b})$ .

**Proof.** We do an induction on the lengths of  $\bar{a}, \bar{b}$ . Start with  $n = 1$ . Let  $a, b$  be **independent**, then

$$\text{tp}(a) = \{\varphi(x) \mid \varphi(x) \text{ has \textbf{cofinitely} many solutions in } \mathcal{M}\}$$

since each **formula** has either finitely many or **cofinitely** many solutions (and  $\neg\varphi(x)$  has the opposite). If  $\varphi(x)$  has finitely many solutions, then  $\mathcal{M} \models \neg\varphi(a)$  because  $a \notin \text{acl}(\emptyset)$ .

**Example.** In ACF,  $\text{tp}(a) = \{p(a) \neq 0 \mid p \in \mathbb{Q}[x], p \neq 0\}$  (or “generated by” these). Also,  $(1+1)x$  being 0 depends on the **characteristic**.

Also, we have

$$\text{tp}(b) = \{\varphi(x) \mid \varphi(x) \text{ has \textbf{cofinitely} many solutions in } \mathcal{N}\}.$$

Since  $\varphi$  has  $k$  non-solutions in  $\mathcal{M}$ ,  $\mathcal{M} \models \exists^{=k}x \neg\varphi(x)$ , so since  $T$  is **complete**,  $T \models \exists^{=k}x \neg\varphi(x)$ , and  $\mathcal{N} \models \exists^{=k}x \neg\varphi(x)$ . So if  $\varphi$  has **cofinitely** many (all but  $k$ ) solutions in  $\mathcal{M}$ , the same is true in  $\mathcal{N}$ . Hence,  $\text{tp}(a) = \text{tp}(b)$ .

**Example.** The **completeness** is important: ACF is a non-example.

For  $n+1$ , let  $\bar{a}a'$  and  $\bar{b}b'$  be **independent**  $(n+1)$ -tuples, so  $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$ . Suppose  $\mathcal{M} \models \varphi(\bar{a}a')$ . Since  $\mathcal{M} \models T$  is **strongly minimal**, and  $a' \notin \text{acl}(\bar{a})$ ,  $\{c \in M \mid \mathcal{M} \models \varphi(\bar{a}, c)\}$  is **cofinite**, with complement of size  $k$ ,  $\mathcal{M} \models \exists^{=k}x (\bar{a}, x)$ , so  $\exists^{=k}x \neg\varphi(\bar{y}, x) \in \text{tp}(\bar{a}) = \text{tp}(\bar{b})$ . So,  $\{c \in N \mid \mathcal{N} \models (\bar{b}, c)\}$  is finite. Since  $b' \notin \text{acl}(\bar{b})$ ,  $\mathcal{N} \models \varphi(\bar{b}, b')$ . ■

**Lemma 4.1.4.** Let  $T$  be a **complete strongly minimal theory** such that and  $\mathcal{M} \models T$ . If  $c \in \text{acl}(A)$ , then there is a **formula**  $\varphi(x, \bar{a}) \in \text{tp}(c/A)$  such that for any  $\psi(x, \bar{a}') \in \text{tp}(c/A)$ ,  $\varphi(x, \bar{a})$  implies  $\psi(x, \bar{a}')$ , i.e.,  $T \models \forall x \forall \bar{y} (\varphi(x, \bar{y}) \rightarrow \psi(x, \bar{y}))$ .<sup>a</sup>

<sup>a</sup>Expanding to get  $\bar{a} = \bar{a}'$ .

**Proof.** There is some **formula**  $\varphi(x, \bar{a}) \in \text{tp}(c/A)$  such that  $\{c' \in M \mid \mathcal{M} \models \varphi(\bar{c}, \bar{a})\}$ , finite. Choose  $\varphi$  such that  $K$  is minimal.

**Claim.** This  $\varphi$  works.

**Proof.** Let  $\psi(x, \bar{a}') \in \text{tp}(c/A)$ . If  $\varphi(x, \bar{a})$  does not imply  $\psi(x, \bar{a}')$ , there is some  $c'$  with  $\mathcal{M} \models \varphi(c', \bar{a}) \wedge \neg\psi(c', \bar{a}')$ . So

$$\{d \mid \mathcal{M} \models \varphi(d, \bar{a}) \wedge \psi(d, \bar{a}')\} \subseteq \{d \mid \mathcal{M} \models \varphi(d, \bar{a})\}.$$

Since  $c'$  is in the right-hand side, but not the left-hand side, but the right-hand side was as small as possible (by choice of  $K$ ), a contradiction, so  $\varphi$  does imply  $\psi$ . ⊗

Most correct...



**Theorem 4.1.7.** Let  $T$  be a **complete strongly minimal theory**. If  $\mathcal{M}, \mathcal{N} \models T$ , then  $\mathcal{N} \cong \mathcal{M}$  if and only if  $\dim \mathcal{N} = \dim \mathcal{M}$ .

**Proof.** Suppose  $\dim \mathcal{M} = \dim \mathcal{N}$ , and let  $A, B$  be **bases** for  $\mathcal{M}, \mathcal{N}$ . Then, we have  $|A| = |B|$ , so there exists a bijection  $f: A \rightarrow B$ , which is a **partial elementary map** since for  $f, \bar{a} \in A$ ,  $\text{tp}(\bar{a}) = \text{tp}(f(\bar{a}))$ .

**Notation** (Partial elementary map). A map  $g: U \subseteq M \rightarrow N$  is a *partial elementary map* if  $\mathcal{M} \models \varphi(\bar{a}) \Leftrightarrow \mathcal{N} \models \varphi(g(\bar{a}))$  for  $\bar{a} \subseteq \text{dom } g$ .<sup>a</sup>

<sup>a</sup>This implies injectivity.

Let  $g$  be a maximal **partial elementary map** from  $\mathcal{M} \rightarrow \mathcal{N}$  extending  $f$  (**Zorn's lemma** shows that such a  $g$  exists).

**Claim.**  $g$  is an **isomorphism**, i.e.,  $\text{dom } g = M$  and  $\text{Im } g = N$ .

**Proof.** Assume that  $c \in M - \text{dom } g$ . We know that  $c \in \text{acl}(A)$ , so let  $\varphi(x, \bar{d})$  be the **formula** from **Lemma 4.1.4** that isolates  $\text{tp}(c/\text{dom } g)$ . Then,  $\mathcal{M} \models \exists x \varphi(x, \bar{d})$ , implying  $\mathcal{N} \models \exists x \varphi(x, g(\bar{d}))$ . Let  $c' \in N$  witness this. Then  $\text{tp}^{\mathcal{M}}(c/\text{dom } g) = \text{tp}^{\mathcal{N}}(c'/\text{Im } g)$  after identifying  $\text{dom } g$  and  $\text{Im } g$ . Then we can define  $g(c) = c'$  to extend  $g$  but remain **partial elementary**, contradicting to the maximality of  $g$ , so  $\text{dom } g = M$ .

Following the same argument, we can show that  $\text{Im } g = N$ , hence  $g$  is an **isomorphism**.  $\otimes$

■

## Lecture 20: Fraïssé Theorem

### 4.1.5 Other Examples of Quantifier Elimination

21 Mar. 14:30

**As previously seen.** ACF, DLO, and also the **random graph theory**.<sup>a</sup>

<sup>a</sup>This is very similar to the argument we gave for DLO.

We discuss some other examples of **theories** which admit **quantifier elimination**.

#### Vector Spaces

On the problem set, we will show that the **theory** of infinite (dimensional) vector spaces admits **quantifier elimination** and is **strongly minimal**.

#### Torsion-Free Abelian Groups

**As previously seen** (Torsion-free). An Abelian group  $G$  is said to be *torsion-free* if no element other than  $e$  has finite order.

**Definition 4.1.9** (Divisible). A **torsion-free** group  $G$  is *divisible* if for every  $n \in \mathbb{N}$  and  $g \in G$ , there is  $h \in G$  such that

$$nh = \underbrace{h + \dots + h}_{n \text{ times}} = g.$$

Since  $G$  is **torsion-free**, for each  $g$ , the corresponding  $h$  is unique.<sup>1</sup> The **theory** DAG of **torsion-free divisible** Abelian groups (i.e.,  $\mathbb{Q}$ -vector spaces in **language**  $\{+, 0, -\}$ ) admits **quantifier elimination**.

#### Ordered Torsion-Free Divisible Abelian Groups

On top of DAG, if we also add an ordering, we get ODAG, the **theory** of ordered **torsion-free divisible** Abelian groups, which also admits **quantifier elimination**.

<sup>1</sup>As if  $nh = g = nh'$ , then  $n(h - h') = 0$  so  $h = h'$ .

## Presburger Arithmetic

Presburger arithmetic is  $\text{Th}(\mathbb{N}, 0, 1, +)$ .

## Real Closed Fields

**As previously seen** (Real-closed field). A *real-closed field* is a field  $F$  that has the same first-order properties as the field of real numbers.

The **theory** of **real-closed fields**, denoted as RCF, does not admit **quantifier elimination** in the **language**  $\mathcal{L} = \{0, 1, +, -, \cdot\}$ . This is because the **formula**

$$\varphi(x, y) = \exists x \ x - y = z^2$$

is not equivalent to a **quantifier-free formula**.

**Intuition.** This **formula** is defining the ordering of the field.

**Claim.**  $\varphi(x, y) = \exists z \ x - y = z^2$  is not equivalent to a **quantifier-free formula** in  $\mathcal{L} = \{0, 1, +, -, \cdot\}$ .

**Proof.** The first way one might try to prove this is to show that there are **real-closed fields**  $\mathcal{M} \subseteq \mathcal{N}$  and  $a, b \in \mathcal{N}$  such that  $\mathcal{N} \models \varphi(a, b)$  but  $\mathcal{M} \models \neg\varphi(a, b)$ . but this strategy will not work, since  $\varphi(x, y)$  is also equivalent, in any **real-closed field**, to the **universal formula**  $\forall z \ y - x \neq z^2$ . this is because only one of  $x - y$  and  $y - x$  can have a square root in a **real-closed field**. Instead, one should think about the test for **quantifier elimination**.

**Proposition 4.1.3.** There is no **quantifier-free formula**  $\psi(x, y)$  such that RCF proves that  $\psi(x, y)$  and  $\varphi(x, y)$  are equivalent.

**Proof.** ■

⊗

But if we add in the ordering as a symbol in the **language**, then in the **language**  $\mathcal{L}_{<}$ , the **theory**  $\text{Th}(\mathbb{R}, 0, 1, +, -, \cdot, <)$  of **real-closed** ordered fields does admit **quantifier elimination**. This was shown by Tarski in the 1940's as part of showing that the **theory** of  $\mathbb{R}$  is **decidable**.

**Example.** An example of eliminating quantifiers from a **formula** that we already know is

$$\text{RCF} \models \exists x \ (a \neq 0 \wedge ax^2 + bx + c = 0) \leftrightarrow a \neq 0 \wedge b^2 - 4ac \geq 0.$$

After proving **quantifier elimination**, we can analyze the **definable** sets like we did for **algebraically closed** fields.

**Remark** (*o-minimal*). The **theory** of **real-closed fields** is not **strongly minimal**, but instead what is called *o-minimal*.<sup>a</sup>

<sup>a</sup>There is a well-developed theory of *o-minimality* which would be an entire course in itself.

**Proof.** The **definable** sets in one variable are finite unions of intervals and points. ⊗

**Remark.** Tarski-Seidenberg algorithm can solve all Euclidean geometry problems.

## 4.2 Ultrapowers


**Intuition.** Fraïssé theorem. Idea is that to build “universal” countable **structures** with “all possible finitary behaviors”.

Assume (for now)  $\mathcal{L}$  is relational.

**Definition 4.2.1 (Age).** For an  $\mathcal{L}$ -structure  $\mathcal{M}$ , its *age* is the class  $\text{Age}(\mathcal{M})$  of all finite  $\mathcal{L}$ -substructure which extend into  $\mathcal{M}$ .

**Definition 4.2.2 (Hereditary property).** A class  $\mathbb{K}$  of finite  $\mathcal{L}$ -structure has the *hereditary property* if for all  $\mathcal{B} \in \mathbb{K}$  and  $\mathcal{A}$  which extends into  $\mathcal{B}$ ,  $\mathcal{A} \in \mathbb{K}$ .

**Definition 4.2.3 (Joint embedding property).** A class  $\mathbb{K}$  of finite  $\mathcal{L}$ -structure has the *joint embedding property* if for all  $\mathcal{A}, \mathcal{B} \in \mathbb{K}$ , there exists  $\mathcal{C} \in \mathbb{K}$  with embedding




**Theorem 4.2.1 (Baby Fraïssé theorem).** A class  $\mathbb{K}$  of finite  $\mathcal{L}$ -structure is  $\text{Age}(\mathcal{M})$  for some countable  $\mathcal{M}$  if and only if

- $\mathbb{K}$  is countable up to isomorphism;
- $\mathbb{K} \neq \emptyset$ ;
- $\mathbb{K}$  has the hereditary property;
- $\mathbb{K}$  has the joint embedding property.

**Proof.** The forward direction is clear. For the backward direction, let  $\mathbb{K} = \{\mathcal{A}_0, \mathcal{A}_1, \dots\}$ , we construct  $\mathcal{M}_0 \subseteq \mathcal{M}_1 \subseteq \mathcal{M}_2 \subseteq \dots$  inductively as follows:

- $\mathcal{M}_0 := \mathcal{A}_0$ .
- $\mathcal{M}_{n+1}$  is chosen from  $\mathbb{K}$  using joint embedding property such that  $\mathcal{M}_n \hookrightarrow \mathcal{M}_{n+1} \hookrightarrow \mathcal{A}_{n+1}$ .



Let  $\mathcal{M} := \bigcup_n \mathcal{M}_n$ . Then,

- $\mathbb{K} \subseteq \text{Age}(\mathcal{M})$ : because each  $\mathcal{A}_n \in \mathbb{K}$  is embedded into  $\mathcal{M}_n \subseteq \mathcal{M}$ , so  $\mathcal{A}_n$  is embedded into  $\mathcal{M}$ ;
- $\text{Age}(\mathcal{M}) \subseteq \mathbb{K}$ : for a finite subset  $\mathcal{N} \subseteq \mathcal{M}$ , we have  $\mathcal{N} \subseteq \mathcal{M}_n$  for some  $n$ , so by hereditary property,  $\mathcal{M}_n \in \mathbb{K}$ , hence  $\mathcal{N} \in \mathbb{K}$ .

■


**Example.**  $\text{Age}(\mathbb{Q}, <) = \text{Age}(\mathbb{N}, <) = \text{Age}(\mathbb{Z}, <) = \{\text{all finite linear orders}\}$ .

**Example.**  $\text{Age}(\text{random graph theory}) = \{\text{all finite graphs}\} = \text{Age}(\coprod \{\text{finite graphs}\})$ .




**Definition 4.2.4** (Ultrahomogeneous). A countable structure  $\mathcal{M}$  is *ultrahomogeneous* if for any finite subsets  $\mathcal{A}, \mathcal{B} \subseteq \mathcal{M}$  and isomorphism  $f: \mathcal{A} \rightarrow \mathcal{B}$ , then there is an isomorphism  $g: \mathcal{M} \cong \mathcal{M}$  extending  $f$ .


**Definition 4.2.5** (Amalgamation property). A class  $\mathbb{K}$  of finite  $\mathcal{L}$ -structure has the *amalgamation property* if for all  $\mathcal{A}$  such that




that  $\tilde{f} \circ f = \tilde{g} \circ g$ , i.e., the following diagram commutes:



**Definition 4.2.6** (Extension property). A countable structure  $\mathcal{M}$  has the *extension property* w.r.t. a class  $\mathbb{K}$  of (finite) structure if for all  $f: \mathcal{A} \hookrightarrow \mathcal{B}$  and  $g: \mathcal{A} \hookrightarrow \mathcal{M}$ , there exists  $\tilde{f}: \mathcal{B} \hookrightarrow \mathcal{M}$  such that  $\tilde{f} \circ f = g$ .



**Note.**  $\mathcal{M}$  ultrahomogeneous implies  $\mathcal{M}$  has extension property w.r.t.  $\text{Age}(\mathcal{M})$ .



**Theorem 4.2.2** (Fraïssé theorem). A class  $\mathbb{K}$  of finite  $\mathcal{L}$ -structure is  $\text{Age}(\mathcal{M})$  for an ultrahomogeneous countable  $\mathcal{M}$  if and only if

- $\mathbb{K}$  is countable up to isomorphism;
- $\mathbb{K} \neq \emptyset$ ;
- $\mathbb{K}$  has the HP;
- $\mathbb{K}$  has the JEP;
- $\mathbb{K}$  has the AP.

Moreover, in that case, there exists a unique-up-to-isomorphism countable ultrahomogeneous  $\mathcal{M}$  such that  $\text{Age}(\mathcal{M}) = \mathbb{K}$ .

**Definition 4.2.7** (Fraïssé class). A class  $\mathbb{K}$  with properties in [Fraïssé theorem](#) is called a *Fraïssé class*.

**Definition 4.2.8** (Fraïssé limit). The countable [ultrahomogeneous](#)  $\mathcal{M}$  such that  $\text{Age}(\mathcal{M}) = \mathbb{K}$  is the *Fraïssé limit* of  $\mathbb{K}$ , denoted as  $\text{Flm}(\mathbb{K})$ .

Now, we try to prove the [Fraïssé theorem](#).

**Proof of Theorem 4.2.2.** To prove the forward direction, we need only to check [AP](#) for  $\text{Age}(\mathcal{M})$ . We use [EP](#) to extend  $\mathcal{A} \hookrightarrow \mathcal{B} \hookrightarrow \mathcal{M}$  along  $\mathcal{A} \hookrightarrow \mathcal{C}$ . Let  $\mathcal{D}$  be the union of images of  $\mathcal{C} \hookrightarrow \mathcal{M}$  and  $\mathcal{B} \hookrightarrow \mathcal{M}$ .

$$\begin{array}{ccc}
 \mathcal{A} & \xrightarrow{f} & \mathcal{B} \\
 \downarrow & \searrow & \downarrow \\
 \text{Age}(\mathcal{M}) \ni \mathcal{C} & \xrightarrow[\exists \text{ by EP}]{} & \mathcal{D} \subseteq \mathcal{M}
 \end{array}$$

For the backward direction,  
 where  $\mathcal{M}'_n$  is obtained from  $\mathcal{M}_n$  by amalgamating ■

# Appendix

# Bibliography

- [HH97] W. Hodges and S.M.S.W. Hodges. *A Shorter Model Theory*. Cambridge University Press, 1997. ISBN: 9780521587136. URL: <https://books.google.com/books?id=S6QYeuo4p1EC>.
- [Hin05] P.G. Hinman. *Fundamentals of Mathematical Logic*. Taylor & Francis, 2005. ISBN: 9781568812625. URL: <https://books.google.com/books?id=xA6D8o72qAgC>.
- [Mar02] D. Marker. *Model Theory : An Introduction*. Graduate Texts in Mathematics. Springer New York, 2002. ISBN: 9780387987606. URL: <https://books.google.com/books?id=gkvogoiEnuYC>.
- [Rob49] Julia Robinson. “Definability and decision problems in arithmetic”. In: *The Journal of Symbolic Logic* 14.2 (1949), pp. 98–114. DOI: [10.2307/2266510](https://doi.org/10.2307/2266510).