

MATH597  
Analysis II

Pingbang Hu

December 16, 2023

## Abstract

This is the graduate-level real analysis course taught by [Jinho Baik](#) at University of Michigan. Everything is built up from measure theory, like integration and differentiation. Topics include measure theory, Hilbert/Banach spaces,  $L^p$  spaces, and some Fourier Analyses.

There are some proofs is **intended** left as assignments, and for completeness, I put them in [Appendix A](#), use it in your **own risks**! You'll lose the chance to practice and really understand the materials.

Additionally, we'll use Folland [[FF99](#)] as our main text, while using Tao [[Tao13](#)] and Axler [[Axl19](#)] as supplementary references.

# Contents

<b>1</b>	<b>Measure</b>	<b>3</b>
1.1	$\sigma$ -Algebra	3
1.2	Measures	7
1.3	Outer Measures	10
1.4	Hahn-Kolmogorov Theorem	16
1.5	Borel Measures on $\mathbb{R}$	20
1.6	Lebesgue-Stieltjes Measure on $\mathbb{R}$	24
1.7	Regularity Properties of Lebesgue-Stieltjes Measures	26
<b>2</b>	<b>Integration</b>	<b>30</b>
2.1	Measurable Function	30
2.2	Integration of Nonnegative Functions	34
2.3	Integration of Complex Functions	39
2.4	$L^1$ Space	41
2.5	Riemann Integrability	44
2.6	Modes of Convergence	45
<b>3</b>	<b>Product Measure</b>	<b>50</b>
3.1	Product $\sigma$ -Algebra	50
3.2	Product Measures	53
3.3	Monotone Class Lemma	54
3.4	Fubini-Tonelli Theorem	55
3.5	Lebesgue Measure on $\mathbb{R}^d$	57
<b>4</b>	<b>Differentiation on Euclidean Space</b>	<b>59</b>
4.1	Hardy-Littlewood Maximal Function	59
4.2	Lebesgue Differentiation Theorem	61
<b>5</b>	<b>Normed Vector Space</b>	<b>64</b>
5.1	Metric Spaces and Normed Spaces	64
5.2	$L^p$ Space	65
5.3	Embedding Properties of $L^p$ Spaces	69
5.4	Banach Spaces	71
5.5	Bounded Linear Transformations	74
5.6	Dual of $L^p$ Spaces	76
<b>6</b>	<b>Signed and Complex Measures</b>	<b>79</b>
6.1	Signed Measures	79
6.2	Absolutely Continuous Measures	84
6.3	Lebesgue Differentiation Theorem for Regular Borel Measures	88
6.4	Monotone Differentiation Theorem	91
6.5	Functions of Bounded Variation	94
<b>7</b>	<b>Hilbert Spaces</b>	<b>101</b>
7.1	Inner Product Spaces	101
7.2	Orthonormal Bases	104

---

<b>8</b>	<b>Introduction to Fourier Analysis</b>	<b>109</b>
8.1	Fourier Series . . . . .	109
<b>A</b>	<b>Additional Proofs</b>	<b>116</b>
A.1	Measure . . . . .	116
A.2	Integration . . . . .	119
A.3	Produce Measure . . . . .	119
A.4	Differentiation on Euclidean Space . . . . .	119
A.5	Normed-Vector-Space . . . . .	119
A.6	Signed and Complex Measures . . . . .	119
A.7	Hilbert Spaces . . . . .	120
A.8	Introduction to Fourier Analysis . . . . .	120

# Chapter 1

## Measure

### Lecture 1: $\sigma$ -Algebra and Motivations

#### 1.1 $\sigma$ -Algebra

05 Jan. 11:00

##### 1.1.1 Motivated Examples

Before we start, we first see some examples.

**Example (Finite power set).** Let  $X = \{a, b, c\}$ . Then

$$\mathcal{P}(X) := \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}\},$$

which is the *power set* of  $X$ . In general, if  $|X| = n$  with  $n < \infty$ ,  $|\mathcal{P}(X)| = 2^n$ .

**Example (Infinite power set).** If  $n = \infty$ , say  $X = \mathbb{N}$ , then  $\mathcal{P}(\mathbb{N})$  is an uncountable set while  $\mathbb{N}$  is a countable set. We can see this as follows. Consider

$$\phi: \mathcal{P}(\mathbb{N}) \rightarrow [0, 1], \quad A \mapsto 0.a_1a_2a_3 \dots (\text{base } 2),$$

where

$$a_i = \begin{cases} 1, & \text{if } i \in A \\ 0, & \text{if } i \notin A, \end{cases}$$

and for example,  $A$  can be  $A = \{2, 3, 6, \dots\} \subseteq \mathbb{N}$ . Note that  $\phi$  is surjective, hence we have

$$|\mathcal{P}(\mathbb{N})| \geq |[0, 1]|.$$

But since  $[0, 1]$  is uncountable, so is  $\mathcal{P}(\mathbb{N})$ .

We like to *measure* the *size* of subsets of  $X$ . Hence, we are intriguing to define a map  $\mu$  such that

$$\mu: \mathcal{P}(X) \rightarrow [0, \infty].$$

**Example.** Let  $X = \{0, 1, 2\}$ . Then we want to define  $\mu: \mathcal{P}(X) \rightarrow [0, \infty]$ , we can have

- $\mu(A) = |A|$ . Then we have
  - $\mu(\{0, 1\}) = 2$
  - $\mu(\{0\}) = 1$
- $\mu(A) = \sum_{i \in A} 2^i$ . Then we have

$$- \mu(\{0, 1\}) = 2^0 + 2^1 = 3$$

**Example.** Let  $X = \{0\} \cup \mathbb{N}$ . Then we want to define  $\mu: \mathcal{P}(\mathbb{N}) \rightarrow [0, \infty]$ , we can have

- $\mu(A) = |A|$ . Then we have
  - $\mu(\{2, 3, 4, 5, \dots\}) = \infty = \mu(\{\text{even numbers}\})$
- $\mu(A) = e^{-1} \sum_{i \in A} \frac{1}{i!}$ . Then we have
  - $\mu(\{0, 2, 4, 6, \dots\}) = e^{-1} (1 + \frac{1}{2!} + \frac{1}{3!} + \dots)$
- $\mu(A) = \sum_{i \in A} a_i$

**Example.** Let  $X = \mathbb{R}$ . Then we want to define  $\mu: \mathcal{P}(\mathbb{R}) \rightarrow [0, \infty]$ , we can have

- $\mu(A) = |A|$
- $\mu((a, b)) = b - a$ .

**Problem.** Can we extend this map to all of  $\mathcal{P}(\mathbb{R})$ ?

**Answer.** No! ⊗

- $\mu((a, b)) = e^b - e^a$ .

**Problem.** Can we extend this map to all of  $\mathcal{P}(\mathbb{R})$ ?

**Answer.** No! ⊗

We immediately see the problems. To extend our native measure method into  $\mathbb{R}$  is hard and will cause something counter-intuitive!<sup>1</sup> Hence, rather than define measurement on *all* subsets in the power set of  $X$ , we only focus on *some* subsets. In other words, we want to define

$$\mu: \mathcal{P}(\mathbb{R}) \supset \mathcal{A} \rightarrow [0, \infty].$$

### 1.1.2 $\sigma$ -Algebra

Now, we start from the definition of the most fundamental element in measure theory.

**Definition 1.1.1 ( $\sigma$ -algebra).** Let  $X$  be a set. A collection  $\mathcal{A}$  of subsets of  $X$ , i.e.,  $\mathcal{A} \subset \mathcal{P}(X)$  is called a  $\sigma$ -algebra on  $X$  if

- $\emptyset \in \mathcal{A}$ .
- $\mathcal{A}$  is closed under complements. i.e., if  $A \in \mathcal{A}$ ,  $A^c = X \setminus A \in \mathcal{A}$ .
- $\mathcal{A}$  is closed under countable unions. i.e., if  $A_i \in \mathcal{A}$ , then  $\bigcup_{i=1}^{\infty} A_i \in \mathcal{A}$ .

**Remark.** There are some easy properties we can immediately derive.

- $X \in \mathcal{A}$  from  $X = X \setminus \emptyset$  and  $\mathcal{A}$  is closed under complement.<sup>a</sup>
- $\bigcap_{i=1}^{\infty} A_i = (\bigcup_{i=1}^{\infty} A_i^c)^c$ , namely  $\mathcal{A}$  is closed under countable intersections.
- $A_1 \cup A_2 \cup \dots \cup A_n = A_1 \cup A_2 \cup \dots \cup A_n \cup \emptyset \cup \emptyset \cup \dots$ , hence  $\mathcal{A}$  is closed under finite unions and intersections.

<sup>a</sup>Recall that  $\emptyset \in \mathcal{A}$ .

<sup>1</sup>E.g., [Banach-Tarski paradox](#).

**Note.** The definition of  $\sigma$ -algebra should remind us the definition of topological basis, and this is indeed the case. We can consider a topological space and put some structure on the  $\sigma$ -algebra  $\mathcal{A}$ , which gives us the following.

**Definition 1.1.2 (Borel set).** Given a topological space  $X$ , a *Borel set* is any set in  $X$  that can be formed from open sets through the operations of countable union, countable intersection and relative complement.

## Lecture 2: Measure

**Example.** We first see some examples.

07 Jan. 11:00

- (a) Let  $\mathcal{A} = \mathcal{P}(X)$ , which is the power  $\sigma$ -algebra.
- (b) Let  $\mathcal{A} = \{\emptyset, X\}$ , which is a trivial  $\sigma$ -algebra.
- (c) Let  $B \subset X$ ,  $B \neq \emptyset$ ,  $B \neq X$ . Then we see that  $\mathcal{A} = \{\emptyset, B, B^c, X\}$  is a  $\sigma$ -algebra.

**Lemma 1.1.1.** Let  $\mathcal{A}_\alpha$ ,  $\alpha \in I$ , be a family of  $\sigma$ -algebra on  $X$ . Then  $\bigcap_{\alpha \in I} \mathcal{A}_\alpha$  is a  $\sigma$ -algebra on  $X$ .

**Proof.** A simple proof can be made as follows. Firstly,  $\emptyset \in \mathcal{A}_\alpha$  for every  $\alpha$  clearly. Moreover, closure under complement and countable unions for every  $\mathcal{A}_\alpha$  implies the same must be true for  $\bigcap_{\alpha \in I} \mathcal{A}_\alpha$ . Hence,  $\bigcap_{\alpha \in I} \mathcal{A}_\alpha$  is a  $\sigma$ -algebra.

**Remark.** Notice that  $I$  may be an uncountable intersection. ■

The above allows us to give the following definition.

**Definition 1.1.3 (Generation of  $\sigma$ -algebra).** Given  $\mathcal{E} \subset \mathcal{P}(X)$ , where  $\mathcal{E}$  is not necessarily a  $\sigma$ -algebra. Let  $\langle \mathcal{E} \rangle$  be the intersection of all  $\sigma$ -algebras on  $X$  containing  $\mathcal{E}$ , then we call  $\langle \mathcal{E} \rangle$  the  $\sigma$ -algebra generated by  $\mathcal{E}$ .

**Remark.** Clearly,  $\langle \mathcal{E} \rangle$  is the smallest  $\sigma$ -algebra containing  $\mathcal{E}$ , and it is unique. To check the uniqueness, we suppose there are two different  $\langle \mathcal{E} \rangle_1$  and  $\langle \mathcal{E} \rangle_2$  generated from  $\mathcal{E}$ . It's easy to show

$$\langle \mathcal{E} \rangle_1 \subseteq \langle \mathcal{E} \rangle_2,$$

and by symmetry, they are equal.

**Example.** We see that  $\{\emptyset, B, B^c, X\} = \langle \{B\} \rangle = \langle \{B^c\} \rangle$ .

**Lemma 1.1.2.** We have the following.

- (a) Given  $\mathcal{A}$  a  $\sigma$ -algebra,  $\mathcal{E} \subset \mathcal{A} \subset \mathcal{P}(X) \Rightarrow \langle \mathcal{E} \rangle \subset \mathcal{A}$ .
- (b)  $\mathcal{E} \subset \mathcal{F} \subset \mathcal{P}(X) \Rightarrow \langle \mathcal{E} \rangle \subset \langle \mathcal{F} \rangle$ .

**Proof.** We'll see that after proving the first claim, the second follows smoothly.

- (a) The first claim is trivial, since we know that  $\langle \mathcal{E} \rangle$  is the smallest  $\sigma$ -algebra containing  $\mathcal{E}$ , then if  $\mathcal{E} \subset \mathcal{A}$ , we clearly have  $\langle \mathcal{E} \rangle \subset \mathcal{A}$  by the definition.

(b) The second claim is also easy. From the first claim and the definition, we have

$$\mathcal{E} \subset \mathcal{F} \subset \langle \mathcal{F} \rangle \Rightarrow \langle \mathcal{E} \rangle \subset \langle \mathcal{F} \rangle.$$

■

At this point, we haven't put any specific structure on  $X$ . Now we try to describe those spaces with good structure, which will give the space some nice properties.

**Definition 1.1.4 (Borel  $\sigma$ -algebra).** For a topological space  $X$ , the *Borel  $\sigma$ -algebra* on  $X$ , denoted as  $\mathcal{B}(X)$ , is the  $\sigma$ -algebra generated by the collection of all open sets in  $X$ .

**Example.** We see that  $\mathcal{B}(\mathbb{R})$  contains

- $\mathcal{E}_1 = \{(a, b) \mid a < b; a, b \in \mathbb{R}\}$ .
- $\mathcal{E}_2 = \{[a, b] \mid a < b; a, b \in \mathbb{R}\}$  since  $[a, b] = \bigcap_{n=1}^{\infty} (a - \frac{1}{n}, b + \frac{1}{n})$ .
- $\mathcal{E}_3 = \{(a, b] \mid a < b; a, b \in \mathbb{R}\}$  since  $(a, b] = \bigcap_{n=1}^{\infty} (a, b + \frac{1}{n})$ .
- $\mathcal{E}_4 = \{[a, b) \mid a < b; a, b \in \mathbb{R}\}$  since  $[a, b) = \bigcap_{n=1}^{\infty} (a - \frac{1}{n}, b)$ .
- $\mathcal{E}_5 = \{(a, \infty) \mid a \in \mathbb{R}\}$  since  $(a, \infty) = \bigcup_{n=1}^{\infty} (a, a + n)$ .
- $\mathcal{E}_6 = \{[a, \infty) \mid a \in \mathbb{R}\}$  since  $[a, \infty) = \bigcup_{n=1}^{\infty} [a, a + n)$ .
- $\mathcal{E}_7 = \{(-\infty, b) \mid b \in \mathbb{R}\}$  since  $(-\infty, b) = \bigcup_{n=1}^{\infty} (b - n, b)$ .
- $\mathcal{E}_8 = \{(-\infty, b] \mid b \in \mathbb{R}\}$  since  $(-\infty, b] = \bigcup_{n=1}^{\infty} (b - n, b]$ .

**Proposition 1.1.1.**  $\mathcal{B}(\mathbb{R}) = \langle \mathcal{E}_i \rangle$  for each  $i = 1, \dots, 8$  in the above example.

**Proof.** Firstly, we see that  $\mathcal{E}_i \subset \mathcal{B}(\mathbb{R}) \Rightarrow \langle \mathcal{E}_i \rangle \subset \mathcal{B}(\mathbb{R})$  by Lemma 1.1.2. Secondly, by definition,  $\mathcal{B}(\mathbb{R}) = \langle \mathcal{E} \rangle$  where

$$\mathcal{E} = \{O \subseteq \mathbb{R} \mid O \text{ is open in } \mathbb{R}\}.$$

It's enough to show  $\mathcal{E} \subset \langle \mathcal{E}_i \rangle$  since if so,  $\langle \mathcal{E} \rangle \subseteq \langle \mathcal{E}_i \rangle$ , and clearly  $\langle \mathcal{E} \rangle \supseteq \langle \mathcal{E}_i \rangle = \mathcal{B}(\mathbb{R})$ , then we will have  $\langle \mathcal{E} \rangle = \langle \mathcal{E}_i \rangle$ . Let  $O \subset \mathbb{R}$  be an open set, i.e.,  $O \in \mathcal{E}$ . We claim that every open set in  $\mathbb{R}$  is a countable union of disjoint open intervals.<sup>a</sup> Thus,

$$O = \bigcup_{j=1}^{\infty} I_j,$$

where  $I_j$  open interval with the form of  $(a, b), (-\infty, b), (a, \infty), (-\infty, \infty)$ . For example,  $\mathcal{E}_1$  is trivially true, and

$$(a, b) = \bigcup_{n=1}^{\infty} \underbrace{\left[ a + \frac{1}{n}, b - \frac{1}{n} \right]}_{\substack{\in \mathcal{E}_2 \\ \in \langle \mathcal{E}_2 \rangle}}$$

shows the case for  $\mathcal{E}_2$  and

$$(a, \infty) = \bigcup_{k=1}^{\infty} (a, a + k)$$

shows the case for  $\mathcal{E}_5$ . It's now straightforward to check open intervals are in  $\langle \mathcal{E}_i \rangle$  for every  $i$ . ■

<sup>a</sup>See [here](#).

Now, to put a structure on a space, we define the following.



**Definition.** Let  $X$  be a space.

**Definition 1.1.5 (Measurable space).** A *measurable space* is a tuple of  $X$  and a  $\sigma$ -algebra  $\mathcal{A}$  on  $X$ , denoted by  $(X, \mathcal{A})$ .

In particular, if the  $\sigma$ -algebra is the *Borel  $\sigma$ -algebra* of  $X$ , then we give it a special name.

**Definition 1.1.6 (Borel space).** A *Borel space* is a tuple of  $X$  and  $\mathcal{B}(X)$ , denoted by  $(X, \mathcal{B}(X))$ .

**Remark.** The definition of *Borel space* means that  $X$  implicitly has some topological structure.

**Definition 1.1.7 (Measurable set).** Given a *measurable space*  $(X, \mathcal{A})$ , every  $E \in \mathcal{A}$  is a so-called  *$\mathcal{A}$ -measurable set*.

## 1.2 Measures

With the definition of *measurable space*, we now can refine our *measure* function  $\mu$  as follows.

**Definition 1.2.1 (Measure).** Given a *measurable space* on  $(X, \mathcal{A})$ , a *measure* is a function

$$\mu: \mathcal{A} \rightarrow [0, \infty]$$

such that

- (null empty set)  $\mu(\emptyset) = 0$ .
- (countable additivity)  $\mu(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \mu(A_i)$  if  $A_1, A_2, \dots \in \mathcal{A}$  are disjoint.

**Definition 1.2.2 (Measure space).** We denote  $(X, \mathcal{A}, \mu)$  as so-called a *measure space* given  $\mu$  is the *measure* on  $(X, \mathcal{A})$ .

**Notation.** We denote  $[0, \infty] := [0, \infty) \cup \{\infty\}$ .

**Remark.** We only want *countable additivity* but not uncountable additivity.

**Proof.** Consider the most intuitive *measure* on  $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ . Since we have

$$(0, 1] = (1/2, 1] \cup (1/4, 1/2] \cup (1/8, 1/4] \cup \dots$$

and also

$$(0, 1] = \bigcup_{x \in (0, 1]} \{x\}.$$

In the first case, we are claiming that

$$1 = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots = \mu((1/2, 1]) + \mu((1/4, 1/2]) + \mu((1/8, 1/4]) + \dots;$$

while in the second case, we are claiming that

$$1 = \sum_{x \in (0, 1]} 0$$

since  $\mu(\{x\}) = 0$  for  $x \in \mathbb{R}$ , which is clearly not what we want. \*

**Example (Counting measure).** For any  $(X, \mathcal{A})$ , we let  $\mu(A) := |A|$ . This is the so-called *counting measure*.

**Example (Dirac-Delta measure).** Let  $x_0 \in X$ . For any  $(X, \mathcal{A})$ , the *Dirac-Delta measure* at  $x_0$  is

$$\mu(A) = \begin{cases} 1, & \text{if } x_0 \in A; \\ 0, & \text{if } x_0 \notin A \end{cases}$$

for every  $A \in \mathcal{A}$ .

**Example.** For  $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$ , given  $A \in \mathcal{P}(\mathbb{N})$ ,

$$\mu(A) = \sum_{i \in A} a_i$$

where  $a_1, a_2, \dots \in [0, \infty)$ .

## Lecture 3: Construct a Measure

After seeing examples of [measures](#), we now want to construct it from ground up.

10 Jan. 11:00

**Note.** If  $A, B \in \mathcal{A}$  and  $A \subset B$ , then

$$\mu(B \setminus A) + \mu(A) = \mu(B) \Rightarrow \mu(B \setminus A) = \mu(B) - \mu(A) \text{ if } \mu(A) < \infty.$$

**Theorem 1.2.1.** Given  $(X, \mathcal{A}, \mu)$  be a [measure space](#). Then the following hold.

(a) Monotonicity.

$$A, B \in \mathcal{A}, A \subset B \Rightarrow \mu(A) \leq \mu(B).$$

(b) Countable subadditivity.

$$A_1, A_2, \dots \in \mathcal{A} \Rightarrow \mu\left(\bigcup_{i=1}^{\infty} A_i\right) \leq \sum_{i=1}^{\infty} \mu(A_i).$$

(c) Continuity from below.

$$\begin{cases} A_1, A_2, \dots \in \mathcal{A} \\ A_1 \subset A_2 \subset A_3 \subset \dots \end{cases} \Rightarrow \mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \lim_{n \rightarrow \infty} \mu(A_n).$$

(d) Continuity from above.

$$\begin{cases} A_1, A_2, \dots \in \mathcal{A} \\ A_1 \supset A_2 \supset A_3 \supset \dots \\ \mu(A_1) < \infty \end{cases} \Rightarrow \mu\left(\bigcap_{i=1}^{\infty} A_i\right) = \lim_{n \rightarrow \infty} \mu(A_n).$$

**Proof.** We prove this theorem one by one.

(a) Since  $A \subset B$ , hence we have

$$\mu(B) = \mu\left(\underbrace{(B \setminus A) \cup A}_{\text{disjoint}}\right) \stackrel{!}{=} \underbrace{\mu(B \setminus A)}_{\geq 0} + \mu(A) \geq \mu(A).$$

(b) This should be trivial from [countable additivity](#) with the fact that  $\mu(A) \geq 0$  for all  $A$ .

DIY!

(c) Let  $B_1 = A_1$ ,  $B_i = A_i \setminus A_{i-1}$  for  $i \geq 2$ , then

$$\bigcup_{i=1}^{\infty} A_i = \bigcup_{i=1}^{\infty} B_i$$

are a disjoint union and  $B_i \in \mathcal{A}$ , hence we see that

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mu(B_i) = \lim_{n \rightarrow \infty} \sum_{i=1}^n \mu(B_i).$$

With  $\mu(\bigcup_{i=1}^n B_i) = \mu(A_n)$ , we have

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \lim_{n \rightarrow \infty} \sum_{i=1}^n \mu(B_i) = \lim_{n \rightarrow \infty} \mu\left(\bigcup_{i=1}^n B_i\right) = \lim_{n \rightarrow \infty} \mu(A_n).$$

(d) Let  $E_i = A_1 \setminus A_i \Rightarrow E_i \in \mathcal{A}$ ,  $E_1 \subset E_2 \subset \dots$ . We then have

$$\bigcup_{i=1}^{\infty} E_i = \bigcup_{i=1}^{\infty} (A_1 \setminus A_i) = A_1 \setminus \left(\bigcap_{i=1}^{\infty} A_i\right),$$

which implies

$$\bigcap_{i=1}^{\infty} A_i = A_1 \setminus \left(\bigcup_{i=1}^{\infty} E_i\right) \Rightarrow \mu\left(\bigcap_{i=1}^{\infty} A_i\right) = \mu(A_1) - \mu\left(\bigcup_{i=1}^{\infty} E_i\right)$$

since  $\mu(\bigcup_{i=1}^{\infty} E_i) \leq \mu(A_1) < \infty$ . Then from [continuity from below](#), we further have

$$\mu\left(\bigcap_{i=1}^{\infty} A_i\right) = \mu(A_1) - \lim_{n \rightarrow \infty} \mu(E_n) = \mu(A_1) - \lim_{n \rightarrow \infty} (\mu(A_1) - \mu(A_n)).$$

From [monotonicity](#), we see that  $\mu(A_n) \leq \mu(A_1) < \infty$ , hence we can split the limit and further get

$$\mu\left(\bigcap_{i=1}^{\infty} A_i\right) = \mu(A_1) - \mu(A_1) + \lim_{n \rightarrow \infty} \mu(A_n) = \lim_{n \rightarrow \infty} \mu(A_n).$$

■

**Note.** Sometimes we also call [continuity from below](#) property as monotone convergence theorem for sets. We'll later see the important [monotone convergence theorem](#) for integral, which is in different content.

**Remark (Condition of continuity from above).** The condition  $\mu(A_1) < \infty$  in [continuity from above](#) is necessary.

**Proof.** Given  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), \mu)$  where  $\mu$  is the [counting measure](#). Then we see

- $A_n = \{n, n+1, n+2, \dots\} \Rightarrow \mu(A_n) = \infty$
- $A_1 \supset A_2 \supset A_3 \supset \dots$
- $\bigcap_{i=1}^{\infty} A_i = \emptyset \Rightarrow \mu(\bigcap_{i=1}^{\infty} A_i) = 0$

We see that in this case, since  $\mu(A_1) \not< \infty$ , hence [continuity from above](#) doesn't hold. ⊗

We now try to characterize some properties of a [measure space](#).

**Definition.** Let  $(X, \mathcal{A}, \mu)$  be a [measure space](#).

**Definition 1.2.3** ( $\mu$ -null set).  $A \subset X$  is a  $\mu$ -null set if  $A \in \mathcal{A}$  and  $\mu(A) = 0$ .

**Definition 1.2.4** ( $\mu$ -subnull set).  $A \subset X$  is a  $\mu$ -subnull set if there exists a  $\mu$ -null set  $B$  such that  $A \subset B$ .

**Definition 1.2.5** (Complete measure space).  $(X, \mathcal{A}, \mu)$  is a complete [measure space](#) if every  $\mu$ -subnull set is  $\mathcal{A}$ -measurable.

**Note.** We see that for a  $\mu$ -subnull set, it's not necessary  $\mathcal{A}$ -measurable if the [measure space](#) is not complete.

**Remark.** From the property of [measure](#), the condition for a [measure space](#)  $(X, \mathcal{A}, \mu)$  being complete is equivalent to saying that every  $\mu$ -subnull set is a  $\mu$ -null set.

**Proof.** This follows from the [monotonicity](#) of a [measure](#) and the fact that a [measure](#) is always non-negative. Finally, a  $\mu$ -subnull set is always in  $\mathcal{A}$ .  $\circledast$

There are some useful terminologies we'll use later relating to  $\mu$ -null.

**Definition 1.2.6** (Almost everywhere). Given a [measure space](#)  $(X, \mathcal{A}, \mu)$ , a statement  $P(x)$ ,  $x \in X$  holds  $\mu$ -almost everywhere (a.e.) if the set  $\{x \in X : P(x) \text{ does not hold}\}$  is  $\mu$ -null.

It's always pleasurable working with finite rather than infinite, hence we give the following notion.

**Definition.** Let  $(X, \mathcal{A}, \mu)$  be a [measure space](#).

**Definition 1.2.7** (Finite measure).  $\mu$  is a finite measure if  $\mu(X) < \infty$ .

**Definition 1.2.8** ( $\sigma$ -finite measure).  $\mu$  is a  $\sigma$ -finite measure if  $X = \bigcup_{n=1}^{\infty} X_n$ ,  $X_n \in \mathcal{A}$ ,  $\mu(X_n) < \infty$ .

**Exercise.** Every [measure space](#) can be completed. Namely, we can always find a bigger  $\sigma$ -algebra to complete the space.

## 1.3 Outer Measures

As we said before, we're now going to construct a [measure](#). And a modern way to do this is to start with something called [outer measure](#).

**Definition 1.3.1** (Outer measure). An outer measure on  $X$  is a function

$$\mu^* : \mathcal{P}(X) \rightarrow [0, \infty]$$

such that

- (null empty set)  $\mu^*(\emptyset) = 0$ .
- (monotonicity)  $\mu^*(A) \leq \mu^*(B)$  if  $A \subset B$ .
- (countable subadditivity)  $\mu^*(\bigcup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \mu^*(A_i)$  for every  $A_i \subset X$ .

**Example.** For  $A \subset \mathbb{R}$ ,

$$\mu^*(A) = \inf \left\{ \sum_{i=1}^{\infty} (b_i - a_i) : \bigcup_{i=1}^{\infty} (a_i, b_i) \supset A \right\}$$

is an **outer measure**.

**Proof.** This follows directly from the **Proposition 1.3.1** we're going to show.  $\circledast$

**Remark.** We see that an **outer measure** need not be a **measure**.

**Proposition 1.3.1.** Let  $\mathcal{E} \subset \mathcal{P}(X)$  such that  $\emptyset, X \in \mathcal{E}$ . Let  $\rho: \mathcal{E} \rightarrow [0, \infty]$  such that  $\rho(\emptyset) = 0$ . Then

$$\mu^*(A) := \inf \left\{ \sum_{i=1}^{\infty} \rho(E_i) : \forall_{i \in \mathbb{N}} E_i \in \mathcal{E}, \bigcup_{i=1}^{\infty} E_i \supset A \right\}$$

is an **outer measure** on  $X$ .

To prove **Proposition 1.3.1**, recall the **Tonelli's theorem for series** first.

**Theorem 1.3.1** (Tonelli's theorem for series). If for all  $i, j \in \mathbb{N}$ ,  $a_{ij} \in [0, \infty]$ , then

$$\sum_{(i,j) \in \mathbb{N}^2} a_{ij} = \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij} = \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} a_{ij}.$$

**Proof.** Read [Tao13, Theorem 0.0.2].  $\blacksquare$

## Lecture 4: Carathéodory Extension Theorem

**As previously seen.** Last time we skip the proof of **Proposition 1.3.1**, which is a quite important theorem for building a **measure**.

12 Jan. 11:00

To see this, we note that from **Proposition 1.3.1**, given a positive function  $\rho$  defined on a subset of the power set of  $X$  with  $\rho(\emptyset) = 0$ , we can induce an **outer measure** from  $\rho$ .

**Note.** Later, we can further induce a natural **measure** from the induced **outer measure**.

We now prove **Proposition 1.3.1**.

**Proof of Proposition 1.3.1.** We need to prove the following.

**Claim.**  $\mu^*$  is well-defined, i.e., inf is taken over a non-empty set.

**Proof.** This is trivial since  $X \in \mathcal{E}$  and  $X \supset A$  for any  $A \in \mathcal{E}$ .  $\circledast$

**Claim.** **Null empty set** holds, i.e.,  $\mu^*(\emptyset) = 0$ .

**Proof.** Since  $\emptyset \in \mathcal{E}$  and

$$\mu^*(\emptyset) = \inf \left\{ \sum_{i=1}^{\infty} \rho(E_i) : \forall_{i \in \mathbb{N}} E_i \in \mathcal{E}, \bigcup_{i=1}^{\infty} E_i \supset \emptyset \right\} = 0$$

since  $\rho(\emptyset) = 0$  for all  $i$  and further, by squeeze theorem,  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \rho(\emptyset) = 0$ .  $\circledast$

**Claim.** **Monotonicity** holds, i.e.,  $A \subset B \Rightarrow \mu^*(A) \leq \mu^*(B)$ .

**Proof.** Suppose for a contradiction that  $A \subset B$  and  $\mu^*(A) > \mu^*(B)$ , then by definition of  $\mu^*$ ,

$$\begin{aligned}\mu^*(A) &= \inf \left\{ \sum_{i=1}^{\infty} \rho(E_i) : \forall_{i \in \mathbb{N}} E_i \in \mathcal{E}, \bigcup_{i=1}^{\infty} E_i \supset A \right\} \\ &> \inf \left\{ \sum_{i=1}^{\infty} \rho(E_i) : \forall_{i \in \mathbb{N}} E_i \in \mathcal{E}, \bigcup_{i=1}^{\infty} E_i \supset B \right\} = \mu^*(B).\end{aligned}$$

Now, let  $B =: (B \setminus A) \cup A$ , then we have

$$\begin{aligned}\mu^*(A) &= \inf \left\{ \sum_{i=1}^{\infty} \rho(E_i) : \forall_{i \in \mathbb{N}} E_i \in \mathcal{E}, \bigcup_{i=1}^{\infty} E_i \supset A \right\} \\ &> \inf \left\{ \sum_{i=1}^{\infty} \rho(E_i) : \forall_{i \in \mathbb{N}} E_i \in \mathcal{E}, \bigcup_{i=1}^{\infty} E_i \supset (B \setminus A) \cup A \right\} = \mu^*(B).\end{aligned}$$

Now, since  $B \setminus A \supseteq \emptyset$ , then this inequality can't hold, hence a contradiction.  $\otimes$

**Claim.** Countable subadditivity holds, i.e.,  $\mu^*(\bigcup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \mu^*(A_i)$  for every  $A_i \subset X$ .

**Proof.** Let  $A_1, A_2, \dots \in X$ . If one of  $\mu^*(A_n) = \infty$ , then result holds. So we may assume  $\mu^*(A_n) < \infty$  for all  $n \in \mathbb{N}$ . Now, fix any  $\epsilon > 0$ , we will show that

$$\mu^*\left(\bigcup_{n=1}^{\infty} A_n\right) \leq \sum_{n=1}^{\infty} \mu^*(A_n) + \epsilon.$$

For each  $n \in \mathbb{N}$ ,  $\exists E_{n,1}, E_{n,2}, \dots \in \mathcal{E}$  such that  $\bigcup_{k=1}^{\infty} E_{n,k} \supset A_n$  and  $\mu^*(A_n) + \epsilon/2^n \geq \sum_{k=1}^{\infty} \rho(E_{n,k})$ .

**Remark.** This is an important trick! We often set the error term as  $\epsilon/2^n$  instead of  $\epsilon$  as in above to accommodate the summation over a countable set.

Then we see that

$$\bigcup_{k=1}^{\infty} A_n \subset \bigcup_{n=1}^{\infty} \bigcup_{k=1}^{\infty} E_{k,n} = \bigcup_{(n,k) \in \mathbb{N}^2} E_{k,n},$$

which implies

$$\mu^*\left(\bigcup_{n=1}^{\infty} A_n\right) \leq \sum_{(n,k) \in \mathbb{N}^2} \rho(E_{k,n}) \stackrel{!}{=} \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \rho(E_{k,n}) \leq \sum_{n=1}^{\infty} \left( \mu^*(A_n) + \frac{\epsilon}{2^n} \right)$$

from the inequality just derived. Now, since the last term is just

$$\sum_{n=1}^{\infty} \left( \mu^*(A_n) + \frac{\epsilon}{2^n} \right) = \sum_{n=1}^{\infty} \mu^*(A_n) + \epsilon,$$

hence we finally have

$$\mu^*\left(\bigcup_{n=1}^{\infty} A_n\right) \leq \sum_{n=1}^{\infty} \mu^*(A_n) + \epsilon$$

for arbitrarily small fixed  $\epsilon > 0$ , hence the subadditivity is proved.  $\otimes$

**Definition 1.3.2** (Carathéodory measurable). Let  $\mu^*$  be an **outer measure** on  $X$ . We say  $A \subset X$  is *Carathéodory measurable with respect to  $\mu^*$*  if

$$\forall E \subset X, \mu^*(E) = \mu^*(E \cap A) + \mu^*(E \setminus A).$$

**Note.** We sometimes write *C-measurable* instead of **Carathéodory measurable** for convenience.

**Lemma 1.3.1.** Let  $\mu^*$  be an **outer measure** on  $X$ . Suppose  $B_1, \dots, B_N$  are disjoint **C-measurable** sets. Then, for all  $E \subset X$ ,

$$\mu^* \left( E \cap \left( \bigcup_{i=1}^N B_i \right) \right) = \sum_{i=1}^N \mu^*(E \cap B_i).$$

**Proof.** Since we have

$$\begin{aligned} \mu^* \left( E \cap \left( \bigcup_{i=1}^N B_i \right) \right) &= \mu^*(E' \cap B_1) + \mu^*(E' \setminus B_1) \\ &= \mu^* \left( E \cap \left( \bigcup_{i=1}^N B_i \cap B_1 \right) \right) + \mu^* \left( E \cap \left( \bigcup_{i=1}^N B_i \right) \cap B_1^c \right) \\ &= \mu^*(E \cap B_1) + \mu^* \left( E \cap \left( \bigcup_{i=2}^N B_i \right) \right) \end{aligned}$$

where the equality comes from the fact that  $B_1$  is **C-measurable** and disjoint from  $B_i, i \neq 1$ . Then, we simply iterate this argument and have the result. Note that in the first inequality, we define  $E' := E \cap \left( \bigcup_{i=1}^N B_i \right)$  for the simplicity of notation. ■

**Remark.** This implies that if we restrict an **outer measure** on a **C-measurable** set, then it becomes finite additive.

**Theorem 1.3.2** (Carathéodory extension theorem). Let  $\mu^*$  be an **outer measure** on  $X$ . Let  $\mathcal{A}$  be the collection of **C-measurable** sets (with respect to  $\mu^*$ ). Then the following holds.

- (a)  $\mathcal{A}$  is a  **$\sigma$ -algebra** on  $X$ .
- (b)  $\mu = \mu^*|_{\mathcal{A}}$  is a **measure** on  $(X, \mathcal{A})$ .
- (c)  $(X, \mathcal{A}, \mu)$  is a **complete measure space**.

**Proof.** We divide the proof in several steps.

- (a) We show  $\mathcal{A}$  is a  **$\sigma$ -algebra** by verifying the following.

- We first show  $\emptyset \in \mathcal{A}$ .

**Claim.**  $\emptyset \in \mathcal{A}$ .

**Proof.** To show this, we simply check that  $\emptyset$  is **C-measurable**. We see that

$$\forall_{E \subset X} \mu^*(E) = \mu^*(E \cap \emptyset) + \mu^*(E \setminus \emptyset) = \mu^*(E),$$

which just shows  $\emptyset \in \mathcal{A}$ . ⊗

- Then we show  $\mathcal{A}$  is closed under complements.

**Claim.**  $\mathcal{A}$  closed under complements.

**Proof.** This is equivalent to say that if  $A$  is **C-measurable**, so is  $A^c$ . We see that if  $A$  is **C-measurable**, then for every  $E \subset X$ ,

$$\mu^*(E) = \mu^*(E \cap A) + \mu^*(E \setminus A).$$

Observing that  $E \cap A = E \setminus A^c$  and  $E \setminus A = E \cap A^c$ , hence

$$\mu^*(E) = \mu^*(E \setminus A^c) + \mu^*(E \cap A^c).$$

We immediately see that above implies  $A^c \in \mathcal{A}$ . ⊗

- We now show  $\mathcal{A}$  is closed under countable unions.

**Note.** To show  $\mathcal{A}$  closed under countable unions, we show that  $\mathcal{A}$  is closed under:

$$\text{finite unions} \xRightarrow{\text{then}} \text{countable } \underline{\text{disjoint}} \text{ unions} \xRightarrow{\text{then}} \text{countable unions}.$$


Hence, we first show  $\mathcal{A}$  is closed under finite unions.

**Claim.**  $A, B \in \mathcal{A} \Rightarrow A \cup B \in \mathcal{A}$ .

**Proof.** Fix  $E \subset X$  arbitrary. We need to show that

$$\mu^*(E) = \mu^*(E \cap (A \cup B)) + \mu^*(E \setminus (A \cup B)),$$

i.e.,  $\mu^*(1 \cup 2 \cup 3 \cup 4) = \mu^*(1 \cup 2 \cup 3) + \mu^*(4)$  given  $A, B \in \mathcal{A}$  and the following figure.



- Since  $A$  is **C-measurable**,
  - \*  $\mu^*(1 \cup 2 \cup 3 \cup 4) = \mu^*(1 \cup 2) + \mu^*(3 \cup 4)$
  - \*  $\mu^*(1 \cup 2 \cup 3) = \mu^*(1 \cup 2) + \mu^*(3)$
- Since  $B$  is **C-measurable**,
  - \*  $\mu^*(3 \cup 4) = \mu^*(3) + \mu^*(4)$

Hence, we have

$$\begin{aligned} \mu^*(1 \cup 2 \cup 3 \cup 4) &= \mu^*(1 \cup 2) + \mu^*(3 \cup 4) \\ &= \mu^*(1 \cup 2) + \mu^*(3) + \mu^*(4) = \mu^*(1 \cup 2 \cup 3) + \mu^*(4). \end{aligned}$$

⊗

We now show  $\mathcal{A}$  is closed under countable disjoint unions.

**Claim.**  $\mathcal{A}$  is closed under countable disjoint unions.



**Proof.** Let  $A_1, A_2, \dots \in \mathcal{A}$  and disjoint. Fix  $E \subset X$  arbitrary. Since  $\mu^*$  is countably subadditive,

$$\mu^*(E) \leq \mu^*\left(E \cap \bigcup_{i=1}^{\infty} A_i\right) + \mu^*\left(E \setminus \bigcup_{i=1}^{\infty} A_i\right),$$

hence we only need to show another way around. Fix  $N \in \mathbb{N}$ , we have  $\bigcup_{n=1}^N A_n \in \mathcal{A}$  since  $N$  is finite, and

$$\begin{aligned} \mu^*(E) &= \mu^*\left(E \cap \left(\bigcup_{n=1}^N A_n\right)\right) + \mu^*\left(E \setminus \left(\bigcup_{n=1}^N A_n\right)\right) \\ &\geq \underbrace{\sum_{n=1}^N \mu^*(E \cap A_n)}_{\stackrel{!}{=} \mu^*(E \cap (\bigcup_{n=1}^N A_n))} + \underbrace{\mu^*\left(E \setminus \bigcup_{n=1}^{\infty} A_n\right)}_{\leq \mu^*(E \setminus (\bigcup_{n=1}^N A_n))}. \end{aligned}$$

Now, take  $N \rightarrow \infty$  then we are done.  $\otimes$

We can then extend this to the case of countable unions.

**Exercise.** Show  $\mathcal{A}$  is closed under countable unions.

Above shows that  $\mathcal{A}$  is a  $\sigma$ -algebra.

The proof will be continued...

## Lecture 5: Hahn-Kolmogorov Theorem

Before proceeding the proof, we first see a stronger version of [Lemma 1.3.1](#) we have seen before.

14 Jan. 11:00

**Lemma 1.3.2.** Let  $\mu^*$  be an outer measure on  $X$ . Suppose  $B_1, B_2, \dots$  are disjoint  $\mathcal{C}$ -measurable sets. Then, for all  $E \subset X$ ,

$$\mu^*\left(E \cap \left(\bigcup_{i=1}^{\infty} B_i\right)\right) = \sum_{i=1}^{\infty} \mu^*(E \cap B_i).$$

**Proof.** Since

$$\sum_{n=1}^{\infty} \mu^*(E \cap B_i) \geq \mu^*\left(E \cap \bigcup_{n=1}^{\infty} B_n\right) \geq \mu^*\left(E \cap \left(\bigcup_{n=1}^N B_n\right)\right) \stackrel{!}{=} \sum_{n=1}^N \mu^*(E \cap B_n),$$

we can just take  $N \rightarrow \infty$  and since  $N \in \mathbb{N}$  is arbitrary, the result follows from squeeze theorem.  $\blacksquare$

Let's continue the proof of [Carathéodory extension theorem](#).

**Proof of Theorem 1.3.2 (cont.)** The (a) is proved, now we prove (b) and (c).

(b) Since from [Definition 1.2.1](#), to show  $\mu$  is a measure, we need to show the following.

- Null empty set property.

**Claim.**  $\mu(\emptyset) = 0$ .

**Proof.** This means that we need to show  $\mu^*|_{\mathcal{A}}(\emptyset) = 0$ . Since  $\emptyset \in \mathcal{A}$  and  $\mu^*$  is an outer measure, hence from the property of outer measure, it clearly holds.  $\otimes$

- Countable additivity property.

**Claim.**  $\mu^*$  on  $\mathcal{A}$  has Countable additivity property.

**Proof.** It follows from [Lemma 1.3.2](#) with  $E = X$  ⊗

(c) The proof is given in [Theorem A.1.1](#). ■

## 1.4 Hahn-Kolmogorov Theorem

We see that we can start with any collection of open sets  $\mathcal{E}$  and any  $\rho$  such that it assigns [measure](#) on  $\mathcal{E}$ , then it induces an [outer measure](#) by [Proposition 1.3.1](#). Then, one can [complete](#) the [outer measure](#) by [Theorem 1.3.2](#). Specifically, we have

$$(\mathcal{E}, \rho) \xrightarrow{\text{Proposition 1.3.1}} (\mathcal{P}(X), \mu^*) \xrightarrow{\text{Theorem 1.3.2}} (\mathcal{A}, \mu)$$

To introduce this concept, we see that we can start with a more general definition compared to [σ-algebra](#) we are working on till now.

**Definition 1.4.1 (Algebra).** Let  $X$  be a set. A collection  $\mathcal{A}$  of subsets of  $X$ , i.e.,  $\mathcal{A} \subset \mathcal{P}(X)$  is called an *algebra on  $X$*  if

- $\emptyset \in \mathcal{A}$ .
- $\mathcal{A}$  is closed under complements. i.e., if  $A \in \mathcal{A}$ ,  $A^c = X \setminus A \in \mathcal{A}$ .
- $\mathcal{A}$  is closed under **finite** unions. i.e., if  $A_i \in \mathcal{A}$ , then  $\bigcup_{i=1}^n A_i \in \mathcal{A}$  for  $n < \infty$ .

**Remark.** The only difference between an [algebra](#) and a [σ-algebra](#) is whether they closed under **countable** unions in the definition.

Now, we can look at a more general setup compared to an [outer measure](#).

**Definition 1.4.2 (Pre-measure).** Let  $\mathcal{A}_0$  be an [algebra](#) on  $X$ . A *pre-measure* on  $X$  with respect to  $\mathcal{A}_0$  is a function

$$\mu_0: \mathcal{A}_0 \rightarrow [0, \infty]$$

such that

- (null empty set)  $\mu_0(\emptyset) = 0$ .
- (finite additivity)  $\mu_0(\bigcup_{i=1}^n A_i) = \sum_{i=1}^n \mu_0(A_i)$  if  $A_1, \dots, A_n \in \mathcal{A}_0$  are disjoint.
- (countable additivity within the [algebra](#)) If  $A \in \mathcal{A}_0$  and  $A = \bigcup_{n=1}^{\infty} A_n$ ,  $A_n \in \mathcal{A}_0$ , disjoint, then

$$\mu_0(A) = \sum_{n=1}^{\infty} \mu_0(A_n).$$

**Lemma 1.4.1.** The [null empty set](#) property and [countable additivity within the algebra](#) implies [finite additivity](#) in [Definition 1.4.2](#).

**Proof.** It's easy to see that since  $\mu_0$  is monotone. ■

**Theorem 1.4.1 (Hahn-Kolmogorov theorem).** Let  $\mu_0$  be a [pre-measure](#) on [algebra](#)  $\mathcal{A}_0$  on  $X$ . Let  $\mu^*$  be the [outer measure](#) induced by  $(\mathcal{A}_0, \mu_0)$  in [Proposition 1.3.1](#). Finally, let  $\mathcal{A}$  and  $\mu$  be the [Carathéodory σ-algebra](#) and [measure](#) for  $\mu^*$ , then  $(\mathcal{A}, \mu)$  extends  $(\mathcal{A}_0, \mu_0)$ , i.e.,

$$\mathcal{A} \supset \mathcal{A}_0, \quad \mu|_{\mathcal{A}_0} = \mu_0.$$

**Proof.** We prove this theorem in two parts. We first show that  $\mathcal{A} \supset \mathcal{A}_0$ .

**Claim.**  $\mathcal{A} \supset \mathcal{A}_0$ .

**Proof.** Let  $A \in \mathcal{A}_0$ , we want to show  $A \in \mathcal{A}$ , i.e.,  $A$  is **C-measurable**, i.e.,

$$\forall E \subset X \quad \mu^*(E) = \mu^*(E \cap A) + \mu^*(E \cap A^c).$$

We first fix an  $E \subset X$ . From **countable subadditivity** of  $\mu^*$ , we have

$$\mu^*(E) \leq \mu^*(E \cap A) + \mu^*(E \cap A^c).$$

Hence, we only need to show another direction. If  $\mu^*(E) = \infty$ , then  $\mu^*(E) = \infty \geq \mu^*(E \cap A) + \mu^*(E \cap A^c)$  clearly. So, assume  $\mu^*(E) < \infty$ .

Fix  $\epsilon > 0$ . By the **Proposition 1.3.1** of  $\mu^*$ ,  $\exists B_1, B_2, \dots \in \mathcal{A}_0$ ,  $\bigcup_{n=1}^{\infty} B_n \supset E$  such that

$$\mu^*(E) + \epsilon \geq \sum_{n=1}^{\infty} \mu_0(B_n) = \sum_{n=1}^{\infty} \left( \underbrace{\mu_0(B_n \cap A)}_{\in \mathcal{A}_0} + \underbrace{\mu_0(B_n \cap A^c)}_{\in \mathcal{A}_0} \right)$$

by the **finite additivity** of  $\mu_0$ . Note that

$$\left\{ \begin{array}{l} \bigcup_{n=1}^{\infty} (B_n \cap A) \supset E \cap A \\ \bigcup_{n=1}^{\infty} (B_n \cap A^c) \subset E \cap A^c \end{array} \right. \Rightarrow \mu^*(E) + \epsilon \geq \mu^*(E \cap A) + \mu^*(E \cap A^c)$$

since

$$\mu^*(E \cap A) \leq \mu^* \left( \bigcup_{n=1}^{\infty} (B_n \cap A) \right) \leq \sum_{n=1}^{\infty} \mu^*(B_n \cap A)$$

and

$$\mu^*(E \cap A^c) \leq \mu^* \left( \bigcup_{n=1}^{\infty} (B_n \cap A^c) \right) \leq \sum_{n=1}^{\infty} \mu^*(B_n \cap A^c).$$

We then see that for any  $\epsilon > 0$ , the inequality

$$\mu^*(E) + \epsilon \geq \mu^*(E \cap A) + \mu^*(E \cap A^c)$$

holds, hence so does

$$\mu^*(E) \geq \mu^*(E \cap A) + \mu^*(E \cap A^c),$$

which implies  $\mathcal{A} \supset \mathcal{A}_0$ . ⊛

*The proof will be continued...*

## Lecture 6: Hahn-Kolmogorov Theorem and Extension

Let's continue the proof of **Hahn-Kolmogorov theorem**.

18 Jan. 11:00

**Proof of Theorem 1.4.1 (cont.)** We proved the first part already, now we prove the part left.

**Claim.**  $\mu|_{\mathcal{A}_0} = \mu_0$ .

**Proof.** Let  $A \in \mathcal{A}_0$ , we want to show that  $\mu(A) = \mu_0(A)$ .

- Firstly, let

$$B_i = \begin{cases} A, & \text{if } i = 1 \\ \emptyset, & \text{if } i \geq 2 \end{cases} \in \mathcal{A}_0,$$

hence  $\bigcup_{i=1}^{\infty} B_i = A$ , then we see that

$$\mu^*(A) \leq \sum_{i=1}^{\infty} \mu_0(B_i) = \mu_0(A)$$

from the [definition](#) of  $\mu^*$  and [countable additivity within the algebra](#) of  $\mu_0$ .

- Secondly, let  $B_i \in \mathcal{A}_0$ ,  $\bigcup_{i=1}^{\infty} B_i \supset A$  be arbitrary. Let  $C_1 = A \cap B_1 \in \mathcal{A}_0$ ,  $C_i = A \cap B_i \setminus \left(\bigcup_{j=1}^{i-1} B_j\right) \in \mathcal{A}_0$  for  $i \geq 2$  since the operations are finite. Then we see

$$A = \bigcup_{i=1}^{\infty} C_i \in \mathcal{A}_0$$

are disjoint countable unions, by [countable additivity within the algebra](#), therefore

$$\mu_0(A) = \sum_{i=1}^{\infty} \mu_0(C_i) \leq \sum_{i=1}^{\infty} \mu_0(B_i) \Rightarrow \mu_0(A) \leq \mu^*(A)$$

by taking the infimum from the [definition](#) of  $\mu^*$ .

Combine these two inequality, we see that

$$\mu^*(A) = \mu_0(A),$$

for every  $A \in \mathcal{A}_0$ , which implies  $\mu(A) = \mu_0(A)$  for every  $A \in \mathcal{A}_0$  from [Theorem 1.3.2](#), where we extend  $\mu^*$  to  $\mu$  respect to  $\mathcal{A}_0$ . ⊗

■

**Definition 1.4.3** (Hahn-Kolmogorov extension).  $(\mathcal{A}, \mu)$  obtained from [Hahn-Kolmogorov theorem](#) is called the *Hahn-Kolmogorov extension* of  $(\mathcal{A}_0, \mu_0)$ .

**Note.** We sometimes say *HK extension* instead of [Hahn-Kolmogorov extensions](#) for simplicity.

We can show the uniqueness of [HK extension](#).

**Theorem 1.4.2** (Uniqueness of HK extension). Let  $\mathcal{A}_0$  be an [algebra](#) on  $X$ ,  $\mu_0$  be a [pre-measure](#) on  $\mathcal{A}_0$ . Let  $(\mathcal{A}, \mu)$  be the [HK extension](#) of  $(\mathcal{A}_0, \mu_0)$ . Finally, let  $(\mathcal{A}', \mu')$  be another extension of  $(\mathcal{A}_0, \mu_0)$ . Then if  $\mu_0$  is [σ-finite](#),  $\mu = \mu'$  on  $\mathcal{A} \cap \mathcal{A}'$ .

**Proof.** First, we note the following.

**Note.** Notice that  $\mathcal{A}_0 \subset \mathcal{A}, \mathcal{A}'$  since they both extend  $\mathcal{A}_0$ .

Let  $A \in \mathcal{A} \cap \mathcal{A}'$ , we need to show

$$\underbrace{\mu(A)}_{\mu^*(A)} = \mu'(A).$$

We'll show this by showing two inequalities. We first show that  $\mu^*(A) \geq \mu'(A)$ .

**Claim.**  $\mu^*(A) \geq \mu'(A)$ .

**Proof.** It's easy to show that  $\mu^*(A) \geq \mu'(A)$  by choosing the arbitrary cover of  $A$  and using the definition of  $\mu^*$ .  $\otimes$

Secondly, we will show that  $\mu(A) \leq \mu'(A)$ .

**Claim.**  $\mu(A) \leq \mu'(A)$ .

**Proof.** We split this into two cases.

- Assume  $\mu(A) < \infty$ , and fix  $\epsilon > 0$ . Then there exists  $B_i \in \mathcal{A}_0$  with  $B := \bigcup_{i=1}^{\infty} B_i \supset A$  such that

$$\mu(A) + \epsilon = \mu^*(A) + \epsilon \geq \sum_{i=1}^{\infty} \mu_0(B_i) \stackrel{!}{=} \sum_{i=1}^{\infty} \mu(B_i) \geq \mu\left(\bigcup_{i=1}^{\infty} B_i\right) = \mu(B).$$

This implies that

$$\mu(B \setminus A) = \mu(B) - \mu(A) \leq \epsilon,$$

where the equality comes from  $A \subset B$  and  $\mu(A) < \infty$ . On the other hand,

$$\mu(B) = \lim_{N \rightarrow \infty} \mu\left(\bigcup_{i=1}^N B_i\right) = \lim_{N \rightarrow \infty} \mu'\left(\bigcup_{i=1}^N B_i\right) = \mu'(B)$$

where the middle equality follows from  $\mu = \mu'$  on  $\mathcal{A}_0$ , hence,

$$\mu(A) \leq \mu(B) = \mu'(B) = \mu'(A) + \mu'(B \setminus A) \leq \mu'(A) + \mu(B \setminus A) \leq \mu'(A) + \epsilon$$

for arbitrary  $\epsilon$ , so we conclude  $\mu(A) \leq \mu'(A)$ .

- Assume  $\mu(A) = \infty$ . Since  $\mu_0$  is  $\sigma$ -finite, so we know  $X = \bigcup_{n=1}^{\infty} X_n$  for some  $X_n \in \mathcal{A}_0$  such that  $\mu_0(X_n) < \infty$ . Replacing  $X_n$  by  $X_1 \cup \dots \cup X_n \in \mathcal{A}_0$ , we may assume that  $X_1 \subset X_2 \subset \dots$ . Then,

$$\forall_{n \in \mathbb{N}} \mu(A \cap X_n) < \infty \stackrel{!}{\Rightarrow} \mu(A \cap X_n) \leq \mu'(A \cap X_n).$$

By continuity from above, we have

$$\mu(A) = \lim_{n \rightarrow \infty} \mu(A \cap X_n) \leq \lim_{n \rightarrow \infty} \mu'(A \cap X_n) = \mu'(A).$$

Combine above two inequalities, the result follows.  $\otimes$

From the uniqueness theorem, we have the following. ■

**Corollary 1.4.1.** Let  $\mu_0$  be a pre-measure on algebra  $\mathcal{A}_0$  on  $X$ . Suppose  $\mu_0$  is  $\sigma$ -finite, then there exists a unique measure  $\mu$  on  $\langle \mathcal{A}_0 \rangle$  that extends  $\mathcal{A}_0$ . Furthermore, we have the following.

(a) The completion of  $(X, \langle \mathcal{A}_0 \rangle, \mu)$  is the HK extension of  $(\mathcal{A}_0, \mu_0)$ .<sup>a</sup>

(b)  $\mu(A) = \inf \left\{ \sum_{i=1}^{\infty} \mu_0(B_i) \mid \forall_{i \in \mathbb{N}} B_i \in \mathcal{A}_0, \bigcup_{i=1}^{\infty} B_i \supset A \right\}$  for all  $A \in \overline{\langle \mathcal{A}_0 \rangle}$ .

<sup>a</sup>This really means the pair of the  $\sigma$ -algebra and the measure in the completion of  $(X, \langle \mathcal{A}_0 \rangle, \mu)$  is the HK extension of  $(\mathcal{A}_0, \mu_0)$ , not the whole tuple of the measurable-space.

## Lecture 7: Borel Measures

### 1.5 Borel Measures on $\mathbb{R}$

21 Jan. 11:00

**As previously seen.** Recall that when we say something is **Borel**, we assume some sort of topological structure on the underlying space implicitly. In our context, we're considering the usual topology on  $\mathbb{R}^n$  specifically. We'll focus on one dimensional case for now.

**Definition 1.5.1 (Distribution function).** If  $F: \mathbb{R} \rightarrow \mathbb{R}$  is increasing<sup>a</sup> and right-continuous, then it's a *distribution function*.

<sup>a</sup>I.e.,  $F(x) \leq F(y)$  for  $x < y$ .

Here are some examples of right-continuous functions.

**Example.**  $F(x) = x$ .

**Example.**  $F(x) = e^x$ .

**Example.**  $F(x) = \begin{cases} 1, & \text{if } x \geq 0; \\ 0, & \text{if } x < 0. \end{cases}$

**Example.** Let  $\mathbb{Q} := \{r_1, r_2, \dots\}$ . Define

$$F_n(x) = \begin{cases} 1, & \text{if } x \geq r_n; \\ 0, & \text{if } x < r_n; \end{cases}, \quad F(x) := \sum_{n=1}^{\infty} \frac{F_n(x)}{2^n}.$$

Then  $F$  is a *distribution function* (hence right-continuous). This is shown in [Lemma A.1.1](#).

**Note.** If  $F$  is increasing, then

$$F(\infty) := \lim_{x \nearrow \infty} F(x), \quad F(-\infty) := \lim_{x \searrow -\infty} F(x)$$

exist in  $[-\infty, \infty]$ . In probability theory, cumulative distribution function (CDF) is a *distribution function* with  $F(\infty) = 1$ ,  $F(-\infty) = 0$ .<sup>a</sup>

<sup>a</sup>There are *distributions* [FF99, §9], but these are different from *distribution functions*.

Now, we see a new definition which is essential to our discussion.

**Definition 1.5.2 (Borel measure).** A *Borel measure* is any *measure*  $\mu$  defined on the  $\sigma$ -algebra of *Borel sets*.

Since we're now considering a topological space, hence it's reasonable to define the following because we have the concept of compact set now.

**Definition 1.5.3 (Locally finite).** Let  $X$  be a Hausdorff topological space,  $\mu$  on  $(X, \mathcal{B}(X))$  is called *locally finite* if  $\mu(K) < \infty$  for every compact set  $K \subset X$ .

**Note.** Some authors will require a *Borel measure* equipped with the *locally finite* property. But formally, this is not so common.

**Lemma 1.5.1.** Let  $\mu$  be a **locally finite Borel measure** on  $\mathbb{R}$ , then

$$F_\mu(x) = \begin{cases} \mu((0, x]), & \text{if } x > 0 \\ 0, & \text{if } x = 0 \\ -\mu((x, 0]), & \text{if } x < 0 \end{cases}$$

is a **distribution function**.

**Proof.** We need to show two things.

**Claim.**  $F_\mu$  is increasing.

**Proof.** To show  $F_\mu$  is increasing, consider  $x < y$  such that

$$F_\mu(x) \leq F_\mu(y)$$

by considering

- $x > 0$ : Then  $F_\mu(x) = \mu((0, x])$  and

$$F_\mu(y) = \mu((0, y]) = \mu((0, x] \cup (x, y]) \geq \mu((0, x]) = F_\mu(x).$$

- $x = 0$ : Then  $F_\mu(x) = 0$  and

$$F_\mu(y) = \mu((0, y]) \geq 0 = F_\mu(0)$$

since  $y > 0$ .

- $x < 0$ : Follows the same argument with  $x > 0$ .

⊗

We now show  $F_\mu$  is right-continuous.

**Claim.**  $F_\mu$  is right-continuous.

**Proof.** Firstly, assume that  $x \geq 0$ , then we see that

$$F_\mu(x) = \mu((0, x]) = \mu((0, x^+])$$

from the fact that a measure is right-continuous.<sup>a</sup> Now, if  $x \leq 0$ , the same argument follows since multiplying  $-1$  will not change the fact that a **measure** is continuous. ⊗

<sup>a</sup>Actually, a measure is always continuous.

■

**Definition 1.5.4 (Half intervals).** We call  $\emptyset, (a, b], (a, \infty), (-\infty, b]$ , and  $(-\infty, \infty)$  *half-intervals*.

**Lemma 1.5.2.** Let  $\mathcal{H}$  be the collection of finite disjoint unions of **half-intervals**. Then,  $\mathcal{H}$  is an **algebra** on  $\mathbb{R}$ .

**Proof.** We observe that  $\emptyset \in \mathcal{H}$  and  $\mathcal{H}$  is closed under finite unions is obvious, hence we only need to show that  $\mathcal{H}$  is closed under complements.

**Claim.**  $\mathcal{H}$  is closed under complements.

**Proof.** We have

- $\emptyset^c = \mathbb{R} = (-\infty, \infty) \in \mathcal{H}$ .
- $(a, b]^c = (-\infty, a] \cup (a, \infty) \in \mathcal{H}$  since it's a two disjoint union of half intervals.
- $(a, \infty)^c = (-\infty, a] \in \mathcal{H}$ .
- $(-\infty, b]^c = (b, \infty) \in \mathcal{H}$ .
- $(-\infty, \infty)^c = \emptyset \in \mathcal{H}$ .

⊛

■

**Proposition 1.5.1** (Distribution function defines a pre-measure). Let  $F: \mathbb{R} \rightarrow \mathbb{R}$  be a **distribution function**. For a **half interval**  $I$ , define

$$\ell(I) := \ell_F(I) = \begin{cases} 0, & \text{if } I = \emptyset; \\ F(b) - F(a), & \text{if } I = (a, b]; \\ F(\infty) - F(a), & \text{if } I = (a, \infty]; \\ F(b) - F(-\infty), & \text{if } I = (-\infty, b]; \\ F(\infty) - F(-\infty), & \text{if } I = (-\infty, \infty). \end{cases}$$

Define  $\mu_0 := \mu_{0,F}: \mathcal{H} \rightarrow [0, \infty]$  by

$$\mu_0(A) = \sum_{k=1}^N \ell(I_k) \text{ if } A = \bigcup_{k=1}^N I_k,$$

where  $A$  is a finite **disjoint** union of **half intervals**  $I_1, \dots, I_N$ . Then,  $\mu_0$  is a **pre-measure** on  $\mathcal{H}$ .

**Proof.** Firstly, we note that  $\mu_0$  is well-defined. And also,  $\mu_0$  satisfies **null empty set** and **Finite additivity** properties clearly. The only nontrivial part needs a proof is the **Countable additivity within  $\mathcal{H}$**  properties. To show that **Countable additivity within  $\mathcal{H}$**  holds, we proceed as follows.

Suppose  $A \in \mathcal{H}$  where  $A = \bigcup_{i=1}^{\infty} A_i$  is a countable disjoint union. It is enough to consider the case that  $A = I$ ,  $A_k = I_k$  are all **half-intervals**.

**Remark.** Since  $\mathcal{H}$  is only a collection of *finite* disjoint **half intervals**, hence after considering  $A = I$ , we can apply the same argument iteratively and stop in finite steps. Formally, we can consider  $H \in \mathcal{H}$ ,  $H = \bigcup_{i=1}^{\infty} A^i$ , where  $A^i$  being a **half interval**. Then by the above argument, we have  $A^i = I^i$  and so on.

Focus on the case  $I = (a, b]$ . Let  $(a, b] = \bigcup_{n=1}^{\infty} (a_n, b_n]$ , which is a disjoint union. Then we only need to check  $F(b) - F(a) = \sum_{n=1}^{\infty} (F(b_n) - F(a_n))$ .

**Claim.**  $F(b) - F(a) \geq \sum_{n=1}^{\infty} (F(b_n) - F(a_n))$ .

**Proof.** Since  $(a, b] \supset \bigcup_{n=1}^N (a_n, b_n]$  for any fixed  $N \in \mathbb{N}$ , hence

$$\forall_{N \in \mathbb{N}} F(b) - F(a) \geq \sum_{n=1}^N (F(b_n) - F(a_n)).$$

By letting  $N \rightarrow \infty$ , we have  $F(b) - F(a) \geq \sum_{n=1}^{\infty} (F(b_n) - F(a_n))$ .

⊛



**Claim.**  $F(b) - F(a) \leq \sum_{n=1}^{\infty} (F(b_n) - F(a_n))$ .

**Proof.** Fix  $\epsilon > 0$ . Since  $F$  is right-continuous,  $\exists a' > a$  such that  $F(a') - F(a) < \epsilon$ . For each  $n \in \mathbb{N}$ ,  $\exists b'_n > b_n$  such that

$$F(b'_n) - F(b_n) < \frac{\epsilon}{2^n}.$$

**Remark.** It's again the  $\epsilon/2^n$  trick we saw before!

Then, we have  $[a', b] \subset \bigcup_{n=1}^{\infty} (a_n, b'_n)$ , hence  $\exists_{N \in \mathbb{N}} [a', b] \subset \bigcup_{n=1}^N (a_n, b'_n)$ , which is only finitely many unions now.<sup>a</sup> In this case, we have

$$F(b) - F(a') \leq \sum_{n=1}^N F(b'_n) - F(a_n).$$

Finally, we see that

$$\begin{aligned} F(b) - F(a) &\leq F(b) - F(a') + \epsilon \\ &\leq \sum_{n=1}^{\infty} (F(b'_n) - F(a_n)) + \epsilon \\ &\leq \sum_{n=1}^{\infty} \left( F(b_n) - F(a_n) + \frac{\epsilon}{2^n} \right) + \epsilon = \sum_{n=1}^{\infty} (F(b_n) - F(a_n)) + 2\epsilon \end{aligned}$$

for any fixed  $\epsilon > 0$ , hence

$$F(b) - F(a) \leq \sum_{n=1}^{\infty} (F(b_n) - F(a_n)).$$

⊗

<sup>a</sup>This essentially follows from the fact that open sets are closed under countable unions, hence the equality will not hold, even after taking the limit.

Combine these two inequalities, we have

$$F(b) - F(a) = \sum_{n=1}^{\infty} (F(b_n) - F(a_n))$$

as we desired. ■

## Lecture 8: Lebesgue-Stieltjes Measure on $\mathbb{R}$

To classify all **measures**, we now see this last theorem to complete the task.

24 Jan. 11:00

**Theorem 1.5.1** (Locally finite Borel measures on  $\mathbb{R}$ ). We have the following.

- (a) If  $F: \mathbb{R} \rightarrow \mathbb{R}$  is a **distribution function**, then there exists a **unique locally finite Borel measure**  $\mu_F$  on  $\mathbb{R}$  such that for every  $a < b$ ,

$$\mu_F((a, b]) = F(b) - F(a).$$

- (b) Suppose  $F, G: \mathbb{R} \rightarrow \mathbb{R}$  are **distribution functions**. Then,  $\mu_F = \mu_G$  on  $\mathcal{B}(\mathbb{R})$  if and only if  $F - G$  is a constant function.

**Proof.** ■

HW.

**Remark.** Theorem 1.5.1 simply states that given a [distribution function](#), if we restrict our attention on [locally finite measures](#) on  $\mathbb{R}$  following our usual convention, then it defines the [measure](#) on  $\mathcal{B}(\mathbb{R})$  uniquely up to a *constant shift*.

## 1.6 Lebesgue-Stieltjes Measure on $\mathbb{R}$

We see that

$$F \text{ distribution function} \xrightarrow{!} \mu_F \text{ on Carathéodory } \sigma\text{-algebra } \mathcal{A}_{\mu_F} \supset \mathcal{B}(\mathbb{R}).$$

Furthermore, we have

$$(\mathcal{A}_{\mu_F}, \mu_F) = \overline{(\mathcal{B}(\mathbb{R}), \mu_F)}.$$

**Definition 1.6.1** (Lebesgue-Stieltjes measure). Given a [distribution function](#)  $F$ , we say  $\mu_F$  on  $\mathcal{A}_{\mu_F}$  is called the *Lebesgue-Stieltjes measure* corresponding to  $F$ .

**Definition.** From [Definition 1.6.1](#), if  $F(x) = x$ , then the induced  $(\mathcal{A}_{\mu_F}, \mu_F)$  is denoted as  $(\mathcal{L}, m)$ .

**Definition 1.6.2** (Lebesgue measure).  $m$  is called *Lebesgue measure*.

**Definition 1.6.3** (Lebesgue  $\sigma$ -algebra).  $\mathcal{L}$  is called *Lebesgue  $\sigma$ -algebra*.

**Remark.** Recall that  $\mathcal{L}$  is induced by [Carathéodory extension theorem](#), so given  $m$ , for all  $A \subset \mathbb{R}$ ,

$$\mathcal{L} := \left\{ A \subset \mathbb{R} \mid \forall_{E \subset \mathbb{R}} m(A) = m(A \cap E) + m(A \setminus E) \right\}.$$

**Note.** We see that since  $F$  is right-continuous and increasing, hence

$$F(x^-) \leq F(x) = F(x^+).$$

Some text will use  $x-$  and  $x+$  instead of  $x^-$  and  $x^+$ , respectively.

We now see some examples.

**Example** (Discrete measure).  $\mu_F((a, b]) = F(b) - F(a)$ . Then

- $\mu_F(\{a\}) = F(a) - F(a^-)$
- $\mu_F([a, b]) = F(b) - F(a^-)$
- $\mu_F((a, b)) = F(b^-) - F(a)$

This is so-called *discrete measure*.

**Example** (Dirac measure). We define

$$F(x) = \begin{cases} 1, & \text{if } x \geq 0; \\ 0, & \text{if } x < 0. \end{cases}$$

Then

- $\mu_F(\{0\}) = 1$

- $\mu_F(\mathbb{R}) = 1$
- $\mu_F(\mathbb{R} \setminus \{0\}) = 0$ . This is easy to see since  $\mathbb{R} \setminus \{0\} = (-\infty, 0) \cup (0, \infty)$ , hence

$$\mu_F(\mathbb{R} \setminus \{0\}) = \mu_F((-\infty, 0) \cup (0, \infty)) = \underbrace{\mu_F((-\infty, 0))}_{0-0} + \underbrace{\mu_F((0, \infty))}_{1-1} = 0,$$

where  $\mu_F((-\infty, 0)) = 0$  follows from  $F(0^-) - F(-\infty) = 0 - 0 = 0$ , while  $\mu_F((0, \infty)) = 0$  follows from  $F(\infty) - F(0) = 1 - 1 = 0$ .

We call  $\mu_F$  the *Dirac measure* at 0.

**Example.** Denote  $\mathbb{Q} = \{r_1, r_2, \dots\}$ , and we define

$$F(x) = \sum_{n=1}^{\infty} \frac{F_n(x)}{2^n} \text{ where } F_n(x) = \begin{cases} 1, & \text{if } x \geq r_n; \\ 0, & \text{if } x < r_n. \end{cases}$$

Then

- $\mu_F(\{r_i\}) > 0$  for all  $r_i \in \mathbb{Q}$ .
- $\mu_F(\mathbb{R} \setminus \mathbb{Q}) = 0$ .

This is shown in [Lemma A.1.2](#).

**Example.** If  $F$  is continuous at  $a$ , then  $\mu_F(\{a\}) = 0$ .

**Example (Lebesgue measure).**  $F(x) = x$ , then recall that we denote  $\mu_F := m$ , and we have

- $m((a, b]) = m((a, b)) = m([a, b]) = b - a$ .

**Example.**  $F(x) = e^x$


- $\mu_F((a, b]) = \mu_F((a, b)) = e^b - e^a$ .

**Example (Middle thirds Cantor set).** Let  $C := \bigcap_{n=1}^{\infty} K_n$ , where we have

$$\begin{aligned} K_0 &:= [0, 1] \\ K_1 &:= K_0 \setminus \left(\frac{1}{3}, \frac{2}{3}\right) \\ K_2 &:= K_1 \setminus \left(\frac{1}{9}, \frac{2}{9}\right) \cup \left(\frac{7}{9}, \frac{8}{9}\right) \\ &\vdots \\ K_n &:= K_{n-1} \setminus \bigcup_{k=1}^{3^{n-1}-1} \left(\frac{3k+1}{3^{n+1}}, \frac{3k+2}{3^{n+1}}\right). \end{aligned}$$

We see that  $C$  is uncountable and with  $m(C) = 0$ . And observe that  $x \in C$  if and only if  $x = \sum_{n=1}^{\infty} \frac{a_n}{3^n}$  for some  $a_n \in \{0, 2\}$ . Hence, we can instead formulate  $K_n$  by

$$K_n = \bigcup_{\substack{a_i \in \{0, 2\} \\ 1 \leq i \leq n}} \left[ \sum_{i=1}^{\infty} \frac{a_i}{3^i}, \sum_{i=1}^{\infty} \frac{a_i}{3^i} + \frac{1}{3^n} \right].$$


 Figure 1.1: The top line corresponds to  $K_0$ , and then  $K_1$ , etc.

The proof of  $m(C) = 0$  is given in [Lemma A.1.3](#).

### Cantor Function

Consider  $F$  as follows. We define a function  $F$  to be 0 to the left of 0, and 1 to the right of 1. Then, define  $F$  to be  $\frac{1}{2}$  on  $(\frac{1}{3}, \frac{2}{3})$ ,  $\frac{1}{4}$  on  $(\frac{1}{9}, \frac{2}{9})$ ,  $\frac{3}{4}$  on  $(\frac{7}{9}, \frac{8}{9})$  and so on. This is so-called *Cantor Function*. We can show  $F$  is continuous and increasing, which makes  $F$  a distribution function. Also, we see that the measure this  $F$  induced is called *Cantor measure*.




Figure 1.2: Cantor Function (Devil's Staircase).

We see that  $F$  is *continuous* and increasing. Furthermore,

Cantor Measure $\mu_F$	Lebesgue Measure $m$
$\mu_F(\mathbb{R} \setminus C) = 0$	$m(\mathbb{R} \setminus C) = \infty > 0$
$\mu_F(C) = 1$	$m(C) = 0$
$\mu_F(\{a\}) = 0$	$m(\{a\}) = 0$

**Remark.**  $\mu_F$  and  $m$  are said to be **singular** to each other.

## 1.7 Regularity Properties of Lebesgue-Stieltjes Measures

We first see a lemma.

**Lemma 1.7.1.** Let  $\mu$  be [Lebesgue-Stieltjes measure](#) on  $\mathbb{R}$ . Then we have

$$\mu(A) \stackrel{!}{=} \inf \left\{ \sum_{i=1}^{\infty} \mu((a_i, b_i]) \mid \bigcup_{i=1}^{\infty} (a_i, b_i] \supset A \right\} = \inf \left\{ \sum_{i=1}^{\infty} \mu((a_i, b_i)) \mid \bigcup_{i=1}^{\infty} (a_i, b_i) \supset A \right\}$$

for every  $A \in \mathcal{A}_\mu$

**Proof.** The second equality follows from the [continuity of the measure](#). ■

**Remark.** This is similar to  $(a, b) = \bigcup_{n=1}^{\infty} (a, b - 1/n]$  and  $(a, b] = \bigcap_{n=1}^{\infty} (a, b + 1/n]$ .

## Lecture 9: Properties of Lebesgue-Stieltjes measure

26 Jan. 11:00

As previously seen. Let  $X \subset [0, \infty]$ . Recall that

- Finite supremum.

$$\alpha = \sup X < \infty \Leftrightarrow \begin{cases} \forall_{x \in X} \alpha \geq x \\ \forall_{\epsilon > 0} \exists_{x \in X} x + \epsilon \geq \alpha. \end{cases}$$

- Infinite supremum.

$$\alpha = \sup X = \infty \Leftrightarrow \forall_{L > 0} \exists_{x \in X} x \geq L.$$

This should be useful latter on.

**Theorem 1.7.1 (Regularity).** Let  $\mu$  be [Lebesgue-Stieltjes measure](#). Then, for all  $A \in \mathcal{A}_\mu$ ,

- (a) (outer regularity)  $\mu(A) = \inf\{\mu(O) \mid O \supset A, O \text{ is open}\}$
- (b) (inner regularity)  $\mu(A) = \sup\{\mu(K) \mid K \subset A, K \text{ is compact}\}$

**Proof.** We check them separately.

(a)

DIY

- (b) Let  $s := \sup\{\mu(K) \mid K \subset A, K \text{ is compact}\}$ , then by [monotonicity](#), we have  $\mu(A) \geq s$ . To show the other direction, we consider

**Claim.** [Inner regularity](#) holds if  $A$  is a bounded set.

**Proof.** Then  $\bar{A} \in \mathcal{B}(\mathbb{R}) \subset \mathcal{A}_\mu$ ,  $\bar{A}$  is also bounded  $\Rightarrow \mu(\bar{A}) < \infty$ . Fix  $\epsilon > 0$ , then by [outer regularity](#), there exists an open  $O \supset \bar{A} \setminus A$ , and  $\mu(O) - \mu(\bar{A} \setminus A) = \mu(O \setminus (\bar{A} \setminus A)) \leq \epsilon$ . Let  $K := \underbrace{A \setminus O}_{K \subset A} = \underbrace{\bar{A} \setminus O}_{\text{compact}}$ , we show that

$$\mu(K) \geq \mu(A) - \epsilon.$$

⊗

DIY

**Claim.** [Inner regularity](#) holds if  $A$  is an unbounded set with  $\mu(A) < \infty$ .

**Proof.** Let  $A = \bigcup_{n=1}^{\infty} A_n$ ,  $A_n = A \cap [-n, n]$  where  $A_1 \subset A_2 \subset \dots$ , then

$$\lim_{n \rightarrow \infty} \mu(A_n) = \mu(A) < \infty.$$

⊗

**Claim.** [Inner regularity](#) holds if  $A$  is an unbounded set with  $\mu(A) = \infty$ .

**Proof.** We can show that

$$\lim_{n \rightarrow \infty} \mu(A_n) = \mu(A) = \infty.$$

Fix  $L > 0$ , then  $\exists N$  such that  $\mu(A_N) \geq L$ .

⊗

■

**Definition.** Let  $X$  be a topological space. Then

**Definition 1.7.1** ( $G_\delta$ -set). A  $G_\delta$ -set is  $G = \bigcap_{i=1}^{\infty} O_i$ ,  $O_i$  open.

**Definition 1.7.2** ( $F_\sigma$ -set). An  $F_\sigma$ -set is  $F = \bigcup_{i=1}^{\infty} F_i$ ,  $F_i$  closed.

**Theorem 1.7.2.** Let  $\mu$  be a Lebesgue-Stieltjes measure. Then the following are equivalent:

- (a)  $A \in \mathcal{A}_\mu$ ;
- (b)  $A = G \setminus M$ ,  $G$  is a  $G_\delta$ -set,  $M$  is a  $\mu$ -null set;
- (c)  $A = F \setminus N$ ,  $F$  is an  $F_\sigma$ -set,  $N$  is a  $\mu$ -null set.

**Proof.** We see that (b)  $\Rightarrow$  (a) and (c)  $\Rightarrow$  (a) are clear.

**Claim.** (a)  $\Rightarrow$  (c).

**Proof.** We consider two cases.

- Assume  $\mu(A) < \infty$ . From the inner regularity, we have

$$\forall n \in \mathbb{N} \exists \text{compact } K_n \subset A \text{ such that } \mu(K_n) + \frac{1}{n} \geq \mu(A).$$

Let  $F = \bigcup_{n=1}^{\infty} K_n$ , then  $N = A \setminus F$  is  $\mu$ -null.

- Assume  $\mu(A) = \infty$ . Let  $A = \bigcup_{k \in \mathbb{Z}} A_k$ ,  $A_k = A \cap (k, k+1]$ . From what we have just shown above,

$$\forall k \in \mathbb{Z} \ A_k = F_k \cup N_k, \quad A = \underbrace{\left( \bigcup_k F_k \right)}_{F_\sigma\text{-set}} \cup \underbrace{\left( \bigcup_k N_k \right)}_{\mu\text{-null}}.$$

⊗

**Claim.** (a)  $\Rightarrow$  (b).

**Proof.** We see that

$$A^c = F \cup N, \quad A = F^c \cap N^c = F^c \setminus N.$$

⊗

■

**Proposition 1.7.1.** Let  $\mu$  be a Lebesgue-Stieltjes measure, and  $A \in \mathcal{A}_\mu$ ,  $\mu(A) < \infty$ . Then we have

$$\forall \epsilon > 0 \exists I = \bigcup_{i=1}^{N(\epsilon)} I_i$$

disjoint open intervals such that  $\mu(A \triangle I) \leq \epsilon$ .

**Proof.** Using outer regularity and the fact that every open set is  $\bigcup_{i=1}^{\infty} I_i$ , where  $I_i$  are disjoint open intervals. ■

We now see some properties of Lebesgue measure.

Check!

DIY

**Theorem 1.7.3.** Let  $A \in \mathcal{L}$ , then we have  $A + s \in \mathcal{L}$ ,  $rA \in \mathcal{L}$  for all  $r, s \in \mathbb{R}$ . i.e.,

$$m(A + s) = m(A), \quad m(rA) = |r| \cdot m(A).$$

**Proof.** ■

DIY

**Example.** We now see some examples.

- (a) Let  $\mathbb{Q} =: \{r_i\}_{i=1}^{\infty}$  which is dense in  $\mathbb{R}$ . Let  $\epsilon > 0$ , and

$$O = \bigcup_{i=1}^{\infty} \left( r_i - \frac{\epsilon}{2^i}, r_i + \frac{\epsilon}{2^i} \right).$$

We see that  $O$  is open and dense<sup>a</sup> in  $\mathbb{R}$ . But we see

$$m(O) \leq \sum_{i=1}^{\infty} \frac{2\epsilon}{2^i} = 2\epsilon.$$

Furthermore,  $\partial O = \overline{O} \setminus O$ ,  $m(\partial O) = \infty$

- (b) There exists uncountable set  $A$  with  $m(A) = 0$ .  
(c) There exists  $A$  with  $m(A) > 0$  but  $A$  contains no non-empty open intervals.  
(d) There exists  $A \notin \mathcal{L}$ . e.g. Vitali set.<sup>b</sup>  
(e) There exists  $A \in \mathcal{L} \setminus \mathcal{B}(\mathbb{R})$ .

<sup>a</sup>[https://en.wikipedia.org/wiki/Dense\\_set](https://en.wikipedia.org/wiki/Dense_set)

<sup>b</sup>[https://en.wikipedia.org/wiki/Vitali\\_set](https://en.wikipedia.org/wiki/Vitali_set)

# Chapter 2

## Integration

### Lecture 10: Integration

#### 2.1 Measurable Function

26 Jan. 11:00

We start with a definition.

**Definition 2.1.1** (Measurable function). Suppose  $(X, \mathcal{A}), (Y, \mathcal{B})$  are measurable spaces. Then we say  $f: X \rightarrow Y$  is  $(\mathcal{A}, \mathcal{B})$ -measurable if

$$\forall_{B \in \mathcal{B}} f^{-1}(B) \in \mathcal{A}.$$

**Remark.** If  $\mathcal{A}$  and  $\mathcal{B}$  are given, we'll sometimes say  $f$  is measurable if it'll not cause any confusions.

**Lemma 2.1.1.** Given two measurable spaces  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$ , and suppose  $\mathcal{B} = \langle \mathcal{E} \rangle$  for some  $\mathcal{E} \subset Y$ . Then,

$$f: X \rightarrow Y \text{ is } (\mathcal{A}, \mathcal{B})\text{-measurable} \Leftrightarrow \forall_{E \in \mathcal{E}} f^{-1}(E) \in \mathcal{A}.$$

**Proof.** We see that the *only if* part ( $\Rightarrow$ ) is clear. On the other direction, we consider the following. Let  $\mathcal{D} = \{E \subset Y \mid f^{-1}(E) \in \mathcal{A}\}$ , then

- $\mathcal{E} \subset \mathcal{D}$  by assumption
- $\mathcal{D}$  is a  $\sigma$ -algebra

Check!

hence, we see that  $\langle \mathcal{E} \rangle = \mathcal{B} \subset \mathcal{D}$  from Lemma 1.1.2. The result then follows from the definition of  $(\mathcal{A}, \mathcal{B})$ -measurable. ■

**Note.** Recall that

- $f^{-1}(E^c) = f^{-1}(E)^c$
- $f^{-1}(\bigcup_{\alpha} E_{\alpha}) = \bigcup_{\alpha} f^{-1}(E_{\alpha})$

**Definition 2.1.2** ( $\mathcal{A}$ -measurable). Let  $(X, \mathcal{A})$  be a measurable space. Then,

$$\left. \begin{array}{l} f: X \rightarrow \mathbb{R} \\ f: X \rightarrow \overline{\mathbb{R}} \\ f: X \rightarrow \mathbb{C} \end{array} \right\} \text{ is } \mathcal{A}\text{-measurable if } \left\{ \begin{array}{l} f \text{ is } (\mathcal{A}, \mathcal{B}(\mathbb{R}))\text{-measurable} \\ f \text{ is } (\mathcal{A}, \mathcal{B}(\overline{\mathbb{R}}))\text{-measurable} \\ \operatorname{Re} f, \operatorname{Im} f: X \rightarrow \mathbb{R} \text{ are } \mathcal{A}\text{-measurable.} \end{array} \right.$$



**Notation.** Notice that

- $\overline{\mathbb{R}} = [-\infty, \infty]$
- $\mathcal{B}(\overline{\mathbb{R}}) = \{E \subset \overline{\mathbb{R}} \mid E \cap \mathbb{R} \in \mathcal{B}(\mathbb{R})\}$ .
- $\operatorname{Re} f$  is the real part of  $f$ , while  $\operatorname{Im} f$  is the imaginary part of  $f$ .

**Example.** We see that

- $\mathcal{A} = \mathcal{P}(X) \Rightarrow$  Every function is  $\mathcal{A}$ -measurable.
- $\mathcal{A} = \{\emptyset, X\} \Rightarrow$  The only  $\mathcal{A}$ -measurable functions are constant functions.

There are two very common kinds of measurable functions are worth mentioning.

**Definition.** Given a measurable function  $f$ , we have the following.

**Definition 2.1.3** (Lebesgue measurable function).  $f$  is a Lebesgue measurable function if  $f: (\mathbb{R}, \mathcal{L}) \rightarrow (\mathbb{C}, \mathcal{B}(\mathbb{C}))$ .

**Definition 2.1.4** (Borel measurable function).  $f$  is a Borel measurable function if  $f: (\mathbb{R}, \mathcal{B}(\mathbb{R})) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))$ .

**Lemma 2.1.2.** Given  $f: X \rightarrow \mathbb{R}$ , then the following are equivalent:

- $f$  is  $\mathcal{A}$ -measurable;
- $\forall a \in \mathbb{R}, f^{-1}((a, \infty)) \in \mathcal{A}$ ;
- $\forall a \in \mathbb{R}, f^{-1}([a, \infty)) \in \mathcal{A}$ ;
- $\forall a \in \mathbb{R}, f^{-1}((-\infty, a)) \in \mathcal{A}$ ;
- $\forall a \in \mathbb{R}, f^{-1}((-\infty, a]) \in \mathcal{A}$ .

**Proof.** The result follows from Lemma 2.1.1 we just saw. ■

**Remark** (Operations preserve  $\mathcal{A}$ -measurability). Given  $f, g: X \rightarrow \mathbb{R}$  and is  $\mathcal{A}$ -measurable, then

- $\phi: \mathbb{R} \rightarrow \mathbb{R}$ ,  $\mathcal{A}$ -measurable, then

$$\phi \circ f: X \rightarrow \mathbb{R}$$

is  $\mathcal{A}$ -measurable.

- $-f, 3f, f^2, |f|$  are all  $\mathcal{A}$ -measurable, and  $\frac{1}{f}$  is  $\mathcal{A}$ -measurable if  $f(x) \neq 0, \forall x \in X$ .

- $f + g$  is  $\mathcal{A}$ -measurable. We see this from

$$(f + g)^{-1}((a, \infty)) = \bigcup_{r \in \mathbb{Q}} (f^{-1}((r, \infty)) \cap g^{-1}((a - r, \infty)))$$

with Lemma 2.1.2.

- $f \cdot g$  is  $\mathcal{A}$ -measurable. We see this from

$$f(x)g(x) = \frac{1}{2} ((f(x) + g(x))^2 - f(x)^2 - g(x)^2).$$

(e) We see that

$$(f \vee g)(x) := \max\{f(x), g(x)\} \text{ and } (f \wedge g)(x) := \min\{f(x), g(x)\}$$

are  $\mathcal{A}$ -measurable.

(f) Let  $f_n: X \rightarrow \overline{\mathbb{R}}$  be  $\mathcal{A}$ -measurable. Then

$$\sup_{n \in \mathbb{N}} f_n, \inf_{n \in \mathbb{N}} f_n, \limsup_{n \rightarrow \infty} f_n, \liminf_{n \rightarrow \infty} f_n$$

are  $\mathcal{A}$ -measurable.

**Proof.** Consider  $\sup_{n \in \mathbb{N}} f_n =: g$ , then

$$g^{-1}((a, \infty]) = \bigcup_{n \in \mathbb{N}} f_n^{-1}((a, \infty])$$

for  $\sup_n f_n(x) = g(x) > a$ . A similar argument can prove the case of  $\inf_{n \in \mathbb{N}} f_n$ .

And notice that  $\limsup_{n \rightarrow \infty} f_n = \inf_{k \in \mathbb{N}} \sup_{n \geq k} f_n$ , then the similar argument also proves this case.  $\otimes$

check

(g) If  $\lim_{n \rightarrow \infty} f_n(x)$  converges for every  $x \in X$ , then  $f$  is  $\mathcal{A}$ -measurable.

(h) If  $f: \mathbb{R} \rightarrow \mathbb{R}$  is continuous  $\Rightarrow f$  is Borel measurable  $\Rightarrow f$  is Lebesgue measurable since the preimage of an open set of a continuous function is open, then we consider  $f^{-1}((a, \infty))$ .

**Definition 2.1.5 (Support).** The *support* of function  $f: X \rightarrow \overline{\mathbb{R}}$  is

$$\text{supp } f := \{x \in X \mid f(x) \neq 0\}.$$

**Definition.** For  $f: X \rightarrow \overline{\mathbb{R}}$ , let  $f^+ := f \vee 0$  and  $f^- := (-f) \vee 0$ , i.e.,  $f^+(x) = \max\{f(x), 0\}$ ,  $f^-(x) = \max\{-f(x), 0\}$ . Then we have the following.

**Definition 2.1.6 (Positive part).**  $f^+$  is the *positive part* of  $f$ .

**Definition 2.1.7 (Negative part).**  $f^-$  is the *negative part* of  $f$ .

**Remark.** If  $\text{supp } f^+ \cap \text{supp } f^- = \emptyset$  and  $f(x) = f^+(x) - f^-(x)$ , then

$$f \text{ is } \mathcal{A}\text{-measurable} \Leftrightarrow f^+, f^- \text{ are } \mathcal{A}\text{-measurable}.$$

**Definition 2.1.8 (Characteristic (Indicator) function).** For  $E \subset X$ , the *characteristic (indicator) function* of  $E$  is

$$\mathcal{X}_E(x) = \mathbb{1}_E(x) = \begin{cases} 1, & \text{if } x \in E; \\ 0, & \text{if } x \in E^c. \end{cases}$$

**Remark.** We see that  $\mathbb{1}_E$  is  $\mathcal{A}$ -measurable  $\Leftrightarrow E \in \mathcal{A}$ .

**Definition 2.1.9 (Simple function).** Let  $(X, \mathcal{A})$  be a measurable space. Then a *simple function*  $\phi: X \rightarrow \mathbb{C}$  that is  $\mathcal{A}$ -measurable and takes only finitely many values.

**Remark.** We see that if  $\phi(X) = \{c_1, \dots, c_N\}$ , then

$$E_i = \phi^{-1}(\{c_i\}) \in \mathcal{A} \Rightarrow \phi = \sum_{i=1}^N \underbrace{c_i}_{\neq \pm\infty} \underbrace{\mathbb{1}_{E_i}}_{\in \mathcal{A}}.$$

## Lecture 11: Integration of nonnegative functions

As previously seen. For a **simple function**  $\phi$ ,  $c_i$  can actually be in  $\mathbb{C}$ .

31 Jan. 11:00

**Theorem 2.1.1.** Given a **measurable space**  $(X, \mathcal{A})$  and let  $f: X \rightarrow [0, \infty]$ , the following are equivalent.

- (a)  $f$  is an  **$\mathcal{A}$ -measurable** function.
- (b) There exists **simple functions**  $0 \leq \phi_1(x) \leq \phi_2(x) \leq \dots \leq f(x)$  such that

$$\forall x \in X \quad \lim_{n \rightarrow \infty} \phi_n(x) = f(x)$$

i.e.,  $f$  is a **pointwise upward** limit of **simple functions**.

**Proof.** We'll prove both directions.

**Claim.** (b)  $\Rightarrow$  (a).

**Proof.** It's clear from the fact that  $f(x) = \sup_n \phi_n(x)$  and **the remark**. ⊗

**Claim.** (a)  $\Rightarrow$  (b).

**Proof.** Assume  $f$  is  **$\mathcal{A}$ -measurable**, and fix  $n \in \mathbb{N}$ . Let  $F_n = f^{-1}([2^n, \infty]) \in \mathcal{A}$ . Also, for  $0 \leq k \leq 2^{2n} - 1$ ,  $E_{n,k} = f^{-1}([\frac{k}{2^n}, \frac{k+1}{2^n}]) \in \mathcal{A}$ . Then, define  $\phi_n$  be

$$\phi_n = \sum_{k=0}^{2^{2n}-1} \frac{k}{2^n} \mathbb{1}_{E_{n,k}} + 2^n \mathbb{1}_{F_n},$$

we have

- $0 \leq \phi_1(x) \leq \phi_2(x) \leq \dots \leq f(x)$  for every  $x \in X$
- $\forall x \in X \setminus F_n$ , we have  $0 \leq f(x) - \phi_n(x) \leq \frac{1}{2^n}$

Furthermore, we see that

$$F_1 \supset F_2 \supset \dots, \quad \bigcap_{n=1}^{\infty} F_n = f^{-1}(\{\infty\}),$$

then

- $x \in f^{-1}([0, \infty]) = X \setminus \bigcap_{n=1}^{\infty} F_n \Rightarrow \lim_{n \rightarrow \infty} \phi_n(x) = f(x)$
- $x \in f^{-1}(\{\infty\}) = \bigcap_{n=1}^{\infty} F_n \Rightarrow f_n(x) \geq 2^n \Rightarrow \lim_{n \rightarrow \infty} \phi_n(x) = \infty = f(x)$

⊗

■

**Corollary 2.1.1.** If  $f$  is bounded on a set  $A \subset \mathbb{R}$ , i.e.,  $\exists L > 0$  such that

$$\forall_{x \in A} |f(x)| \leq L,$$

then there exists a sequence of **simple functions**  $\{\phi_n\}$  such that  $\phi_n \rightarrow f$  **uniformly** on  $A$ .

**Proof.** \_\_\_\_\_ ■

DIY

**Corollary 2.1.2.** If  $f: X \rightarrow \mathbb{C}$  is a **measurable function** if and only if there exists **simple functions**  $\phi_n: X \rightarrow \mathbb{C}$  such that

$$0 \leq |\phi_1(x)| \leq |\phi_2(x)| \leq \cdots \leq |f(x)|$$

with

$$\forall_{x \in X} \lim_{n \rightarrow \infty} \phi_n(x) = f(x).$$

**Proof.** \_\_\_\_\_ ■

DIY

## 2.2 Integration of Nonnegative Functions

We start with our first definition about integral.

**Definition 2.2.1** (Integration of nonnegative function). Let  $(X, \mathcal{A}, \mu)$  be a **measure space**, and  $\phi: X \rightarrow [0, \infty]$  such that

$$\phi = \sum_{i=1}^N c_i \mathbb{1}_{E_i}$$

be a **simple function**. Define

$$\int \phi = \int \phi \, d\mu = \int_X \phi \, d\mu = \sum_{i=1}^N c_i \mu(E_i).$$

Furthermore, for  $A \in \mathcal{A}$ ,

$$\int_A \phi = \int_A \phi \, d\mu = \int \phi \mathbb{1}_A \, d\mu.$$

**Note.** Note that

- In the expression  $\sum_{i=1}^N c_i \mu(E_i)$ , we're using the convention  $0 \cdot \infty = 0$ .
- The function  $\phi \mathbb{1}_A$  is also a **simple function** since both  $\phi$  and  $\mathbb{1}_A$  are **simple function**.

**Proposition 2.2.1.** Suppose we have  $\phi, \psi \geq 0$  be two **simple functions**. Then,

- Definition 2.2.1** is well-defined.
- $\int c\phi = c \int \phi$  for  $c \in [0, \infty)$ .
- $\int \phi + \psi = \int \phi + \int \psi$ .
- $\phi(x) \geq \psi(x)$  for all  $x \Rightarrow \int \phi \geq \int \psi$ .
- $\nu(A) = \int_A \phi \, d\mu$  is a **measure** on  $(X, \mathcal{A})$ .

**Proof.** \_\_\_\_\_ ■

DIY

**Definition 2.2.2** (Generalization of Integration of nonnegative function). Given a **measure space**  $(X, \mathcal{A}, \mu)$  with  $f: X \rightarrow [0, \infty]$  be  **$\mathcal{A}$ -measurable**. Define

$$\int f = \int f \, d\mu = \sup \left\{ \int \phi : 0 \leq \phi \leq f \text{ such that } \phi \text{ is simple} \right\}.$$

**Note.** Note that

- If  $f$  is a **simple function**, the **Definition 2.2.1** and **Definition 2.2.2** of  $\int f$  are the same.
- $\int cf = c \int f$  for  $c \in [0, \infty)$ .
- If  $f \geq g \geq 0 \Rightarrow \int f \geq \int g$ .
- But  $\int f + g = \int f + \int g$  is not trivial.

**Theorem 2.2.1** (Monotone convergence theorem). Given  $(X, \mathcal{A}, \mu)$  be a **measure space**. Then if

- $f_n: X \rightarrow [0, \infty]$  be  **$\mathcal{A}$ -measurable** for every  $n \in \mathbb{N}$ ;
- $0 \leq f_1(x) \leq f_2(x) \leq \dots$  for every  $x \in X$ ;
- $\lim_{n \rightarrow \infty} f_n(x) = f(x)$  for every  $x \in X$ ,

we have

$$\lim_{n \rightarrow \infty} \int f_n = \int f.$$

**Proof.** Note that if  $\lim_{n \rightarrow \infty} \int f_n$  exists, then it's equal to  $\sup_n \int f_n$ . Then

- $f_n \leq f \Rightarrow \int f_n \leq \int f \Rightarrow \lim_{n \rightarrow \infty} \int f_n \leq \int f$ .
- Fix a **simple function**  $0 \leq \phi \leq f$ , then it's enough to show  $\lim_{n \rightarrow \infty} \int f_n \geq \int \phi$ . We first fix  $\alpha = (0, 1)$ , then it's also enough to show

$$\lim_{n \rightarrow \infty} \int f_n \geq \alpha \int \phi.$$

Let  $A_n := \{x \in X \mid f_n(x) \geq \alpha \phi(x)\}$ , then since  $f_n$  is **measurable**,

- $A_n \in \mathcal{A}$
- $A_1 \subset A_2 \subset A_3 \subset \dots$
- $\bigcup_{n=1}^{\infty} A_n = X$

We then have

$$\int f_n \geq \int f_n \mathbb{1}_{A_n} \geq \int \alpha \phi \mathbb{1}_{A_n} = \alpha \int_{A_n} \phi = \alpha \nu(A_n)$$

where  $\nu(A) = \int_A \phi$  is a **measure**. This implies

$$\lim_{n \rightarrow \infty} \int f_n \geq \alpha \lim_{n \rightarrow \infty} \nu(A_n) \stackrel{!}{=} \alpha \nu(X) = \alpha \int \phi.$$

■

**Corollary 2.2.1** (Linearity of nonnegative integral). Let  $f, g \geq 0$  be **measurable**, then

$$\int f + g = \int f + \int g.$$

**Proof.** There exists [simple functions](#)  $\phi_n$  and  $\psi_n$  such that

- $0 \leq \phi_1 \leq \phi_2 \leq \dots$  and  $\phi_n \rightarrow f$  [pointwise](#)
- $0 \leq \psi_1 \leq \psi_2 \leq \dots$  and  $\psi_n \rightarrow g$  [pointwise](#)

Then,

$$\int (f + g) \stackrel{!}{=} \lim_{n \rightarrow \infty} \int (\phi_n + \psi_n) = \lim_{n \rightarrow \infty} \int \phi_n + \int \psi_n \stackrel{!}{=} \int f + \int g.$$

■

## Lecture 12: Fatou's Lemma

We start with a useful corollary.

2 Feb. 11:00

**Corollary 2.2.2** (Tonelli's theorem for nonnegative series and integrals). Given  $g_n \geq 0$  for every  $n \in \mathbb{N}$  and let  $g_n$  be [measurable](#), then

$$\int \sum_{n=1}^{\infty} g_n = \sum_{n=1}^{\infty} \int g_n.$$

**Proof.** Let  $f_N := \sum_{n=1}^N g_n$  such that  $\lim_{N \rightarrow \infty} f_N = \sum_{n=1}^{\infty} g_n =: f$ , then since  $g_n \geq 0$ , we have  $0 \leq f_1 \leq f_2 \leq \dots$  with

$$\lim_{N \rightarrow \infty} f_N(x) = \sum_{n=1}^{\infty} g_n(x).$$

By [Theorem 2.2.1](#), we have

$$\lim_{N \rightarrow \infty} \underbrace{\int \sum_{n=1}^N g_n}_{f_N} = \underbrace{\int \sum_{n=1}^{\infty} g_n}_f.$$

Now, since the terms in the limit on the left-hand side is just a finite sum, by [Corollary 2.2.1](#), we have

$$\underbrace{\lim_{N \rightarrow \infty} \sum_{n=1}^N \int g_n}_{\sum_{n=1}^{\infty} \int g_n} = \lim_{N \rightarrow \infty} \int \sum_{n=1}^N g_n = \int \sum_{n=1}^{\infty} g_n,$$

hence

$$\int \sum_{n=1}^{\infty} g_n = \sum_{n=1}^{\infty} \int g_n.$$

■

**Remark.** Recall that we have seen [two series case](#) before. We'll later see two integrals cases.

**Theorem 2.2.2** (Fatou's Lemma). Suppose  $f_n \geq 0$  and [measurable](#), then

$$\int \liminf_{n \rightarrow \infty} f_n \leq \liminf_{n \rightarrow \infty} \int f_n.$$

**Proof.** Before we start we note the following.

**Remark.** Recall that

$$\liminf_{n \rightarrow \infty} f_n := \lim_{k \rightarrow \infty} \inf_{n \geq k} f_n = \sup_{k \in \mathbb{N}} \inf_{n \geq k} f_n$$

and

$$\exists \lim_{n \rightarrow \infty} a_n \Leftrightarrow \limsup_{n \rightarrow \infty} a_n = \liminf_{n \rightarrow \infty} a_n.$$

Let  $g_k = \inf_{n \geq k} f_n$ , then  $g_k$  is measurable and  $0 \leq g_1 \leq g_2 \leq \dots$ . Now, from Theorem 2.2.1, we have

$$\int \lim_{k \rightarrow \infty} g_k = \lim_{k \rightarrow \infty} \int g_k.$$

Notice that the left-hand side is just  $\int \liminf_{n \rightarrow \infty} f_n$ , while the right-hand side is just  $\lim_{k \rightarrow \infty} \int \inf_{n \geq k} f_n$ , i.e.,

$$\int \liminf_{n \rightarrow \infty} f_n = \lim_{k \rightarrow \infty} \int \inf_{n \geq k} f_n.$$

We see that we want to take the inf outside the integral on the right-hand side. Observe that

$$\forall_{m \geq k} \inf_{n \geq k} f_n \leq f_m \Rightarrow \forall_{m \geq k} \int \inf_{n \geq k} f_n \leq \int f_m \Rightarrow \int \inf_{n \geq k} f_n \leq \inf_{m \geq k} \int f_m.$$

Then, we have

$$\int \liminf_{n \rightarrow \infty} f_n = \lim_{k \rightarrow \infty} \int \inf_{n \geq k} f_n \leq \lim_{k \rightarrow \infty} \inf_{m \geq k} \int f_m = \liminf_{m \rightarrow \infty} \int f_m.$$

■

**Example (Escape to horizontal infinity).** Given  $(\mathbb{R}, \mathcal{L}, m)$ , let  $f_n := \mathbb{1}_{(n, n+1)}$ . We immediately see that

- $f_n \rightarrow 0$  pointwise
- $\int f_n = 1$  for every  $n$
- $\int f = 0$

From Theorem 2.2.2, we have a strict inequality

$$0 = \int \liminf_{n \rightarrow \infty} f_n, \liminf_{n \rightarrow \infty} \int f_n = 1.$$

**Example (Escape to width infinity).** Given  $(\mathbb{R}, \mathcal{L}, m)$ , let  $f_n := \frac{1}{n} \mathbb{1}_{(0, n)}$ .

**Example (Escape to vertical infinity).** Given  $(\mathbb{R}, \mathcal{L}, m)$ , let  $f_n := n \mathbb{1}_{(0, \frac{1}{n})}$ .

**Lemma 2.2.1** (Markov's inequality). Let  $f \geq 0$  be measurable. Then

$$\forall_{c \in (0, \infty)} \mu(\{x \mid f(x) \geq c\}) \leq \frac{1}{c} \int f.$$

**Proof.** Denote  $\{x \mid f(x) \geq c\} =: E$ , then

$$f(x) \geq c \mathbb{1}_E(x) \Rightarrow \int f \geq c \int \mathbb{1}_E = c \cdot \mu(E).$$

■

**Remark.** Notice that  $E = f^{-1}([c, \infty])$ , hence  $E$  is measurable.

**Proposition 2.2.2.** Let  $f \geq 0$  be measurable. Then,

$$\int f = 0 \Leftrightarrow f = 0 \text{ a.e.}$$

i.e.,

$$\int f d\mu = 0 \Leftrightarrow \mu(A) = 0$$

where  $A = \{x \mid f(x) > 0\} = f^{-1}((0, \infty])$ .

**Proof.** Firstly, assume that  $f = \phi$  is a simple function. We may write

$$\phi = \sum_{i=1}^N c_i \mathbb{1}_{E_i}$$

where  $E_i$  are disjoint and  $c_i \in (0, \infty)$ . Then,

$$\int \phi = \sum_{i=1}^N c_i \mu(E_i) = 0 \Leftrightarrow \mu(E_1) = \dots = \mu(E_N) = 0 \Leftrightarrow \mu(A) = 0, \quad A = \bigcup_{i=1}^N E_i.$$

Now, assume that  $f$  is a general function where  $f \geq 0$  is the only constraint, and we consider two cases.

- Assume  $\mu(A) = 0$  (i.e.,  $f = 0$  a.e.). Let  $0 \leq \phi \leq f$ , where  $\phi$  is simple. Then

$$\forall_{x \in A^c} \phi(x) = 0$$

since  $f(x) = 0, \forall x \in A^c$ . This implies that  $\phi = 0$  a.e. since  $\mu(A) = 0$ , so  $\int \phi = 0$ . We then have

$$\int f = 0$$

from Definition 2.2.2.

- Assume  $\int f = 0$ . Let  $A_n = f^{-1}([\frac{1}{n}, \infty])$ . Then we see that
  - $A_1 \subset A_2 \subset \dots$
  - $\bigcup_{n=1}^{\infty} A_n = f^{-1}(\bigcup_{n=1}^{\infty} [\frac{1}{n}, \infty]) = f^{-1}((0, \infty)) = A$ .

We then have

$$\mu(A_n) = \mu\left(\left\{x \mid f(x) \geq \frac{1}{n}\right\}\right) \stackrel{!}{\leq} n \int f = 0,$$

which further implies

$$\mu(A) = \lim_{n \rightarrow \infty} \mu(A_n) = 0$$

from the continuity of measure from below. ■

**Corollary 2.2.3.** If  $f, g \geq 0$  are both measurable and  $f = g$  a.e., then

$$\int f = \int g.$$

**Proof.** Let  $A = \{x \mid f(x) \neq g(x)\}^a$ . Then by assumption,  $\mu(A) = 0$ , hence

$$f \mathbb{1}_A = 0 \text{ a.e.}, \quad g \mathbb{1}_A = 0 \text{ a.e.}$$



This further implies that

$$\begin{aligned}\int f &= \int f(\mathbb{1}_A + \mathbb{1}_{A^c}) \stackrel{!}{=} \int f\mathbb{1}_A + \int f\mathbb{1}_{A^c} \\ &= \int f\mathbb{1}_{A^c} = \int g\mathbb{1}_{A^c} = \int g\mathbb{1}_{A^c} + \int g\mathbb{1}_A = \int g.\end{aligned}$$

■

<sup>a</sup> $A$  is measurable indeed.

**Corollary 2.2.4.** Let  $f_n \geq 0$  be measurable. Then

- (a)  $\left. \begin{array}{l} 0 \leq f_1 \leq f_2 \leq \dots \leq f \text{ a.e.} \\ \lim_{n \rightarrow \infty} f_n = f \text{ a.e.} \end{array} \right\} \Rightarrow \lim_{n \rightarrow \infty} \int f_n = \int f.$
- (b)  $\lim_{n \rightarrow \infty} f_n = f \text{ a.e.} \Rightarrow \int f \leq \liminf_{n \rightarrow \infty} \int f_n.$

**Proof.**

■

DIY

**Remark.** Almost all the theorems we've proved can be replaced by theorems dealing with **almost everywhere** condition.

## Lecture 13: Integration of Complex Functions

### 2.3 Integration of Complex Functions

4 Feb. 11:00

As usual, we start with a definition.

**Definition 2.3.1 (Integrable).** Let  $(X, \mathcal{A}, \mu)$  be a **measure space** and let  $f: X \rightarrow \overline{\mathbb{R}}$  and  $g: X \rightarrow \mathbb{C}$  be **measurable**.<sup>a</sup>

Then  $f, g$  are called *integrable* if  $\int |f| < \infty$  and  $\int |g| < \infty$ , and we define

$$\int f = \int f^+ - \int f^-, \quad \int g = \int \operatorname{Re} g + i \int \operatorname{Im} g.$$

Furthermore, for  $f: X \rightarrow \overline{\mathbb{R}}$ , we define

$$\int f = \begin{cases} \infty, & \text{if } \int f^+ = \infty, \int f^- < \infty; \\ -\infty, & \text{if } \int f^+ < \infty, \int f^- = \infty. \end{cases}$$

<sup>a</sup>Recall that for a complex-valued function like  $g$ , this means that both  $\operatorname{Re} g$  and  $\operatorname{Im} g$  are measurable.

We now see a lemma.

**Lemma 2.3.1.** Let  $f, g: X \rightarrow \overline{\mathbb{R}}$  or  $\mathbb{C}$  **integrable**. Assume that  $f(x) + g(x)$  is well-defined for all  $x \in X$ .<sup>a</sup> Then we have

- (a)  $f + g, cf$  for all  $c \in \mathbb{C}$  are **integrable**.
- (b)  $\int f + g = \int f + \int g$ . This is not trivial since  $(f + g)^+ \neq f^+ + g^+$ .
- (c)  $|\int f| \leq \int |f|$ .

<sup>a</sup>That is, we never see  $\infty + (-\infty)$  or  $(-\infty) + \infty$ .

**Proof.** Check [FF99] page 53. ■

**Lemma 2.3.2.** Let  $(X, \mathcal{A}, \mu)$  be a **measure space** and let  $f$  be an **integrable** function on  $X$ . Then

- (a)  $f$  is finite **a.e.**, i.e.,  $\{x \in X \mid |f(x)| = \infty\}$  is a **null set**.
- (b) The set  $\{x \in X \mid f(x) \neq 0\}$  is  **$\sigma$ -finite**.

**Proof.** ■

HW 5  
Q8 by  
Lemma 2.2.1

**Proposition 2.3.1.** Let  $(X, \mathcal{A}, \mu)$  be a **measure space**, then

- (a) If  $h$  is **integrable** on  $X$ , then

$$\forall_{E \in \mathcal{A}} \int_E h = 0 \Leftrightarrow \int |h| = 0 \Leftrightarrow h = 0 \text{ a.e.}$$

- (b) If  $f, g$  are **integrable** on  $X$ , then

$$\forall_{E \in \mathcal{A}} \int_E f = \int_E g \Leftrightarrow f = g \text{ a.e.}$$

**Proof.** We prove this one by one.

- (a) We see that the second equivalence is done in **Proposition 2.2.2**, hence we prove the first equivalence only. Since we have

$$\int |h| = 0 \Rightarrow \left| \int_E h \right| \leq \int_E |h| \leq \int |h| = 0,$$

which shows one implication. Now assume that  $\int_E h = 0$  for all  $E \in \mathcal{A}$ , then we can write  $h$  as

$$h = u + iv = (u^+ - u^-) + i(v^+ - v^-).$$

Let  $B := \{x \in X \mid u^+(x) > 0\}$ , then by assumption, we have

$$0 = \int_B h = \operatorname{Re} \int_B h = \int_B u = \int_B u^+ = \int_B u^+ + \int_{B^c} u^+ = \int u^+,$$

hence  $u^+ = 0$  **almost everywhere**. Similarly, we have  $u^-, v^+, v^-$  are all zero **almost everywhere**. This gives us that  $h$  is zero **almost everywhere** as desired.

- (b) ■

DIY

**Theorem 2.3.1** (Dominated convergence theorem). Let  $(X, \mathcal{A}, \mu)$  be a **measure space**, and

- Let  $f_n$  be **integrable** on  $X$ .
- $\lim_{n \rightarrow \infty} f_n(x) = f(x)$  **almost everywhere**.
- There is a  $g: X \rightarrow [0, \infty]$  such that  $g$  is **integrable** and

$$\forall_{n \in \mathbb{N}} |f_n(x)| \leq g(x) \text{ a.e.}$$

Then we have

$$\lim_{n \rightarrow \infty} \int f_n = \int f = \int \lim_{n \rightarrow \infty} f_n.$$

**Proof.** Let  $F$  be the countable union of [null set](#) on which the three conditions may fail. Then we see that after modifying the definition of  $f_n, f$  and  $g$  on  $F$ , we may assume that all three conditions hold everywhere since modifying on a [null set](#) does not change the integral.

We now consider the  $\mathbb{R}$ -valued case only. Note that the second and the third conditions imply that  $f$  is [integrable](#) since  $|f| \leq g(x)$ . We then see that  $g + f_n \geq 0$  and  $g - f_n \geq 0$  because  $-g \leq f_n \leq g$ . From [Theorem 2.2.2](#), we have

$$\int g + f \leq \liminf_{n \rightarrow \infty} \int g + f_n, \quad \int g - f \leq \liminf_{n \rightarrow \infty} \int g - f_n.$$

From the [linearity of integral](#), we have

$$\int g + \int f \leq \int g + \liminf_{n \rightarrow \infty} \int f_n, \quad \int g - \int f \leq \int g - \liminf_{n \rightarrow \infty} \int f_n.$$

Now, since  $\int g < \infty$ , we can cancel it, which gives

$$\int f \leq \liminf_{n \rightarrow \infty} \int f_n, \quad -\int f \leq \liminf_{n \rightarrow \infty} \int -f_n = -\limsup_{n \rightarrow \infty} \int f_n,$$

which implies

$$\int f \leq \liminf_{n \rightarrow \infty} \int f_n \leq \limsup_{n \rightarrow \infty} \int f_n \leq \int f.$$

This shows that the limit exists, and the desired result indeed holds. ■

**Corollary 2.3.1** (Tonelli's theorem for series and integrals). Suppose  $f_n$  are [integrable](#) functions such that

$$\sum_{n=1}^{\infty} \int |f_n| < \infty,$$

then we have

$$\int \sum_{n=1}^{\infty} f_n = \sum_{n=1}^{\infty} \int f_n.$$

**Proof.** Take  $G(x)$  to be

$$G(x) := \sum_{n=1}^{\infty} |f_n(x)|,$$

then we see  $G(x) \geq |F_N(x)|$  where  $F_N(x) := \sum_{n=1}^N f_n(x)$ . By [Corollary 2.2.2](#), we have

$$\int G(x) = \sum_{n=1}^{\infty} \int |f_n(x)| < \infty.$$

Lastly, from [Theorem 2.3.1](#), the result follows. ■

**Remark.** Compare to [Corollary 2.2.2](#), we see that we further generalize the result!

## Lecture 14: $L^1$ Space

### 2.4 $L^1$ Space

7 Feb. 11:00

We now introduce another space called  $L^p$  spaces, which are function spaces defined using a natural generalization of the [p-norm](#) for finite-dimensional vector spaces. We sometimes call it Lebesgue spaces also. Before we start, we need to define a *norm*.

**Definition 2.4.1 (Norm).** Let  $V$  be a vector space over field  $\mathbb{R}$  or  $\mathbb{C}$ . A *norm* is a [seminorm](#)

**Definition 2.4.2 (Seminorm).** A *seminorm* on  $V$  is

$$\|\cdot\| : V \rightarrow [0, \infty)$$

such that

- $\|cv\| = |c| \|v\|$  for every  $v \in V$  and every scalar  $c$ .
- $\|v + w\| \leq \|v\| + \|w\|$  for every  $v, w \in V$ .

with an additional condition

- $\|v\| = 0 \Leftrightarrow v = 0$ .

**Lemma 2.4.1.** A [normed](#) vector space is a metric space with metric  $\rho(v, w) = \|v - w\|$ .

**Proof.** ■

DIY


**Example ( $p$ -norm).**  $V = \mathbb{R}^d$  with

$$\|x\|_p = \begin{cases} \left( \sum_{i=1}^d |x_i|^p \right)^{1/p}, & \text{if } p \in [0, \infty); \\ \max_{1 \leq i \leq d} |x_i|, & \text{if } p = \infty \end{cases}$$

is a [normed](#) vector space. The unit ball

$$\{x \in \mathbb{R}^d \mid \|x\|_p \leq 1\}$$

for different  $p$  has the following figures.



**Remark.** All  $p$ -norms induce the same topology. i.e., if  $U$  is open in  $p$ -norm, it is open in  $p'$ -norm as well.

**Note.** Recall that we say  $f$  is [integrable](#) means

$$\int |f| < \infty,$$

and if  $f = g$  a.e., then

$$\int f = \int g$$

**Definition 2.4.3** ( $L^1$  Space). Given  $(X, \mathcal{A}, \mu)$ ,

$$f \in L^1(X, \mathcal{A}, \mu) (= L^1(X, \mu) = L^1(X) = L^1(\mu))$$

means that  $f$  is an **integrable** function on  $X$ .

**Lemma 2.4.2.**  $L^1(X, \mathcal{A}, \mu)$  is a vector space with **seminorm**

$$\|f\|_1 = \int |f|.$$

**Proof.** ■

Check this is indeed a **seminorm**.

**Definition 2.4.4** ( $L^1$  Space with equivalence class). Define  $f \sim g$  if  $f = g$  a.e., then

$$L^1(X, \mathcal{A}, \mu) / \sim = L^1(X, \mathcal{A}, \mu),$$

i.e., we simply denote the collection of equivalence classes by itself.<sup>a</sup>

<sup>a</sup>By some abusing of notation of  $L^1$ .

**Remark.** We have

- With **Definition 2.4.4**,  $L^1(X, \mathcal{A}, \mu)$  is a **normed** vector space.
- We say that the  $L^1$ -metric  $\rho(f, g)$  is simply

$$\rho(f, g) = \int |f - g|.$$

## Dense Subsets of $L^1$

**Note.** Recall the definition of a **dense set**.

**Definition 2.4.5** (Step function). A *step function* on  $\mathbb{R}$  is

$$\psi = \sum_{i=1}^N c_i \mathbb{1}_{I_i},$$

where  $I_i$  is an interval.

**Notation.** We denote the collection of continuous functions with compact support by  $C_c(\mathbb{R})$ .

**Theorem 2.4.1.** We have the following.

- $\{\text{integrable simple functions}\}$  is dense in  $L^1(X, \mathcal{A}, \mu)$  (with respect to  $L^1$ -metric).
- $(X, \mathcal{A}, \mu) = (\mathbb{R}, \mathcal{A}_\mu, \mu)$ , where  $\mu$  is a **Lebesgue-Stieltjes-measure**. Then the set of **integrable simple** functions is dense in  $L^1(\mathbb{R}, \mathcal{A}_\mu, \mu)$ .
- $C_c(\mathbb{R})$  is dense in  $L^1(\mathbb{R}, \mathcal{L}, m)$ .

**Proof.** We prove this one by one.

- (a) Since there exists **simple functions**  $0 \leq |\phi_1| \leq |\phi_2| \leq \dots \leq |f|$ , where  $\phi_n \rightarrow f$  **pointwise**. Then by **Theorem 2.3.1**, we have

$$\lim_{n \rightarrow \infty} \int \underbrace{|f_n - f|}_{\leq |\phi_n| + |f| \leq 2|f|} = 0$$

where  $2|f|$  is in  $L^1$ .

- (b) Let  $\mathbb{1}_E$  approximate by  $\sum_{i=1}^{\infty} c_i \mathbb{1}_{I_i}$ . From **Theorem 1.7.1** for **Lebesgue-Stieltjes-measure**,

$$\forall \epsilon' > 0 \exists I = \bigcup_{i=1}^N I_i \text{ such that } \mu(E \triangle I) \leq \epsilon'.$$

- (c) To approximate  $\mathbb{1}_{(a,b)}$ , we simply consider function  $g \in C_c(\mathbb{R})$  such that

$$\int |\mathbb{1}_{(a,b)} - g| \leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

■

## Lecture 15: Riemann Integral

### 2.5 Riemann Integrability

9 Feb. 11:00

We are now working in  $(\mathbb{R}, \mathcal{L}, m)$ . Let's first revisit the definition of Riemann Integral. Let  $P$  be a partition of  $[a, b]$  as

$$P = \{a = t_0 < t_1 < \dots < t_k = b\}.$$

Then the *lower Riemann sum* of  $f$  using  $P$  is equal to  $L_P$ , which is defined as

$$L_P = \sum_{i=1}^K \left( \inf_{[t_{i-1}, t_i]} f \right) (t_i - t_{i-1}),$$

and the *upper Riemann sum* of  $f$  using  $P$  is equal to  $U_P$ , which is defined as

$$U_P = \sum_{i=1}^K \left( \sup_{[t_{i-1}, t_i]} f \right) (t_i - t_{i-1}).$$

Then we call

- *Lower Riemann integral* of  $f = \underline{I} = \sup_P L_P$
- *Upper Riemann integral* of  $f = \bar{I} = \inf_P U_P$

**Definition 2.5.1** (**Riemann (Darboux) integrable**). A **bounded** function  $f: [a, b] \rightarrow \mathbb{R}$  is called *Riemann (Darboux) integrable* if  $\underline{I} = \bar{I}$ . If so, then we write

$$\underline{I} = \bar{I} = \int_a^b f(x) dx.$$

**Note.** We see that

- If  $P \subset P'$ , then

$$L_P \leq L_{P'}, \quad U_{P'} \leq U_P.$$

- Recall that continuous functions on  $[a, b]$  are **Riemann integrable** on  $[a, b]$ .

**Theorem 2.5.1.** Let  $f: [a, b] \rightarrow \mathbb{R}$  be a bounded function. Then

- (a) If  $f$  is **Riemann integrable**, then  $f$  is **Lebesgue measurable**, thus **Lebesgue integrable**. Further,

$$\int_a^b f(x) dx = \int_{[a,b]} f dm.$$

- (b) If  $f$  is **Riemann integrable**  $\Leftrightarrow f$  is continuous Lebesgue a.e.<sup>a</sup>

<sup>a</sup>Here, we mean that the set where  $f$  is discontinuous is a **null set** under **Lebesgue measure**.

**Proof.** There exists  $P_1 \subset P_2 \subset \dots$  such that  $L_{P_n} \nearrow \underline{I}$  and  $U_{P_n} \searrow \bar{I}$ .

**Note.** Here, we took refinements of  $P_n$  if needed.

Now, define **simple (step) functions**

- $\phi_n = \sum_{i=1}^K (\inf_{[t_{i-1}, t_i]} f) \mathbb{1}_{(t_{i-1}, t_i]}$
- $\psi_n = \sum_{i=1}^K (\sup_{[t_{i-1}, t_i]} f) \mathbb{1}_{(t_{i-1}, t_i]}$

if  $P_n = \{a = t_0 < t_1 < \dots < t_K\}$ . Let  $\phi := \sup_n \phi_n$  and  $\psi := \inf_n \psi_n$ . We then see that  $\phi, \psi$  are **Lebesgue (Borel) measurable function**.

**Note.** Note that

- $\exists M > 0$  such that  $\forall_{n \in \mathbb{N}} |\phi_n|, |\psi_n| \leq M \mathbb{1}_{[a,b]}$
- $\int \phi_n dm = L_{P_n}, \int \psi_n dm = U_{P_n}$

By **Theorem 2.3.1** and the fact that  $M \mathbb{1}_{[a,b]} \in L^1(\mathbb{R}, \mathcal{L}, m)$ , we have

$$\underline{I} = \lim_{n \rightarrow \infty} \int \phi_n dm = \int \phi dm, \quad \bar{I} = \lim_{n \rightarrow \infty} \int \psi_n dm = \int \psi dm.$$

Thus,

$$f \text{ is Riemann integrable} \Leftrightarrow \int \phi = \int \psi \Leftrightarrow \int (\psi - \phi) = 0 \Leftrightarrow \psi = \phi \text{ Lebesgue a.e.}$$

■

## 2.6 Modes of Convergence

As we should already see, there are different *modes* of convergence. Let's formalize them.

**Definition.** Let  $f_n, f: X \rightarrow \mathbb{C}$ , and  $S \subset X$ . Then we have the following definitions.

**Definition 2.6.1 (Pointwise convergence).**  $f_n \rightarrow f$  *pointwise* on  $S$  if

$$\forall_{x \in S} \forall_{\epsilon > 0} \exists N \in \mathbb{N} \forall_{n \geq N} |f_n(x) - f(x)| < \epsilon.$$

**Definition 2.6.2** (Uniformly convergence).  $f_n \rightarrow f$  *uniformly* on  $S$  if

$$\forall \epsilon > 0 \exists N \in \mathbb{N} \forall x \in S \forall n \geq N |f_n(x) - f(x)| < \epsilon.$$

**Remark.** We see that we can replace  $\forall \epsilon > 0$  by  $\forall k \in \mathbb{N}$  with  $\epsilon$  changing to  $\frac{1}{k}$ .

**Lemma 2.6.1.** Let  $B_{n,k}$  be

$$B_{n,k} := \left\{ x \in X \mid |f_n(x) - f(x)| < \frac{1}{k} \right\}.$$

Then

(a)  $f_n \rightarrow f$  *pointwise* on  $S$  if and only if

$$S \subset \bigcap_{k=1}^{\infty} \bigcup_{N=1}^{\infty} \bigcap_{n=N}^{\infty} B_{n,k}.$$

(b)  $f_n \rightarrow f$  *uniformly* on  $S$  if and only if  $\exists N_1, N_2, \dots \in \mathbb{N}$  such that

$$S \subset \bigcap_{k=1}^{\infty} \bigcap_{n=N_k}^{\infty} B_{n,k}.$$

**Proof.** This essentially follows from Definition 2.6.1. ■

**Definition.** Let  $(X, \mathcal{A}, \mu)$  be a *measure space*. Assuming that  $f_n, f$  are *measurable functions*, then we have the following.

**Definition 2.6.3** (Converge almost everywhere).  $f_n \rightarrow f$  *almost everywhere* means

$\exists$  *null set*  $E$  such that  $f_n \rightarrow f$  *pointwise* on  $E^c$ .


**Definition 2.6.4** (Converge in  $L^1$ ).  $f_n \rightarrow f$  *in  $L^1$*  means

$$\lim_{n \rightarrow \infty} \|f_n - f\| = 0.$$

**Example.** Given  $(\mathbb{R}, \mathcal{L}, m)$  and let  $f = 0$ . Consider the following functions.

- (a)  $f_n = \mathbb{1}_{(n, n+1)}$ .
- (b)  $f_n = \frac{1}{n} \mathbb{1}_{(0, n)}$ .
- (c)  $f_n = n \mathbb{1}_{(0, \frac{1}{n})}$ .
- (d) The *typewriter function*.





**Proof.** We see that different function sequences converge in different senses.

**Exercise.** Classify in what senses do (a), (b), (c) and the typewriter function converge.

⊛

## Lecture 16: Modes of Convergence

Let's start with a proposition.

11 Feb. 11:00

**Proposition 2.6.1** (Fast  $L^1$  convergence leads to a.e. convergence). Let  $(X, \mathcal{A}, \mu)$  be a **measure space**, and  $f_n, f$  are all **measurable** functions on  $X$ . Then

$$\sum_{n=1}^{\infty} \|f_n - f\|_1 < \infty \Rightarrow f_n \rightarrow f \text{ a.e.}$$

**Proof.** Let

$$E := \bigcup_{k=1}^{\infty} \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} B_{n,k}^c = \{x \in X \mid f_n(x) \not\rightarrow f(x)\}.$$

By Lemma 2.2.1, we see that

$$\forall_k \forall_N \mu(B_{n,k}^c) \leq k \int |f_n - f| \Rightarrow \forall_k \mu\left(\bigcup_{n=N}^{\infty} B_{n,k}^c\right) \leq \sum_{n=N}^{\infty} k \|f_n - f\|_1 \rightarrow 0$$

as  $N \rightarrow \infty$ . Now, by continuity of measure from above,

$$\forall_k \mu\left(\bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} B_{n,k}^c\right) = \lim_{N \rightarrow \infty} \mu\left(\bigcup_{n=N}^{\infty} B_{n,k}^c\right) = 0 \Rightarrow \mu(E) = 0$$

since  $f_n \rightarrow f$  pointwise on  $E^c$ . ■

**Corollary 2.6.1.** Given  $\{f_n\}_n$  such that  $f_n \rightarrow f$  in  $L^1$ , there exists a subsequence  $\{f_{n_j}\}_{n_j}$  where  $f_{n_j} \rightarrow f$  a.e.

**Proof.** Since

$$\forall_{j \in \mathbb{N}} \forall_{n_j \in \mathbb{N}} \|f_{n_j} - f\|_1 \leq \frac{1}{j^2}.$$

Then,

$$\sum_{j=1}^{\infty} \|f_{n_j} - f\|_1 < \infty.$$

From Proposition 2.6.1, we have the desired result. ■

**Definition 2.6.5** (Converge in measure). Let  $f_n, f$  be measurable functions on  $(X, \mathcal{A}, \mu)$ . Then  $f_n \rightarrow f$  in measure if

$$\forall_{\epsilon > 0} \lim_{n \rightarrow \infty} \mu(\{x \in X \mid |f_n(x) - f(x)| \geq \epsilon\}) = 0.$$

**Example.** Let  $f_n = n \mathbb{1}_{(0, \frac{1}{n})}$  and  $f = 0$  on  $(\mathbb{R}, \mathcal{B}(\mathbb{R}), m)$ , then  $f_n \rightarrow f$  in measure.


**Proof.** We see that

$$\forall \epsilon > 0 \quad \{x \in X \mid |f_n(x) - f(x)| > \epsilon\} = \left(0, \frac{1}{n}\right),$$

$f_n \rightarrow 0$  in measure. (Recall that  $f_n \not\rightarrow 0$  in  $L^1$ )

⊛

**Remark.** We see that



Finally, we have the following.

**Definition.** Let  $f_n, f$  be measurable functions on  $(X, \mathcal{A}, \mu)$ .

**Definition 2.6.6** (Uniformly almost everywhere).  $f_n \rightarrow f$  uniformly almost everywhere if  $\exists$  null set  $F$  such that  $f_n \rightarrow f$  uniformly on  $F^c$ .

**Definition 2.6.7** (Almost uniformly).  $f_n \rightarrow f$  *almost uniformly* if  $\forall \epsilon > 0 \exists F \in \mathcal{A}$  such that  $\mu(F) < \epsilon$ ,  $f_n \rightarrow f$  **uniformly** on  $F^c$ .

**Lemma 2.6.2.** We have

$$f_n \rightarrow f \text{ uniformly on } S \Leftrightarrow \exists N_1, N_2, \dots \in \mathbb{N} \ S \subset \bigcap_{k=1}^{\infty} \bigcup_{n=N_k}^{\infty} B_{n,k}.$$

**Theorem 2.6.1** (Egorov's theorem). Let  $f_n, f$  be measurable functions on  $(X, \mathcal{A}, \mu)$ . Suppose  $\mu(X) < \infty$ , then

$$f_n \rightarrow f \text{ a.e.} \Leftrightarrow f_n \rightarrow f \text{ almost uniformly.}$$

**Proof.** We prove two directions.

( $\Leftarrow$ )

( $\Rightarrow$ ) Fix  $\epsilon > 0$ . We see that

$$f_n \rightarrow f \text{ a.e.} \Rightarrow \mu \left( \bigcup_{k=1}^{\infty} \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} B_{n,k}^c \right) = 0 \Rightarrow \forall_k \mu \left( \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} B_{n,k}^c \right) = 0.$$

From continuity of measure from above and  $\mu(X) < \infty$ , we further have

$$\forall_k \lim_{N \rightarrow \infty} \mu \left( \bigcup_{n=N}^{\infty} B_{n,k}^c \right) = 0 \Rightarrow \forall_k \exists N_k \in \mathbb{N} \mu \left( \bigcup_{n=N_k}^{\infty} B_{n,k}^c \right) < \frac{\epsilon}{2^k}.$$

Now, let

$$F := \bigcup_{k=1}^{\infty} \bigcup_{n=N_k}^{\infty} B_{n,k}^c,$$

we see that  $\mu(F) < \epsilon$ , hence  $f_n \rightarrow f$  **uniformly**. ■

DIY

## Chapter 3

# Product Measure

### 3.1 Product $\sigma$ -Algebra

Before we start, we see the setup.

- Product space.

$$X = \prod_{\alpha \in I} X_{\alpha}$$

where  $x = (x_{\alpha})_{\alpha \in I} \in X$ .

- Coordinate map.

$$\pi_{\alpha}: X \rightarrow X_{\alpha}.$$

Now we see the formal definition.

**Definition 3.1.1** (Product  $\sigma$ -algebra). Let  $(X_{\alpha}, \mathcal{A}_{\alpha})$  be a measurable space for all  $\alpha \in I$ . Then a product  $\sigma$ -algebra on  $X = \prod_{\alpha \in I} X_{\alpha}$  is

$$\bigotimes_{\alpha \in I} \mathcal{A}_{\alpha} = \left\langle \bigcup_{\alpha \in I} \pi_{\alpha}^{-1}(\mathcal{A}_{\alpha}) \right\rangle,$$

where  $\pi_{\alpha}^{-1}(\mathcal{A}_{\alpha}) = \{\pi_{\alpha}^{-1}(E) \mid E \in \mathcal{A}_{\alpha}\}$ .

**Notation.** We denote  $I = \{1, \dots, d\} \Rightarrow X = \prod_{i=1}^d X_i, x = (x_1, \dots, x_d)$ . Also,

$$\bigotimes_{i=1}^d \mathcal{A}_i = \mathcal{A}_1 \otimes \dots \otimes \mathcal{A}_d.$$

**Lemma 3.1.1.** If  $I$  is countable, then

$$\bigotimes_{i=1}^{\infty} \mathcal{A}_i = \left\langle \left\{ \prod_{i=1}^{\infty} E_i \mid \forall_i E_i \in \mathcal{A}_i \right\} \right\rangle.$$

**Proof.** If  $E_i \in \mathcal{A}_i$ , then  $\pi_i^{-1}(E_i) = \prod_{j=1}^{\infty} E_j$ , where  $E_j = X_j$  for  $j \neq i$ . On the other hand, since

$$\prod_{i=1}^{\infty} E_i = \bigcap_{i=1}^{\infty} \pi_i^{-1}(E_i),$$

from Lemma 1.1.2, the result follows. ■

## Lecture 17: Product Measure

We now see a lemma.

14 Feb. 11:00

**Lemma 3.1.2.** Suppose  $\mathcal{A}_\alpha = \langle \mathcal{E}_\alpha \rangle$  for every  $\alpha \in I$ . Then

- (a)  $\pi_\alpha^{-1}(\mathcal{A}_\alpha) = \langle \pi_\alpha^{-1}(\mathcal{E}_\alpha) \rangle$
- (b)  $\bigotimes_\alpha \mathcal{A}_\alpha = \langle \bigcup_\alpha \pi_\alpha^{-1}(\mathcal{E}_\alpha) \rangle$
- (c) If  $I$  is countable, then

$$\bigotimes_{i=1}^{\infty} \mathcal{A}_i = \left\langle \left\{ \prod_{i=1}^{\infty} E_i \mid \forall_i E_i \in \mathcal{E}_i \right\} \right\rangle$$

**Proof.** We prove this one by one.

- (a) Note that for  $f: Y \rightarrow Z$ , and  $\mathcal{B}$  be a  $\sigma$ -algebra on  $Z$ , then  $f^{-1}(\mathcal{B})$  is also a  $\sigma$ -algebra since  $f^{-1}(\emptyset) = \emptyset$ ,  $f^{-1}(B)^c = f^{-1}(B^c)$ , and  $\bigcup_n f^{-1}(B_n) = f^{-1}(\bigcup_n B_n)$ . Hence,  $\pi_\alpha^{-1}$  is a  $\sigma$ -algebra on  $X$ , i.e.,

$$\pi_\alpha^{-1}(\mathcal{E}_\alpha) \subset \pi_\alpha^{-1}(\mathcal{A}_\alpha) \stackrel{!}{\Rightarrow} \langle \pi_\alpha^{-1}(\mathcal{E}_\alpha) \rangle \subset \pi_\alpha^{-1}(\mathcal{A}_\alpha).$$

To show the other direction, let  $\mathcal{M}$  being

$$\mathcal{M} = \{B \subset X_\alpha \mid \pi_\alpha^{-1}(B) \in \langle \pi_\alpha^{-1}(\mathcal{E}_\alpha) \rangle\}.$$

We now check

- $\mathcal{M}$  is a  $\sigma$ -algebra.
- $\mathcal{E}_\alpha \subset \mathcal{M}$ . This is true by definition of  $\mathcal{M}$ .

Thus,  $\langle \mathcal{E}_\alpha \rangle = \mathcal{A}_\alpha \subset \mathcal{M}$ . Hence, if  $E \in \mathcal{A}_\alpha$ ,  $E \in \mathcal{M}$ , implying

$$\pi_\alpha^{-1}(E) \in \langle \pi_\alpha^{-1}(\mathcal{E}_\alpha) \rangle,$$

i.e.,  $\mathcal{A}_\alpha \subset \langle \pi_\alpha^{-1}(\mathcal{E}_\alpha) \rangle$ .

(b)

(c)

Check  
(Easy)!

DIY

DIY

■

**Theorem 3.1.1.** Suppose  $X_1, \dots, X_d$  are metric spaces. Let  $X = \prod_{i=1}^d X_i$  with product metric defined as

$$\rho(x, y) = \sum_{i=1}^d \rho_i(x_i, y_i).$$

Then,

- (a)  $\bigotimes_{i=1}^d \mathcal{B}(X_i) \subset \mathcal{B}(X)$
- (b) If in addition, each  $X_i$  has a countable dense subset,

$$\bigoplus_{i=1}^d \mathcal{B}(X_i) = \mathcal{B}(X).$$

**Proof.**

DIY

■

**Remark.** We see that

- $\mathcal{B}(\mathbb{R}^d) = \mathcal{B}(\mathbb{R}) \otimes \cdots \otimes \mathcal{B}(\mathbb{R})$
- let  $f = u + iv: X \rightarrow \mathbb{C}$ , and  $\mathcal{A}$  be a  $\sigma$ -algebra on  $X$ . Then

$$\bigvee_{E \in \mathcal{B}(\mathbb{R})} u^{-1}(E), v^{-1}(E) \in \mathcal{A} \Leftrightarrow f^{-1}(F) \in \mathcal{A}, \forall F \in \mathcal{B}(\mathbb{C})$$

with  $\mathcal{B}(\mathbb{C}) = \mathcal{B}(\mathbb{R}^2) = \mathcal{B}(\mathbb{R}) \otimes \mathcal{B}(\mathbb{R})$ .

We first focus on 2 dimensional case. Specifically, we think of our coordinate is  $x$  and  $y$  on  $\mathbb{R}^2$ .

**Definition.** Let  $X, Y$  be two sets, then we have the following.

**Definition 3.1.2** (*x-section, y-section for set*). For  $E \subset X \times Y$ ,

$$E_x = \{y \in Y \mid (x, y) \in E\}, \quad E^y = \{x \in X \mid (x, y) \in E\},$$

where  $E_x$  is called the *x-section* of  $E$ , while  $E_y$  is called the *y-section* of  $E$ .

**Definition 3.1.3** (*x-section, y-section for function*). For  $f: X \times Y \rightarrow \mathbb{C}$ , define  $f_x: Y \rightarrow \mathbb{C}$ ,  $f^y: X \rightarrow \mathbb{C}$  by

$$f_x(y) = f(x, y) = f^y(x),$$

where  $f_x(y)$  is called the *x-section* of  $f$ , while  $f_y(x)$  is called the *y-section* of  $f$ .

**Example.** We see that

$$(\mathbb{1}_E)_x = \mathbb{1}_{E_x}$$

and

$$(\mathbb{1}_E)^y = \mathbb{1}_{E^y}.$$

**Proposition 3.1.1.** Given two measurable spaces  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$ , then

- (a) If  $E \in \mathcal{A} \otimes \mathcal{B}$ , then

$$\bigvee_{x \in X} \bigvee_{y \in Y} E_x \in \mathcal{B}, E^y \in \mathcal{A}.$$

- (b) If  $f: X \times Y \rightarrow \mathbb{C}$  is  $\mathcal{A} \otimes \mathcal{B}$ -measurable, then

$$\bigvee_{x \in X} \bigvee_{y \in Y} f_x \text{ is } \mathcal{B}\text{-measurable, } f^y \text{ is } \mathcal{A}\text{-measurable.}$$

**Proof.** We prove this one by one.

- (a) Let  $\mathcal{F} := \left\{ E \subset X \times Y \mid \bigvee_{x \in X} \bigvee_{y \in Y} E_x \in \mathcal{B}, E^y \in \mathcal{A} \right\}$ , then

- $\mathcal{F}$  is a  $\sigma$ -algebra.
  - $\emptyset \in \mathcal{F}$ .
  - $(E^c)_x = E_x^c$ .
  - $\left( \bigcup_{j=1}^{\infty} E_j \right)_x = \bigcup_{j=1}^{\infty} (E_j)_x$ .

And the same is true for  $y$ .

- Let  $\mathcal{R}_0 := \{A \times B \mid A \in \mathcal{A}, B \in \mathcal{B}\} \subset \mathcal{F}$ , which is again easy to show from definition.

Hence, we see that  $\langle \mathcal{R}_0 \rangle = \mathcal{A} \otimes \mathcal{B} \subset \mathcal{F}$ .

(b) Since

$$(f_x)^{-1}(B) = (f^{-1}(B))_x$$

and

$$(f^y)^{-1}(B) = (f^{-1}(B))^y,$$

the result follows from (a). ■

## 3.2 Product Measures

We start with the definition.

**Definition 3.2.1 (Rectangle).** Given two measurable spaces  $(X, \mathcal{A}), (Y, \mathcal{B})$ , a (measurable) rectangle is  $R = A \times B$  where  $A \in \mathcal{A}$  and  $B \in \mathcal{B}$ . Furthermore, we let

$$\mathcal{R}_0 := \{R = A \times B \mid A \in \mathcal{A}, B \in \mathcal{B}\},$$

and

$$\mathcal{R} := \left\{ \bigcup_{i=1}^N R_i \mid N \in \mathbb{N}, R_1, \dots, R_N \text{ disjoint rectangles} \right\}.$$

**Note.** Whenever we're talking about rectangle, they're always measurable.

**Lemma 3.2.1.**  $\mathcal{R}$  is an algebra, and

$$\langle \mathcal{R}_0 \rangle = \langle \mathcal{R} \rangle = \mathcal{A} \otimes \mathcal{B}.$$

**Proof.** Simply observe that

$$(A \times B)^c = (A^c \times Y) \cup (X \times B^c).$$

DIY

## Lecture 18: Monotone Class

Let's start with a theorem.

16 Feb. 11:00

**Theorem 3.2.1.** Let  $(X, \mathcal{A}, \mu), (Y, \mathcal{B}, \nu)$  be measure spaces. Then

(a) There is a measure  $\mu \times \nu$  on  $\mathcal{A} \otimes \mathcal{B}$  satisfying

$$(\mu \times \nu)(A \times B) = \mu(A)\nu(B)$$

for every  $A \in \mathcal{A}, B \in \mathcal{B}$ .

(b) If  $\mu, \nu$  are  $\sigma$ -finite, then  $\mu \times \nu$  is unique.

**Proof.** We prove this one by one.

(a) Define  $\pi: \mathcal{R} \rightarrow [0, \infty]$  by  $\pi(A \times B) = \mu(A)\nu(B)$ , and extending linearly, we have

$$\pi(A \times B) = \mu(A)\nu(B),$$

hence

$$\pi\left(\bigcup_{i=1}^N A_i \times B_i\right) = \sum_{i=1}^n \pi(A_i \times B_i).$$

We claim that  $\pi$  is a **pre-measure**. To show this, it's enough to check that  $\pi(A \times B) = \sum_{n=1}^{\infty} \pi(A_n \times B_n)$  if  $A \times B = \coprod_n A_n \times B_n$ . Since  $A_n \times B_n$  are disjoint, so

$$\mathbb{1}_{A \times B}(x, y) = \sum_{n=1}^{\infty} \mathbb{1}_{A_n \times B_n}(x, y).$$

Thus,

$$\mathbb{1}_A(x) \mathbb{1}_B(y) = \sum_{n=1}^{\infty} \mathbb{1}_{A_n}(x) \mathbb{1}_{B_n}(y).$$

Integrating with respect to  $x$ , and applying [Theorem 1.3.1](#), we have

$$\int_X \mathbb{1}_A(x) \mathbb{1}_B(y) d\mu(x) = \sum_{n=1}^{\infty} \int_X \mathbb{1}_{A_n}(x) \mathbb{1}_{B_n}(y) d\mu(x),$$

which implies

$$\mu(A) \mathbb{1}_B(y) = \sum_{n=1}^{\infty} \mu(A_n) \mathbb{1}_{B_n}(y)$$

for every  $y$ . We can then integrate again with respect to  $y$  and apply [Theorem 1.3.1](#), we have

$$\int_Y \mu(A) \mathbb{1}_B(y) d\nu(y) = \sum_{n=1}^{\infty} \int_Y \mu(A_n) \mathbb{1}_{B_n}(y) d\nu(y),$$

which gives us

$$\mu(A) \nu(B) = \sum_{n=1}^{\infty} \mu(A_n) \nu(B_n).$$

Hence, we see that  $\mu$  is indeed a **pre-measure**, so [Theorem 1.4.1](#) gives  $\mu \times \nu$  on  $\langle \mathcal{R} \rangle = \mathcal{A} \otimes \mathcal{B}$  extending  $\pi$  on  $\mathcal{R}$ .

- (b) If  $\mu, \nu$  are  **$\sigma$ -finite**, then  $\pi$  is  **$\sigma$ -finite** on  $\mathcal{R}$ , then [Theorem 1.4.2](#) applies. Moreover, we have that

$$(\mu \times \nu)(E) = \inf \left\{ \sum_{i=1}^{\infty} \mu(A_i) \nu(B_i) \mid E \subset \bigcup_{i=1}^{\infty} A_i \times B_i, A_i \in \mathcal{A}, B_i \in \mathcal{B} \right\}.$$

■

### 3.3 Monotone Class Lemma

Let's start with a definition.

**Definition 3.3.1 (Monotone Class).** If  $X$  is a set, and  $C \subset \mathcal{P}(X)$ , we say that  $C$  is a *monotone class* on  $X$  if

- $C$  is closed under countable increasing unions.
- $C$  is closed under countable decreasing intersections.

**Example.** Every  **$\sigma$ -algebra** is a **monotone class**.

**Example.** If  $C_\alpha$  are (arbitrarily many) **monotone classes** on a set  $X$ , then  $\bigcap_\alpha C_\alpha$  is a **monotone class**. Furthermore, if  $\mathcal{E} \subset \mathcal{P}(X)$ , there is a unique smallest **monotone class** containing  $\mathcal{E}$ , denoted by  $\langle \mathcal{E} \rangle$ , which follows the same idea as in [Definition 1.1.3](#).



**Theorem 3.3.1 (Monotone Class Lemma).** Suppose  $\mathcal{A}_0$  is an algebra on  $X$ . Then  $\langle \mathcal{A}_0 \rangle^a$  is the monotone class generated by  $\mathcal{A}_0$ .

<sup>a</sup> $\langle \mathcal{A}_0 \rangle$  is the  $\sigma$ -algebra generated by  $\mathcal{A}_0$  by Definition 1.1.3.

**Proof.** Let  $\mathcal{A} = \langle \mathcal{A}_0 \rangle$  and let  $\mathcal{C}$  be the monotone class generated by  $\mathcal{A}_0$ . Since  $\mathcal{A}$  is a  $\sigma$ -algebra, it's a monotone class. Note that it contains  $\mathcal{A}_0$ , hence  $\mathcal{A} \supset \mathcal{C}$ .

To show  $\mathcal{C} \supset \mathcal{A}$ , it's enough to show that  $\mathcal{C}$  is a  $\sigma$ -algebra. We check that

1.  $\emptyset \in \mathcal{A}_0 \subseteq \mathcal{C}$ .
2. Let  $\mathcal{C}' = \{E \subset X \mid E^c \in \mathcal{C}\}$ .
  - $\mathcal{C}'$  is a monotone class.
  - $\mathcal{A}_0 \subset \mathcal{C}'$  because if  $E \in \mathcal{A}_0$ , then  $E^c \in \mathcal{A}_0$ , so  $E^c \in \mathcal{C}$ , thus  $E \in \mathcal{C}'$ .

We see that  $\mathcal{C}' \subset \mathcal{C}'$ , so  $\mathcal{C}$  is closed under complements.

3. For  $E \subset X$ , let  $\mathcal{D}(E) = \{F \in \mathcal{C} \mid E \cup F \in \mathcal{C}\}$ .
  - $\mathcal{D}(E) \subset \mathcal{C}$ .
  - $\mathcal{D}(E)$  is a monotone class.
  - If  $E \in \mathcal{A}_0$ , then  $\mathcal{A}_0 \subset \mathcal{D}(E)$ . We see this by picking  $F \in \mathcal{A}_0$ , then  $E \cup F \in \mathcal{A}_0 \subset \mathcal{C}$ .

Hence,  $\mathcal{C} = \mathcal{D}(E)$  if  $E \in \mathcal{A}_0$ .

4. Let  $\mathcal{D} = \{E \in \mathcal{C} \mid \mathcal{D}(E) = \mathcal{C}\}$ . That is  $\mathcal{D} = \{E \in \mathcal{C} \mid E \cup F, \forall F \in \mathcal{C}\}$ . Then we have
  - $\mathcal{A}_0 \subset \mathcal{D}$  by 3.
  - $\mathcal{D}$  is a monotone class.
  - $\mathcal{D} \subset \mathcal{C}$  by definition.

Thus,  $\mathcal{D} = \mathcal{C}$ , so if  $E, F \in \mathcal{C}$ , then  $E \cup F \in \mathcal{C}$ . This implies that  $\mathcal{C}$  is closed under finite unions.

5. Now to show that  $\mathcal{C}$  is closed under countable unions, let  $E_1, E_2, \dots \in \mathcal{C}$ . We may then define

$$F_N = \bigcup_{n=1}^N E_n \in \mathcal{C}.$$

Then we see that  $F_1 \subset F_2 \subset \dots$ , hence  $\bigcup_N F_N \in \mathcal{C}$ . But this simply implies

$$\bigcup_N F_N = \bigcup_n E_n,$$

so we're done. ■

## Lecture 19: Fubini-Tonelli's Theorem

**As previously seen.** If  $E \in \mathcal{A} \otimes \mathcal{B} \Rightarrow E_x \in \mathcal{B}, E^y \in \mathcal{A} \forall x \in X, \forall y \in Y$ . Note that the reverse is not true.

18 Feb. 11:00

### 3.4 Fubini-Tonelli Theorem

We start with a theorem.

**Theorem 3.4.1** (Tonelli's theorem for characteristic functions). Given  $(X, \mathcal{A}, \mu)$  and  $(Y, \mathcal{B}, \nu)$  be  $\sigma$ -finite measure space. Suppose  $E \in \mathcal{A} \otimes \mathcal{B}$ , then

- (a)  $\alpha(x) := \nu(E_x): X \rightarrow [0, \infty]$  is a  $\mathcal{A}$ -measurable function.
- (b)  $\beta(y) := \mu(E^y): Y \rightarrow [0, \infty]$  is a  $\mathcal{B}$ -measurable function.
- (c)  $(\mu \times \nu)(E) = \int_X \nu(E_x) d\mu(x) = \int_Y \mu(E^y) d\nu(y)$ .

**Proof.** We prove this in two cases.

1. Assume that  $\mu, \nu$  are finite measures. Let

$$C := \{E \in \mathcal{A} \otimes \mathcal{B} \mid \text{Conditions (a), (b), (c) hold}\}.$$

It's enough to prove that  $\langle \mathcal{R} \rangle = \mathcal{A} \otimes \mathcal{B} \subset C$ . We further observe that from the Theorem 3.3.1 and the fact that  $\mathcal{R}$  is an algebra, it's also enough to show that

- $\mathcal{R} \subset C$ .
- $C$  is a monotone class.

From condition (a),

$$\alpha(x) = \nu((A \times B)_x) = \begin{cases} \nu(B), & \text{if } x \in A; \\ 0, & \text{if } x \notin A \end{cases} = \nu(B) \mathbb{1}_A.$$

And from condition (b),

$$(\mu \times \nu)(A \times B) = \mu(A)\nu(B)$$

and

$$\int_X \nu((A \times B)_x) d\mu(x) = \nu(B)\mu(A).$$

Let  $E_n \in C$ ,  $E_1 \subset E_2 \subset \dots$ . We need to show  $E = \bigcup_{n=1}^{\infty} E_n \in C$ . We now see that

$$E_x = \bigcup_{n=1}^{\infty} (E_n)_x, (E_1)_x \subset (E_2)_x \subset \dots \Rightarrow \alpha(x) = \nu(E_x) \stackrel{!}{=} \lim_{n \rightarrow \infty} \nu((E_n)_x) \quad \forall x \in X.$$

This implies that (a) is proved. For (c), we see that

$$(\mu \times \nu)(E) \stackrel{!}{=} \lim_{n \rightarrow \infty} (\mu \times \nu)(E_n) = \lim_{n \rightarrow \infty} \int_X \nu((E_n)_x) d\mu(x) \stackrel{!}{=} \int_X \nu(E_x) d\mu(x).$$

Now let  $F_n \in C$ ,  $F_1 \supset F_2 \supset \dots$ . We need to show that  $F = \bigcap_{n=1}^{\infty} F_n \in C$ . Instead of using Theorem 2.2.1, we now want to use Theorem 2.3.1, which is applicable since  $\mu(X), \nu(Y) < \infty$  by assumption.

2. Assume  $\mu$  and  $\nu$  are  $\sigma$ -finite measures. We then have a sequence  $\{X_n \times Y_n\}$  of rectangles of with only finite measure. Now, just consider if  $E \in \mathcal{A} \otimes \mathcal{B}$ , (a) applies to  $E \cap (X_n \times Y_n)$  for each  $n$ , with

$$X \times Y = \bigcup_{n=1}^{\infty} (X_n \times Y_n), \begin{cases} X_1 \subset X_2 \subset \dots, & \mu(X_k) < \infty \\ Y_1 \subset Y_2 \subset \dots, & \nu(Y_k) < \infty, \end{cases}$$

we have

$$\mu \times \nu(E \cap (X_n \times Y_n)) = \int \mathbb{1}_{X_n}(x) \cdot \nu(E_x \cap Y_n) d\mu(x) = \int \mathbb{1}_{Y_n}(y) \mu(E^y \cap X_n) d\nu(y).$$

By applying Theorem 2.2.1, the result follows. ■

**Theorem 3.4.2** (Fubini-Tonelli's theorem). Given two  $\sigma$ -finite measure space  $(X, \mathcal{A}, \mu), (Y, \mathcal{B}, \nu)$ , we have the following.

**Theorem** (Tonelli's theorem). If  $f: X \times Y \rightarrow [0, \infty]$  is  $\mathcal{A} \otimes \mathcal{B}$ -measurable, then

- (a)  $g(x) := \int_Y f(x, y) d\nu(y), X \rightarrow [0, \infty]$  is a  $\mathcal{A}$ -measurable function.
- (b)  $h(x) := \int_X f(x, y) d\mu(x), Y \rightarrow [0, \infty]$  is a  $\mathcal{B}$ -measurable function.
- (c) We have

$$\int_{X \times Y} f d(\mu \times \nu) = \int_X \left( \int_Y f(x, y) d\nu(y) \right) d\mu(x) = \int_Y \left( \int_X f(x, y) d\mu(x) \right) d\nu(y).$$

**Theorem** (Fubini's theorem). If  $f \in L^1(X \times Y, \mu \times \nu)$ , then

- (a)  $f_x \in L^1(Y, \nu)$  for  $\mu$ -a.e.  $x$ , and  $g(x) \in L^1(X, \mu)$  defined  $\mu$ -a.e.
- (b)  $f_y \in L^1(X, \mu)$  for  $\nu$ -a.e.  $y$ , and  $h(y) \in L^1(Y, \nu)$  defined  $\nu$ -a.e.
- (c) The iterated integral formulas hold. Namely, we have

$$\int_{X \times Y} f d(\mu \times \nu) = \int_X \left( \int_Y f(x, y) d\nu(y) \right) d\mu(x) = \int_Y \left( \int_X f(x, y) d\mu(x) \right) d\nu(y).$$

**Proof.** Read [FF99]. ■

**Remark.** The Fubini and Tonelli's theorem are frequently used in tandem. Say that if one want to reverse the order of integration in a double integral  $\iint f d\mu d\nu$ . We first verify that  $\int |f| d(\mu \times \nu) < \infty$  by using Tonelli's theorem to evaluate this integral as an iterated integral. Then, we apply Fubini theorem to conclude that

$$\iint f d\mu d\nu = \iint f d\nu d\mu.$$

## Lecture 20: Lebesgue Measure on $\mathbb{R}^d$

### 3.5 Lebesgue Measure on $\mathbb{R}^d$

21 Feb. 11:00

**Example.**  $(\mathbb{R}^2, \mathcal{L} \otimes \mathcal{L}, m \times m)$  is not complete.

**Proof.** We see that

- Let  $A \in \mathcal{L}, A \neq \emptyset, m(A) = 0$ .
- Let  $B \subset [0, 1], B \notin \mathcal{L}$  (Vital set for example).
- Let  $E = A \times B, F = A \times [0, 1]$ .

We see that  $E \subset F, F \in \mathcal{L} \otimes \mathcal{L}, (m \times m)(F) = m(A)m([0, 1]) = 0$ , i.e.,  $F$  is a null set. But  $E$  is not  $\mathcal{L} \otimes \mathcal{L}$ -measurable since otherwise, its sections are all measurable. ⊛

**Definition 3.5.1.** Let  $(\mathbb{R}^d, \mathcal{L}^d, m^d)$  be the completion of

$$(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d), m \times \cdots \times m),$$

which is same as the *completion* of

$$(\mathbb{R}^d, \mathcal{L} \otimes \cdots \otimes \mathcal{L}, m \times \cdots \times m).$$

**Remark.** We see that

$$\mathcal{L}^d \supsetneq \mathcal{L} \otimes \cdots \otimes \mathcal{L} = \left\langle \left\{ \prod_{i=1}^d E_i \mid E_i \in \mathcal{L} \right\} \right\rangle.$$

**Definition 3.5.2** (Rectangle in  $\mathbb{R}^d$ ). A *rectangle* in  $\mathbb{R}^d$  is  $R = \prod_{i=1}^d E_i$  where  $E_i \in \mathcal{B}(\mathbb{R})$ .

**Definition 3.5.3** (Lebesgue measure in  $\mathbb{R}^d$ ). We let the *Lebesgue measure* in  $\mathbb{R}^d$ , denoted as  $m^d$ , defined as

$$m^d(E) := \inf \left\{ \sum_{k=1}^{\infty} m^d(R_k) \mid E \subset \bigcup_{k=1}^{\infty} R_k, R_k \text{ is rectangles} \right\}.$$

**Theorem 3.5.1.** Let  $E \subset \mathcal{L}^d$ . Then

- (a)  $m^d(E) = \inf \{m^d(O) \mid \text{open } O \supset E\} = \sup \{m^d(K) \mid \text{compact } K \subset E\}$ .
- (b)  $E = A_1 \cup N_1 = A_2 \setminus N_2$ , where  $A_1$  is  $F_\sigma$ ,  $A_2$  is  $G_\delta$ , and  $N_i$  are null.
- (c) If  $m^d(E) < \infty$ ,  $\forall \epsilon > 0$ ,  $\exists R_1, \dots, R_m$  rectangles whose sides are intervals such that

$$m^d \left( E \Delta \left( \bigcup_{i=1}^m R_i \right) \right) < \epsilon.$$

**Proof.** Similar to  $d = 1$  case. ■

**Theorem 3.5.2.** Integrable step functions and  $C_c(\mathbb{R}^d)$ , the collection of continuous functions, are dense in  $L^1(\mathbb{R}^d, \mathcal{L}^d, m^d)$

**Proof.** See [FF99]. ■

**Theorem 3.5.3.** Lebesgue measure in  $\mathbb{R}^d$  is translation-invariant.

**Proof.** See [FF99]. ■

**Theorem 3.5.4** (Effect of linear transformation on Lebesgue measure). If  $T \in \text{GL}(\mathbb{R}^d)$ ,  $e \in \mathcal{L}^d$ , then  $T(E)$  is measurable and

$$m(T(E)) = |\det T| \cdot m(E).$$

**Proof.** See [FF99]. ■

## Chapter 4

# Differentiation on Euclidean Space

**As previously seen.** Given  $f: [a, b] \rightarrow \mathbb{R}$ , there are two versions of **fundamental theorem of calculus**:

(a)

$$\int_a^b f'(x) dx = f(b) - f(a).$$

(b)

$$\frac{d}{dx} \int_a^x f(t) dt = f(x),$$

which follows from

$$\lim_{r \rightarrow 0^+} \frac{1}{r} \int_x^{x+r} f(t) dt = f(x) = \lim_{r \rightarrow 0^+} \frac{1}{r} \int_{x-r}^x f(t) dt.$$

**Remark.** We see that

$$\lim_{r \rightarrow 0^+} \frac{1}{r} \int_x^{x+r} (f(t) - f(x)) dt = 0 = \lim_{r \rightarrow 0^+} \frac{1}{r} \int_{x-r}^x (f(t) - f(x)) dt,$$

where we have

$$f(x) = \frac{1}{r} \int_x^{x+r} f(t) dt.$$

This generalized to  $f: \mathbb{R}^d \rightarrow \mathbb{R}$ , namely

$$\lim_{r \rightarrow 0^+} \frac{1}{\text{vol}(B(x, r))} \int_{B(x, r)} (f(t) - f(x)) \underbrace{dt}_{\mathbb{R}^d} \stackrel{?}{=} 0.$$

### 4.1 Hardy-Littlewood Maximal Function

We first see our notation.

**Notation.** Given a(n) (open) ball in  $\mathbb{R}^d$ ,  $B = B(a, r)$ , denote  $cB = B(a, cr)$  for  $c > 0$ .

**Lemma 4.1.1 (Vitali-type covering lemma).** Let  $B_1, \dots, B_k$  be a finite collection of open balls in  $\mathbb{R}^d$ . Then there exists a sub-collection  $B'_1, \dots, B'_m$  of disjoint open balls such that

$$\bigcup_{i=1}^m (3B'_i) \supset \bigcup_{i=1}^k B_i.$$

**Proof.** Greedy Algorithm. ■

## Lecture 21: Hardy-Littlewood Maximal Function and Inequality

**Notation.** We let

$$\int_E f \, dm = \int_E f(x) \, dx.$$

25 Feb. 11:00

The problem we're working on is

$$\frac{1}{m(B(w, r))} \int_{B(w, r)} f(y) \, dy \xrightarrow[?]{r \rightarrow 0} f(x).$$

**Definition 4.1.1 (Locally integrable).** Given  $f: \mathbb{R}^d \rightarrow \mathbb{C}$  be [Lebesgue measurable](#) function. Then we say  $f$  is *locally integrable* if for every compact  $K \subset \mathbb{R}^d$ ,

$$\int_K |f| \, dm < \infty.$$

In this case, we write  $f \in L^1_{\text{loc}}(\mathbb{R}^d)$ .

**Definition 4.1.2 (Hardy-Littlewood maximal function).** Given  $f \in L^1_{\text{loc}}(\mathbb{R}^d)$ , the *Hardy-Littlewood maximal function* for  $f$  is defined as

$$Hf(x) := \sup \{A_r(x) \mid r > 0\},$$

where

$$A_r(x) := \frac{1}{m(B(x, r))} \int_{B(x, r)} |f(y)| \, dy.$$

**Note.** We note that  $A_r(\cdot)$  means *averaging function* over an open ball with radius being  $r$ .

**Lemma 4.1.2.** Let  $f \in L^1_{\text{loc}}(\mathbb{R}^d)$ , then

- (a)  $A_r(x)$  is jointly continuous for  $(x, r) \in \mathbb{R}^d \times (0, \infty)$ .
- (b)  $Hf(x)$  is [Borel measurable](#).

**Proof.** We outline the proof.

- (a) Let  $(x, r) \rightarrow (x^*, r^*) \Rightarrow A_r(x) \rightarrow A_{r^*}(x^*)$ . Let  $(x_n, r_n)$  be any sequence which converges to  $x^*, r^*$ , then we consider  $\lim_{n \rightarrow \infty} A_{r_n}(x_n)$  and we can calculate

$$\int \underbrace{|f(y)| \mathbb{1}_{B(x_n, r_n)}(y)}_{:= h_n(y)},$$

then we apply [Theorem 2.3.1](#) to  $h_n$ .

- (b) Observe that

$$(Hf)^{-1}(\underbrace{(a, \infty)}_{\text{open}}) = \bigcup_{r>0} A_r^{-1}((a, \infty))$$

is open, since  $A_r^{-1}((a, \infty))$  is open from the 1. Note that the equality comes from the fact that  $Hf = \sup_r A_r$ .

**Theorem 4.1.1** (Hardy-Littlewood maximal inequality). There exists  $C_d > 0$  such that for every  $f \in L^1(\mathbb{R}^d)$ ,

$$\forall_{\alpha > 0} m(\{x \in \mathbb{R}^d \mid Hf(x) > \alpha\}) \leq \frac{C_d}{\alpha} \int |f(x)| \, dx.$$

**Proof.** We first fix  $f \in L^1$  and  $\alpha > 0$ . We define

$$E := \{x \mid Hf(x) > \alpha\},$$

which is a Borel measurable set by Lemma 4.1.2. Then

$$x \in E \Rightarrow \exists_{r_x > 0} A_{r_x}(x) > \alpha \Rightarrow m(B(x, r_x)) < \frac{1}{\alpha} \int_{B(x, r_x)} |f(y)| \, dy.$$

From inner regularity, we have

$$m(E) = \sup \{m(K) \mid \text{compact } K \subset E\}.$$

Let  $K \subset E$  be compact, then

$$K \subset \bigcup_{x \in K} B(x, r_x) \stackrel{K \text{ compact}}{\Rightarrow} K \subset \bigcup_{i=1}^N B_i \stackrel{!}{\Rightarrow} K \subset \bigcup_{i=1}^m \{3B'_j\}.$$

From here, we further have

$$m(K) \leq \sum_{i=1}^m m(3B'_j) = 3^d \sum_{j=1}^m m(B'_j) \leq \frac{3^d}{\alpha} \sum_{j=1}^m \int_{B'_j} |f(y)| \, dy.$$

Now, since  $B'_1, \dots, B'_m$  are disjoint, hence we finally have

$$m(K) \leq \frac{3^d}{\alpha} \int_{\mathbb{R}^d} |f(y)| \, dy.$$

## Lecture 22: Lebesgue Differentiation Theorem

We should compare the Hardy-Littlewood maximal inequality to Markov's inequality. Namely, there exists  $C_d > 0$  (can take  $3^d$ ) such that for all  $f \in L^1(\mathbb{R}^d)$ ,  $\alpha > 0$ , we have

$$\begin{cases} m(\{x \mid Hf(x) > \alpha\}) \leq \frac{C_d}{\alpha} \int |f|; \\ m(\{x \mid |f(x)| > \alpha\}) \leq \frac{1}{\alpha} \int |f|. \end{cases}$$

### 4.2 Lebesgue Differentiation Theorem

We start with a theorem!

**Theorem 4.2.1** (Lebesgue differentiation theorem). Let  $f \in L^1$ , then

$$\lim_{r \rightarrow 0} \frac{1}{m(B(x, r))} \int_{B(x, r)} |f(y) - f(x)| \, dy = 0$$

for a.e.  $x$ .

**Proof.** The result holds for  $f \in C_c(\mathbb{R}^d)$ , namely for those continuous functions with **compact support**. This is because for any  $\epsilon > 0$ , if  $r$  is small and  $|f(y) - f(x)| < \epsilon$ , then

$$\frac{1}{m(B(x, r))} \int_{B(x, r)} |f(y) - f(x)| \, dy < \epsilon.$$

Now, let  $f \in L^1(\mathbb{R}^d)$  and fix  $\epsilon > 0$ . By density, there exists  $g \in C_c(\mathbb{R}^d)$  with  $\|f - g\|_1 < \epsilon$ . We then have

$$\int_{B_r(x)} |f(y) - f(x)| \, dy \leq \int_{B_r(x)} |f(y) - g(y)| \, dy + \int_{B_r(x)} |g(y) - g(x)| \, dy + \int_{B_r(x)} |g(x) - f(x)| \, dy.$$

**Note.** We use  $B_r(x)$  above to denote  $B(x, r)$  for spacing reason only. Nothing tricky here.

Divide all of these by  $m(B(x, r))$ , and take  $\limsup_{r \rightarrow \infty}$ , we need to understand the error terms, namely

$$\frac{1}{m(B(x, r))} \int_{B(x, r)} |f(x) - g(x)| \, dy = |g(x) - f(x)|$$

and

$$\frac{1}{m(B(x, r))} \int_{B(x, r)} |f(y) - g(y)| \, dy \leq (H(f - g))(x).$$

We define

$$Q(x) := \limsup_{r \rightarrow \infty} \frac{1}{m(B(x, r))} \int_{B(x, r)} |f(y) - f(x)| \, dy.$$

We want to show  $m(\{x \in X \mid Q(x) > 0\}) = 0$ . Let  $E_\alpha = \{x \in X \mid Q(x) > \alpha\}$ . It is enough to show  $m(E_\alpha) = 0$  for all  $\alpha > 0$  because  $\{x \in X \mid Q(x) > 0\} = \bigcup_n E_{\frac{1}{n}}$ . We know by the above that

$$Q(x) \leq (H(f - g))(x) + 0 + |g(x) - f(x)|.$$

Therefore,

$$E_\alpha \subset \{x \in X \mid (H(f - g))(x) > \alpha/2\} \cup \{x \in X \mid |g(x) - f(x)| > \alpha/2\}.$$

By the **Hardy-Littlewood maximal inequality** and **Markov's inequality**, we have

$$\begin{cases} m(\{x \mid (H(f - g))(x) > \alpha/2\}) \leq \frac{2C_d}{\alpha} \int |f - g|; \\ m(\{x \mid |g(x) - f(x)| > \alpha/2\}) \leq \frac{2}{\alpha} \int |f - g|. \end{cases}$$

Thus,

$$0 \leq m(E_\alpha) \leq \frac{2C_d}{\alpha} \|f - g\|_1 + \frac{2}{\alpha} \|f - g\|_1 \leq \frac{2(C_d + 1)}{\alpha} \epsilon.$$

Taking  $\epsilon \rightarrow 0$ ,  $m(E_\alpha)$  does not depend on  $\epsilon$  and  $g$ , hence  $m(E_\alpha) = 0$ . ■

**Corollary 4.2.1.** **Theorem 4.2.1** also holds for  $f \in L^1_{\text{loc}}(\mathbb{R}^d)$ .

**Proof.** Using the fact that  $m^d$  is  $\sigma$ -finite, and apply **Theorem 4.2.1**. Specifically, partition  $\mathbb{R}^d$  into countably many compact sets  $K_i$  and apply **Theorem 4.2.1** to  $f \mathbb{1}_{K_i}$  for all  $i$ . ■

**Corollary 4.2.2.** For  $f \in L^1_{\text{loc}}$ , we have

$$\lim_{r \rightarrow 0} \frac{1}{m(B(x, r))} \int_{B(x, r)} f(y) \, dy = f(x)$$

for **a.e.**  $x$ .



**Proof.** Use that

$$f(x) = \frac{1}{m(B(x, r))} \int_{B(x, r)} f(y) \, dy$$

and the triangle inequality. ■

DIY

**Definition 4.2.1 (Lebesgue point).** Let  $f \in L^1_{\text{loc}}(\mathbb{R}^d)$ , the point  $x \in \mathbb{R}^d$  is called a *Lebesgue point* of  $f$  if

$$\lim_{r \rightarrow 0} \frac{1}{m(B(x, r))} \int_{B(x, r)} |f(y) - f(x)| \, dy = 0.$$

**Remark.** Corollary 4.2.1 tells us that almost all points in  $\mathbb{R}^d$  are *Lebesgue points* for  $f$ .

**Definition 4.2.2 (Shrink nicely).** We say that  $\{E_r\}_{r>0}$  *shrinks nicely* to  $x$  as  $r \rightarrow 0$  if  $E_r \subset B(x, r)$  and

$$\exists_{c>0} c \cdot m(B(x, r)) \leq m(E_r).$$

**Note.** Here,  $c$  is independent of  $r$ .

**Corollary 4.2.3.** Suppose  $E_r$  *shrink nicely* to 0, and  $f \in L^1_{\text{loc}}(\mathbb{R}^d)$ , and  $x$  is a *Lebesgue point*. Then

$$\lim_{r \rightarrow 0} \frac{1}{m(E_r)} \int_{E_r} |f(y) - f(x)| \, dy = 0 \text{ and } \lim_{r \rightarrow 0} \frac{1}{m(E_r)} \int_{E_r} |f(y)| \, dy = f(x).$$

**Corollary 4.2.4.** If  $f \in L^1_{\text{loc}}(\mathbb{R})$ , then  $F(x) = \int_0^x f(y) \, dy$  is differentiable and  $F'(x) = f(x)$  *almost everywhere*.

# Chapter 5

## Normed Vector Space

### Lecture 23: Metric, normed and $L^p$ Spaces

#### 5.1 Metric Spaces and Normed Spaces

09 Mar. 11:00

We have seen the definition of a [norm](#) before, now we formally introduce the concept of *metric*.

**Definition 5.1.1 (Metric).** Let  $Y$  be a set, a function  $\rho: Y \times Y \rightarrow [0, \infty)$  is a *metric* on  $Y$  if

- $\rho(x, y) = \rho(y, x)$  for all  $x, y \in Y$ .
- $\rho(x, z) \leq \rho(x, y) + \rho(y, z)$  for all  $x, y, z \in Y$ .
- $\rho(x, y) = 0$  if and only if  $x = y$ .

**Note.** The following make sense in a [metric](#) space.

- (a) Open/closed balls.
- (b) Open/closed sets.
- (c) Convergence sequences ( $x_n \rightarrow x$  with respect to  $\rho$  if and only if  $\lim_{n \rightarrow \infty} \rho(x_n, x) = 0$ ).
- (d) Continuous functions.

**Example.** We have the following [metric](#) spaces.

- (a)  $\mathbb{Q}$  with  $\rho(x, y) = |x - y|$ .
- (b)  $\mathbb{R}$  with  $\rho(x, y) = |x - y|$ .
- (c)  $\mathbb{R}_+$  with  $\rho(x, y) = |\ln(y/x)|$ .
- (d)  $\mathbb{R}^d$  with

$$\rho_p(x, y) = \left( \sum_{i=1}^d |x_i - y_i|^p \right)^{1/p}$$

and

$$\rho_\infty(x, y) = \max_{1 \leq i \leq d} |x_i - y_i|.$$

These all give the same open sets, hence they are topologically equivalent.

(e)  $C([0, 1])$  with

$$\rho_p(f, g) = \left( \int_0^1 |f - g|^p \right)^{1/p}$$

and

$$\rho_\infty(f, g) = \max_{x \in [0, 1]} |f(x) - g(x)|.$$

(f) Let  $(X, \mathcal{A}, \mu)$  be a **measure space** with  $\mu(X) < \infty$ . Let  $Y$  be the set of **measurable functions** on  $X$ , then

$$\rho(f, g) = \int \min\{|f(x) - g(x)|, 1\} d\mu(x)$$

is a **metric** and  $f_n \rightarrow f$  in  $\rho$  if and only if  $f_n \rightarrow f$  in **measure**.

Let  $V$  be a vector space over scalar field  $K = \mathbb{R}$  or  $K = \mathbb{C}$ .

**As previously seen** (Metric induced by a norm). Recall the definition of **seminorm** and **norm**. We see that a **norm** induces a **metric**

$$\rho(v, w) := \|v - w\|,$$

and we have

$$v_n \rightarrow v \Leftrightarrow \lim_{n \rightarrow \infty} \|v_n - v\| = 0.$$

**Example.** We first see some common examples of **normed** vector space.

(a)  $L^1(X, \mathcal{A}, \mu)$  with  $\|f\|_1 := \int |f| d\mu$ .

(b)  $C([0, 1])$  with  $\|f\|_1 := \int_0^1 |f(x)| dx$ ,  $\|f\|_\infty := \max_{0 \leq x \leq 1} |f(x)|$ .

(c) For  $\mathbb{R}^d$  and  $0 < p < \infty$ , we have

$$\|x\|_p := \left( \sum_{i=1}^d |x_i|^p \right)^{1/p}, \quad \|x\|_\infty := \max_{1 \leq i \leq d} |x_i|.$$

## 5.2 $L^p$ Space

It turns out that we can generalize  $L^1$  into  $L^p$ .

**Definition 5.2.1** ( $L^p$  space). Given a **measure space**  $(X, \mathcal{A}, \mu)$  and a **measurable function**  $f$  and  $p$  such that  $0 < p < \infty$ , we define a **seminorm**  $\|\cdot\|_p$  such that

$$\|f\|_p := \left( \int_X |f|^p d\mu \right)^{1/p},$$

which induces the so-called  $L^p$  space, denoted as  $L^p(X, \mathcal{A}, \mu)$  where

$$L^p(X, \mathcal{A}, \mu) := \{f \mid \|f\|_p < \infty\}.$$

**Remark.** Note that  $\|\cdot\|_p$  is only a **seminorm**. But if we identify functions which are equal **almost everywhere**, then it's indeed a **norm**.

**Example.**  $(\mathbb{R}, \mathcal{L}, m)$  has  $f(x) = x^{-\alpha} \mathbb{1}_{(1, \infty)}(x) \in L^p$  if and only if  $\alpha p > 1$ . In contrast,  $g(x) = x^{-\beta} \mathbb{1}_{(0, 1)}(x) \in L^p$  if and only if  $\beta p < 1$ .

Similar to **Definition 5.2.1**, we have the following.

**Definition 5.2.2** ( $\ell^p$  space). If  $(X, \mathcal{P}(X), \nu)$  is equipped with the [counting measure](#), then we say it's an  $\ell^p$  space such that

$$\ell^p(X) := L^p(X, \mathcal{P}(X), \nu).$$

**Remark.** We are interested in  $\ell^p(\mathbb{N})$  in particular. We have

$$\ell^p := \ell^p(\mathbb{N}) = \left\{ a = (a_1, a_2, \dots) \mid \|a\|_p = \left( \sum_{i=1}^{\infty} |a_i|^p \right)^{1/p} < \infty \right\}.$$

**Lemma 5.2.1.**  $L^p(X, \mathcal{A}, \nu)$  is a vector space for all  $p \in (0, \infty)$ .

**Proof.** We verify the following.

- $c \cdot f \in L^p(X, \mathcal{A}, \mu)$  for  $c \in \mathbb{R}$ . Indeed, since

$$\|cf\|_p = \left( \int |cf|^p d\mu \right)^{1/p} = |c| \|f\|_p < \infty \Leftrightarrow \|f\|_p < \infty,$$

which implies  $c \cdot f \in L^p(X, \mathcal{A}, \mu)$ .

- $f + g \in L^p(X, \mathcal{A}, \mu)$ . Indeed, since for any real numbers  $\alpha, \beta$ , we have

$$(\alpha + \beta)^p \leq (2 \cdot \max\{|\alpha|, |\beta|\})^p = 2^p \cdot \max\{|\alpha|^p, |\beta|^p\} \leq 2^p (|\alpha|^p + |\beta|^p),$$

which implies that for  $f, g \in L^p(X, \mathcal{A}, \mu)$ , we have

$$\|f + g\|_p < \infty \Leftrightarrow \|f + g\|_p^p = \int |f + g|^p d\mu \leq 2^p \int (|f|^p + |g|^p) d\mu < \infty.$$

This further implies

$$\|f + g\|_p < \infty \Leftrightarrow \|f\|_p, \|g\|_p < \infty,$$

which is what we want. ■

We see that in the above derivation, it doesn't give us the triangle inequality, namely

$$\|f + g\|_p \leq \|f\|_p + \|g\|_p,$$

hence we need some new results.

**Theorem 5.2.1** (Hölder's inequality). Let  $1 < p < \infty$ , and let  $q := p/(p-1)$  so that  $1/p + 1/q = 1$ . Then we have

$$\|f \cdot g\|_1 \leq \|f\|_p \|g\|_q.$$

**Proof.** We prove this in steps.

**Claim.** We have

$$t \leq \frac{t^p}{p} + 1 - \frac{1}{p} = \frac{t^p}{p} + \frac{1}{q}$$

for all  $t \geq 0$ .

**Proof.** By taking  $F(t) := t - t^p/p$  and  $t \geq 0$ , we see that the maximum of  $F$  implies the above inequality. ⊗

**Claim (Young's Inequality).** We have

$$\alpha\beta \leq \frac{\alpha^p}{p} + \frac{\beta^q}{q}$$

for  $\alpha, \beta > 0$ .<sup>a</sup>

<sup>a</sup>[https://en.wikipedia.org/wiki/Young's\\_inequality\\_for\\_products](https://en.wikipedia.org/wiki/Young's_inequality_for_products)

**Proof.** This follows by taking  $t := \alpha/\beta^{q-1}$  in the first inequality we obtained.  $\otimes$

Then, without loss of generality, we can assume that  $0 < \|f\|_p, \|g\|_q < \infty$ . Now, consider  $F(x) = f(x)/\|f\|_p$ ,  $G(x) = g(x)/\|g\|_q$ . We know that  $\|F\|_p = 1 = \|G\|_q$ . Then by Young's Inequality, we have

$$\int |F(x)G(x)| \, d\mu \leq \int \frac{|F(x)|^p}{p} + \frac{|G(x)|^q}{q} \Rightarrow \frac{\|fg\|_1}{\|f\|_p \|g\|_q} \leq \frac{1}{p} + \frac{1}{q} = 1,$$

which implies our desired result.  $\blacksquare$

**Example.** For  $p = q = 2$ ,  $X = \{1, \dots, d\}$  with  $\mu$  being the **counting measure**, then for any  $x, y \in \mathbb{R}^d$ , we have

$$\sum_{i=1}^d |x_i y_i| \leq \sqrt{\sum_{i=1}^d x_i^2} \sqrt{\sum_{i=1}^d y_i^2}$$

We now see how we can obtain the desired triangle inequality.

**Theorem 5.2.2 (Minkowski's Inequality).** Let  $1 \leq p < \infty$ , then for  $f, g \in L^p$ ,

$$\|f + g\|_p \leq \|f\|_p + \|g\|_p.$$

**Proof.** For  $p = 1$ , it's easy since it's just triangle inequality. Now, we assume that  $1 < p < \infty$ , and we may assume also that  $\|f + g\| \neq 0$  without loss of generality. Then

$$\begin{aligned} \int |f(x) + g(x)|^p &\leq \int |f(x) + g(x)|^{p-1} (|f(x)| + |g(x)|) \\ &\leq \left( \int |f + g|^{(p-1)q} \right)^{1/q} \left[ \left( \int |f|^p \right)^{1/p} + \left( \int |g|^p \right)^{1/p} \right] \\ &\leq \left( \int |f + g|^p \right)^{1/q} (\|f\|_p + \|g\|_p). \end{aligned}$$

We then see that

$$\underbrace{(|f(x) + g(x)|^p)^{1-1/q}}_{(|f(x)+g(x)|^p)^{1/p}} \leq \|f\|_p + \|g\|_p,$$

which is just  $\|f + g\|_p \leq \|f\|_p + \|g\|_p$ .  $\blacksquare$

## Lecture 24: Embedding $L^p$ Space

**Definition 5.2.3 (Essential supremum).** For a **measurable function**  $f$  on  $(X, \mathcal{A}, \mu)$ , we define

$$S := \{\alpha \geq 0 \mid \mu(\{x \mid |f(x)| > \alpha\}) = 0\} = \{\alpha \geq 0 \mid |f(x)| \leq \alpha \text{ a.e.}\}.$$

11 Mar. 11:00

Then, we say that the *essential supremum* of  $f$ , denoted as  $\|f\|_\infty$ , is defined as

$$\|f\|_\infty := \begin{cases} \inf S, & \text{if } S \neq \emptyset; \\ \infty, & \text{if } S = \emptyset. \end{cases}$$

**Definition 5.2.4** ( $L^\infty$  space). Let  $L^\infty(X, \mathcal{A}, \mu)$  be

$$L^\infty(X, \mathcal{A}, \mu) = \{f \mid \|f\|_\infty < \infty\}.$$

**Definition 5.2.5** ( $\ell^\infty$  space). We let  $\ell^\infty$  be defined as

$$\ell^\infty = L^\infty(\mathcal{N}, \mathcal{P}(\mathcal{N}), \nu),$$

where  $\nu$  is the [counting measure](#).

**Example.** Consider  $(\mathbb{R}, \mathcal{L}, m)$ . Then

$$\begin{aligned} f(x) &= \frac{1}{x} \mathbb{1}_{(0, \infty)}(x) \notin L^\infty; \\ g(x) &= x \mathbb{1}_{\mathbb{Q}}(x) + \frac{1}{1+x^2} \in L^\infty. \end{aligned}$$

If  $f$  is continuous on  $(\mathbb{R}, \mathcal{L}, m)$ , then  $\|f\|_\infty = \sup_{x \in \mathbb{R}} |f(x)|$ . For  $a \in \ell^\infty$ , we have  $\|a\|_\infty = \sup_{i \in \mathbb{N}} |a_i|$ , and sequences in  $\ell^\infty$  are exactly the bounded sequences.

**Lemma 5.2.2.** We have the following.

(a) Suppose  $f \in L^\infty(X, \mathcal{A}, \mu)$ . Then,

$$\begin{cases} \mu(\{x \mid |f(x)| > \alpha\}) = 0, & \text{if } \alpha \geq \|f\|_\infty; \\ \mu(\{x \mid |f(x)| > \alpha\}) > 0, & \text{if } \alpha < \|f\|_\infty. \end{cases}$$

(b)  $|f(x)| \leq \|f\|_\infty$  [almost everywhere](#).

(c)  $f \in L^\infty$  if and only if there exists a bounded [measurable function](#)  $g$  such that  $f = g$  [almost everywhere](#).

**Proof.** ■

DIY

**Theorem 5.2.3.** We have the following.

(a)  $\|fg\|_1 \leq \|f\|_1 \|g\|_\infty$ .

(b)  $\|f + g\|_\infty \leq \|f\|_\infty + \|g\|_\infty$ .

(c)  $f_n \rightarrow f$  in  $L^\infty$  if and only if  $f_n \rightarrow f$  [uniformly almost everywhere](#).

**Proof.** We'll do one implication in (c). Let  $A_n = \{x \mid |f_n(x) - f(x)| > \|f_n - f\|_\infty\}$ . Then  $\mu(A_n) = 0$ . Let  $A = \bigcup_n A_n$ , we see that  $\mu(A) = 0$  as well.

For  $x \in A^c$  and for every  $n$ , we have

$$|f_n(x) - f(x)| \leq \|f_n - f\|_\infty.$$

Given  $\epsilon > 0$ , there is an  $N$  so that  $\|f_n - f\|_\infty < \epsilon$  for all  $n \geq N$ . But then for all  $x \in A^c$ ,  $|f_n(x) - f(x)| < \epsilon$  as well. ■

DIY (a)  
and (b)

**Remark.** The motivation for (a) is that

$$\frac{1}{1} + \frac{1}{\infty} = 1,$$

and we want to have the similar result as in Theorem 5.2.1.

**Proposition 5.2.1.** We have the following.

- (a) For  $1 \leq p < \infty$ , the collection of simple functions with finite measure support is dense in  $L^p(X, \mathcal{A}, \mu)$ .
- (b) For  $1 \leq p < \infty$ , the collection of step functions with finite measure support is dense in  $L^p(\mathbb{R}, \mathcal{L}, m)$ , so is  $C_c(\mathbb{R})$ .
- (c) For  $p = \infty$ , the collection of simple functions is dense in  $L^\infty(X, \mathcal{A}, \mu)$ .

**Proof.** ■

DIY

**Remark.** Note that  $C_c(\mathbb{R})$  is **not** dense in  $L^\infty(\mathbb{R}, \mathcal{L}, m)$ .

### 5.3 Embedding Properties of $L^p$ Spaces

**Definition 5.3.1** (Equivalent norm). Two norms  $\|\cdot\|, \|\cdot\|'$  on  $V$  are *equivalent* if there exists  $c_1, c_2 > 0$ , such that

$$c_1 \|v\| \leq \|v\|' \leq c_2 \|v\|$$

for all  $v \in V$ .

**Note.** We see that

- (a) These norms gives the same topological properties (open sets, closed sets, convergence, etc.).
- (b) Definition 5.3.1 is an equivalence relation on norms.

**Example.** For  $\mathbb{R}^d$  we have the norms  $\|\cdot\|_p$  for  $1 \leq p \leq \infty$ . All of these are equivalent. We see that for  $1 \leq p < \infty$ ,


$$\|x\|_p = \left( \sum_{i=1}^d |x_i|^p \right)^{1/p} \leq (d \|x\|_\infty^p)^{1/p} = d^{1/p} \|x\|_\infty.$$

Also,

$$\|x\|_p = \left( \sum_{i=1}^d |x_i|^p \right)^{1/p} \geq (\|x\|_\infty^p)^{1/p} = \|x\|_\infty.$$

Thus,  $\|\cdot\|_p$  is equivalent to  $\|\cdot\|_\infty$  for every  $1 \leq p < \infty$ , and transitivity gives that they are all equivalent.

Another way of thinking of this, by assuming  $v \neq 0$ , and scaling by some  $t$ , we may assume  $v$  lies on the unit circle in one of the norms. Then we are squeezing a unit circle in  $\|\cdot\|'$  between two circles of radius  $c_1, c_2$  in  $\|\cdot\|$ . In picture, we have to show that  $\|\cdot\|_2$  and  $\|\cdot\|_\infty$  are equivalent, we have



since the circles in  $\|\cdot\|_\infty$  are squares.

**Example.** For  $1 \leq p, q \leq \infty$ , we have  $L^p(\mathbb{R}, m)$ -norm and  $L^q(\mathbb{R}, m)$ -norm are not equivalent, even worse, we have that

$$L^p(\mathbb{R}, m) \not\subseteq L^1(\mathbb{R}, m), \quad L^p(\mathbb{R}, m) \not\supseteq L^1(\mathbb{R}, m).$$

## Lecture 25: Banach Spaces

**Proposition 5.3.1.** Suppose  $\mu(X) < \infty$ , then for every  $0 < p < q \leq \infty$ ,  $L^q \subseteq L^p$ .

14 Mar. 11:00

**Proof.** Suppose  $q < \infty$ , then

$$\int |f|^p \leq \left( \int (|f|^p)^{q/p} \right)^{p/q} \left( \int 1^{q/(q-p)} \right)^{1-p/q} = \left( \int |f|^q \right)^{p/q} \mu(X)^{1-p/q} < \infty$$

where we split  $\int |f|^p$  into  $\int |f|^p \cdot 1$ . From Hölder's inequality with  $q/p > 1$ , we have

$$\|f\|_p \leq \|f\|_q \mu(X)^{1/p-1/q} < \infty.$$

The case that  $q = \infty$  is left as an exercise. ■

DIY

**Proposition 5.3.2.** If  $0 < p < q \leq \infty$ , then  $\ell^p \subseteq \ell^q$ .

**Proof.** We consider two cases.

- When  $q = \infty$ , we have

$$\|a\|_\infty^p = \left( \sup_i |a_i| \right)^p = \sup_i |a_i|^p \leq \sum_{i=1}^{\infty} |a_i|^p.$$

Thus,  $\|a\|_\infty \leq \|a\|_p$ .

- When  $q < \infty$ , we see that

$$\sum_{i=1}^{\infty} |a_i|^q = \sum_{i=1}^{\infty} |a_i|^p \cdot |a_i|^{q-p} \leq \|a\|_\infty^{q-p} \sum_{i=1}^{\infty} |a_i|^p \leq \|a\|_\infty^{q-p} \cdot \|a\|_p^p = \|a\|_p^q.$$

Therefore,

$$\|a\|_q \leq \|a\|_p.$$



**Proposition 5.3.3.** For all  $0 < p < q < r \leq \infty$ ,  $L^p \cap L^r \subseteq L^q$ .

**Proof.** ■

DIY

## 5.4 Banach Spaces

Let's start with a definition.

**Definition 5.4.1 (Cauchy sequence).** Let  $Y, \rho$  be a [metric](#) space. We call  $x_n$  a *Cauchy sequence* if for every  $\epsilon > 0$ , there exists an  $N \in \mathbb{N}$  such that for all  $n, m \geq N$ ,  $\rho(x_n, x_m) < \epsilon$ .

**Note.** Convergent sequence are [Cauchy](#).

**Definition 5.4.2 (Complete).** A [metric](#) space  $(Y, \rho)$  is called *complete* if every [Cauchy sequence](#) in  $Y$  converges.

**Example.** We first see some examples.

- (a) We see that  $\mathbb{Q}$  with  $\rho(x, y) = |x - y|$  is **not** [complete](#), but  $\mathbb{R}$  with the same [metric](#) is [complete](#).
- (b)  $C([0, 1])$  with  $\rho(f, g) = \|f - g\|_\infty$  is [complete](#), but with  $\rho(f, g) = \int |f - g|$  is not.

**Definition 5.4.3 (Banach space).** A *Banach space* is a [complete normed](#) vector space.

**Remark.** Namely, a vector space equipped with a [norm](#) whose [metric induced by the norm](#) is [complete](#).

**Theorem 5.4.1.** Let  $(V, \|\cdot\|)$  be a [normed](#) space. Then,

$V$  is [complete](#)  $\Leftrightarrow$  every absolutely convergent series is convergent.

i.e., if  $\sum_{i=1}^{\infty} \|v_i\| < \infty$ , then  $\left\{ \sum_{i=1}^N v_i \right\}_{N \in \mathbb{N}}$  converges to some  $s \in V$ .

Before we prove [Theorem 5.4.1](#), we first see one of the result based on this theorem.<sup>1</sup>

**Theorem 5.4.2 (Riesz-Fischer theorem).** For every  $1 \leq p \leq \infty$ , we have  $L^p(X, \mathcal{A}, \mu)$  is [complete](#), hence a [Banach space](#).

**Proof.** We prove this in two cases.

- We first prove this for  $1 \leq p < \infty$ . Suppose  $f_n \in L^p$  and  $\sum_{n=1}^{\infty} \|f_n\|_p < \infty$ .

We need to show that there is an  $F \in L^p$  such that  $\left\| \sum_{n=1}^N f_n - F \right\|_p \rightarrow 0$  as  $N \rightarrow \infty$ . i.e., we need to show the following.

1.  $\sum_{n=1}^{\infty} f_n(x)$  is convergent [a.e.](#) In fact, we can show this by showing the following.

**Claim.** We have

$$\int \sum_{n=1}^{\infty} |f_n(x)| < \infty.$$

<sup>1</sup>The proof can be found in [here](#).

**Proof.** Let  $G(x) = \sum_{n=1}^{\infty} |f_n(x)| = \sup_N \sum_{n=1}^N |f_n(x)|$ ,  $G: X \rightarrow [0, \infty]$ . Also, let  $G_N(x) = \sum_{n=1}^N |f_n(x)|$ . Then, we have

$$0 \leq G_1 \leq G_2 \leq \dots \leq G,$$

and  $G_N \rightarrow G$ . Furthermore,

$$0 \leq G_1^p \leq G_2^p \leq \dots \leq G^p,$$

and  $G_N^p \rightarrow G^p$ . From [monotone convergence theorem](#),

$$\int G^p = \lim_{N \rightarrow \infty} \int G_N^p.$$

From [Minkowski inequality](#), we further have

$$\|G_N\|_p \leq \sum_{n=1}^N \|f_n\|_p \leq \sum_{n=1}^{\infty} \|f_n\|_p := B < \infty.$$

Thus,

$$\int G(x)^p = \lim_{N \rightarrow \infty} \int G_N^p = \lim_{N \rightarrow \infty} \|G_N\|_p^p \leq B^p < \infty.$$

We see that  $G$  is finite [a.e.](#) as desired. This implies that  $\sum_{n=1}^{\infty} |f_n(x)| < \infty$  [a.e.](#), so  $\sum_{n=1}^{\infty} f_n(x)$  converges [a.e.](#) Now, we simply let

$$F(x) = \begin{cases} \sum_{n=1}^{\infty} f_n(x), & \text{if it converges;} \\ 0, & \text{otherwise.} \end{cases}$$

⊗

2.  $F \in L^p$ , where  $F(x) := \sum_{n=1}^{\infty} f_n(x)$  [a.e.](#) and say is zero elsewhere.

**Claim.**  $F$  defined in this way is indeed in  $L^p$ .

**Proof.** This is clear since

$$|F(x)| \leq G(x) \Rightarrow \int |F|^p \leq \int G^p < \infty,$$

hence  $F \in L^p$ .

⊗

3. We then show the last condition we need to check.

**Claim.**  $\left\| \sum_{n=1}^N f_n - F \right\|_p \rightarrow 0$  as  $N \rightarrow \infty$ .

**Proof.** We now see that

$$\left| \sum_{n=1}^N f_n(x) - F(x) \right|^p \leq \left( \sum_{n=1}^{\infty} |f_n(x)| + |F(x)| \right)^p \leq (2G(x))^p.$$

Since  $2G \in L^p$ , so  $2G^p \in L^1$ . Thus, by [dominated convergence theorem](#), we have

$$\lim_{N \rightarrow \infty} \int \left| \sum_{n=1}^N f_n(x) - F(x) \right|^p dx = 0.$$

This implies

$$\left\| \sum_{n=1}^N f_n - F \right\|_p \rightarrow 0$$

as  $N \rightarrow \infty$ . ⊗

- Now assume  $1 \leq p \leq \infty$ . DIY

■

## Lecture 26: Bounded Linear Transformations

We now prove [Theorem 5.4.1](#), completing the proof of [Theorem 5.4.2](#) since the latter relies on this result. 16 Mar. 11:00

**Proof of Theorem 5.4.1.** We prove it by proving two directions.

( $\Rightarrow$ ) Suppose  $V$  is [complete](#), and fix an absolutely convergent series  $\sum_n v_n$ . Define  $s_N = \sum_{n=1}^N v_n$ . It suffices to show the partial sums are a [Cauchy Sequence](#).

Fix  $\epsilon > 0$ , then because  $\sum_{n=1}^{\infty} \|v_n\| < \infty$ , there is a  $K \in \mathbb{N}$  so that

$$\sum_{n=K}^{\infty} \|v_n\| < \epsilon.$$

Now let  $M > N > K$ , we see that

$$\|s_M - s_N\| = \left\| \sum_{n=N+1}^M v_n \right\| \leq \sum_{n=N+1}^M \|v_n\| \leq \sum_{n=N}^{\infty} \|v_n\| < \epsilon,$$

so this is [Cauchy](#).

( $\Leftarrow$ ) Now suppose  $v_n, n \in \mathbb{N}$  is a [Cauchy sequence](#). For all  $j \in \mathbb{N}$ , there exists an  $N_j \in \mathbb{N}$  such that

$$\|v_n - v_m\| < \frac{1}{2^j}$$

for all  $n, m \geq N_j$ . Without loss of generality, we may assume  $N_1 < N_2 < \dots$

Let  $w_1 = v_{N_1}$ ,  $w_j = v_{N_j} - v_{N_{j-1}}$  for  $j \geq 2$ . Therefore,

$$\sum_{j=1}^{\infty} \|w_j\| \leq \|v_{N_1}\| + \sum_{j=2}^{\infty} \frac{1}{2^{j-1}} < \infty.$$

Thus,  $\sum_{j=1}^k w_j \rightarrow s \in V$  as  $k \rightarrow \infty$ . But by telescoping, we have

$$v_{N_k} = \sum_{j=1}^k w_j \rightarrow s.$$

Now we claim that since  $v_n$  is **Cauchy**, so  $v_n \rightarrow s$ .

Explicitly, take  $\epsilon > 0$ , and let  $k$  be large enough so that  $\|v_{N_k} - s\| < \epsilon$  and  $1/2^k < \epsilon$ . Then if  $n > N_k$  then

$$\|v_n - s\| \leq \|v_n - v_{N_k}\| + \|v_{N_k} - s\| < \epsilon + \epsilon = 2\epsilon.$$

Thus,  $v_n \rightarrow s$ . ■

## 5.5 Bounded Linear Transformations

**Definition 5.5.1** (Bounded linear transformation). Given two **normed** vector spaces  $(V, \|\cdot\|)$ ,  $(W, \|\cdot\|')$ , a linear map  $T: V \rightarrow W$  is called a *bounded map* if there exists  $c \geq 0$  such that

$$\|Tv\|' \leq c\|v\|$$

for all  $v \in V$ .

**Proposition 5.5.1.** Suppose  $T: (V, \|\cdot\|) \rightarrow (W, \|\cdot\|')$  is a linear map. Then the following are equivalent.

- (a)  $T$  is continuous.
- (b)  $T$  is continuous at 0.
- (c)  $T$  is a **bounded map**.

**Proof.** (a)  $\Rightarrow$  (b) is clear.

**Claim.** (b)  $\Rightarrow$  (c).

**Proof.** Take  $\epsilon = 1$ , then there exists a  $\delta > 0$  such that  $\|Tu\|' < 1$  if  $\|u\| < \delta$ .

Now take an arbitrary  $\|v\| \in V$ ,  $v \neq 0$ . Let  $u = \frac{\delta}{2\|v\|}v$ . Then  $\|u\| < \delta$ . Therefore,

$$\|Tu\|' < 1 \Rightarrow \frac{\delta}{2\|v\|} \|Tv\|' < 1 \Rightarrow \|Tv\|' < \frac{2}{\delta} \|v\|.$$

Then  $2/\delta$  is our constant. ⊗

**Claim.** (c)  $\Rightarrow$  (a).

**Proof.** Fix  $v_0 \in V$ . Then for some constant  $c$

$$\|Tv - Tv_0\|' = \|T(v - v_0)\|' \leq c\|v - v_0\|.$$

Thus,  $T$  is continuous, as when  $v \rightarrow v_0$  the right-hand side goes to zero, and so  $Tv \rightarrow Tv_0$ . ⊗

**Example.** Let's see some examples.

- (a) We can look at

$$\begin{aligned} T: \ell^1 &\rightarrow \ell^1 \\ (a_1, a_2, \dots) &\mapsto (a_2, a_3, \dots). \end{aligned}$$

Then clearly  $\|Ta\|_1 \leq \|a\|_1$ , so  $T$  is a **bounded linear transformation**.

- (b) We can also look at  $S: (C([-1, 1]), \|\cdot\|_1) \rightarrow \mathbb{C}$ , where  $Sf = f(0)$ .  $S$  is not a **bounded linear**

transformation, because we can make

$$\begin{cases} \|Sf\| &= |f(0)| = n \\ \|f\|_1 &= 1 \end{cases}$$

for every  $n \in \mathbb{N}$  (take  $f$ 's graph to be a skinny triangle shooting up to  $n$  at 0).

(c) But  $U: (C([-1, 1]), \|\cdot\|_\infty) \rightarrow \mathbb{C}$  defined by  $Uf = f(0)$  is a **bounded linear transformation**, because  $|f(0)| \leq \|f\|_\infty$ .

(d) Let  $A$  be an  $n \times m$  matrix. Then  $T: \mathbb{R}^m \rightarrow \mathbb{R}^n$  defined by  $v \mapsto Av$  is a **bounded linear transformation**.

Explicitly this is

$$(Tv)_i = (Av)_i = \sum_{j=1}^m A_{ij}v_j.$$

(e) Let  $K(x, y)$  be a continuous function on  $[0, 1] \times [0, 1]$ . We'll define

$$T: (C[0, 1], \|\cdot\|_\infty) \rightarrow (C[0, 1], \|\cdot\|_\infty)$$

by

$$(Tf)(x) = \int_0^1 K(x, y)f(y) dy.$$

This is an analogue of matrix multiplication ( $K$  is like a continuous matrix). This is a **bounded linear transformation**.

(f) Let us look at  $T: L^1(\mathbb{R}) \rightarrow (C(\mathbb{R}), \|\cdot\|_\infty)$  defined by

$$(Tf)(t) = \int_{-\infty}^{\infty} e^{-itx} f(x) dx$$

that is the Fourier transform of  $f$ .

(g)  $T: (C^\infty[0, 1], \|\cdot\|_\infty) \rightarrow (C^\infty[0, 1], \|\cdot\|_\infty)$ . Define

$$(Tf)(x) = f'(x).$$

This is not a **bounded linear transformation**. In contrast,  $S$ , defined on the same spaces

$$(Sf)(x) = \int_0^x f(t) dt$$

is bounded.

**Definition 5.5.2 (Operator norm).** Let  $L(V, W)$  be defined as a vector space such that

$$L(V, W) := \{T: V \rightarrow W \mid T \text{ is a bounded linear transformation}\}.$$

Then for  $T \in L(V, W)$ , the *operator norm* of  $T$  is

$$\begin{aligned} \|T\| &:= \inf\{c \geq 0 \mid \|Tv\|'' \leq c\|v\|' \text{ for all } v \in V\} \\ &= \sup\left\{\frac{\|Tv\|''}{\|v\|'} \mid v \neq 0, v \in V\right\} \\ &= \sup\{\|Tv\|'' \mid \|v\|' = 1, v \in V\}. \end{aligned}$$

**Lemma 5.5.1.** We have that

- (a) The [three definitions](#) of  $\|T\|$  above are all equal.  
 (b)  $(L(V, W), \|\cdot\|)$  is indeed a [normed](#) space.

**Proof.** ■

DIY

## Lecture 27: Dual Space

As previously seen. From [Definition 5.5.2](#), we have that

18 Mar. 11:00

$$\|Tv\|'' \leq \|T\| \|v\|'.$$

**Remark.** Notice that this [Definition 5.5.2](#) is only for [bounded linear transformation](#).

**Theorem 5.5.1.** If  $W$  is [complete](#), then  $L(V, W)$  is [complete](#).

**Proof.** Suppose  $T_n$  is a [Cauchy sequence](#) in  $L(V, W)$ . Fix  $v \in V$  and let  $w_n := T_n v \in W$ , we then have

$$\|w_n - w_m\| = \|T_n v - T_m v\| = \|(T_n - T_m)v\| \leq \underbrace{\|T_n - T_m\|}_{\rightarrow 0} \underbrace{\|v\|}_{\text{fixed value}}.$$

Thus,  $w_n$  is [Cauchy](#), so it converges since  $W$  is [complete](#). We call its unique limit  $Tv$ . This makes  $T: V \rightarrow W$  into a function. We must show it is a [bounded linear transformation](#) and that  $\|T_n - T\| \rightarrow 0$ . ■

DIY

## 5.6 Dual of $L^p$ Spaces

**Example.** Let  $w \in \mathbb{R}^d$ , and denote the [inner product](#) between  $w$  and  $v \in \mathbb{R}^d$  by

$$v \cdot w := \langle v, w \rangle.$$

Then we can consider

$$\max\{v \cdot w \mid \|v\|_2 = 1\} = \|w\|_2.$$

If  $w \in \mathbb{C}^d$ , this is similar since

$$\max\{|v \cdot w| \mid \|v\|_2 = 1\} = \|w\|_2.$$

These maximums are achieved by  $v = \frac{\overline{w}}{\|w\|}$  if  $w \neq 0$ .

**Proposition 5.6.1.** Let  $1/p + 1/q = 1$  with  $1 \leq q < \infty$ . For every  $g \in L^q$ ,

$$\|g\|_q = \sup \left\{ \left| \int fg \right| \mid \|f\|_p = 1 \right\}.$$

Suppose  $\mu$  is [σ-finite](#), then the result also holds for  $q = \infty$ ,  $p = 1$ .

**Proof.** By [Hölder's inequality](#), we know that

$$\left| \int fg \right| \leq \int |fg| = \|fg\|_1 \leq \|f\|_p \|g\|_q = \|g\|_q.$$

Thus, the supremum is less or equal to  $\|g\|_q$ .

(a) Let

$$f(x) = \frac{|g(x)|^{q-1} \cdot \overline{\operatorname{sgn}(g(x))}}{\|g\|_q^{q-1}}$$

Then  $\int |f|^p = 1$ , and  $\int fg = \|g\|_q$ .

Check

**Note.** For  $\alpha \in \mathbb{C}$ ,  $\operatorname{sgn}(\alpha) := e^{i\theta}$  where  $\alpha = |\alpha| e^{i\theta}$ .

(b) The case that  $\mu$  is  $\sigma$ -finite and  $q = \infty, p = 1$  can be shown.

DIY

■

**Remark.** One could use the above to prove [Minkowski's inequality](#) (as it only uses [Hölder's inequality](#)).

**Definition 5.6.1** (Dual space). For a [normed](#) space  $(V, \|\cdot\|)$ , its *dual space* is  $V^* = L(V, \mathbb{R})$  or  $V^* = L(V, \mathbb{C})$ .

**Remark.** Namely, the [dual space](#) of  $V$  contains [bounded linear transformations](#) with codomain being the scalar field.

**Definition 5.6.2** (Linear functional). Given a [normed](#) space  $(V, \|\cdot\|)$ ,  $\ell \in V^*$  is called a *linear functional* on  $V$ . i.e.,

- $\ell: V \rightarrow \mathbb{R}$  (or  $\mathbb{C}$ ).
- $\ell$  is linear.
- There exists a  $c \geq 0$  such that  $|\ell(v)| = c\|v\|$ .

**Note.**  $V^*$  is always a [Banach space](#) (even if  $V$  is not [complete](#)).

**Corollary 5.6.1.** We have the following.

(a) Let  $1/p + 1/q = 1, 1 \leq q < \infty$ . For  $g \in L^q$  define  $\ell_g \in L^p \rightarrow \mathbb{C}$  by

$$\ell_g(f) = \int fg.$$

Then  $\ell_g \in (L^p)^*$ . Furthermore,  $\|\ell_g\| = \|g\|_q$ .

(b) If  $\mu$  is  $\sigma$ -finite, then this also holds for  $q = \infty, p = 1$ .

**Proof.**  $\ell_g$  is clearly linear in  $f$  because the integral is linear. Then [Proposition 5.6.1](#) gives in both (a) and (b) that

$$\|g\|_q = \sup\{|\ell_g(f)| \mid \|g\|_p = 1\} = \|\ell_g\|$$

and so  $\ell_g$  is a [bounded linear transformation](#) with the desired properties. ■

**Theorem 5.6.1.** We have the following.

(a) Let  $1/p + 1/q = 1, 1 \leq q < \infty$ . The map  $T: L^q \rightarrow (L^p)^*$  given by  $Tg = \ell_g$  is an isometric<sup>a</sup> linear isomorphism. This means that

- $T$  is a [bounded linear transformation](#).

- $T$  is bijective.
- $T$  is **norm**-preserving.

(b) If  $\mu$  is  **$\sigma$ -finite** then this also holds for  $q = \infty, p = 1$ .

<sup>a</sup>A map  $T$  is called isometric if for a given  $g$ ,  $\|Tg\| = \|g\|$ .

**Proof.** We have already proved this is isometric in [Corollary 5.6.1](#), it is clearly linear, and isometry implies injectivity.

We will prove that it is surjective later. ■

Fix!!!

**Note.** Even if  $\mu$  is  **$\sigma$ -finite** we might not have  $L^1 \cong (L^\infty)^*$ . Also note that  $L^2 \cong (L^2)^*$ , and for all  $1 < p < \infty$  we have  $(L^p)^{**} \cong L^p$ .



# Chapter 6

## Signed and Complex Measures

### Lecture 28: Signed Measure

21 Mar. 11:00

**As previously seen.** Suppose  $f: X \rightarrow [0, \infty]$  is a measurable function on  $(X, \mathcal{A}, \mu)$ .

We can define  $\nu(E) = \int_E f d\mu$  for  $E \in \mathcal{A}$ , and  $\nu$  is a measure on  $(X, \mathcal{A})$ . This gives a map from the set of non-negative measurable functions on  $X$  to measures on  $X$ . This is injective if we identify functions which are equal almost everywhere. But it is not necessarily surjective. We can then think of measures as a generalization of functions.

For an example, think of a Dirac-Delta measure on  $\mathbb{R}$ . This is not the Lebesgue integral of any non-negative measurable function.

What if instead we took  $f: X \rightarrow \mathbb{R}, \overline{\mathbb{R}}$  or  $\mathbb{C}$ ? We could take the same construction to get  $\nu(E) = \int_E f d\mu$ , but this is no longer a measure as it can take  $\mathbb{R}, \overline{\mathbb{R}}$  or  $\mathbb{C}$  values.

### 6.1 Signed Measures

**Definition 6.1.1.** Let  $(X, \mathcal{A})$  be a measurable space. A signed measure is a function

$$\nu: \mathcal{A} \rightarrow [-\infty, \infty) \text{ or } \nu: \mathcal{A} \rightarrow (-\infty, \infty]$$

such that

- $\nu(\emptyset) = 0$ .
- If  $A_1, A_2, \dots \in \mathcal{A}$  are disjoint then

$$\nu\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \nu(A_i)$$

where the series on the right-hand side converges absolutely if  $\nu(\bigcup_{i=1}^{\infty} A_i) \in (-\infty, \infty)$ .

**Remark.** This means the series does not depend on rearrangement if our function  $\nu$  takes finite value on the set  $\bigcup_i A_i$ .

**Example.** Consider

- $\nu$  is a positive measure (i.e., measure), then  $\nu$  is a signed measure.
- If we have positive measures  $\mu_1, \mu_2$  such that either  $\mu_1(X) < \infty$  or  $\mu_2(X) < \infty$ , then  $\nu = \mu_1 - \mu_2$  is a signed measure.
- If  $f: X \rightarrow \overline{\mathbb{R}}$  on a measure space  $(X, \mathcal{A}, \mu)$  such that  $\int_X f^+ d\mu < \infty$  or  $\int_X f^- d\mu < \infty$ , we can

define

$$\nu(E) = \int_E f \, d\mu$$

and this will be a [signed measure](#).

**Note.** The following weird things happen with [signed measures](#).

- (a)  $A \subseteq B$  does not imply  $\nu(A) \leq \nu(B)$ , as  $\nu(B) = \nu(A) + \nu(B \setminus A)$ , and  $\nu(B \setminus A)$  may be negative.
- (b) If  $A \subseteq B$  and  $\nu(A) = \infty$ , then  $\nu(B) = \infty$ , because  $\nu(B \setminus A) \in (-\infty, \infty]$ .
- (c) Similarly, if  $A \subseteq B$  and  $\nu(A) = -\infty$  then  $\nu(B) = -\infty$ .

**Lemma 6.1.1.** If  $\nu$  is a [signed measure](#) on  $(X, \mathcal{A})$ , then we have the following.

- (a) Continuity from below. If  $E_n \in \mathcal{A}$  and  $E_1 \subseteq E_2 \subseteq \dots$  then

$$\nu\left(\bigcup_{n=1}^{\infty} E_n\right) = \lim_{N \rightarrow \infty} \nu(E_N).$$

- (b) Continuity from above. If  $E_n \in \mathcal{A}$ ,  $E_1 \supseteq E_2 \supseteq \dots$ , and  $-\infty < \nu(E_1) < \infty$  then

$$\nu\left(\bigcap_{n=1}^{\infty} E_n\right) = \lim_{N \rightarrow \infty} \nu(E_N).$$

**Proof.** Read [FF99]. ■

**Definition.** Let  $\nu$  be a [signed measure](#) on  $(X, \mathcal{A})$ . Let  $E \in \mathcal{A}$ , then we say that

**Definition 6.1.2** (Positive set for a signed measure).  $E$  is *positive* for  $\nu$  if for all  $F \subseteq E$ ,  $\nu(F) \geq 0$ .

**Definition 6.1.3** (Negative set for a signed measure).  $E$  is *negative* for  $\nu$  if for all  $F \subseteq E$ ,  $\nu(F) \leq 0$ .

**Definition 6.1.4** (Null set for a signed measure).  $E$  is *null* for  $\nu$  if for all  $F \subseteq E$ ,  $\nu(F) = 0$ .

**Note.** We see that

- (a) If  $E$  is a [positive set](#),  $F \subseteq E$ , then  $\nu(F) \leq \nu(E)$ .
- (b) If  $E$  is a [negative set](#),  $F \subseteq E$ , then  $\nu(F) \geq \nu(E)$ .

**Lemma 6.1.2.** Let  $\nu$  be a [signed measure](#) on  $(X, \mathcal{A})$ , then

- (a) If  $E$  is [positive](#),  $G \subseteq E$  is [measurable](#), then  $G$  is [positive](#).
- (b) If  $E$  is [negative](#),  $G \subseteq E$  is [measurable](#), then  $G$  is [negative](#).
- (c) If  $E$  is [null](#),  $G \subseteq E$  is [measurable](#), then  $G$  is [null](#).
- (d)  $E_1, E_2, \dots$  are [positive](#) sets, then  $\bigcup_{i=1}^{\infty} E_i$  is [positive](#).

**Proof.** The first three are trivial from their definition. For 4., if  $E_1, \dots$  are [positive sets](#), let

$F_n := E_n \setminus \bigcup_{j=1}^{n-1} E_j$ . Then  $F_n \subset E_n$ , so  $F_n$  is **positive sets** from 1., hence if  $E \subset \bigcup_{j=1}^{\infty} E_j$ , then

$$\nu(E) = \sum_{j=1}^{\infty} \nu(E \cap F_j) \geq 0$$

as desired. ■

**Lemma 6.1.3.** Suppose that  $\nu$  is a **signed measure** with  $\nu: \mathcal{A} \rightarrow [-\infty, \infty)$ . Suppose  $E \in \mathcal{A}$  and  $0 < \nu(E) < \infty$ , then there exists a **measurable**  $A \subseteq E$  such  $A$  is a **positive set** and  $\nu(A) > 0$ .

**Proof.** If  $E$  is **positive**, we're done. Otherwise, there exist **measurable** subsets with **negative** measure. Let  $n_1 \in \mathbb{N}$  be the least such  $n_1$  such that there exists  $E_1 \subseteq E$  with  $\nu(E_1) < -1/n_1$ .

If  $E \setminus E_1$  is **positive**, we're done. Else we can inductively define  $n_2, n_3, \dots$  as well as  $E_2, E_3, \dots$

Explicitly, if  $E \setminus \bigcup_{i=1}^{k-1} E_i$  is not **positive**, let  $n_k$  be the least such that there exists  $E_k \subseteq E \setminus \bigcup_{i=1}^{k-1} E_i$  with  $\nu(E_k) < -1/n_k$ .

Note then that if  $n_k \geq 2$ , for all  $B \subseteq E \setminus \bigcup_{i=1}^{k-1} E_i$  we have that  $\nu(B) \geq -\frac{1}{n_k-1}$ .

Now let  $A = E \setminus \bigcup_{i=1}^{\infty} E_i$ . Since  $E = A \cup (\bigcup_i E_i)$  we have by **countable additivity** that

$$0 < \nu(E) = \nu(A) + \sum_{k=1}^{\infty} \nu(E_k) < \nu(A).$$

Furthermore,  $\nu(E), \nu(A)$  are both in  $(0, \infty)$ , and we see that

$$0 < \nu(E) \leq \nu(A) - \sum_{k=1}^{\infty} \frac{1}{n_k}.$$

Therefore, the sum on the right-hand side must converge, meaning that  $1/n_k \rightarrow 0$  as  $k \rightarrow \infty$ . That is  $\lim_{k \rightarrow \infty} n_k = \infty$ .

Now if  $B \subseteq A$ , then  $B \subseteq E \setminus \bigcup_{i=1}^{\infty} E_i$ . Therefore,  $B \subseteq E \setminus \bigcup_{i=1}^{k-1} E_i$ . By the note above, for large enough  $k$  such that  $n_k \geq 2$  we have

$$\nu(B) \geq \frac{-1}{n_k - 1},$$

then taking  $k \rightarrow \infty$  we have  $\nu(B) \geq 0$ , and so  $A$  is a **positive** set as desired. ■


**Remark.** Notice that the same is true for  $-\infty < \nu(E) < 0$ .

**Theorem 6.1.1 (Hahn decomposition theorem).** If  $\nu$  is a **signed measure** on  $(X, \mathcal{A})$ , then there exist  $P, N \in \mathcal{A}$  such that

$$P \cap N = \emptyset, \quad P \cup N = X,$$

where  $P$  is **positive for  $\nu$** ,  $N$  is **negative for  $\nu$** .

Furthermore, if  $P', N'$  are another such pair, then  $P \triangle P' (= N \triangle N')$  is **null** for  $\nu$ .



## Lecture 29: Hahn and Jordan Decomposition Theorem

We now prove [Theorem 6.1.1](#).

23 Mar. 11:00

**Proof of Theorem 6.1.1.** We first show the uniqueness. We see that  $P \setminus P' \subseteq P, P \setminus P' \subseteq N'$ . Thus,  $P \setminus P' \subseteq P \cap N'$  is both [positive](#) and [negative](#), hence  $P \setminus P'$  is [null](#). Similarly, for  $P' \setminus P$ , and then their union  $P \triangle P'$  is [null](#) as well.

To show the existence, without loss of generality suppose  $\nu: \mathcal{A} \rightarrow [-\infty, \infty)$ . If not, consider  $-\nu$ . Let

$$s := \sup\{\nu(E) \mid E \in \mathcal{A} \text{ is a } \text{positive set}\},$$

which is a nonempty supremum because  $\emptyset$  is [positive](#). Then there exist  $P_1, P_2, \dots$  [positive sets](#) such that  $\lim_{n \rightarrow \infty} \nu(P_n) = s$ .

Then we have that  $P = \bigcup_n P_n$  is [positive](#) by [Lemma 6.1.2](#). We then have  $\nu(P) \leq s$  and  $\nu(P) = \nu(P_n) + \nu(P \setminus P_n) \geq \nu(P_n)$ . Thus,

$$\nu(P) \geq \lim_{n \rightarrow \infty} \nu(P_n) = s.$$

Hence,  $\nu(P) = s$  and the supremum is in fact a max. We then know that  $s = \nu(P) < \infty$  because  $\nu$  does not attain the value infinity.

Now let  $N = X \setminus P$ . We claim that  $N$  is [negative](#). If not then there exists a [measurable](#)  $E \subseteq N$  with  $\nu(E) > 0$ . By assumption,  $\nu(E) < \infty$ . Then  $0 < \nu(E) < \infty$ , so by [Lemma 6.1.3](#) there exists a [measurable](#)  $A \subseteq E$  such that  $A$  is [positive](#) and  $\nu(A) > 0$ .

But we then know that

$$\nu(P \cup A) = \nu(P) + \nu(A) > \nu(P)$$

which is a contradiction since  $P \cup A$  is a [positive set](#), and  $\nu(P)$  is maximal. Therefore,  $N$  is [negative](#), and the theorem holds.

Finally, if  $P', N'$  is another pair of sets as in the statement of the theorem, we have

$$P \setminus P' \subseteq P, \quad P \setminus P' \subseteq N',$$

so that  $P \setminus P'$  is both positive and negative, hence null; likewise for  $P' \setminus P$ . ■

**Definition 6.1.5 (Singular).** If  $\mu, \nu$  are [signed measures](#) on  $(X, \mathcal{A})$ , then we say  $\mu$  and  $\nu$  are *singular to each other*, denoted as  $\mu \perp \nu$ , if there exists  $E, F \in \mathcal{A}$  such that  $E \cap F = \emptyset, E \cup F = X$ ,  $F$  is [null](#) for  $\mu$ ,  $E$  is [null](#) for  $\nu$ .

**Example.** Consider  $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$  with

- The [Lebesgue measure](#)  $m$ .
- The [Cantor measure](#)  $\mu_C$  induced by the [Cantor function](#).
- The [discrete measure](#)  $\mu_D = \delta_1 + 2\delta_{-1}$ .

We then see that (a)  $m \perp \mu_D$ . (b)  $m \perp \mu_C$ . (c)  $\mu_C \perp \mu_D$ .

**Proof.** We see them as follows.

- Take  $E = \mathbb{R} \setminus \{-1, 1\}, F = \{-1, 1\}$  to see that  $m \perp \mu_D$ .
- Take  $E = \mathbb{R} \setminus K$  and  $F = K$  where  $K$  is the [Cantor set](#) to see that  $m \perp \mu_C$ .
- We can also see that  $\mu_C \perp \mu_D$ .

⊛

**Theorem 6.1.2 (Jordan decomposition theorem).** Let  $\nu$  be a [signed measure](#) on  $(X, \mathcal{A})$ . Then there

exists unique **positive measures**  $\nu^+, \nu^-$  on  $(X, \mathcal{A})$  such that for all  $E \in \mathcal{A}$  we have

$$\nu(E) = \nu^+(E) - \nu^-(E)$$

and  $\nu^+ \perp \nu^-$ .

**Proof.** For existence, we take  $\nu^+(E) := \nu(E \cap P)$ ,  $\nu^-(E) := -\nu(E \cap N)$  where  $P, N$  is the **Hahn decomposition** of  $X$ .

If there exists  $\mu^+, \mu^-$  such that  $\nu = \mu^+ + \mu^-$  and  $\mu^+ \perp \mu^-$ , let  $E, F \in \mathcal{A}$  be such that  $E \cap F = \emptyset$ ,  $E \cup F = X$ , and  $\mu^+(F) = \mu^-(E) = 0$ . Then we have that  $X = E \cup F$  is another **Hahn decomposition** for  $\nu$ , so  $P \triangle E$  is  $\nu$ -null. Therefore, for any  $A \in \mathcal{A}$ ,  $\mu^+(A) = \mu^+(A \cap E) = \nu(A \cap E) = \nu(A \cap P) = \nu^+(A)$ , hence  $\mu^+ = \nu^+$ . Likewise, we have  $\nu^- = \mu^-$ . ■

## Lecture 30: Absolutely Continuous Measures

**Example.** For an example of **Theorem 6.1.2**, let  $(X, \mathcal{A}, \mu)$  be a **measure space**,  $f: X \rightarrow \overline{\mathbb{R}}$ , and  $\nu(E) = \int_E f d\mu$ . Then

$$\nu^+(E) = \int_E f^+ d\mu, \quad \nu^-(E) = \int_E f^- d\mu.$$

25 Mar. 11:00

**Definition.** Given a **signed measure**  $\nu$  on  $(X, \mathcal{A})$  and its **Jordan decomposition**  $\nu = \nu^+ - \nu^-$ .

**Definition 6.1.6** (Positive variation). We call  $\nu^+$  the *positive variation* of  $\nu$ .

**Definition 6.1.7** (Negative variation). We call  $\nu^-$  the *negative variation* of  $\nu$ .

**Definition 6.1.8** (Total variation). The *total variation measure* of  $\nu$ , denoted as  $|\nu|$ , is defined as  $|\nu| := \nu^+ + \nu^-$ .

**Remark.** There is always a **positive measure** on  $X$ .

**Proof.** Consider the **total variation**  $|\nu|$  for an arbitrary **signed measure**  $\nu$ . ⊛

**Example.** In the above **example**,  $|\nu|(E) = \int_E |f| d\mu$ .

**Lemma 6.1.4.** We have the following

- (a)  $|\nu(E)| \leq |\nu|(E)$ .
- (b)  $E$  is  $\nu$ -null if and only if  $E$  is  $|\nu|$ -null.
- (c) If  $\kappa$  is another **signed measure**, then  $\kappa \perp \nu$  if and only if  $\kappa \perp |\nu|$  if and only if  $\kappa \perp \nu^+$  and  $\kappa \perp \nu^-$ .

**Proof.** ■

DIY

**Definition 6.1.9** (Finite signed measure). A **signed measure**  $\nu$  is *finite* if  $|\nu|$  is a **finite measure**, and similarly for  $\sigma$ -finite.

**Remark.** This holds if and only if  $\nu^+, \nu^-$  are both **finite** (resp.  $\sigma$ -finite) **measures**.

## 6.2 Absolutely Continuous Measures

**Definition 6.2.1** (Absolutely continuous measure). Let  $\mu$  be a **positive measure**,  $\nu$  be a **signed measure**, both on  $(X, \mathcal{A})$ . We say that  $\nu$  is *absolutely continuous with respect to*  $\mu$ , denoted as  $\nu \ll \mu$ , provided that for all  $E \in \mathcal{A}$ ,  $\mu(E) = 0$  implies  $\nu(E) = 0$ .

**Remark.** This is equivalent to every  $\mu$ -null set being  $\nu$ -null.

**Example.** If  $(X, \mathcal{A}, \mu)$ ,  $f: X \rightarrow \overline{\mathbb{R}}$ ,  $\nu(E) = \int_E f d\mu$ , then  $\nu \ll \mu$ .

**Notation.**  $d\nu = f d\mu$  means  $\nu$  is a **signed measure** defined by

$$\nu(E) = \int_E f d\mu.$$

**Lemma 6.2.1.** If  $\mu$  is a **positive measure**,  $\nu$  is a **signed measure** on  $(X, \mathcal{A})$ , then

- (a)  $\nu \ll \mu$  if and only if  $|\nu| \ll \mu$  if and only if  $\nu^+ \ll \mu$  and  $\nu^- \ll \mu$ .
- (b)  $\nu \ll \mu$  and  $\nu \perp \mu$  implies  $\nu = 0$ .

**Proof.** For (b), write  $X = A \cup B$ ,  $A \cap B = \emptyset$ ,  $A$   $\mu$ -null,  $B$   $\nu$ -null. Then

$$\nu(E) = \nu(E \cap A) + \nu(E \cap B) = \nu(E \cap A).$$

Then  $E \cap A \subseteq A$ , so  $\nu(E \cap A) = 0$ . By **absolute continuity**,  $\nu(E \cap A) = 0$ , thus  $\nu(E) = 0$ . ■

DIY (a)

**Theorem 6.2.1** (Radon-Nikodym theorem). Suppose  $\mu$  is a  **$\sigma$ -finite positive measure**,  $\nu$  is a  **$\sigma$ -finite signed measure**, and suppose  $\nu \ll \mu$ . Then there exists  $f: X \rightarrow \overline{\mathbb{R}}$  such that  $d\nu = f d\mu$ , in other words  $\nu(E) = \int_E f d\mu$ .

If  $g$  is another such function with  $d\nu = g d\mu$  then  $f = g$   $\mu$ -a.e..

**Proof.** We'll prove a more general form called **Lebesgue Radon Nikodym theorem**, which is a more general theorem compare to this theorem. ■

**Definition 6.2.2** (Radon-Nikodym derivative). Suppose  $\nu \ll \mu$ . The *Radon-Nikodym derivative of  $\nu$  with respect to  $\mu$*  is a function

$$\frac{d\nu}{d\mu}: X \rightarrow \overline{\mathbb{R}}$$

such that

$$\nu(E) = \int_E \frac{d\nu}{d\mu} d\mu$$

for all  $E \in \mathcal{A}$ .

**Remark.** i.e. we have  $d\nu = \frac{d\nu}{d\mu} d\mu$  in our notation.

**Note.** By **Theorem 6.2.1**, such a function exists and is unique up to equivalence  $\mu$ -a.e. in the  **$\sigma$ -finite** case.

**Example.** Let  $F(x) = e^{2x}: \mathbb{R} \rightarrow \mathbb{R}$ , then

$$\frac{dF}{dm} = 2e^{2x}.$$

**Proof.** Since  $F$  is continuous and strictly increasing, so we may define a [Lebesgue-Stieltjes measure](#)  $\mu_F$  on  $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ .

This is defined to be the unique [locally finite measure](#) satisfying

$$\mu_F([a, b]) = F(b) - F(a) = e^{2b} - e^{2a}.$$

Then one can check that

$$\mu_F(E) = \int_E 2e^{2x} dx.$$

By uniqueness and the classical [fundamental theorem of calculus](#), since the right-hand side is a [locally finite Borel measure](#), and  $\kappa([a, b]) = e^{2b} - e^{2a}$ , thus  $\mu_F = \kappa$ . Therefore,  $\mu_F \ll m$  and we have

$$\frac{d\mu_F}{dm} = 2e^{2x} = \frac{dF}{dx}.$$

⊛

**Example.** Let  $\mu_C = C(X): \mathbb{R} \rightarrow \mathbb{R}$  be the [Cantor function](#). Then  $C'(x) = 0$  outside the [Cantor set](#), but we don't always have

$$\mu_C(E) \neq \int_E 0 dx.$$

So the candidate derivative is 0, but this fails.

**Proof.** In particular,

$$C(b) - C(a) \neq \int_a^b C'(x) dx.$$

In fact,  $\mu_C \not\ll m$  because  $\mu_C \perp m$  and  $\mu_C \neq 0$ .

Thus, the existence of a derivative [almost everywhere](#) and continuity is not enough to guarantee a version of the [fundamental theorem of calculus](#) to hold. ⊛

## Lecture 31: Lebesgue-Radon-Nikodym Theorem

**Lemma 6.2.2.** Let  $\mu, \nu$  be [finite positive measures](#) on  $(X, \mathcal{A})$ . Then either

- $\nu \perp \mu$ .
- There exists an  $\epsilon > 0$ , an  $F \in \mathcal{A}$  such that  $\mu(F) > 0$  and  $F$  is a [positive set for the measure](#)  $\nu - \epsilon\mu$ , i.e., for all  $G \subseteq F$ ,  $\nu(G) \geq \epsilon\mu(G)$ .

**Proof.** Let  $\kappa_n = \nu - (1/n)\mu$ . By [Theorem 6.1.1](#) we have  $X = P_n \cup N_n$  for  $P_n$  [positive](#),  $N_n$  [negative](#) for  $\kappa_n$ . Also, we let  $P = \bigcup_n P_n$ ,  $N = \bigcap_n N_n = X \setminus P$ , then  $X = P \cup N$ .

We see that for any  $n$  we have  $\kappa_n(N) \leq 0$  because  $N \subseteq N_n$ . Thus,

$$0 \leq \nu(N) \leq \frac{1}{n}\mu(N),$$

which implies  $\nu(N) = 0$ . Because  $\nu$  is [positive](#) for any  $N' \subseteq N$  we have  $0 \leq \nu(N') \leq \nu(N)$ , and thus  $\nu(N') = 0$ . This shows  $N$  is [null](#) for  $\nu$ . Now, we see that

- If  $\mu(P) = 0$ , then  $\nu \perp \mu$ .
- If  $\mu(P) \neq 0$ , then we have  $\mu(P) > 0$  hence  $\mu(P_n) > 0$  for some  $n$ . With  $F = P_n$  and  $\epsilon = 1/n$ , then  $F$  is a [positive set](#) for  $\kappa_n = \nu - (1/n)\mu$  as desired. ■

28 Mar. 11:00

**Theorem 6.2.2** (Lebesgue-Radon-Nikodym theorem). Let  $\mu$  be a  $\sigma$ -finite positive measure,  $\nu$  a  $\sigma$ -finite signed measure on  $(X, \mathcal{A})$ . Then there are unique  $\sigma$ -finite signed measures  $\lambda, \rho$  on  $(X, \mathcal{A})$  such that

$$\lambda \perp \mu, \quad \rho \ll \mu, \quad \nu = \lambda + \rho.$$

Furthermore, there exists a measurable function  $f: X \rightarrow \mathbb{R}$  such that  $d\rho = f d\mu$ .<sup>a</sup> And if there is another  $g$  such that  $d\rho = g d\mu$ , then  $f = g$   $\mu$ -a.e.

<sup>a</sup>That is for all  $E \in \mathcal{A}$ ,  $\rho(E) = \int_E f d\mu$ .

**Proof.** We prove it step by step.

- (a) Assume  $\mu, \nu$  are finite positive measures. We first prove the existence of  $\lambda, f$ , and  $d\rho = f d\mu$ . Let

$$\begin{aligned} \mathcal{F} &= \left\{ g: X \rightarrow [0, \infty] \mid \int_E g d\mu \leq \nu(E), \forall E \in \mathcal{A} \right\} \\ &= \{g: X \rightarrow [0, \infty] \mid d\nu - g d\mu \text{ is a positive measure}\}. \end{aligned}$$

This set is nonempty since  $g = 0 \in \mathcal{F}$ . Let  $s = \sup\{\int_X g d\mu \mid g \in \mathcal{F}\}$ .

**Claim.** There is an  $f \in \mathcal{F}$  such that  $s = \int_X f d\mu$ .

**Proof.** If  $g, h \in \mathcal{F}$ , we can define  $u(x) = \max\{g(x), h(x)\}$ , then  $u \in \mathcal{F}$ . This can be seen by letting  $A = \{x \mid g(x) \geq h(x)\}$ , then

$$\int_E u d\mu = \int_{E \cap A} g d\mu + \int_{E \cap A^c} h d\mu \leq \nu(E \cap A) + \nu(E \cap A^c) = \nu(E).$$

There exist measurable functions  $g_1, g_2, \dots \in \mathcal{F}$  such that

$$\lim_{n \rightarrow \infty} \int_X g_n d\mu = s.$$

We can replace  $g_2$  by  $\max(g_1, g_2)$ ,  $g_3$  by  $\max(g_1, g_2, g_3)$ . Generally,

$$g_n \leftarrow \max(g_1, g_2, \dots, g_n),$$

so that we may assume  $0 \leq g_1 \leq g_2 \leq \dots$

Then we still know that  $\lim_{n \rightarrow \infty} \int_X g_n d\mu = s$ , as all the relevant integrals are bounded above by  $s$ . Now let  $f(x) = \sup_n g_n(x) = \lim_{n \rightarrow \infty} g_n(x)$ , by monotone convergence theorem,

$$\int_E f d\mu = \lim_{n \rightarrow \infty} \int_E g_n d\mu \leq \nu(E).$$

Thus,  $f \in \mathcal{F}$ , and when  $E = X$  we get  $\int_X f d\mu = s$  as desired. ⊛

Let  $\rho(E) := \int_E f d\mu$ , then we of course have  $\rho \ll \mu$ , and also, we know

$$0 \leq \rho(X) = \int_X f d\mu \leq \nu(X) < \infty.$$

Thus,  $\rho$  is a finite positive measure, so we can define  $\lambda(E) := \nu(E) - \rho(E)$ , then

$$\lambda(E) = \nu(E) - \int_E f d\mu \geq 0$$

because  $f \in \mathcal{F}$ . Thus,  $\lambda$  is also a positive measure, and  $\lambda(X) \leq \nu(X) < \infty$ . It remains to show the following.

**Claim.**  $\lambda \perp \mu$ .



**Proof.** Suppose not, by Lemma 6.2.2, there exists  $\epsilon > 0$ ,  $F \in \mathcal{A}$  such that  $\mu(F) > 0$  and  $F$  is a positive set for  $\lambda - \epsilon\mu$ .

Then this says that  $d\lambda - \epsilon\mathbb{1}_F d\mu$  is a positive measure, that is,

$$d\nu - f d\mu - \epsilon\mathbb{1}_F d\mu$$

is a positive measure. But, this will break maximality of  $f$ , specifically, let  $g(x) = f(x) + \epsilon\mathbb{1}_F(x)$ . Then for all  $E \in \mathcal{A}$  we have

$$\begin{aligned} \int_E g d\mu &= \int_E f d\mu + \epsilon\mu(E \cap F) \\ &= \nu(E) - \lambda(E) + \epsilon\mu(E \cap F) \\ &\leq \nu(E) - \lambda(E \cap F) + \epsilon\mu(E \cap F) \leq \nu(E) \end{aligned}$$

since  $\lambda(E \cap F) - \epsilon\mu(E \cap F) \geq 0$ . Thus,  $g \in \mathcal{F}$ . We then see that

$$s \geq \int_X g d\mu = \int_X f d\mu + \int_X \epsilon\mathbb{1}_F d\mu = s + \epsilon\mu(F) > s,$$

which is a contradiction.  $\otimes$

We see that the existence of  $\lambda$ ,  $f$ , and  $d\rho = f d\mu$  is proved. As for uniqueness, if there are  $\lambda'$  and  $f'$  such that  $d\nu = d\lambda' + f' d\mu$ , we then have

$$d\lambda - d\lambda' = (f' - f) d\mu.$$

But we see that  $\lambda - \lambda' \perp \mu$  while  $(f' - f) d\mu \ll d\mu$ , hence

$$d\lambda - d\lambda' = (f' - f) d\mu = 0,$$

so  $\lambda = \lambda'$  and  $f = f'$   $\mu$ -a.e. by Proposition 2.3.1.

- (b) Suppose that  $\mu$  and  $\nu$  are  $\sigma$ -finite measures. Then  $X$  is a countable disjoint union of  $\mu$ -finite sets and a countable disjoint union of  $\nu$ -finite sets. By taking intersections of these we obtain a disjoint sequence  $\{A_j\} \subset \mathcal{A}$  such that  $\mu(A_j)$  and  $\nu(A_j)$  are finite for all  $j$  and  $X = \bigcup_j A_j$ . Define  $\mu_j(E) = \mu(E \cap A_j)$  and  $\nu_j(E) = \nu(E \cap A_j)$ , then by the reasoning above, for each  $j$  we have

$$d\nu_j = d\lambda_j + f_j d\mu_j$$

where  $\lambda_j \perp \mu_j$ . Since  $\mu_j(A_j^c) = \nu_j(A_j^c) = 0$ , we have

$$\lambda_j(A_j^c) = \nu_j(A_j^c) - \int_{A_j^c} f_j d\mu_j = 0,$$

and we may assume that  $f_j = 0$  on  $A_j^c$ . Let  $\lambda = \sum_j \lambda_j$  and  $f = \sum_j f_j$ , we then have

$$d\nu = d\lambda + f d\mu, \quad \lambda \perp \mu,$$

and  $d\lambda$  and  $f d\mu$  are  $\sigma$ -finite, as desired. As for uniqueness, it's the same as for the first case.

- (c) We now consider the general case. If  $\nu$  is a signed measure, we apply the preceding argument to  $\nu^+$  and  $\nu^-$  and subtract the results.

■

**Remark.** Notationally, we may write  $d\nu = d\lambda + f d\mu$ , where  $d\lambda$  and  $d\mu$  are singular to each other.

## Lecture 32: Lebesgue Differentiation Theorem for Regular Borel Measures

We now do an example to illustrate [Theorem 6.2.2](#).

30 Mar. 11:00

**Example.** Let  $\mu = m$ ,  $\nu = \mu_F$  ([Lebesgue-Stieltjes measure](#) for  $F$ ). We'll define  $F(x)$  by

$$F(x) = \begin{cases} e^{3x}, & \text{if } x \leq 0; \\ 1, & \text{if } 0 < x < 1; \\ 5, & \text{if } x \geq 1. \end{cases}$$

Then we have

$$\mu_F(E) = \int_{E \cap \mathbb{R}_{<0}} 3e^{3x} dx + 4\delta_1(E).$$

**Proof.** It is enough to check on  $(-\infty, x]$  because these are [locally finite Borel measures](#) on  $\mathbb{R}$ . Then we have

$$\mu_F = d\rho + d\lambda = f dm + d\lambda$$

where  $f = \mathbb{1}_{\mathbb{R}_{<0}} 3e^{3x}$  and  $\lambda = 4\delta_1$ ,  $\lambda \perp m$ . \*

Specifically, we call such a decomposition *Lebesgue decomposition* of  $\nu$  with respect to  $\mu$ . Now, with the condition  $\nu \ll \mu$ , [Theorem 6.2.2](#) implies that  $d\nu = f d\mu$  for some  $f$ , which is exactly the statement of [Theorem 6.2.1](#). And, it should be clear now that the definition of [Radon Nikodym derivative](#) of  $\nu$  with respect to  $\mu$ , denoted as  $d\nu/d\mu$ , is just  $f$  in this case.

**As previously seen.** If  $\nu = \nu^+ - \nu^-$ , we defined the [total variation](#)  $|\nu| = \nu^+ + \nu^-$ . Then we have  $|\nu(E)| \leq |\nu|(E)$ .

### 6.3 Lebesgue Differentiation Theorem for Regular Borel Measures

**Definition 6.3.1** (Regular). A Borel [signed measure](#)  $\nu$  on  $\mathbb{R}^d$  is called *regular* if

- (compact finite)  $|\nu|(K) < \infty$  for all compact  $K$ .
- (outer regularity) We have [outer regularity](#)

$$|\nu|(E) = \inf\{|\nu|(U) \mid \text{open } U \supseteq E\}$$

for every [Borel set](#)  $E$ .

**Example.** We see that

- Any [Lebesgue-Stieltjes measure](#) on  $\mathbb{R}$  has this property from [Theorem 1.7.1](#), so is the difference between two of them (at least if one of them is [finite](#)).
- The [Lebesgue measure](#) on  $\mathbb{R}^d$  is [regular](#).

**Note.** From [compact finiteness](#), if  $\nu$  is [regular](#) then it is  $\sigma$ -finite.

**Lemma 6.3.1.**  $f \in L^1_{\text{loc}}(\mathbb{R}^d)$  if and only if  $d\nu = f dm$  is [regular](#).

**Proof.** We prove this in two directions.

( $\Leftarrow$ ) Suppose  $d\nu = f dm$  is **regular**. Then

$$|\nu|(K) = \int_K |f| dm < \infty$$

for all compact  $K$ , thus  $f \in L^1_{\text{loc}}(\mathbb{R}^d)$ .

( $\Rightarrow$ ) Suppose  $f \in L^1_{\text{loc}}(\mathbb{R}^d)$ . This condition is clearly equivalent to **compact finiteness**. If this holds, then the **outer regularity** may be verified directly as follows. Suppose that  $E$  is a bounded **Borel set**. Given  $\delta > 0$ , by **Theorem 3.5.1**, there is a bounded open  $U \supset E$  such that  $m(U) < m(E) + \delta$  and hence  $m(U \setminus E) < \delta$ . But then, given  $\epsilon > 0$ , there is<sup>a</sup> an open  $U \supset E$  such that

$$\int_{U \setminus E} f dm < \epsilon$$

and hence

$$\int_U f dm < \int_E f dm + \epsilon.$$

The case of unbounded  $E$  follows easily by writing  $E = \bigcup_{j=1}^{\infty} E_j$  where  $E_j$  is bounded and finding an open  $U_j \supset E_j$  such that

$$\int_{U_j \setminus E_j} f dm < \epsilon 2^{-j}.$$

■

<sup>a</sup>This follows from [FF99] Corollary 3.6.

**As previously seen.** Recall the **Lebesgue differentiation theorem**, here we had that if  $f \in L^1_{\text{loc}}(\mathbb{R}^d)$  implies that for Lebesgue **almost every**  $x$ ,

$$\lim_{r \rightarrow 0} \frac{1}{m(E_r)} \int_{E_r} f(y) dy = f(x)$$

for any  $\{E_r\}$  **shrinks nicely** to  $x$ .

**Corollary 6.3.1.** Let  $\rho$  be a **regular signed Borel measure** on  $\mathbb{R}^d$ . Suppose  $\rho \ll m$ . Then  $d\rho = f dm$  for some  $f \in L^1_{\text{loc}}(\mathbb{R}^d)$ , so then for Lebesgue **almost every**  $x$  we have

$$\lim_{r \rightarrow 0} \frac{1}{m(E_r)} \int_{E_r} f(y) dy = f(x).$$

Writing this nicely, using established notation, this is

$$\lim_{r \rightarrow 0} \frac{\rho(E_r)}{m(E_r)} = \frac{d\rho}{dm}(x)$$

for every  $\{E_r\}$  **shrinks nicely** to  $x$ .

**Proposition 6.3.1.** Let  $\lambda$  be a **regular positive Borel measure** on  $\mathbb{R}^d$ . Suppose  $\lambda \perp m$ . Then for **Lebesgue almost every**  $x$ , we have

$$\lim_{r \rightarrow 0} \frac{\lambda(E_r)}{m(E_r)} = 0$$

for every  $\{E_r\}$  **shrinking to  $x$  nicely** (equivalently, **shrinking to 0 nicely**).

**Proof.** It is enough to consider  $E_r = B(x, r)$ . We wish to prove that

$$G := \left\{ x \mid \limsup_{r \rightarrow 0} \frac{\lambda(E_r)}{m(E_r)} \neq 0 \right\} = \bigcup_{n=1}^{\infty} G_n$$

where

$$G_n := \left\{ x \mid \limsup_{r \rightarrow 0} \frac{\lambda(E_r)}{m(E_r)} > \frac{1}{n} \right\}$$

such that  $m(G) = 0$ . We see that it suffices to show  $m(G_n) = 0$  for all  $n$ . Since  $\lambda \perp m$ , so we know there exists  $A, B$  such that  $\mathbb{R}^d = A \cup B$  disjoint with  $\lambda(A) = 0$ ,  $m(B) = 0$ . Thus, it suffices to show  $m(G_n \cap A) = 0$ .

**Note.** Alternatively, we can simply define  $G_n$  over  $A$  instead of  $\mathbb{R}^d$ , as in Folland[FF99].

**Claim.** Given a  $A$  and  $B$  defined above induced from Theorem 6.2.2,  $m(G_n \cap A) = 0$  for all  $n$

**Proof.** Fix  $\epsilon > 0$ , since  $\lambda$  is regular, there exists an open set  $U \supseteq A$  such that  $\lambda(U) \leq \lambda(A) + \epsilon = \epsilon$ . We see that for every  $x \in G_n \cap A$ , there is an  $r_x > 0$  such that  $\lambda(B(x, r_x))/m(B(x, r_x)) > 1/n$  where  $B(x, r_x) \subseteq U$ .

Let  $K \subseteq G_n \cap A$ , compact. Then  $K \subseteq \bigcup_{x \in K} B(x, r_x)$ . By compactness, we can take a finite sub-cover, and then use Lemma 4.1.1 to find disjoint  $B_1, B_2, \dots, B_N$  such that each of  $B_i$  is in the form of  $B(x_i, r_{x_i})$  and  $K \subseteq \bigcup_i 3B_i$ . Therefore,

$$m(K) \leq 3^d \sum_{i=1}^N m(B_i) \leq 3^d n \sum_{i=1}^N \lambda(B_i) = 3^d n \lambda \left( \bigcup_{i=1}^N B_i \right) \leq 3^d n \lambda(U) = 3^d n \epsilon.$$

By inner regularity,  $m(G_n \cap A) \leq 3^d n \epsilon$  for any  $\epsilon > 0$ . Taking  $\epsilon \rightarrow 0$  yields  $m(G_n \cap A) = 0$ , so then  $m(G_n) = 0$  as desired.  $\otimes$

## Lecture 33: Monotone Differentiation Theorem

As previously seen. We have that if  $\rho \ll m$  is regular then

$$\lim_{r \rightarrow 0} \frac{\rho(E_r)}{m(E_r)} = \frac{d\rho}{dm}(x)$$

for Lebesgue almost every  $x$ , where  $\{E_r\}$  shrinks nicely to  $x$ . Likewise, if  $\lambda \perp m$  regular (positive measure) then

$$\lim_{r \rightarrow 0} \frac{\lambda(E_r)}{m(E_r)} = 0$$

for Lebesgue almost every  $x$ , where  $\{E_r\}$  shrinks nicely to  $x$ .

From this, we can easily deduce the following important result.

**Theorem 6.3.1** (Lebesgue differentiation theorem for regular measures). Let  $\nu$  be a regular Borel signed measure on  $\mathbb{R}^d$ . Then  $d\nu = d\lambda + f dm$ ,  $\lambda \perp m$  by Theorem 6.2.2. Then for Lebesgue almost every  $x$ ,

$$\lim_{r \rightarrow 0} \frac{\nu(E_r)}{m(E_r)} = f(x)$$

for every  $\{E_r\}$  shrinks nicely to  $x$ .

1 Apr. 11:00

**Proof.** It must be checked that  $\nu$  **regular** implies  $\lambda, f \, dm$  are **regular**. In particular, since  $f \in L^1_{\text{loc}}$ , so from **Theorem 4.2.1** and its corollary (**Corollary 4.2.1**, **Corollary 4.2.2**), we see that it suffices to show that if  $\lambda$  is **regular** and  $\lambda \perp m$ , then for Lebesgue a.e.  $x$ ,

Check!

$$\lim_{r \rightarrow 0} \frac{\lambda(E_r)}{m(E_r)} \rightarrow 0$$

when  $\{E_r\}$  **shrinks nicely to  $x$** . It also suffices to take  $E_r = B(r, x)$  and to assume that  $\lambda$  is **positive**, since for some  $\alpha > 0$ , we have

$$\left| \frac{\lambda(E_r)}{m(E_r)} \right| \leq \frac{|\lambda|(E_r)}{m(E_r)} \leq \frac{|\lambda|(B(r, x))}{m(E_r)} \leq \frac{|\lambda|(B(r, x))}{\alpha m(B(r, x))}.$$

Therefore, if  $|\lambda|(E_r)/m(E_r) \rightarrow 0$ , so does  $|\lambda(E_r)/m(E_r)|$ , hence  $\lambda(E_r)/m(E_r)$ . We see that the result then follows directly from **Proposition 6.3.1**. ■

## 6.4 Monotone Differentiation Theorem

We first formalize one ambiguous notation we used long time ago with discussing **distribution function**. Namely,  $F(x^+)$ ,  $F(x^-)$ .

**Definition.** For an  $F: \mathbb{R} \rightarrow \mathbb{R}$  that is monotonically increasing, we have the following.

**Definition 6.4.1** ( $F(x^+)$ ). We define  $F(x^+) = \lim_{y \rightarrow x^+} F(y)$ .

**Definition 6.4.2** ( $F(x^-)$ ). We define  $F(x^-) = \lim_{y \rightarrow x^-} F(y)$ .

**Remark.** If  $F$  is monotonically increasing, then  $F(x^+)$ ,  $F(x^-)$  exist and are

$$\inf_{y > x} F(y), \quad \sup_{y < x} F(y)$$

respectively since it's bounded below/above respectively by  $F(x)$ .

**Lemma 6.4.1.** If  $F: \mathbb{R} \rightarrow \mathbb{R}$  is monotonically increasing, then

$$D = \{x \in \mathbb{R} \mid F \text{ is discontinuous at } x\}$$

is a countable set.

**Proof.** Since  $F$  is monotonically increasing,  $x \in D$  if and only if  $F(x^+) > F(x^-)$ . For each  $x \in D$ , let  $I_x = (F(x^-), F(x^+))$ , not empty. Also, if  $x, y \in D$ ,  $x \neq y$ , then  $I_x, I_y$  are disjoint. Now, for  $|x| < N$ ,  $I_x$  lies in the interval  $(F(-N), F(N))$ . Hence,

$$\sum_{|x| < N} [F(x^+) - F(x^-)] \leq F(N) - F(-N) < \infty,$$

which implies that

$$D \cap (-N, N) = \{x \in (-N, N) \mid F(x^+) \neq F(x^-)\}$$

is countable. Since this is true for all  $N$ , the result follows. ■

**Theorem 6.4.1** (Monotone differentiation theorem). Let  $F$  be an increasing function, then

- $F$  is differentiable **Lebesgue almost everywhere**.

- $G(x) := F(x^+)^a$  is differentiable **almost everywhere**.
- $G' = F'$  **almost everywhere**

<sup>a</sup>Observe that  $G$  is increasing and right-continuous.

**Proof.** Start with  $G(x) := F(x^+)$ , which is increasing and right-continuous on  $\mathbb{R}$ . There is then a **Lebesgue-Stieltjes measure**  $\mu_G$  on  $\mathbb{R}$ , thus it is **regular** on  $\mathbb{R}$ . We see

$$\frac{G(x+h) - G(x)}{h} = \begin{cases} \frac{\mu_G((x, x+h])}{m((x, x+h])}, & \text{if } h > 0; \\ \frac{\mu_G((x+h, x])}{m((x+h, x])}, & \text{if } h < 0. \end{cases}$$

Note that both  $\{(x, x+h]\}$  and  $\{(x+h, x]\}$  **shrink nicely** to  $x$  as  $|h| \rightarrow 0$ . By **Theorem 6.3.1** (since these **shrink nicely**), we then know that these both converge for **Lebesgue almost every**  $x$  to some common limit  $f(x)$ . Hence,  $G'$  exists **Lebesgue almost everywhere**. We now show that by defining  $H := G - F$ ,  $H'$  exists and equals zero **a.e.**

Observe that  $H(x) = G(x) - F(x) \geq 0$ , and we see that

$$\{x \mid H(x) > 0\} \subseteq \{x \mid F \text{ is discontinuous at } x\}.$$

The latter set is then countable by **Lemma 6.4.1**, hence we can write  $\{x \mid H(x) > 0\} = \{x_n\}$ . Then let

$$\mu := \sum_n H(x_n) \delta_{x_n}.$$

This is a **Borel measure**, so we check if it is **locally finite**. Indeed, since

$$\mu((-N, N)) = \sum_{-N < x_n < N} H(x_n) \leq G(N) - F(-N) < \infty,$$

where checking the inequality just consists of seeing that the intervals  $(F(x_n), G(x_n))$  are disjoint and is a subset of  $(F(-N), G(N))$ , so

$$\sum_{-N < x_n < N} H(x_n) = \mu \left( \bigcup_n (F(x_n), G(x_n)) \right) \leq \mu((F(-N), G(N))).$$

Thus,  $\mu$  is a **Lebesgue-Stieltjes measure** on  $\mathbb{R}$ , so it is **regular**.

**Remark.** Special to  $\mathbb{R}$ , we have that for a **measure**,

$$\text{locally finite Borel} \Rightarrow \text{Lebesgue-Stieltjes} \Rightarrow \text{regular} \Rightarrow \text{outer regularity}.$$

Also, we have  $\mu \perp m$  since  $m(E) = \mu(E^c) = 0$  where  $E = \{x_n\}$ . Then we have that

$$\left| \frac{H(x+h) - H(x)}{h} \right| \leq \frac{H(x+h) + H(x)}{|h|} \leq \frac{\mu((x-2h, x+2h))}{|h|},$$

which goes to 0 for **Lebesgue almost every**  $x$  by **Theorem 6.3.1** and that  $\mu \perp m$ .

Thus,  $H$  is differentiable **almost everywhere** and  $H' = 0$  **almost everywhere**, which implies  $F$  is differentiable **almost everywhere** and  $F' = G'$  **almost everywhere**. ■

**Proposition 6.4.1.** Suppose  $F$  is an increasing function, then  $F'$  exists **almost everywhere** and is **measurable**, then

$$\int_a^b F'(x) dx \leq F(b) - F(a).$$

**Example.** The inequality can't be made into equality in **Proposition 6.4.1** by the given condition.

**Proof.** Take  $F(x)$  as

$$F(x) := \begin{cases} 0, & \text{if } x \leq 0; \\ 1, & \text{if } x > 0. \end{cases}$$

Then  $F'(x) = 0$  almost everywhere. So

$$\int_{-1}^1 F'(x) \, dx = 0 < 1 = F(1) - F(-1).$$

\*)

**Example.** The inequality can't be made into equality in Proposition 6.4.1 even if  $F$  is continuous in addition.

**Proof.** Take  $F(x)$  to be the Cantor function. Then  $F'(x) = 0$  almost everywhere, but

$$\int_0^1 F'(x) \, dx = 0 < 1 = F(1) - F(0).$$

\*)

## Lecture 34: Functions of Bounded Variation

4 Apr. 11:00

**Proof of Proposition 6.4.1.** Let

$$G(x) := \begin{cases} F(a), & \text{if } x < a; \\ F(x), & \text{if } a \leq x \leq b; \\ F(b), & \text{if } x > b. \end{cases}$$

Then  $G$  is increasing. We define

$$g_n(x) = \frac{G(x + 1/n) - G(x)}{1/n} \rightarrow F'(x)$$

for almost every  $x \in [a, b]$ . We note that  $g_n(x) \geq 0$ . Theorem 2.2.2 tells us that

$$\int_a^b F'(x) \, dx = \int_a^b \liminf_{n \rightarrow \infty} g_n(x) \, dx \leq \liminf_{n \rightarrow \infty} \int_a^b g_n(x) \, dx.$$

We then evaluate

$$\begin{aligned} \int_a^b g_n(x) \, dx &= n \left( \int_{a+1/n}^{b+1/n} G(x) \, dx - \int_a^b G(x) \, dx \right) \\ &= n \left( \int_b^{b+1/n} G(x) \, dx - \int_a^{a+1/n} G(x) \, dx \right) \\ &\leq n \left( G\left(b + \frac{1}{n}\right) \cdot \frac{1}{n} - G(a) \cdot \frac{1}{n} \right) = F(b) - F(a). \end{aligned}$$

Therefore,

$$\int_a^b F'(x) \, dx \leq F(b) - F(a).$$

■

## 6.5 Functions of Bounded Variation

**Definition 6.5.1** (Total variation function). For  $F: \mathbb{R} \rightarrow \mathbb{R}$ , the *total variation function* of  $F$  is  $T_F: \mathbb{R} \rightarrow [0, \infty]$  defined by

$$T_F(x) = \sup \left\{ \sum_{i=1}^n |F(x_i) - F(x_{i-1})| \mid n \in \mathbb{N}, -\infty < x_0 < x_1 < \cdots < x_n = x \right\}.$$

**Lemma 6.5.1.**  $T_F(b)$  is equal to

$$T_F(a) + \sup \left\{ \sum_{i=1}^n |F(x_i) - F(x_{i-1})| \mid n \in \mathbb{N}, a = x_0 < x_1 < \cdots < x_n = b \right\}$$

if  $a < b$ .

**Proof.** The idea is that the sums in the Definition 6.1.8 of  $T_F$  are made bigger if the additional subdivision points  $x_j$  are added. Hence, if  $a < b$ ,  $T_F(b)$  is unaffected if we assume that  $a$  is always one of the subdivision points. ■

**Remark.**  $T_F$  is increasing.

**Definition 6.5.2** (Bounded variation). We say that  $F$  is of *bounded variation*, denoted as  $F \in \text{BV}$ , provided that

$$T_F(\infty) = \lim_{x \rightarrow \infty} T_F(x) < \infty.$$

Similarly,  $F \in \text{BV}([a, b])$  means that

$$\sup \left\{ \sum_{i=1}^n |F(x_i) - F(x_{i-1})| \mid n \in \mathbb{N}, a = x_0 < x_1 < \cdots < x_n = b \right\} < \infty.$$

**Remark.** We see the following.

- (a) If  $F$  is of *bounded variation*, then  $F$  is bounded.
- (b)  $F(x) = \sin x$  is not of *bounded variation*, but it is of *bounded variation* over any  $[a, b]$ .
- (c) For  $F(x)$  defined as

$$F(x) = \begin{cases} \frac{\sin x}{x}, & \text{if } x \neq 0; \\ 1, & \text{if } x = 0 \end{cases}$$

is not of *bounded variation* of  $[a, b]$  if  $a < 0 < b$  because the harmonic series does not converge.

Before we see more properties of *bounded variation* function, we introduce a useful characterization of a function.

**Definition 6.5.3** (Lipschitz). A function  $F: [a, b] \rightarrow \mathbb{C}$  is called *Lipschitz* if there exists an  $M \geq 0$  such that

$$|F(x) - F(y)| \leq M |x - y|.$$

**Remark.** We have the following.

- (a) If  $F, G$  are of *bounded variation*,  $\alpha F + \beta G$  are of *bounded variation* for  $\alpha, \beta \in \mathbb{C}$ .
- (b) If  $F$  is increasing and bounded, then  $F$  is a function of *bounded variation*.



(c) If  $F$  is **Lipschitz** on  $[a, b]$ , then  $F \in \text{BV}([a, b])$ .

(d) If  $F$  is differentiable, and  $F'$  is bounded on  $[a, b]$ , then  $F$  is **Lipschitz** from mean value theorem, so it is in  $\text{BV}([a, b])$ .

In particular, we have the following.

**Remark.** If  $F(x) = \int_{-\infty}^x f(t) dt$  for  $f \in L^1(\mathbb{R})$ , then  $F \in \text{BV}$ .

**Proof.** We see this by

$$\sum_{i=1}^n |F(x_i) - F(x_{i-1})| = \sum_{i=1}^n \left| \int_{x_{i-1}}^{x_i} f(t) dt \right| \leq \sum_{i=1}^n \int_{x_{i-1}}^{x_i} |f(t)| dt = \int_{x_0}^{x_n} |f(t)| dt \leq \int_{-\infty}^{\infty} |f(t)| dt,$$

which is finite since  $f \in L^1(\mathbb{R})$ . \*

**Lemma 6.5.2.** If  $F \in \text{BV}$ , then  $T_F$  is bounded, increasing,  $T_F(-\infty) = 0$ .

**Proof.** Fix  $\epsilon > 0$  and for a  $x \in \mathbb{R}$ , there exists  $x_0 < \dots < x_n = x$  such that

$$\sum_{i=1}^n |F(x_i) - F(x_{i-1})| \geq T_F(x) - \epsilon.$$

From **Lemma 6.5.1**, we see that  $T_F(x) - T_F(x_0) \geq T_F(x) - \epsilon$ , and hence  $T_F(y) \leq \epsilon$  for  $y \leq x_0$ , so  $T_F(-\infty) = 0$ . ■

**Lemma 6.5.3.**  $F \in \text{BV}$ , then  $T_F \pm F$  are increasing and bounded.

**Proof.** Let  $x < y$  and fix  $\epsilon > 0$ , then there are points  $x_0 < x_1 < \dots < x_n = x$  such that

$$T_F(x) \leq \sum_{i=1}^n |F(x_i) - F(x_{i-1})| + \epsilon.$$

Furthermore,

$$T_F(y) \geq \sum_{i=1}^n |F(x_i) - F(x_{i-1})| + |F(y) - F(x)|.$$

Then, since  $\pm(F(y) - F(x)) \leq |F(y) - F(x)|$ , we have<sup>a</sup>

$$T_F(y) \pm (F(y) - F(x)) \geq \sum_{i=1}^n |F(x_i) - F(x_{i-1})| \geq T_F(x) - \epsilon,$$

hence

$$T_F(y) \pm F(y) \geq T_F(x) \pm F(x) - \epsilon.$$

Taking  $\epsilon \rightarrow 0$  yields the result. ■

<sup>a</sup>To be precise, we note that  $\pm(F(y) - F(x)) \geq -|F(y) - F(x)|$  and proceed.

With this, we have the following useful decomposition for any  $F \in \text{BV}$ .

**Definition 6.5.4 (Jordan decomposition).** For any  $F \in \text{BV}$ , the *Jordan decomposition* of  $F$  is

$$F = \frac{T_F + F}{2} - \frac{T_F - F}{2},$$

which is a difference of increasing and bounded functions.

Specifically, we give special names for the terms in the **Jordan decomposition** of  $F$ .

**Definition 6.5.5** (Positive variation). The *positive variation* of  $F$  is  $\frac{1}{2}(T_F + F)$ .

**Definition 6.5.6** (Negative variation). The *negative variation* of  $F$  is  $\frac{1}{2}(T_F - F)$ .

**Proof.** It's clear that the above decomposition holds, we just need to check that both terms are increasing and bounded functions. They are increasing from [Lemma 6.5.3](#). To show they are bounded, for  $y > x$ ,

$$T_F(y) \pm F(y) \geq T_F(x) \pm F(x)$$

implies that

$$|F(y) - F(x)| \leq T_F(y) - T_F(x) \leq T_F(\infty) - T_F(-\infty) < \infty,$$

so that  $F$ , and hence  $T_F \pm F$  is all bounded.  $\circledast$

**Note.** Do not be confused between [Definition 6.5.4](#) and [Theorem 6.1.2](#) for *signed measure*! Also, the *positive variation of a function* and *positive variation of a measure* is confusing, same for the negative case.

**Theorem 6.5.1.**  $F$  is of *bounded variation* if and only if  $F = F_1 - F_2$  for  $F_1, F_2$  increasing and bounded.

**Proof.** The forward implication is given by [Lemma 6.5.3](#) and [the Jordan decomposition](#), and the other direction is easy to check.  $\blacksquare$

Check!

**Corollary 6.5.1.**  $F \in \text{BV}$  implies that  $F$  is differentiable *almost everywhere*. Furthermore,

- (a)  $F(x^+), F(x^-)$  exist for all  $x$  as do  $F(-\infty), F(\infty)$ .
- (b) The set of discontinuities of  $F$  is countable.
- (c)  $G(x) = F(x^+)$  is differentiable and  $G' = F'$  *almost everywhere*.
- (d)  $F' \in L^1(\mathbb{R}, m)$  (i.e.  $F \in L^1_{\text{loc}}(\mathbb{R})$ ) for every  $a < b$ .

**Proof.**  $\blacksquare$

DIY

## Lecture 35: Continue on Functions of Bounded Variation

**Definition 6.5.7** (Normalized bounded variation). A function  $G \in \text{BV}$  is said to have *normalized bounded variation*, denoted as  $G \in \text{NBV}$  provided that  $G$  is right continuous and  $G(-\infty) = 0$ .

6 Apr. 11:00

**Example.** If  $F$  is increasing and bounded,  $F$  right continuous,  $F(-\infty) = 0$ .

$F(x) = \int_{-\infty}^x f(t) dt, f \in L^1(\mathbb{R})$ . Midterm gave  $F$  is uniformly continuous.

**Lemma 6.5.4.** If  $F \in \text{BV}$  is right continuous, then  $T_F \in \text{NBV}$ .

**Proof.**  $T_F$  is bounded, increasing, and satisfies  $T_F(-\infty) = 0$  by [Lemma 6.5.2](#). Thus,  $T_F \in \text{BV}$ . Hence, we just need to check that  $T_F$  is right continuous. Suppose not, then there is a point  $a \in \mathbb{R}$  such that  $c := T_F(a^+) - T_F(a) > 0$  since  $T_F$  is increasing.

Fix  $\epsilon > 0$ , since  $F(x)$  and  $g(x) := T_F(x^+)$  are right continuous, there exists a  $\delta > 0$  such that for  $y \in (a, a + \delta]$  we have  $|F(y) - F(a)| < \epsilon$  and  $|g(y) - g(a)| < \epsilon$ . We then have that

$$T_F(y) - T_F(a^+) \leq \underbrace{T_F(y^+) - T_F(a^+)}_{|g(y) - g(a)|} < \epsilon.$$

There exist  $a = x_0 < x_1 < \dots < x_n = a + \delta$  such that

$$\sum_{i=1}^n |F(x_i) - F(x_{i-1})| \geq T_F(a + \delta) - T_F(a) - \frac{c}{4} \geq T_F(a^+) - T_F(a) - \frac{c}{4} = \frac{3c}{4}.$$

From  $|F(x_1) - F(a)| < \epsilon$ , we have

$$\sum_{i=2}^n |F(x_i) - F(x_{i-1})| \geq \frac{3}{4}c - \epsilon.$$

From the same reason, there exist  $a = t_0 < \dots < t_k = x_1$  such that

$$\sum_{i=1}^k |F(t_i) - F(t_{i-1})| \geq T_F(x_1) - T_F(a) - \frac{c}{4} \geq \frac{3}{4}c$$

since  $T_F$  is increasing and  $x_1 > a$  strictly. Then as  $[a, a + \delta] = [a, x_1] \cup [x_1, a + \delta]$  we see that

$$T_F(a + \delta) - T_F(a) \geq \sum_{j=1}^k |F(t_j) - F(t_{j-1})| + \sum_{i=2}^n |F(x_i) - F(x_{i-1})| \geq \frac{3}{4}c + \left(\frac{3}{4}c - \epsilon\right) = \frac{3}{2}c - \epsilon.$$

Combine these all together, we get

$$\epsilon + c \geq \underbrace{T_F(a + \delta) - T_F(a^+)}_{\leq \epsilon} + \underbrace{T_F(a^+) - T_F(a)}_{\leq c} = \underbrace{T_F(a + \delta) - T_F(a)}_{\text{above}} \geq \frac{3}{2}c - \epsilon \Rightarrow c \leq 4\epsilon.$$

By taking  $\epsilon \rightarrow 0$  yields  $c = 0$ , which is a contradiction. ■

**Corollary 6.5.2.**  $F \in \text{NBV}$  if and only if  $F = F_1 - F_2$ ,  $F_1, F_2 \in \text{NBV}$  and increasing.

**Proof.** Consider the [Jordan decomposition](#)

$$F = \frac{T_F + F}{2} - \frac{T_F - F}{2}$$

and proceed similar as in [Theorem 6.5.1](#). ■

**Theorem 6.5.2.** We have the following.

- (a) Suppose that  $\mu$  is a [finite signed Borel measure](#) on  $\mathbb{R}$ , then  $F(x) = \mu((-\infty, x]) \in \text{NBV}$ .
- (b)  $F \in \text{NBV}$  implies that there exists a unique [finite signed Borel measure](#) on  $\mathbb{R}$  satisfying  $\mu_F((-\infty, x]) = F(x)$ .

**Proof.** We prove this one by one.

- (a) Let  $\mu = \mu^+ - \mu^-$ , then  $F = F^+ - F^-$ , where  $F^\pm(x) = \mu^\pm((-\infty, x])$ , which are bounded, right continuous and  $F^\pm(-\infty) = 0$ , so  $F^\pm \in \text{NBV}$ .
- (b) Let  $F \in \text{NBV}$ , then  $F = F_1 - F_2$ ,  $F_1, F_2 \in \text{NBV}$  and increasing from [Corollary 6.5.2](#). Then define  $\mu_{F_1}, \mu_{F_2}$  as the [Lebesgue-Stieltjes measure](#) correspond to  $F_1$  and  $F_2$ , and set  $\mu_F := \mu_{F_1} - \mu_{F_2}$ . Uniqueness is shown in [Appendix A.6.1](#). ■

**Remark (Jordan decomposition).** We see that why both  $\mu^\pm$  and  $F^\pm$  all has the name *Jordan decomposition* from the above proof of (a).

The next natural arising question is the following.

**Problem 6.5.1.** Which functions in NBV correspond to measures  $\mu$  such that  $\mu \perp m$  or  $\mu \ll m$ ?

One answer is given in the following proposition.

**Proposition 6.5.1.** We have the following.

- (a) If  $F \in \text{NBV}$ , then  $F$  is differentiable **almost everywhere** and  $F' \in L^1(\mathbb{R}, m)$ .
- (b)  $\mu_F = \lambda + F' m$  for some measure  $\lambda$  satisfying  $\lambda \perp m$ .
- (c)  $\mu_F \perp m$  if and only if  $F' = 0$  Lebesgue **almost everywhere**.
- (d)  $\mu_F \ll m$  if and only if  $\int_{-\infty}^x F'(t) dt = F(x) - F(-\infty) = F(x)$ .

**Proof.** For (d), we have

$$\begin{aligned}
 \mu_F \ll m &\Leftrightarrow \lambda = 0 \\
 &\Leftrightarrow \mu_F = F' m \\
 &\Leftrightarrow \mu_F(E) = \int_E F' m \quad \forall \text{ Borel } E \\
 &\Leftrightarrow F(x) = \mu_F((-\infty, x]) = \int_{-\infty}^x F'(t) dt, \quad \forall x \in \mathbb{R}.
 \end{aligned}$$

The last converse comes from the uniqueness of [Theorem 6.5.2](#) above. ■

Check (a),  
(b), (c)

## Lecture 36: Absolutely Continuous Functions

### 6.5.1 Absolutely Continuous Functions

8 Apr. 11:00

We start with a definition.

**Definition 6.5.8** (Absolutely continuous function). We say that  $F: \mathbb{R} \rightarrow \mathbb{R}$  is *absolutely continuous*, denoted as  $F \in AC$ , if for all  $\epsilon > 0$ , there exists a  $\delta > 0$  such that if  $(a_1, b_1), \dots, (a_N, b_N)$  are finitely many disjoint open intervals satisfying  $\sum_{n=1}^N (b_n - a_n) < \delta$ , then

$$\sum_{n=1}^N |F(b_n) - F(a_n)| < \epsilon.$$

**Note.** Do not be confused between [Definition 6.5.8](#) and [Definition 6.2.1](#).

**Lemma 6.5.5.** We have the following.

- (a) If  $F$  is **absolutely continuous**, then it is uniformly continuous.
- (b) If  $F$  is **Lipschitz**, then  $F$  is **absolutely continuous**.
- (c) Given  $f \in L^1$ ,  $F(x) = \int_{-\infty}^x f(t) dt$  is **absolutely continuous**.

**Proof.** We prove this one by one.

- (a) We simply take  $N = 1$  in [Definition 6.5.8](#).
- (b) This is trivial.

(c) We write this out as

$$\sum_{n=1}^N |F(b_n) - F(a_n)| = \sum_{n=1}^N \left| \int_{a_n}^{b_n} f(t) dt \right| \leq \sum_{n=1}^N \int_{a_n}^{b_n} |f(t)| dt = \int_E |f(t)| dt$$

where  $E = \bigcup_{n=1}^N (a_n, b_n)$ , so  $m(E) = \sum_{n=1}^N (b_n - a_n)$ . From Midterm Q1, if  $f \in L^1(X, \mu)$ , for all  $\epsilon > 0$ , there is a  $\delta > 0$  such that  $\mu(E) < \delta$  implies  $\int_E |f| < \epsilon$ . This directly implies that this function is **absolutely continuous**. ■

**Example.** The **Cantor function**  $F$  is uniformly continuous. However, we will see that it is not **absolutely continuous**.

**Proposition 6.5.2.** Suppose  $F \in \text{NBV}$ , then  $F$  is **absolutely continuous** if and only if  $\mu_F \ll m$ .

**Proof.** We prove two directions.

( $\Leftarrow$ ) Suppose  $\mu_F \ll m$ . Then  $F(x) = \int_{-\infty}^x F'(t) dt$ , and  $F' \in L^1(\mathbb{R}, m)$ , by **Proposition 6.5.1**. Therefore,  $F \in \text{AC}$ .

( $\Rightarrow$ ) Now suppose  $F \in \text{AC}$ . Note that since  $F$  is continuous,

$$\mu_F((a, b)) = \lim_{n \rightarrow \infty} \mu_F((a, b - 1/n]) = \lim_{n \rightarrow \infty} F(b - 1/n) - F(a) = F(b) - F(a).$$

We let  $E$  be a **Borel set** with  $m(E) = 0$ . Fix  $\epsilon > 0$ , we will show  $|\mu_F(E)| \leq \epsilon$ . Let  $\delta > 0$  be the constant from  $F \in \text{AC}$ .

We know that there exists open  $U_1 \supseteq U_2 \supseteq \cdots \supseteq E$  such that  $\lim_{n \rightarrow \infty} m(U_n) = m(E) = 0$ , and open  $V_1 \supseteq V_2 \supseteq \cdots \supseteq E$  such that  $\lim_{n \rightarrow \infty} \mu_F(V_n) = \mu_F(E)$  by regularity.

Let  $O_n = U_n \cap V_n$ , then  $O_1 \supseteq O_2 \supseteq \cdots \supseteq E$ , and by monotonicity (for  $\mu_F$  decomposing into pos/neg first)

$$\lim_{n \rightarrow \infty} m(O_n) = m(E) = 0, \quad \lim_{n \rightarrow \infty} \mu_F(O_n) = \mu_F(E).$$

Thus without loss of generality, we may assume  $m(O_1) < \delta$ . Each  $O_n$  is a countable union of disjoint intervals

$$O_n = \bigcup_{k=1}^{\infty} (a_k^n, b_k^n).$$

For any  $N$  we also have

$$\sum_{k=1}^N (b_k^n - a_k^n) \leq m(O_n) \leq m(O_1) < \delta.$$

Therefore,

$$\left| \mu_F \left( \bigcup_{k=1}^N (a_k^n, b_k^n) \right) \right| = \left| \sum_{k=1}^N \mu_F((a_k^n, b_k^n)) \right| \leq \sum_{k=1}^N |\mu_F((a_k^n, b_k^n))| \leq \sum_{k=1}^N |F(b_k^n) - F(a_k^n)| < \epsilon,$$

which implies

$$|\mu_F(O_n)| = \lim_{N \rightarrow \infty} \left| \mu_F \left( \bigcup_{k=1}^N (a_k^n, b_k^n) \right) \right| \leq \epsilon$$

and

$$|\mu_F(E)| = \lim_{n \rightarrow \infty} |\mu_F(O_n)| \leq \epsilon.$$

By taking  $\epsilon \rightarrow 0$  we have  $\mu_F(E) = 0$ , hence  $\mu_F \ll m$ . ■

**Corollary 6.5.3.**  $F \in \text{NBV} \cap \text{AC}$  if and only if  $F(x) = \int_{-\infty}^x f(t) dt$  for some  $f \in L^1(\mathbb{R}, m)$ . If this holds, we have  $f = F'$  **Lebesgue almost everywhere**.

**Lemma 6.5.6.** If  $F \in \text{AC}([a, b])$ , then  $F \in \text{NBV}([a, b])$ .

**Proof.** ■

DIY

We then have the following.

**Theorem 6.5.3** (Fundamental theorem of calculus). For  $F \in [a, b] \rightarrow \mathbb{R}$ , the following are equivalent.

- (a)  $F \in \text{AC}([a, b])$ .
- (b)  $F(x) - F(a) = \int_a^x f(t) dt$  for some  $f \in L^1([a, b], m)$ .
- (c)  $F$  is differentiable **almost everywhere** on  $[a, b]$  and  $F(x) - F(a) = \int_a^x F'(t) dt$ .

**Definition.** Let  $\mu$  be a **finite signed Borel measure** on  $\mathbb{R}$ .

**Definition 6.5.9** (Discrete measure).  $\mu$  is called a *discrete measure* if there is a countable set  $\{x_n\}$  and  $c_n \neq 0$  such that  $\sum_{n=1}^{\infty} |c_n| < \infty$  and  $\mu = \sum_n c_n \delta_{x_n}$ .

**Note.**  $\delta_{x_n}$  is the **Dirac delta measure** at  $x_n$ .

**Definition 6.5.10** (Continuous measure).  $\mu$  is called a *continuous measure* if  $\mu(\{a\}) = 0$  for all  $a \in \mathbb{R}$ .

**Lemma 6.5.7.** Given a **finite signed Borel measure**  $\mu$ ,

- (a) Any  $\mu = \mu_d + \mu_c$ , where  $\mu_d$  is **discrete**,  $\mu_c$  is **continuous** are uniquely determined.
- (b)  $\mu$  **discrete** implies  $\mu \perp m$ .
- (c)  $\mu \ll m$  implies  $\mu$  is **continuous**.

**Proof.** We see that (b) and (c) are obvious, so we prove (a) only. Let  $E = \{x \in \mathbb{R} \mid \mu(\{x\}) = 0\}$ , then for any countable subset  $F$  of  $E$ , the series  $\sum_{x \in F} \mu(\{x\})$  converges absolutely to  $\mu(F)$ , so

$$\{x \in E \mid |\mu(\{x\})| > k^{-1}\}$$

is finite for all  $k$ , and it follows that  $E$  itself is countable. Hence,  $\mu_d(A) = \mu(A \cap E)$  is **discrete** and  $\mu_c(A) = \mu(A \setminus E)$  is **continuous**. ■

**Corollary 6.5.4.** For  $\mu$  a **finite signed Borel measure** on  $\mathbb{R}$ , we have that

$$\mu = \mu_d + \mu_{ac} + \mu_{sc}$$

where  $\mu_d$  is **discrete**,  $\mu_{ac}$  is **absolutely continuous** respect to  $m$ , and  $\mu_{sc}$  is **continuous** and **singularly** to  $m$ .

# Chapter 7

## Hilbert Spaces

### Lecture 37: Hilbert Spaces

#### 7.1 Inner Product Spaces

11 Apr. 11:00

**Definition 7.1.1 (Inner product).** Let  $V$  be a (complex) vector space. An *inner product* is a function

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{C}$$

such that

- $\langle \alpha x + \beta y, z \rangle = \alpha \langle x, z \rangle + \beta \langle y, z \rangle$  for all  $x, y, z \in V$ , and  $\alpha, \beta \in \mathbb{C}$ .
- $\langle x, y \rangle = \overline{\langle y, x \rangle}$  for every  $x, y \in V$ .
- $\langle x, x \rangle \in [0, \infty)$ , and  $\langle x, x \rangle = 0$  if and only if  $x = 0$ .

**Note.** Note that we have conjugate linearity in the second argument, i.e.,

$$\langle x, \alpha y + \beta z \rangle = \overline{\alpha} \langle x, y \rangle + \overline{\beta} \langle x, z \rangle$$

for any  $x, y, z \in V$  and  $\alpha, \beta \in \mathbb{C}$ .

**Proof.** This follows from

$$\langle x, \alpha y + \beta z \rangle = \overline{\langle \alpha y + \beta z, x \rangle} = \overline{\alpha \langle y, x \rangle + \beta \langle z, x \rangle} = \overline{\alpha} \overline{\langle y, x \rangle} + \overline{\beta} \overline{\langle z, x \rangle} = \overline{\alpha} \langle x, y \rangle + \overline{\beta} \langle x, z \rangle.$$

⊛

**Example.** We have the following examples.

- $\mathbb{R}^d$  with  $\langle x, y \rangle = x \cdot y = \sum_{i=1}^d x_i y_i$ .
- $\mathbb{C}^d$  with  $\langle x, y \rangle = \sum_{i=1}^d x_i \overline{y_i}$ .
- $L^2(X, \mu)$  with  $\langle f, g \rangle = \int_X f \overline{g} d\mu$ . Note by [Theorem 5.2.1](#),

$$\left| \int_X f \overline{g} \right| \leq \|f \overline{g}\|_1 \leq \|f\|_2 \|g\|_2 < \infty$$

because  $1/2 + 1/2 = 1$ .

- A special case is  $\ell^2$ , where we have

$$\langle x, y \rangle = \sum_{i=1}^{\infty} x_i \overline{y_i}.$$

**Note.** Note that

$$\|x + y\|^2 = \langle x + y, x + y \rangle = \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle = \|x\|^2 + 2 \operatorname{Re} \langle x, y \rangle + \|y\|^2$$

**Theorem 7.1.1 (Cauchy-Schwarz Inequality).** Given an inner product space,  $|\langle x, y \rangle| \leq \|x\| \|y\|$ .

**Proof.** This is clear if  $\langle x, y \rangle = 0$ . Assume  $\langle x, y \rangle \neq 0$ , then for every  $\alpha \in \mathbb{C}$ , we know that

$$0 \leq \|\alpha x - y\|^2 = |\alpha|^2 \|x\|^2 - 2 \operatorname{Re} \alpha \langle x, y \rangle + \|y\|^2.$$

Write  $\langle x, y \rangle = |\langle x, y \rangle| e^{i\theta}$ , and take  $\alpha = e^{-i\theta} t$  for arbitrary  $t \in \mathbb{R}$ . Then, the right-hand side gives

$$0 \leq \|x\|^2 t^2 - 2 |\langle x, y \rangle| t + \|y\|^2.$$

Note this is a real quadratic function of  $t$ , with at most one real root. Thus, the discriminant  $\Delta \leq 0$ . Specifically, we have

$$\Delta = 4 |\langle x, y \rangle|^2 - 4 \|x\|^2 \|y\|^2 \leq 0 \Leftrightarrow |\langle x, y \rangle|^2 \leq \|x\|^2 \|y\|^2 \Leftrightarrow |\langle x, y \rangle| \leq \|x\| \|y\|.$$

■

**Definition 7.1.2 (Induced norm from inner product).** Given an inner product space  $V$ , let

$$\|x\| := \sqrt{\langle x, x \rangle},$$

which is so-call the *norm induced from the inner product*.

**Proof.** We need to check that this actually defines a **norm**. We check the following.

**Claim.**  $\|x\| = 0 \Leftrightarrow x = 0$  for all  $x \in V$ .

**Proof.** This follows from the **definition of an inner product**. ⊗

**Claim.**  $\|\alpha x\| = |\alpha| \|x\|$  for all  $x \in V$ .

**Proof.** This follows from

$$\|\alpha x\| = \sqrt{\langle \alpha x, \alpha x \rangle} = \sqrt{\alpha \overline{\alpha} \langle x, x \rangle} = |\alpha| \|x\|.$$

⊗

**Claim (Triangle inequality).**  $\|x + y\| \leq \|x\| + \|y\|$  for all  $x, y \in V$ .



**Proof.** The triangle inequality is less obvious, and comes from [Theorem 7.1.1](#). Namely,

$$\begin{aligned}\|x + y\|^2 &= \|x\|^2 + 2 \operatorname{Re} \langle x, y \rangle + \|y\|^2 \\ &\leq \|x\|^2 + 2 |\langle x, y \rangle| + \|y\|^2 \\ &\leq \|x\|^2 + 2 \|x\| \|y\| + \|y\|^2 = (\|x\| + \|y\|)^2\end{aligned}$$

Taking square root on both sides, we have

$$\|x + y\| \leq \|x\| + \|y\|.$$

⊗

■

**Theorem 7.1.2** (Parallelogram law). Let  $V$  be a [normed vector space](#). Then,  $\|\cdot\|$  is [induced by an inner product](#) if and only if

$$\|x + y\|^2 + \|x - y\|^2 = 2 \|x\|^2 + 2 \|y\|^2$$

for all  $x, y \in V$ .

**Proof.** We show two directions.

( $\Rightarrow$ ) This follows from

$$\|x \pm y\|^2 = \|x\|^2 \pm 2 \operatorname{Re} \langle x, y \rangle + \|y\|^2$$

and

$$\|x \pm iy\|^2 = \|x\|^2 \pm 2 \operatorname{Im} \langle x, y \rangle + \|y\|^2.$$

( $\Leftarrow$ ) Firstly, we define

$$\langle x, y \rangle = \frac{1}{4} \left( \|x + y\|^2 - \|x - y\|^2 + \|x + iy\|^2 - i \|x - iy\|^2 \right)$$

as motivated by the above relationship.

**Exercise.** Check this [inner product](#) is indeed inducing the desired norm.

■

**Example.** Consider  $L^p(\mathbb{R}, m)$ ,  $f = \mathbb{1}_{(0,1)}$ ,  $g = \mathbb{1}_{(1,2)}$ . We see the [parallelogram law](#) is satisfied only when  $p = 2$ .

**Remark.** Hence,  $L^p(\mathbb{R}, m)$  is only an [inner product](#) space when  $p = 2$ .

Since we're doing real analysis, we want to deal with limits. It turns out that with an [inner product](#) space, we can say something more compare to the case of a [normed](#) vector space. We now illustrate this.

**Definition.** Given a vector space  $V$  with either a [norm](#) or an [inner product](#), we have the followings.

**Definition 7.1.3 (Strong convergence).** We say that  $x_n \in V$  converges to  $x \in V$  strongly if

$$\lim_{n \rightarrow \infty} \|x_n - x\| = 0.$$

**Definition 7.1.4 (Weak convergence).** We say that  $x_n \in V$  converges to  $x \in V$  weakly if for any fixed  $y \in V$ ,

$$\lim_{n \rightarrow \infty} \langle x_n - x, y \rangle = 0.$$

**Lemma 7.1.1 (Strong convergence implies weak convergence).** Suppose  $V$  is an inner product space. If  $x_n \rightarrow x$  strongly, then  $x_n \rightarrow x$  weakly.

**Proof.** By Cauchy-Schwarz inequality,

$$0 \leq |\langle x_n - x, y \rangle| \leq \|x_n - x\| \cdot \|y\|.$$

Since  $\|x_n - x\| \rightarrow 0$  and  $\|y\|$  is constant in  $n$ , from the squeeze theorem, we have

$$\langle x_n - x, y \rangle \rightarrow 0$$

as  $n \rightarrow \infty$ . ■

**Example.** Consider  $\ell^2$ ,  $x_n = (0, \dots, 0, 1, 0, \dots)$  and  $x = 0$ . Then  $x_n$  does not converge strongly to any vector, but it does converge to 0 weakly.

**Proof.** If we fix  $y \in \ell^2$ , then

$$\langle x_n - x, y \rangle = \bar{y}_n$$

which goes to 0 as  $n \rightarrow \infty$  because  $\sum_n |y_n|^2 < \infty$ . Therefore,  $x_n \rightarrow 0$  weakly, but we see that

$$\|x_n - 0\| = \|x_n\| = 1.$$

Thus,  $x_n \not\rightarrow 0$  strongly. ⊛

## 7.2 Orthonormal Bases

**Definition 7.2.1 (Orthogonal).** Two vectors  $x, y$  are orthogonal if  $\langle x, y \rangle = 0$ , denoted as  $x \perp y$ .

**Remark.** Do not confuse between this notation and Definition 6.1.5.

**Lemma 7.2.1 (Pythagorean theorem).** If  $x_1, \dots, x_n \in V$ ,  $\langle x_i, x_j \rangle = 0$  for all  $i \neq j$ , then

$$\left\| \sum_{i=1}^n x_i \right\|^2 = \sum_{i=1}^n \|x_i\|^2.$$

**Proof.** Use that  $\|x + y\|^2 = \|x\|^2 + 2 \operatorname{Re} \langle x, y \rangle + \|y\|^2$  and induct. ■

**Definition 7.2.2 (Orthonormal set).** We call  $\{e_i\}_{i \in I}$  an orthonormal set if

$$\langle e_i, e_j \rangle = \begin{cases} 0, & \text{if } i \neq j; \\ 1, & \text{if } i = j. \end{cases}$$

## Lecture 38: Orthonormal Basis

Let's start with a lemma.

13 Apr. 11:00

**Lemma 7.2.2** (Best approximation). Let  $e_1, \dots, e_N$  be **orthonormal**, then for  $x \in V$ , let  $\alpha_i = \langle x, e_i \rangle$ , we have

$$\left\| x - \sum_{i=1}^N \alpha_i e_i \right\| \leq \left\| x - \sum_{i=1}^N \beta_i e_i \right\|$$

for all  $\beta_1, \dots, \beta_N \in \mathbb{C}$ . Namely,  $\sum_{i=1}^N \alpha_i e_i$  is the *best approximation* to  $x$  within the span of  $e_1, \dots, e_N$ .

**Note.** We can also think of it as an **orthogonal** projection.

**Proof.** Let  $z = x - \sum_{i=1}^N \alpha_i e_i$ ,  $w = \sum_{i=1}^N (\alpha_i - \beta_i) e_i$  for any  $\{\beta_i\}$ . Note that for all  $n = 1, \dots, N$ , we have

$$\langle z, e_n \rangle = \langle x, e_n \rangle - \alpha_n = 0.$$

Thus,  $\langle z, w \rangle = 0$ . So by the **Pythagorean theorem**,

$$\|z + w\|^2 = \|z\|^2 + \|w\|^2 \geq \|z\|^2,$$

which proves the result. ■

**Lemma 7.2.3** (Bessel's inequality). Let  $\{e_i\}_1^\infty$  be an **orthonormal set**. For  $x \in V$ , let  $\alpha_i = \langle x, e_i \rangle$ . Then we have

$$\sum_{i=1}^{\infty} |\alpha_i|^2 \leq \|x\|^2,$$

which is so-called the *Bessel's inequality*. More generally, we have

$$\|x\|^2 = \left\| x - \sum_{i=1}^N \alpha_i e_i \right\|^2 + \sum_{i=1}^N |\alpha_i|^2$$

for all  $N \in \mathbb{N}$ .

**Proof.** We see that

$$\begin{aligned} \left\| x - \sum_{i=1}^N \alpha_i e_i \right\|^2 &= \|x\|^2 - 2 \operatorname{Re} \left\langle x, \sum_{i=1}^N \alpha_i e_i \right\rangle + \left\| \sum_{i=1}^N \alpha_i e_i \right\|^2 \\ &= \|x\|^2 - 2 \sum_{i=1}^N \operatorname{Re} \overline{\alpha_i} \langle x, e_i \rangle + \sum_{i=1}^N |\alpha_i|^2 \\ &= \|x\|^2 - 2 \sum_{i=1}^N |\alpha_i|^2 + \sum_{i=1}^N |\alpha_i|^2 \\ &= \|x\|^2 - \sum_{i=1}^N |\alpha_i|^2. \end{aligned}$$

From this, we see that the inequality holds trivially. ■

**Remark.** Lemma 7.2.3 actually holds even for an uncountable collection.

**Definition 7.2.3** (Orthonormal basis). An **orthonormal set**  $\{e_i\}$  is said to be an *orthonormal basis* of

$V$  if  $\overline{W} = V$ , where

$$W = \left\{ \sum_{i=1}^N \beta_i e_i \mid N \in \mathbb{N}, \beta_1, \dots, \beta_N \in \mathbb{C} \right\}$$

is the subspace of finite linear combinations. In other words, for all  $x \in V$  and for every  $\epsilon > 0$ , there exists  $w \in W$  such that  $\|x - w\| < \epsilon$ .

**Example.** We have the following.

- (a) For  $\mathbb{C}^d$ , the orthonormal basis is  $e_i = (0, \dots, 0, 1, 0, \dots, 0)$  for  $i = 1, \dots, d$ .
- (b) For  $\ell^2$ , the orthonormal basis is the countably many  $e_i = (0, \dots, 0, 1, 0, \dots)$  for  $i \in \mathbb{N}$ .

**Definition 7.2.4 (Hilbert Space).** A Hilbert space is a complete inner product space.

**Note.** Namely, a Hilbert space is a Banach space with an inner product.

**Example.**  $\mathbb{R}^d$ ,  $\mathbb{C}^d$ ,  $L^2(X, \mathcal{A}, \mu)$ ,  $\ell^2$  are all Hilbert spaces.

**Example.**  $C([0, 1]) \subseteq L^2(X, \mathcal{A}, \mu)$  is not a Hilbert space.

**Proof.** We see this by the fact that it is not complete. Take a function  $f_n$  so that  $f_n$  is zero from 0 to  $1/2$  and 1 from  $1/2 + 1/n$  to 1, connected continuously by a line. We then see that  $f_n$  is Cauchy, but its natural limit is discontinuous.  $\otimes$

**Theorem 7.2.1.** Let  $\mathcal{H}$  be a Hilbert space. Let  $\{e_i\}_{i=1}^\infty$  be an orthonormal set. The following are equivalent.

- (a)  $\{e_i\}_{i=1}^\infty$  is an orthonormal basis.
- (b) If  $x \in \mathcal{H}$  and  $\langle x, e_i \rangle = 0$  for all  $i$ , then  $x = 0$ .
- (c) If  $x \in \mathcal{H}$ , then  $s_N := \sum_{i=1}^N \alpha_i e_i \rightarrow x$  strongly where  $\alpha_i = \langle x, e_i \rangle$ .
- (d) If  $x \in \mathcal{H}$ , then  $\|x\|^2 = \sum_{i=1}^\infty |\alpha_i|^2$ .

**Proof.** We prove this as follows.

**Claim.** (c)  $\Rightarrow$  (d).

**Proof.** From Lemma 7.2.3, we have

$$\|x\|^2 = \|x - s_N\|^2 + \sum_{i=1}^N |\alpha_i|^2.$$

Taking  $N \rightarrow \infty$  and noting  $s_N \rightarrow x$  strongly gives

$$\|x\|^2 = \lim_{N \rightarrow \infty} \sum_{i=1}^N |\alpha_i|^2 = \sum_{i=1}^\infty |\alpha_i|^2.$$

$\otimes$

**Claim.** (d)  $\Rightarrow$  (a).

**Proof.** Using the same equality from Lemma 7.2.3, we have  $\|x\|^2 = \|x - s_N\|^2 + \sum_{i=1}^N |\alpha_i|^2$ . Now, take  $N \rightarrow \infty$  yields  $\|x - s_N\|^2 \rightarrow 0$  so  $\|x - s_N\| \rightarrow 0$ . Therefore,  $s_N \rightarrow x$  strongly, yielding that  $x$  can be approximated by finite linear combinations as desired.  $\otimes$

**Claim.** (a)  $\Rightarrow$  (b).

**Proof.** Fix  $x \in \mathcal{H}$ , and fix  $\epsilon > 0$ . Then by (a), there exists a  $y = \sum_{i=1}^k \beta_i e_i$  such that  $\|x - y\| < \epsilon$ .

By the best approximation lemma,  $\|x - s_k\| \leq \|x - y\| < \epsilon$ . If  $\langle x, e_i \rangle = 0$  for all  $i$ , then  $s_k = 0$ , so  $\|x\| < \epsilon$ . Taking  $\epsilon \rightarrow 0$  would yield  $\|x\| = 0$ , implying  $x = 0$ .  $\otimes$

**Claim.** (b)  $\Rightarrow$  (c).

**Proof.** Bessel's inequality gives

$$\sum_{i=1}^{\infty} |\alpha_i|^2 \leq \|x\|^2 < \infty.$$

We now see that for  $N > M$ ,

$$\|s_N - s_M\|^2 = \left\| x - \sum_{i=M+1}^N \alpha_i e_i \right\|^2 = \sum_{i=M+1}^N |\alpha_i|^2 \rightarrow 0$$

as  $N > M \rightarrow \infty$ , by convergence of the series. This implies that  $\{s_N\}_{N=1}^{\infty}$  is a Cauchy sequence in  $\mathcal{H}$ . Since  $\mathcal{H}$  is complete, there is a vector  $y$  such that  $s_N \rightarrow y$  strongly.

**Problem.** Is  $y = x$ ?

**Answer.** Fix  $i \in \mathbb{N}$ , consider  $\langle y - x, e_i \rangle$ . We see that

$$\langle y - x, e_i \rangle = \langle y - s_N, e_i \rangle + \langle s_N - x, e_i \rangle.$$

We can compute that for  $N > i$ , that

$$\langle s_N - x, e_i \rangle = \alpha_i - \langle x, e_i \rangle = 0.$$

Hence,  $\langle y - x, e_i \rangle = \langle y - s_N, e_i \rangle$ . Because strong convergence implies weak convergence, taking  $N \rightarrow \infty$  yields that  $\langle y - x, e_i \rangle = 0$  for all  $i \in \mathbb{N}$ . Therefore, by the assumption of (b),  $y - x = 0$ , so  $x = y$  and we're done.  $\otimes$

$\otimes$

**Remark.** Note that we did not use the Hilbert space properties for everything except (b)  $\Rightarrow$  (c). When  $\mathcal{H}$  is replaced by any inner product space  $V$  we only have

$$(c) \Rightarrow (d) \Rightarrow (a) \Rightarrow (b).$$

**Note** (Planchenel identity). The (d) in Theorem 7.2.1 is called the *Planchenel identity*.

**Definition 7.2.5** (Separable). A metric space is called *separable* if there exists a countable dense subset.

---

**Example.**  $\mathbb{R}^d \supseteq \mathbb{Q}^d$  and  $\ell^p$  with  $1 \leq p < \infty$  are all [separable](#), but not for  $p = \infty$ .

**Proof.** To do this, consider sequences of rational numbers. ⊛

**Example.**  $L^p(\mathbb{R}, m)$  is [separable](#) for  $1 \leq p < \infty$ .

**Proof.** Take [step functions](#) with rational heights and rational endpoints to intervals. ⊛

**Theorem 7.2.2.** Every [separable Hilbert space](#) has a countable [orthonormal basis](#).

**Proof.** Gram-Schmidt algorithm<sup>a</sup> will construct such an [orthonormal basis](#) explicitly. ■

---

<sup>a</sup>[https://en.wikipedia.org/wiki/Gram-Schmidt\\_process](https://en.wikipedia.org/wiki/Gram-Schmidt_process)

**Note** (Dimension of a Hilbert space). The cardinality of an [orthonormal basis](#) is determined by the space, and we can call this the *dimension* of a [Hilbert space](#).

# Chapter 8

## Introduction to Fourier Analysis

### Lecture 39: Introduction to Fourier Analysis

As previously seen. [Planchenel identity](#) says that if  $e_i$  is an [orthonormal basis](#), then

15 Apr. 11:00

$$\langle x, y \rangle = \sum_{i=1}^{\infty} \alpha_i \bar{\beta}_i$$

where  $\alpha_n = \langle x, e_n \rangle$ ,  $\beta_n = \langle y, e_n \rangle$ .

### 8.1 Fourier Series

We'll be considering the [Hilbert space](#)  $L^2([-\pi, \pi])$ , which by scaling is equivalent to any finite interval. Let's start with a lemma.

**Lemma 8.1.1.** The set

$$\left\{ e_n(x) = \frac{1}{\sqrt{2\pi}} e^{inx} = \frac{1}{\sqrt{2\pi}} (\cos(nx) + i \sin(nx)) \right\}$$

is an [orthonormal set](#) in  $L^2([-\pi, \pi])$ .

**Proof.** We see that

$$\langle e_n, e_m \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(m-n)x} dx = \begin{cases} 1, & \text{if } m = n; \\ 0, & \text{if } m \neq n, \end{cases}$$

hence, the result follows directly. ■

Now, a natural problem arises:

**Problem 8.1.1.** Is  $\{e_n\}$  defined in [Lemma 8.1.1](#) an [orthonormal basis](#)?

**Answer.** By [Hölder's inequality](#) on  $L^2([-\pi, \pi])$ , we have

$$\|f\|_1 \leq \|1\|_2 \|f\|_2 = \sqrt{2\pi} \cdot \|f\|_2 < \infty.$$

Similarly, we have

$$\|f\|_2 = \sqrt{\int_{-\pi}^{\pi} |f(t)|^2 dt} \leq \sqrt{\|f\|_{\infty}^2 2\pi},$$

which implies

$$\|f\|_1 \leq \sqrt{2\pi} \|f\|_2 \leq 2\pi \|f\|_{\infty}.$$

⊛

**Definition 8.1.1** (Fourier coefficient). For  $f \in L^1([-\pi, \pi])$ , its  $n^{\text{th}}$  Fourier coefficient is

$$\hat{f}_n := \langle f, e_n \rangle = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} f(y) e^{-iny} dy.$$

We can show that Definition 8.1.1 form an orthonormal basis, we show that

$$\sum_{n=-M}^N \hat{f}_n e_n(x)$$

converge strongly to  $f(x)$  as  $M, N \rightarrow \infty$ . Explicitly, this is true since

$$\sum_{n=-M}^N \hat{f}_n e_n(x) = \frac{1}{\sqrt{2\pi}} \sum_{n=-M}^N \left[ \int_{-\pi}^{\pi} f(y) e^{-iny} dy \right] e^{inx} = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} f(y) \left( \sum_{n=-M}^N e^{in(x-y)} \right) dy.$$

**Definition 8.1.2** (Poisson kernel). For  $0 \leq r < 1$ , the Poisson kernel is defined as

$$P_r(t) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} e^{int} r^{|n|}.$$

Explicitly, summing as a geometric series gives us

$$P_r(t) = \frac{1 - r^2}{2\pi(1 - 2r \cos t + r^2)}.$$

**Lemma 8.1.2.** For  $f \in L^1([-\pi, \pi])$  and  $0 \leq r < 1$ , for  $x \in [-\pi, \pi]$ ,

$$\sum_{n=-\infty}^{\infty} \hat{f}_n e_n(x) r^{|n|}$$

converge absolutely and uniformly to

$$\int_{-\pi}^{\pi} P_r(x - y) f(y) dy.$$

**Proof.** We have

$$\sum_{n=-\infty}^{\infty} \left| \hat{f}_n e_n(x) r^{|n|} \right| \leq \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} \left( \int_{-\pi}^{\pi} |f(y) e^{-iny}| dy \right) |e_n(x)| r^{|n|} = \frac{\|f\|_1}{2\pi} \sum_{n=-\infty}^{\infty} r^{|n|} < \infty.$$

Therefore, Corollary 2.3.1 applies, and we further have

$$\begin{aligned} \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} \left( \int_{-\pi}^{\pi} f(y) e^{-iny} dy \right) e^{inx} r^{|n|} &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) \left( \sum_{n=-\infty}^{\infty} e^{in(x-y)} r^{|n|} \right) dy \\ &= \int_{-\pi}^{\pi} P_r(x - y) f(y) dy. \end{aligned}$$

Uniform convergence takes a small bit more work. ■

DIY

**Note.** Note that

$$P_r(0) = \frac{1 - r^2}{2\pi(1 - r)^2} = \frac{1 + r}{2\pi(1 - r)} \rightarrow 0 \text{ as } r \rightarrow 1.$$



For any  $t \neq 0$  we have

$$\frac{1 - r^2}{1\pi(1 - 2r \cos t + r^2)} \rightarrow 0$$

as the bottom is always nonzero and finite.

**Definition 8.1.3 (Good kernel).** A kernel  $K_r$  is said to be in a *family<sup>a</sup> of good kernels* if

- (a)  $K_r(t) \geq 0$ .
- (b)  $\int_{-\pi}^{\pi} K_r(t) dt = 1$ .
- (c) For every  $\delta > 0$ ,

$$\int_{[-\pi, \pi] \setminus [-\delta, \delta]} K_r(t) dt \rightarrow 0$$

as  $r \rightarrow 1$ .<sup>b</sup>

<sup>a</sup>Respect to  $r$ .

<sup>b</sup>This definition is not universal. Some text uses other convention.

**Lemma 8.1.3.**  $P_r(t)$  form a *family of good kernels*.

**Proof.** For (b), we use the first formula with [Fubini's theorem](#). For (a) and (c), we use the second formula, namely

$$\int_{[-\pi, \pi] \setminus [-\delta, \delta]} P_r(t) dt \leq \frac{1 - r^2}{2\pi(1 - 2r \cos \delta + r^2)} 2\pi \rightarrow 0$$

as  $r \rightarrow 1$ . ■

**Lemma 8.1.4.** For  $f \in C([-\pi, \pi])$  satisfying  $f(-\pi) = f(\pi)$ , we have

$$\lim_{r \nearrow 1} \int_{-\pi}^{\pi} P_r(x - y) f(y) dy = f(x)$$

uniformly for  $x \in [-\pi, \pi]$ .

**Proof.** We extend  $f$  to  $f: \mathbb{R} \rightarrow \mathbb{C}$  by setting  $f(x + 2\pi) = f(x)$ , then  $f$  is uniformly continuous and bounded. Also, we have

$$\begin{aligned} & \int_{-\pi}^{\pi} P_r(x - y) f(y) dy - f(x) \\ &= \int_{-\pi}^{\pi} P_r f(x - y) dy - f(x) \\ &= \int_{-\pi}^{\pi} P_r(y) f(x - y) dy - f(x) \int_{-\pi}^{\pi} P_r(y) dy \\ &= \int_{-\delta}^{\delta} P_r(y) (f(x - y) - f(x)) dy + \int_{[-\pi, \pi] \setminus [-\delta, \delta]} P_r(y) (f(x - y) - f(x)) dy. \end{aligned}$$

Now, fix  $\epsilon > 0$ , then  $f$  is uniformly continuous, so we can choose a  $\delta > 0$  so that  $|f(x - y) - f(x)| < \epsilon$  for any choice of  $x$ . Then, the left-hand side is bounded by  $\epsilon$ . As for the right-hand side, we note that  $f$  is bounded, so for some  $M$ , we have

$$\left| \int_{[-\pi, \pi] \setminus [-\delta, \delta]} P_r(y) (f(x - y) - f(x)) dy \right| \leq M \int_{[-\pi, \pi] \setminus [-\delta, \delta]} P_r(y) dy.$$

Therefore, sending  $r \nearrow 1$  will give

$$\int_{-\pi}^{\pi} P_r(x-y)f(y) \, dy - f(x) \rightarrow 0,$$

as we desired. ■

## Lecture 40: Fourier Analysis

**As previously seen.** We have from last time that for  $0 \leq r < 1$ ,

18 Apr. 11:00

$$e_n(x) := \frac{1}{\sqrt{2\pi}} e^{inx}$$

$$P_r(t) := \sum_{n=-\infty}^{\infty} e_n(t) r^{|n|}$$

$$f \in L^1([-\pi, \pi]) \Rightarrow \sum_{n=-\infty}^{\infty} \hat{f}_n e_n(x) r^{|n|} = \int_{-\pi}^{\pi} P_r(x-y)f(y) \, dy$$

uniformly in  $x \in [-\pi, \pi]$ . Furthermore, if  $f \in C([-\pi, \pi])$  and  $f(\pi) = f(-\pi)$  then

$$\lim_{r \rightarrow 1} \int_{-\pi}^{\pi} P_r(x-y)f(y) \, dy = f(x)$$

uniformly in  $x \in [-\pi, \pi]$ .

Finally, we also have

$$\|f\|_1 \leq \sqrt{2\pi} \|f\|_2 \leq 2\pi \|f\|_{\infty}.$$

**Theorem 8.1.1 (Fourier Series).** The set of  $e_n(x) = \frac{1}{\sqrt{2\pi}} e^{inx}$  for  $n \in \mathbb{Z}$  is an **orthonormal basis** of  $L^2([-\pi, \pi])$ .

**Proof.** From **Lemma 8.1.1**, this is an **orthonormal set**. Thus, we just need to show that for all  $f \in L^2([-\pi, \pi])$ , that there exists  $h = \sum_{n=-M}^N \beta_n e_n(x)$  such that  $\|f - h\|_2 < \epsilon$ . That is we need to show that the span of the  $e_n$  is dense.

Let  $f \in L^2([-\pi, \pi])$  and fix an  $\epsilon > 0$ . Then there is a function  $g \in C([-\pi, \pi])$  with  $g(\pi) = g(-\pi)$  such that  $\|f - g\|_2 < \frac{\epsilon}{3}$  which follows from a simple density argument.

Let  $g_r(x) = \int_{-\pi}^{\pi} P_r(x-y)g(y) \, dy$ . By the above, there exists an  $r \in [0, 1)$  such that  $\|g_r - g\|_{\infty} < \frac{\epsilon}{3\sqrt{2\pi}}$ .

Therefore,  $\|g_r - g\|_2 < \frac{\epsilon}{3}$ . Consider  $g_{r,N}(x) = \sum_{n=-N}^N \hat{g}_n e_n(x) r^{|n|}$ . By the above there exists an  $N \in \mathbb{N}$  such that  $\|g_{r,N} - g_r\|_{\infty} < \frac{\epsilon}{3\sqrt{2\pi}}$ , thus  $\|g_{r,N} - g_r\|_2 < \frac{\epsilon}{3}$ . Therefore,

$$\|f - g_{r,N}\| < \epsilon$$

where  $g_{r,N}$  is a finite linear combination of  $e_n$ 's, so these form an **orthonormal basis** as desired. ■

**Example (Plancherel's identity).** We have  $\|f\|_2^2 = \sum_{n=-\infty}^{\infty} |\hat{f}_n|^2$ .

**Proof.** For  $f(x) = x$ , we have

$$\hat{f}_n = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} x e^{-inx} \, dx = \begin{cases} 0, & \text{if } n = 0; \\ \frac{(-1)^n i \sqrt{2\pi}}{n}, & \text{if } n \geq 0. \end{cases}$$

Together these imply that

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}.$$

⊛

**Example (Isoperimetric inequality).** Consider a parametric curve  $(x(t), y(t)) \in \mathbb{R}^2$  with  $t \in [-\pi, \pi]$ . Assume that

- (a) This is a closed curve, so  $(x(-\pi), y(-\pi)) = (x(\pi), y(\pi))$ .
- (b) We assume these are smooth, but in fact we just need  $x, y$  are  $C^1$  functions.
- (c) The curve is simple.

Suppose that

$$L = \int_{-\pi}^{\pi} \sqrt{x'(t)^2 + y'(t)^2} dt = 2\pi,$$

then the largest area  $A$  this curve can enclose is  $\pi$ .

**Proof.** By **Green's theorem**,

$$A = \frac{1}{2} \oint_C (x dy - y dx) = \frac{1}{2} \int_{-\pi}^{\pi} (x(t)y'(t) - x'(t)y(t)) dt.$$

Arc length parametrization, so that  $x'(t)^2 + y'(t)^2 = 1$  for all  $t$ . Then the condition  $L = 2\pi$  is automatically satisfied and can be written as

$$\int_{-\pi}^{\pi} (x'(t)^2 + y'(t)^2) dt = 2\pi.$$

For ease of computation, rewrite using complex numbers, i.e.,  $z(t) = x(t) + iy(t)$ . This is then subject to

$$\int_{-\pi}^{\pi} |z'(t)|^2 dt = 2\pi.$$

Rewriting our above formula for area, we need to find the maximum of

$$A = \frac{1}{4i} \int_{-\pi}^{\pi} (\overline{z(t)} z'(t) - z(t) \overline{z'(t)}) dt.$$

Note  $z$  is  $C^1$  and  $z(\pi) = z(-\pi)$ . Denote  $\hat{z}_n = \alpha_n$ , then we now have

$$\hat{z}'_n = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} z'(t) e^{-int} dt = (z(t) e^{-int})_{\pi}^{\pi} + in \int_{-\pi}^{\pi} z(t) e^{-int} dt = in \alpha_n.$$

By **Plancherel's identity**, we have that

$$\int_{-\pi}^{\pi} |z'(t)|^2 dt = \sum_{n=-\infty}^{\infty} |in \alpha_n|^2 = \sum_{n=-\infty}^{\infty} n^2 |\alpha_n|^2 = 2\pi.$$

We know **Plancherel's identity** says  $\langle f, g \rangle = \sum_{n=-\infty}^{\infty} \hat{f}_n \overline{\hat{g}_n}$ . Therefore,

$$A = \frac{1}{4i} \sum_{n=-\infty}^{\infty} \overline{\alpha_n} (in \alpha_n) - \alpha_n \overline{(in \alpha_n)} = \frac{1}{2} \sum_{n=-\infty}^{\infty} n |\alpha_n|^2.$$

Then the question becomes what is the maximum of  $\frac{1}{2} \sum_{n=-\infty}^{\infty} n |\alpha_n|^2$  subject to  $\sum_{n=-\infty}^{\infty} n^2 |\alpha_n|^2 = 2\pi$ .

We will show that

$$2\pi - \sum_{-\infty}^{\infty} n |\alpha_n|^2 \geq 0$$

as we guess the maximum should be  $\pi$ , by virtue of the circle being theoretically optimal. Then we have

$$2\pi - \sum_{-\infty}^{\infty} n |\alpha_n|^2 = \sum_{n=-\infty}^{\infty} (n^2 - n) |\alpha_n|^2 \geq 0$$

because term by term every term is non-negative.

Thus, the area cannot be more than  $\pi$ . We have  $A = \pi$  if and only if equality holds above, that is  $\alpha_n = 0$  for  $n \neq 0, 1$ . This means  $z(t) = \alpha_0 + \alpha_1 e_1(x)$ , that is

$$z(t) = \alpha_0 + C e^{it}.$$

This means that  $z(t)$  is a circle about  $\alpha_0$ .

⊛

**Note.** Professor Baik likes Dym and McKean's Fourier Series & Integrals [DM72]!

# Appendix

# Appendix A

## Additional Proofs

### A.1 Measure

This section gives all additional proofs in [Section 1](#).

**Theorem A.1.1** ([Theorem 1.3.2 \(c\)](#)). Under the setup of [Carathéodory extension theorem](#),  $(X, \mathcal{A}, \mu)$  is a [complete measure space](#).

**Proof.** We see this in two parts.

**Claim.** If  $A \subset X$  satisfies  $\mu^*(A) = 0$ , then  $A$  is [Carathéodory measurable](#) with respect to  $\mu^*$ .

**Proof.** If  $A \subset X$  and  $\mu^*(A) = 0$ , where  $\mu^*$  is an outer measure on  $X$ , we'll show that  $A$  is [Carathéodory measurable](#) with respect to  $\mu^*$ . Equivalently, we want to show that for any  $E \subset X$ ,

$$\mu^*(E) = \mu^*(E \cap A) + \mu^*(E \setminus A).$$

Firstly, noting that  $(E \cap A) \subset A$ , and by [monotonicity](#) of  $\mu^*$ , we see that

$$\mu^*(E \cap A) \leq \mu^*(A) = 0,$$

and since  $\mu^* \geq 0$ , hence  $\mu^*(E \cap A) = 0$ . Now, we only need to show that

$$\mu^*(E) = \mu^*(E \setminus A).$$

Since  $E \setminus A = E \cap A^c$ , and hence we have  $E \cap A^c \subset E$ , so

$$\mu^*(E) \geq \mu^*(E \setminus A).$$

To show another direction, we note that

$$\mu^*(E) \leq \mu^*(E \cup A) = \mu^*((E \setminus A) \cup A) \leq \mu^*(E \setminus A),$$

hence we conclude that  $A$  is [Carathéodory measurable](#) with respect to  $\mu^*$  if  $\mu^*(A) = 0$ . ⊗

**Claim.** If  $A$  is [μ-subnull](#), then  $A \in \mathcal{A}$ .

**Proof.** Let  $\mathcal{A}$  denotes the [Carathéodory  \$\sigma\$ -algebra](#), and  $\mu := \mu^*|_{\mathcal{A}}$ . We want to show if  $A$  is  $\mu$ -subnull, then  $A \in \mathcal{A}$ . Firstly, if  $A$  is  $\mu$ -subnull, then there exists a  $B \in \mathcal{A}$  such that  $A \subset B$  and  $\mu(B) = 0$ . But since from the [monotonicity](#) of  $\mu^*$ , we further have

$$0 = \mu(B) = \mu^*(B) \geq \mu^*(A),$$

hence  $\mu^*(A) = 0$ . From the first claim, we immediately see that  $A$  is [Carathéodory measurable](#) with respect to  $\mu^*$ , which implies  $A$  is in [Carathéodory  \$\sigma\$ -algebra](#), hence  $A \in \mathcal{A}$ .  $\otimes$

We see that the second claim directly proves that  $(X, \mathcal{A}, \mu)$  is a [complete measure space](#).  $\blacksquare$

**Lemma A.1.1.** The function  $F$  defined in [this example](#) is a [distribution function](#)

**Proof.** We define

$$F_n(x) = \begin{cases} 1, & \text{if } x \geq r_n; \\ 0, & \text{if } x < r_n \end{cases}$$

where  $\{r_1, r_2, \dots\} = \mathbb{Q}$ , and

$$F(x) = \sum_{n=1}^{\infty} \frac{F_n(x)}{2^n} = \sum_{n; r_n \leq x} \frac{1}{2^n}$$

is both increasing and right-continuous.

- Increasing. Consider  $x < y$ . We see that

$$F(x) = \sum_{n; r_n \leq x} \frac{1}{2^n} \leq \sum_{n; r_n \leq y} \frac{1}{2^n} = F(y)$$

clearly.<sup>a</sup>

- Right-continuous. We want to show  $F(x^+) = F(x)$ . Let  $x^+(\epsilon) := x + \epsilon$  with  $\epsilon > 0$ , we'll show that

$$\lim_{\epsilon \rightarrow 0} F(x^+(\epsilon)) = \lim_{\epsilon \rightarrow 0} F(x + \epsilon) = F(x).$$

Firstly, we have

$$F(x^+(\epsilon)) - F(x) = \sum_{n; r_n \leq x+\epsilon} \frac{1}{2^n} - \sum_{n; r_n \leq x} \frac{1}{2^n} = \sum_{n; x < r_n \leq x+\epsilon} \frac{1}{2^n},$$

and we want to show

$$\lim_{\epsilon \rightarrow 0} F(x^+(\epsilon)) - F(x) = \lim_{\epsilon \rightarrow 0} \sum_{n; x < r_n \leq x+\epsilon} \frac{1}{2^n} = 0.$$

**Remark.** The strict is crucial to show the result, since if  $x = r_k$  for some fixed  $k$ , then we can't make the summation arbitrarily small.

<sup>a</sup>This is trivial since we're always going to sum more strictly positive terms in  $F(y)$  than in  $F(x)$ .

Before we show how we choose  $\epsilon$ ,<sup>b</sup> we see that

$$\sum_{n=k}^{\infty} \frac{1}{2^n} = 2^{1-k}.$$

Now, we observe that

$$\sum_{n; x < r_n \leq x + \epsilon} \frac{1}{2^n} \leq \sum_{n = \arg \min_k x < r_k \leq x + \epsilon}^{\infty} \frac{1}{2^n} = 2^{1-k}.$$

With this observation, it should be fairly easy to see that we can choose  $\epsilon$  based on how small we want to make  $2^{1-k}$  be,<sup>c</sup> and we indeed see that

$$\lim_{k \rightarrow \infty} 2^{1-k} = 0,$$

which implies that  $F$  is right-continuous by squeeze theorem. ■

<sup>b</sup>To be precise, how  $\epsilon$  depends on  $r_n$ .

<sup>c</sup>We're referring to  $\epsilon - \delta$  proof approach.

**Lemma A.1.2.** The function  $F$  defined in [this example](#) satisfies

- $\mu_F(\{r_i\}) > 0$  for all  $r_i \in \mathbb{Q}$ .
- $\mu_F(\mathbb{R} \setminus \mathbb{Q}) = 0$

given in [this example](#).

**Proof.** We prove them one by one. And notice that  $F$  is indeed a distribution function as we proved in [Lemma A.1.1](#).

1. To show  $\mu_F(\{r\}) > 0$  for every  $r \in \mathbb{Q}$ , we first note that

$$\{r\} = \bigcap_{a-1 \leq x < r} (x, r].$$

Then, we see that

$$\mu_F(\{r\}) = \mu_F\left(\bigcap_{a-1 \leq x < a} (x, r]\right),$$

where each  $(x, r] \in \mathcal{A}$  and  $(x, r] \supset (y, r]$  whenever  $r-1 \leq x \leq y < r$ . Notice that we implicitly assign the order of the index by the order of  $x$ . Then, we see that  $\mu_F(r-1, r] < \infty$ .<sup>a</sup> Then, from [continuity from above](#), we see that

$$\mu_F(\{r\}) = \lim_{i \rightarrow \infty} \mu_F((x_i, r]),$$

where we again implicitly assign an order to  $x$  as the usual order on  $\mathbb{R}$  by given index  $i$ . It's then clear that as  $i \rightarrow \infty$ ,  $x_i \rightarrow r$ . From the definition of  $F$ , we see that

$$F((x_i, r]) = F(r) - F(x_i) = \sum_{n; r_n \leq r} \frac{1}{2^n} - \sum_{n; r_n \leq x_i} \frac{1}{2^n} = \sum_{n; x_i < r_n \leq r} \frac{1}{2^n}.$$

It's then clear that since  $r \in \mathbb{Q}$ , there exists an  $i'$  such that  $r_{i'} = r$ . Then, we immediately see that no matter how close  $x_i \rightarrow r$ , this sum is at least

$$\frac{1}{2^{i'}}$$

for a fixed  $i'$ . Hence, we conclude that  $\mu_F(\{r\}) > 0$  for every  $r \in \mathbb{Q}$ .



2. Now, we show  $\mu_F(\mathbb{R} \setminus \mathbb{Q}) = 0$ . Firstly, we claim that

$$\mu_F(\mathbb{Q}) = 1$$

and

$$\mu_F(\mathbb{R}) = 1$$

as well. Since  $\mu_F(\mathbb{Q}) = 1$  is clear, we note that the second one essentially follows from the fact that we can write

$$\mathbb{R} = \lim_{N \rightarrow \infty} \bigcup_{i=1}^N (a - i, a + i]$$

for any  $a \in \mathbb{R}$ , say 0. From [continuity from below](#), we have

$$\mu_F \left( \bigcup_{i=1}^{\infty} (-i, +i] \right) = \lim_{n \rightarrow \infty} \mu_F((-n, n]) = \sum_{n; r_n \in \mathbb{Q}} \frac{1}{2^n} = 1.$$

Given the above, from countable additivity of  $\mu_F$ , we have

$$\mu_F(\mathbb{R} \setminus \mathbb{Q}) + \underbrace{\mu_F(\mathbb{Q})}_1 = \underbrace{\mu_F(\mathbb{R})}_1 \Rightarrow \mu_F(\mathbb{R} \setminus \mathbb{Q}) = 0$$

as we desired. ■

---

<sup>a</sup>This will be  $\mu(A_1)$  in the condition of [continuity from above](#). Furthermore, since  $\mathbb{Q}$  is countable, hence  $F(x) < \infty$  is promised.

**Lemma A.1.3** (Cantor set has measure 0). Let  $C$  denotes the [middle thirds Cantor set](#), then the [Lebesgue measure](#) of  $C$  is 0. i.e.,

$$m(C) = 0.$$

**Proof.** Since we're removing  $\frac{1}{3}$  of the whole interval at each  $n$ , we see that the measure of those removing parts, denoted by  $r$ , is

$$m(r) = \sum_{n=1}^{\infty} \frac{2^{n-1}}{3^n} = \frac{1}{2} \sum_{n=1}^{\infty} \left(\frac{2}{3}\right)^n = 1.$$

Then, by [countable additivity](#) of  $m$ , we see that

$$m(C) = m([0, 1]) - m(r) = 1 - 1 = 0. \quad \blacksquare$$

## A.2 Integration

## A.3 Product Measure

## A.4 Differentiation on Euclidean Space

## A.5 Normed-Vector-Space

## A.6 Signed and Complex Measures

### A.6.1 Uniqueness of [Theorem 6.5.2](#)

**Claim.** We show that if  $F \in \text{NBV}$ , then the unique

**Proof.** We observe that

$$\begin{aligned}\mu((-\infty, x]) &= \mu^+((-\infty, x]) - \mu^-((-\infty, x]) \\ &= \nu^+((-\infty, x]) - \nu^-((-\infty, x]) = \nu((-\infty, x]).\end{aligned}$$

Then, since we know that  $\mu^\pm$  is uniquely defined, hence we define

$$F^\pm(x) := \mu^\pm((-\infty, x]), \quad G^\pm(x) := \nu^\pm((-\infty, x]).$$

Then we see that

$$F^+ + G^- = G^+ + F^-,$$

while both left-hand side and right-hand sides are right-continuous and increasing. We hence associate it with the Lebesgue-Stieljes measure, which is uniquely defined, denoted it as  $\lambda$ . Hence, we see that

$$\lambda((a, b]) = (F^+ + G^-)(b) - (F^+ + G^-)(a) = (G^+ + F^-)(b) - (G^+ + F^-)(a)$$

which implies

$$\begin{aligned}(\mu^+((-\infty, b]) + \nu^-((-\infty, b])) - (\mu^+((-\infty, a]) + \nu^-((-\infty, a])) \\ = (\nu^+((-\infty, b]) + \mu^-((-\infty, b])) - (\nu^+((-\infty, a]) + \mu^-((-\infty, a])),\end{aligned}$$

i.e.,

$$\mu^+((a, b]) + \nu^-((a, b]) = \nu^+((a, b]) + \mu^-((a, b])$$

which further implies

$$\underbrace{\mu^+((a, b]) - \mu^-((a, b])}_{\mu((a, b])} = \underbrace{\nu^+((a, b]) - \nu^-((a, b])}_{\nu((a, b])}.$$

Since this holds for all  $(a, b]$ , while  $(a, b]$  generates  $\mathcal{B}(\mathbb{R})$ , hence we have that  $\mu = \nu$ . ⊛

## A.7 Hilbert Spaces

## A.8 Introduction to Fourier Analysis

# Bibliography

- [Axl19] S. Axler. *Measure, Integration & Real Analysis*. Graduate Texts in Mathematics. Springer International Publishing, 2019. ISBN: 9783030331429. URL: <https://books.google.com/books?id=8hCDyQEACAAJ>.
- [DM72] H. Dym and H.P. McKean. *Fourier Series and Integrals*. Probability and Mathematical Statistics. Elsevier Science, 1972. ISBN: 9780122264511. URL: <https://books.google.com/books?id=atkQAQAAIAAJ>.
- [FF99] G.B. Folland and G.B.A. FOLLAND. *Real Analysis: Modern Techniques and Their Applications*. A Wiley-Interscience publication. Wiley, 1999. ISBN: 9780471317166. URL: <https://books.google.com/books?id=uPkYAQAAIAAJ>.
- [Tao13] T. Tao. *An Introduction to Measure Theory*. Graduate studies in mathematics. American Mathematical Society, 2013. ISBN: 9781470409227. URL: <https://books.google.com/books?id=SPGJjwEACAAJ>.