

MATH681  
Mathematical Logic

Pingbang Hu

January 10, 2023

## Abstract

This is a graduate-level mathematical logic course taught by [Matthew Harrison-Trainor](#), aiming to obtain insights into all other branches of mathematics, such as algebraic geometry, analysis, etc. Specifically, we will cover model theory beyond the basic foundational ideas of logic.

While there are no required textbooks, some books do cover part of the material in the class. For example, Marker's *Model Theory: An Introduction* [[Mar02](#)], Hodges's *A Shorter Model Theory* [[HH97](#)], and Hinman's *Fundamentals of Mathematical Logic* [[Hin05](#)].



# Contents

<b>1</b>	<b>Language, Logic, and Structures</b>	<b>2</b>
1.1	Syntax and Semantics . . . . .	3
1.2	First-Order Logic . . . . .	5

# Chapter 1

## Language, Logic, and Structures

### Lecture 1: Introduction to Mathematical Logic

The goal of mathematical logic is to obtain insights into other areas of mathematics – algebra, analysis, combinatorics, and so on, by formalizing the **process** of mathematics. 5 Jan. 14:30

**Remark.** More concretely, there are different branches:

- (a) Model Theory: Study subsets of an object defined by a formula (i.e., first-order logic).
- (b) Computability Theory / Recursion Theory: Formalizing what it means to have an algorithm and studying relative computability.
- (c) Set Theory: Study the structure of the mathematical universe.
- (d) Proof Theory: Study the syntactic nature of proofs.

In this class, we study model theory in nature; specifically, we will cover

- basic definitions of logic:
  - What is a formula?
  - What does it mean for a formula to be true?
  - What is a proof?
- Soundness & completeness theorems:
  - Anything provable is true.
  - Anything true is provable.
- Compactness theorem:
  - Non-standard objects exist.
- Using compactness theorem for applications:
  - Chevalley's theorem

The main theme of this course will be *syntax* v.s. *semantics*:

Syntax	v.s.	Semantics
proofs		truth
form of a formula		mathematical structures
number and type of quantifiers		isomorphisms, embeddings

## 1.1 Syntax and Semantics

### 1.1.1 Languages and Structures

Let's start with the fundamental object, [language](#).

**Definition 1.1.1 (Language).** A *language*  $\mathcal{L}$  consists of:

- a set  $\mathcal{F}$  of function symbols  $f$  with arities  $n_f$ ;
- a set  $\mathcal{R}$  of relation symbols  $R$  with arities  $n_R$ ;
- a set  $\mathcal{C}$  of constant symbols  $c$ .

A [language](#) is also sometimes called a *signature*, in which case we use  $\sigma$  rather than  $\mathcal{L}$ .

**Note.** A constant is the same as a 0-ary function.

**Remark.** Any or all sets in [Definition 1.1.1](#) might be empty.

**Example (Graph).** The [language](#) of graphs,  $\mathcal{L}_{\text{graph}} = \{\mathcal{E}\}$  where  $\mathcal{E}$  is a binary (2-ary) relation symbol.

**Example (Ring).** The [language](#) of rings,  $\mathcal{L}_{\text{ring}} = \{0, 1, +, \cdot, -\}$ , where  $0, 1$  are constants,  $+, \cdot$  are binary functions, and  $-$  is a unary function.

**Example (Ordered ring).** The [language](#) of ordered rings,  $\mathcal{L}_{\text{ord}} = \mathcal{L}_{\text{ring}} \cup \{\leq\}$  where  $\leq$  is the binary relation for an ordered ring.

Then, given a [language](#), we can now interpret it in the following way.

**Definition 1.1.2 (Structure).** Given a [language](#)  $\mathcal{L}$ , an  $\mathcal{L}$ -*structure*  $\mathcal{M}$  consists of:

- a non-empty set  $M$  called the *universe*, *domain*, or *underlying set* of  $\mathcal{M}$ ;
- for each function symbol  $f \in \mathcal{F}$ , a function  $f^{\mathcal{M}}: M^{n_f} \rightarrow M$ ;
- for each relation symbol  $R \in \mathcal{R}$ , a relation  $R^{\mathcal{M}} \subseteq M^{n_R}$ ;
- for each constant symbol  $c \in \mathcal{C}$ , an element  $c^{\mathcal{M}} \in M$ .

**Note (Interpretation).** We call  $f^{\mathcal{M}}, R^{\mathcal{M}}, c^{\mathcal{M}}$  the *interpretation in  $\mathcal{M}$*  of symbols  $f, R, c$ , respectively.

Basically, a [structure](#) gives meaning to the symbols from the [language](#), and we often write

$$\mathcal{M} = (M, f^{\mathcal{M}}, \dots, R^{\mathcal{M}}, \dots, c^{\mathcal{M}}, \dots) = (M, f^{\mathcal{M}}, R^{\mathcal{M}}, c^{\mathcal{M}}: f \in \mathcal{F}, R \in \mathcal{R}, c \in \mathcal{C}).$$

**Notation.** We usually use  $\mathcal{M}, \mathcal{N}, \dots, \mathcal{A}, \mathcal{B}, \dots$  to refer to [structures](#), and  $M, N, \dots, A, B, \dots$  for the domains.<sup>a</sup>

<sup>a</sup>Some people use  $|\mathcal{M}|$  for the domain of  $\mathcal{M}$ .

It's time to look at some examples.

**Example.** The rationals  $\mathbb{Q}$  and integers  $\mathbb{Z}$  are both  $\mathcal{L}_{\text{ring}}$ -structures.

**Proof.** Clearly, the domain is the set of rationals, and naively, we let  $+^{\mathbb{Q}} = +$  in  $\mathbb{Q}$ ,  $0^{\mathbb{Q}} = 0$  in  $\mathbb{Q}$ ,  $1^{\mathbb{Q}} = 1$  in  $\mathbb{Q}$ , etc. In this way,  $\mathbb{Q} = (\mathbb{Q}, 0, 1, +, \cdot, -)$  is an  $\mathcal{L}_{\text{ring}}$ -structure. Similarly,  $\mathbb{Z} =$

$(\mathbb{Z}, 0, 1, +, \cdot, -)$  is as well. ⊗

While the [language](#) we have seen are all intuitively correct with their name, i.e.,  $\mathcal{L}_{\text{ring}}$ ,  $\mathcal{L}_{\text{ord}}$ , and  $\mathcal{L}_{\text{graph}}$ , they are really just the high-level abstraction of the objects in the subscript.

**Example.** Nothing forces an  $\mathcal{L}_{\text{ring}}$ -structure to be a ring.

**Proof.** Since an  $\mathcal{L}_{\text{ring}}$ -structure is just any [structure](#) with two binary functions, a unary function, and two constants interpreting the symbols of the [language](#); hence we can define an  $\mathcal{L}_{\text{ring}}$ -structure  $\mathcal{M}$  as

- $\mathcal{M} = \{0, 5, 11\}$ ;
- $0^{\mathcal{M}} = 5$ ;
- $1^{\mathcal{M}} = 11$ ;
- $+^{\mathcal{M}}$  is the constant function 0;
- $\cdot^{\mathcal{M}}$  is the function 5;
- $-^{\mathcal{M}}$  is the identity.

This is clearly not a ring since it fails nearly every axiom of a ring. ⊗

**Note.** Later, we will talk about theories that let us restrict to structures we want.

### 1.1.2 Embeddings and Isomorphisms

We can now consider the relation between [structures](#).

**Definition 1.1.3 (Embedding).** Given a [language](#)  $\mathcal{L}$  and let  $\mathcal{M}$  and  $\mathcal{N}$  be  $\mathcal{L}$ -structures. A map  $\eta: \mathcal{M} \rightarrow \mathcal{N}$  is an  $\mathcal{L}$ -embedding if it is one-to-one and preserves the interpretation of all symbols of  $\mathcal{L}$ :

- (a) for each  $f \in \mathcal{F}$  of arity  $n_f$ , and  $a_1, \dots, a_{n_f} \in \mathcal{M}$ ,

$$\eta(f^{\mathcal{M}}(a_1, \dots, a_{n_f})) = f^{\mathcal{N}}(\eta(a_1), \dots, \eta(a_{n_f}));$$

- (b) for each relation  $R \in \mathcal{R}$  of arity  $n_R$ , and  $a_1, \dots, a_{n_R} \in \mathcal{M}$ ,

$$(a_1, \dots, a_{n_R}) \in R^{\mathcal{M}} \Leftrightarrow (\eta(a_1), \dots, \eta(a_{n_R})) \in R^{\mathcal{N}};$$

- (c) for each constant  $c \in \mathcal{C}$ ,  $\eta(c^{\mathcal{M}}) = c^{\mathcal{N}}$ .

From the definition, an  $\mathcal{L}$ -embedding is an injection, and naturally, we have the following.

**Definition 1.1.4 (Isomorphism).** An  $\mathcal{L}$ -isomorphism is a bijective  $\mathcal{L}$ -embedding.

**Definition.** Given a [language](#)  $\mathcal{L}$  and let  $\mathcal{M}$  and  $\mathcal{N}$  be  $\mathcal{L}$ -structures. Suppose  $M \subseteq N$  and the inclusion map  $\iota: M \hookrightarrow N$  is an  $\mathcal{L}$ -embedding.

**Definition 1.1.5 (Substructure).**  $\mathcal{M}$  is a *substructure* of  $\mathcal{N}$ .

**Definition 1.1.6 (Extension).**  $\mathcal{N}$  is an *extension* of  $\mathcal{M}$ .

**Example.** Ring [embeddings](#) are  $\mathcal{L}_{\text{ring}}$ -embeddings.

This generalizes the notions of embedding and isomorphism for many kinds of mathematical structures.

**Remark.** Asking that  $\eta$  be injective is the same as (b) in Definition 1.1.3 for the relation  $=$  of equality since

$$a = b \in \mathcal{M} \Leftrightarrow \eta(a) = \eta(b) \in \mathcal{N}.$$

However, the notion of **substructure** is **language** sensitive. For groups, there are two possible **languages**:

- (a)  $\mathcal{L}_1 = \{e, \cdot\}$ ;
- (b)  $\mathcal{L}_2 = \{e, \cdot, {}^{-1}\}$ , i.e., with the unary inverse operation.

While both seem OK at first glance, we should use the second one.

**Remark.** Using  $\mathcal{L}_2$ , the **substructure** of a group is the same thing as a subgroup. But if we use  $\mathcal{L}_1$ , then  $(\mathbb{N}, +, 0)$  is a **substructure** of  $(\mathbb{Z}, +, 0)$ , while  $\mathbb{N}$  is not a group for sure.

**Proof.** Simply observe that both  $(\mathbb{N}, 0, +)$ ,  $(\mathbb{Z}, 0, +)$  are  $\mathcal{L}_1$ -structures. ⊗

Similarly, we include  $-$  in  $\mathcal{L}_{\text{ring}}$  for a similar reason as in the previous **example**.

**Example.** An  $\mathcal{L}_{\text{ring}}$ -**substructure** of a field will be a subring, not a subfield. If we want subfields, use  $\mathcal{L}_{\text{ring}} \cup \{{}^{-1}\}$ .<sup>a</sup>

<sup>a</sup>We can set  $0^{-1} = 0$ , but never use this.

## Lecture 2: Formulas and First-Order Logic

We start by asking that given a function symbol  $f$  of arity  $n$ , could we replace  $f$  with an  $(n+1)$ -ary  $R$  relation for its graph? 10 Jan. 14:30

**Example.** Let  $\mathcal{L}$  be a **language** with only relation symbols. Let  $\mathcal{A}$  be an  $\mathcal{L}$ -**structure**. For any  $B \subseteq A$ , there is a **substructure**  $B$  of  $A$  with domain  $B$ .

What is  $B$ ? For each relation symbol  $R$ , let  $R^B = R^A \cap B^{n_R}$ , and this will make  $B$  a **substructure** of  $A$ .

The above is not true for function symbols though.

**Example.** If  $G = (\mathbb{Z}, 0, +)$ , then  $\mathbb{N}$  is not the domain of a subgroup. So if we took  $\mathcal{L} = \{0, +, {}^{-1}\}$ , where  $0$  is the unary relation,  $+$  is the ternary relation, and  ${}^{-1}$  is the binary relation, an  $\mathcal{L}$ -**substructure** of a group might not be a subgroup.

## 1.2 First-Order Logic

### 1.2.1 Terms, Formulas, and Truths

Intuitive, an  $\mathcal{L}$ -**formula** is an expression built using the symbols in a **language**  $\mathcal{L}$ ,  $=$ , the logical connectives  $\wedge, \vee, \neg$ , and variable symbols  $v_1, v_2, \dots, x, y, z$ , and also quantifiers  $\exists$  and  $\forall$ .

**Definition 1.2.1 (Term).** Given a **language**  $\mathcal{L}$ , the set of  $\mathcal{L}$ -*terms* are defined inductively by:

- (a) Each constant symbol is a *term*.
- (b) Each variable symbol  $v_1, \dots$  is a *term*.
- (c) If  $f$  is a function symbol, and  $t_1, \dots, t_{n_f}$  are **terms**, then  $f(t_1, \dots, t_{n_f})$  is a *term*.

If  $\mathcal{M}$  is an  $\mathcal{L}$ -structure, and  $t$  is a term involving only variables among  $v_1, \dots, v_n$ , then  $t$  has an interpretation  $t^{\mathcal{M}}: M^n \rightarrow M$ .

Then, we define  $t^{\mathcal{M}}$  inductively as follows: On input  $a_1, \dots, a_n \in M$

- (a) If  $t$  is a constant  $c$ ,

$$t^{\mathcal{M}}(a_1, \dots, a_n) = c^{\mathcal{M}}.$$

- (b) If  $t$  is a variable  $v_i$ ,

$$t^{\mathcal{M}}(a_1, \dots, a_n) = a_i.$$

- (c) If  $t$  is  $f(s_1, \dots, s_k)$ , then

$$t^{\mathcal{M}}(a_1, \dots, a_n) = f^{\mathcal{M}}(s_1^{\mathcal{M}}(a_1, \dots, a_n), \dots, s_k^{\mathcal{M}}(a_1, \dots, a_n)).$$

**Intuition.** We are basically substituting for variables and evaluating the expression.

**Example.** In  $(\mathbb{R}, 0, 1, +, \cdot, -)$ , technically, a term looks like

$$\cdot(+ (1, 1), + (x, y)),$$

but we will write terms the natural way, i.e.,

$$(1 + 1)(x + y).$$

Also, we will use  $\underline{n}$  or  $n$  to represent the term

$$\underline{n} = \underbrace{1 + 1 + \dots + 1}_{n \text{ times}}.$$

So we could write the above term as

$$2 \cdot (x + y)$$

Then, what do the terms in the ring language look like? They are the polynomials with integer coefficients, assuming we interpret them in a ring.

**Definition 1.2.2 (Formula).** Given a language  $\mathcal{L}$ , the  $\mathcal{L}$ -formulas are defined inductively:

- (a) If  $s, t$  are terms,  $s = t$  is a formula.
- (b) If  $R$  is a relation symbol of arity  $n_R$ , and  $s_1, \dots, s_{n_R}$  are term, then  $R(s_1, \dots, s_{n_R})$  is a formula.
- (c) If  $f$  is a formula, then  $\neg f$  is a formula.
- (d) If  $\varphi$  and  $\psi$  are formulas, then  $\varphi \wedge \psi$  and  $\varphi \vee \psi$  are formulas.
- (e) If  $\varphi$  is a formula, and  $v_1$  is a variable,  $\exists v_1 \varphi$  and  $\forall v_1 \varphi$  are formulas.

**Notation** (Atomic formula). Formulas of the form (a) and (b) in Definition 1.2.2 are called *atomic formulas*.

**Notation** (Quantifier-free formula). Formulas of the form (a), (b), (c), and (d) in Definition 1.2.2 are called *quantifier-free formulas*.

**Example.** We can say that an element  $x$  of a ring has a square root by

$$\exists y \, y^2 = x$$



**Example.** A group is torsion of order 2 can be said by

$$\forall x \, x \cdot x = e.$$

**Example.** We can write down all the field/group/... axioms as **formulas**.

Notice that for the first example, the **formula**  $\exists y \, y^2 = x$  only has meaning if we assign what  $x$  is. In this case, we say that  $y$  is *bound* by  $\exists y$ . But this is local:

**Example.** Consider

$$y = 1 \wedge \exists y \, y^2 = x,$$

while the first appearance of  $y$  is free, the second appearance of  $y$  is bound by  $\exists y$ , or we say that  $y$  is in the scope of  $\exists y$ .

While our definitions work perfectly fine with the above example, but sometimes we don't want this to happen. In such a case, we simply replace the bound instances of  $y$  with a new variable  $z$ .

**Definition 1.2.3 (Free variable).** The *free variables*  $\text{FV}(\varphi)$  of a **formula**  $\varphi$  are defined inductively:

- (a)  $\text{FV}(s = t)$  is the set of variables showing up in  $s$  or  $t$ .
- (b)  $\text{FV}(R(s_1, \dots, s_{n_R}))$  is the set of variables showing up in  $s_1, \dots, s_{n_R}$ .
- (c)  $\text{FV}(\neg\varphi) = \text{FV}(\varphi)$ .
- (d)  $\text{FV}(\varphi \wedge \psi) = \text{FV}(\varphi \vee \psi) = \text{FV}(\varphi) \cup \text{FV}(\psi)$ .
- (e)  $\text{FV}(\exists x \, \varphi) = \text{FV}(\forall x \, \varphi) = \text{FV}(\varphi) \setminus \{x\}$ .

**Example.**  $\text{FV}(\exists y \, y^2 = x) = \{x\}$ .

**Example.**  $\text{FV}(\forall x \, x \cdot x = e) = \emptyset$ .

**Definition 1.2.4 (Sentence).** A **formula**  $\varphi$  is called a *sentence* if it has no **free variables**.

**Notation.** If  $\varphi$  is a **formula** with **free variables** among  $x_1, \dots, x_n$  we often write  $\varphi(x_1, \dots, x_n)$ .

**Remark.** So given  $\varphi(x_1, \dots, x_n)$ , we know that  $\varphi$  has no other **free variables**.

**Example.** It's valid to write  $\varphi(x, y, z) := x = y$ .

**Definition 1.2.5 (Truth).** Given a **language**  $\mathcal{L}$  and an  $\mathcal{L}$ -**structure**  $\mathcal{M}$ , let  $\varphi(x_1, \dots, x_n)$  be an  $\mathcal{L}$ -**formula**. Let  $a_1, \dots, a_n \in \mathcal{M}$ . Define  $\mathcal{M} \models \varphi(\bar{a})^a$  as follows:

- (a) If  $\varphi$  is  $s = t$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if  $s^{\mathcal{M}}(\bar{a}) = t^{\mathcal{M}}(\bar{a})$ .
- (b) If  $\varphi$  is  $R(t_1, \dots, t_{n_R})$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if  $(t_1^{\mathcal{M}}(\bar{a}), \dots, t_{n_R}^{\mathcal{M}}(\bar{a})) \in R^{\mathcal{M}}$ .
- (c) If  $\varphi$  is  $\neg\psi$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if  $\mathcal{M} \not\models \psi(\bar{a})$ .
- (d) If  $\varphi$  is  $\psi_1 \wedge \psi_2$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if  $\mathcal{M} \models \psi_1(\bar{a})$  and  $\mathcal{M} \models \psi_2(\bar{a})$ .
- (e) If  $\varphi$  is  $\psi_1 \vee \psi_2$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if  $\mathcal{M} \models \psi_1(\bar{a})$  or  $\mathcal{M} \models \psi_2(\bar{a})$ .
- (f) If  $\varphi$  is  $\exists y \, \psi(\bar{x}, y)$ ,<sup>b</sup> then  $\mathcal{M} \models \varphi(\bar{a})$  if there's  $b \in \mathcal{M}$  such that  $\mathcal{M} \models \psi(\bar{a}, b)$ .

---

(g) If  $\varphi$  is  $\forall y \psi(\bar{x}, y)$ , then  $\mathcal{M} \models \varphi(\bar{a})$  if for all  $b \in \mathcal{M}$  such that  $\mathcal{M} \models \psi(\bar{a}, b)$ .

<sup>a</sup>We read this as  $\varphi$  is true of  $\bar{a}$  in  $\mathcal{M}$ .

<sup>b</sup>Recall that  $\bar{x} = (x_1, \dots, x_n)$ .

# Appendix

# Bibliography

- [HH97] W. Hodges and S.M.S.W. Hodges. *A Shorter Model Theory*. Cambridge University Press, 1997. ISBN: 9780521587136. URL: <https://books.google.com/books?id=S6QYeuo4p1EC>.
- [Hin05] P.G. Hinman. *Fundamentals of Mathematical Logic*. Taylor & Francis, 2005. ISBN: 9781568812625. URL: <https://books.google.com/books?id=xA6D8o72qAgC>.
- [Mar02] D. Marker. *Model Theory : An Introduction*. Graduate Texts in Mathematics. Springer New York, 2002. ISBN: 9780387987606. URL: <https://books.google.com/books?id=gkvogoiEnuYC>.