Local exploration laws and regular driving characteristics

Autonomous mathematical investigation, with independent critical checks

26 October 2026

Abstract

We investigate the corrected source-neighborhood locality axiom. For continuous semimartingale Loewner drivers with absolutely continuous, right-continuous drift, locality itself gives an initial quadratic-variation rate. Two shrinking mirrored slits show that local boundedness of the initial drift forces zero drift and variance six. Strong domain Markov then identifies SLE_6, without assuming conformal covariance or Brownian driving. The exact instability rates are inverse-linear and inverse-cubic in the slit parameter. Separately, exact scaling proves that a fixed nonconformal affine SLE_6 deformation cannot have the stated tame drift structure in every bounded domain. These are conditional results: the analytic hypotheses have not been derived from RSW or from the project axioms, and the corrected classification and universality remain open.

1 Current locality and scope

Two colored marked domains that agree near their common source must have identical initial exploration laws until exit from that neighborhood, even if their remote targets differ. This is the user's corrected locality axiom at commit `0e47018`. We fix one color orientation throughout. The terminal-germ construction for the earlier fixed-target axiom is excluded; its original proofs are retained in `archive/original-locality-results.tex` and the detailed research record.

The first theorem below needs only a deletion consequence of corrected locality. It does not claim that locality supplies regular driving drift. For rough restart domains we state the prime-end neighborhood or admissible test-deletion requirement explicitly. All driving clocks use $\partial_t g = 2/(g - W)$ and hence $\mathrm{hcap}(K_t) = 2t$. The standard conformal-map comparison for Loewner chains is derived below; ordinary SLE facts used for affine examples are described in [1, 2]. Continuous Loewner generation and infinite lifetime of affine SLE images were proved in the companion research record [3].

2 A near-source drift-boundedness criterion

Status. This is a new conditional rigidity result. It does not assume conformal invariance, Brownian driving, or local-martingale driving. Instead it shows that a locally bounded *initial driving drift*, together with an absolutely continuous drift part and locality, already forces zero drift and variance six. No regularity of quadratic variation is assumed. Strong DMP then identifies the whole driving law. The boundedness and characteristic hypotheses are not derived from RSW or from the three project axioms. A finite-dimensional bulk-jet closure is given as one explicit sufficient hypothesis.

Only a particular deletion consequence of the corrected source-neighborhood locality is needed. Arbitrary target changes and the full force of corrected locality are not used in this theorem. Thus the result is compatible with, but does not exhaust, the corrected research objective.

2.1 Chart-dependent initial characteristics and their covariance

Let $f : \mathbb{H} \to D$ be an inverse chordal conformal chart, with source $f(0) = p$ and target $f(\infty) = q$, both interpreted as prime ends. For the physical ensemble in D, let W^f be its driving process in this chart, with half-plane capacity normalization

$$\partial_t g_t(z) = \frac{2}{g_t(z) - W_t^f}, \qquad W_0^f = 0.$$

Assume that it is a continuous semimartingale in its natural trace filtration, with decomposition

$$W_t^f = M_t^f + \int_0^t b_s^f \, ds, \qquad C_t^f = \langle M^f \rangle_t. \tag{DR1}$$

The drift-density paths b^f are assumed finite and right-continuous, after local stopping if necessary. No absolute continuity or regularity of C^f is assumed. At the initial time suppose

$$b_0^f = B(f) \in \mathbb{R} \tag{DR2}$$

is deterministic. It may depend on the entire physical marked domain and its conformal chart. In particular no Markov property of the single real driver is assumed. A finite deterministic initial variance rate $V(f)$ will be *derived* below from locality and (DR1)–(DR2).

Let $A \subset \overline{\mathbb{H}}$ be a half-plane hull disjoint from 0, let $h : \mathbb{H} \setminus A \to \mathbb{H}$ be hydrodynamically normalized, and set

$$\bar{h}(z) = h(z) - h(0), \qquad f_A = f \circ \bar{h}^{-1}. \tag{DR3}$$

The source is still the same physical prime end, since $\bar{h}(0) = 0$; the target is also unchanged. Put $d = h'(0) > 0$, $e = h''(0)$. Source locality, together with the usual deletion-chart calculation, will give a finite deterministic initial variance rate $V(f)$ and the covariance

$$\boxed{V(f_A) = V(f), \qquad B(f_A) = \frac{B(f)}{d} + \left(\frac{V(f)}{2} - 3\right)\frac{e}{d^2}.} \tag{DR4}$$

These are identities for *initial* characteristics and do not identify the two complete driving laws.

Here are the details without assuming a quadratic-variation density. If h_t is the evolving deletion map, then $du = h_t'(W_t)^2 dt$, and Itô's formula gives

$$d\widehat{W}_{u(t)} = h_t'(W_t)\, dM_t + h_t'(W_t) b_t \, dt + \frac{1}{2} h_t''(W_t)\, dC_t - 3h_t''(W_t)\, dt. \tag{DR5}$$

The -3 comes from

$$\partial_t h_t(z) = \frac{2h_t'(W_t)^2}{h_t(z) - h_t(W_t)} - \frac{2h_t'(z)}{z - W_t}, \qquad (\partial_t h_t)(W_t) = -3h_t''(W_t).$$

The comparison of filtrations is justified by geometric invertibility: from a deleted-domain initial trace one reconstructs its original trace and capacity, so the inverse capacity clock is adapted and its inverse is a stopping-time clock. Locality therefore compares the same trace filtration, not an unrelated enlargement. Adaptedness of an arbitrary increasing clock would not suffice for this step.

Local bracket-regularization lemma (proved). Suppose $h''(0) \neq 0$. Then on a positive initial interval, C_t is absolutely continuous and has a finite right-continuous density. In particular its initial right derivative $V(f)$ exists and is deterministic.

Proof. The deleted-domain driver also has an absolutely continuous drift, $\int \widehat{b}_u \, du$, by (DR1). Uniqueness of the continuous semimartingale decomposition applied to (DR5), after the justified capacity time change, gives the signed-measure identity

$$h_t''(W_t)(dC_t - 6\,dt) = 2\big[h_t'(W_t)^2 \widehat{b}_{u(t)} - h_t'(W_t) b_t\big] dt. \tag{DR5a}$$

The second derivative starts nonzero and remains so on a positive initial interval. On that interval the bracket is absolutely continuous, with

$$v_t = 6 + 2\frac{h_t'(W_t)^2 \widehat{b}_{u(t)} - h_t'(W_t) b_t}{h_t''(W_t)}. \tag{DR5b}$$

The derivative paths of h_t along the continuous driving point are continuous, while the two drift-density paths are right-continuous and finite. Thus v is right-continuous and finite there. In particular

$$V(f) = v_0 = 6 + 2\frac{d^2 B(f_A) - dB(f)}{e} \tag{DR5c}$$

is deterministic. This is an intrinsic initial rate of C, so it does not depend on which nonzero-e deletion was used to derive it. Applying the same time change to bracket gives $V(f_A) = V(f)$. Rearranging (DR5c) gives the drift identity (DR4). □

There is always an available nonzero-e test deletion, for example one vertical slit at any fixed real point different from 0. Right-continuity makes the initial drift and variance rates intrinsic, rather than arbitrary versions of densities at a single time. The bracket density has only been proved locally at the start; its global regularity is not being presumed.

Source-neighborhood convention. In half-plane coordinates these slit deletions preserve a full neighborhood of the source end. Under f this is a prime-end neighborhood. Thus (DR4) is justified on arbitrary rough domains if the corrected locality assumption is formulated for such neighborhoods, or if these particular deletions are explicitly admissible. For regular analytic source boundaries it is ordinary Euclidean neighborhood locality. If locality is instead stated only using Euclidean balls at non-singleton source impressions, the all-rough-domain extension must remain an explicit additional hypothesis.

2.2 Two vanishing slits force both characteristics

For $\epsilon > 0$ take the reflected vertical slits

$$A_\epsilon^{\pm} = \{\, \pm\epsilon^2 + iy : 0 \le y \le 2\epsilon \,\}. \tag{DR6}$$

Their hydrodynamic maps are

$$h_\epsilon^{\pm}(z) = \pm\epsilon^2 + \sqrt{(z \mp \epsilon^2)^2 + 4\epsilon^2}, \tag{DR7}$$

with the square-root branch asymptotic to $z \mp \epsilon^2$ at infinity. Let $f_\epsilon^{\pm} = f \circ (h_\epsilon^{\pm} - h_\epsilon^{\pm}(0))^{-1}$. Both sequences are conformal charts of bounded simply connected domains and

$$f_\epsilon^{\pm} \longrightarrow f \quad \text{locally uniformly in } \mathbb{H}. \tag{DR8}$$

Indeed the slits have diameter 2ϵ and approach 0; directly from (DR7), their maps and inverse maps converge to the identity on compact subsets of $\mathbb{H}$, while $h_\epsilon^\pm(0) \to 0$. Crucially their boundary derivatives do *not* converge to the identity jet:

$$d_\epsilon = (h_\epsilon^\pm)'(0) = \frac{\epsilon}{\sqrt{\epsilon^2+4}} \longrightarrow 0, \qquad e_\epsilon^\pm = (h_\epsilon^\pm)''(0) = \mp\frac{4}{\epsilon(\epsilon^2+4)^{3/2}}. \tag{DR9}$$

In particular

$$\frac{e_\epsilon^\pm}{d_\epsilon^2} = \mp\frac{4}{\epsilon^3\sqrt{\epsilon^2+4}}. \tag{DR10}$$

Thus compact convergence of inverse charts permits very large second-jet changes precisely at the source boundary.

Initial rigidity theorem (proved). Fix f and assume (DR1)–(DR4) for f and the charts $f_\epsilon^\pm$. If

$$\sup_{0<\epsilon<\epsilon_0} \max\{|B(f_\epsilon^+)|, |B(f_\epsilon^-)|\} < \infty \tag{DR11}$$

for some $\epsilon_0 > 0$, then

$$B(f) = 0, \qquad V(f) = 6. \tag{DR12}$$

No continuity of B is required.

Proof. Write $b = B(f)$, $v = V(f)$ and $b_\epsilon^\pm = B(f_\epsilon^\pm)$. Equation (DR4) gives

$$b_\epsilon^\pm = \frac{b}{d_\epsilon} + \left(\frac{v}{2} - 3\right)\frac{e_\epsilon^\pm}{d_\epsilon^2}.$$

Since $e_\epsilon^- = -e_\epsilon^+$,

$$b_\epsilon^+ + b_\epsilon^- = \frac{2b}{d_\epsilon}, \qquad b_\epsilon^+ - b_\epsilon^- = (v-6)\frac{e_\epsilon^+}{d_\epsilon^2}. \tag{DR13}$$

The first right side is bounded only if $b = 0$, since $d_\epsilon \to 0$. The second is bounded only if $v = 6$, since its multiplier has magnitude asymptotic to $2\epsilon^{-3}$. This proves (DR12). □

Exact instability and weaker rate conditions. For real numbers U, C one has $\max\{|U + C|, |U - C|\} = |U| + |C|$. Hence the two drift covariances give the exact identity

$$\boxed{\max\{|b_\epsilon^+|, |b_\epsilon^-|\} = \frac{|b|\sqrt{\epsilon^2+4}}{\epsilon} + \frac{4|v/2-3|}{\epsilon^3\sqrt{\epsilon^2+4}}.} \tag{DR14}$$

There is no possible cancellation between the two reflected tests. In particular, if the displayed maximum is $o(\epsilon^{-3})$, then $v = 6$. If it is $o(\epsilon^{-1})$, then $b = 0$ and $v = 6$. Boundedness is more than sufficient. Conversely, if $v \neq 6$, its leading blow-up is $|v-6|\epsilon^{-3}$; if $v = 6$ and $b \neq 0$, its leading blow-up is $2|b|\epsilon^{-1}$. This is a concrete, quantitative obstruction to any alternative semimartingale driving law with tame initial drift under source perturbations.

2.3 Local boundedness in marked-chart topology suffices

Equip inverse marked charts with locally uniform convergence on $\mathbb{H}$, retaining the fixed preimages $0, \infty$ of source and target. If B is locally bounded at every f in this topology, then (DR8) implies (DR11). In particular continuity of B is sufficient, but unnecessarily strong. This is a hypothesis about the driving characteristic itself; weak convergence of curve laws alone does not imply it.

Conditional classification theorem (proved). Suppose a Borel family of continuous Loewner-generating exploration laws on all bounded simply connected marked prime-end domains has strong DMP and source-neighborhood locality (including the test deletions (DR6) in each chart). Suppose its drivers satisfy (DR1)–(DR2), with finite right-continuous drift-density paths and the usual restart compatibility of characteristic kernels, with no initial assumption on quadratic variation. If the initial drift map B is locally bounded in marked inverse- chart topology, then every driving process is

$$W_t^f = \sqrt{6}\, B_t,$$

and the family is chordal SLE_6 in these conformal coordinates.

Proof. The initial rigidity theorem gives $B(f) = 0$ and the derived $V(f) = 6$ for every marked inverse chart. At any deterministic rational time r, condition on the initial exploration and use strong DMP in its remaining marked domain and current conformal chart. The centered future driver is exactly the restarted driver; its initial drift is b_r^f, by right-continuity. The restarted law has initial drift zero, so $b_r^f = 0$ almost surely. Countably many rational times and right-continuity give $b_t^f = 0$ throughout each localized lifetime. Therefore all domain chart drivers are continuous local martingales.

For completeness the no-density local-martingale locality argument then identifies their brackets. With both drifts zero, (DR5a) becomes $h_t''(W_t)(dC_t - 6\,dt) = 0$. A nonzero-$e$ vertical deletion yields $C_t = 6t$ on a positive initial interval. Let

$$T = \inf\{t \geq 0 : C_t - 6t \neq 0\}.$$

This is an entrance stopping time of a continuous adapted process into an open set, in the usual right-continuous filtration. The initial zero interval gives $T > 0$. If $T < \infty$, continuity gives $C_T = 6T$ and the first-deviation definition gives nonzero values arbitrarily soon after T. Strong DMP restarted at T gives another positive zero interval, a contradiction. Thus $C_t = 6t$ throughout. Lévy's characterization gives Brownian driving, and Loewner generation identifies SLE_6. □

The Borel-kernel condition means that the family admits the standard strong conditional-kernel interpretation on random slit domains, and that the characteristics can be chosen consistently on that family. It is not a claim that deterministic-domain convergence of discrete explorations proves this property. Translation invariance is not needed in this conditional theorem. If its hypotheses hold only after one fixed physical affine normalization, the conclusion is the corresponding fixed-affine SLE_6 family; an affine change of physical coordinates is not presumed to preserve semimartingale driving in an arbitrary preselected complex structure.

2.4 A finite-dimensional closure that is completely classified

For example, fix finitely many interior points $z_j \in \mathbb{H}$ and finitely many nonnegative derivative orders m_j. Let

$$J(f) = (f^{(k)}(z_j) : 1 \leq j \leq N, 0 \leq k \leq m_j).$$

Suppose the initial drift has a representation

$$B(f) = \mathcal{B}(J(f)), \tag{DR15}$$

where $\mathcal{B}$ is locally bounded at every realized jet vector. The bracket has no assumed density or finite-dimensional representation; its initial rate is supplied by the local regularization lemma. Locally uniform convergence of conformal maps implies convergence of all these interior jets by Cauchy's

formula, so (DR15) implies local boundedness of B in marked-chart topology. The conditional classification theorem therefore applies: within (DR1)–(DR2), every source-local strong-DMP family with such a locally bounded finite-dimensional *bulk-jet drift closure* is SLE_6.

This statement does not cover a closure depending on unbounded source boundary derivatives such as $f'(0)$ or a boundary pre-Schwarzian. Those quantities can change without control under (DR8), and can carry exactly the singular terms in (DR13). Nor does it show that true percolation limits have semimartingale driving in a fixed physical complex structure. The new rigorous reduction is that any non-SLE candidate with regular characteristic rates must violate the displayed near-source drift bound, in the explicit ϵ^{-1} or ϵ^{-3} manner above.

2.5 The criterion also detects a wrong affine coordinate system

Corollary (proved). Let A be an orientation-preserving real-linear map which is not a Euclidean similarity, and let the physical ensemble be the fixed-affine pullback

$$P_{D,p,q} = (A^{-1})_* \,\mathrm{Law}(\mathrm{SLE}_6(AD; Ap, Aq)).$$

Although this family satisfies corrected source-neighborhood locality and strong DMP, in the original Euclidean complex structure it cannot satisfy both the regular semimartingale characteristic assumptions (DR1)–(DR2) in all marked domains and the locally bounded initial-drift condition above.

Proof. If both properties held, the conditional classification theorem would make this ensemble ordinary SLE_6 in the original complex structure. The single-domain affine identifiability theorem would then make A a similarity, a contradiction. □

Thus the drift criterion selects a conformal structure. It is not a coordinate-free consequence of coarse crossing bounds: bounded affine changes preserve such bounds, whereas a wrong affine structure necessarily violates at least one of the criterion's additional analytic hypotheses. The corollary does not decide whether the failure is semimartingality, right-continuity of the drift density, deterministic initial drift, or boundedness of the initial drift; distinguishing these failures is a separate question.

2.6 What corrected locality says at a regular analytic source germ

Here is a separate necessary local structure statement that uses the full corrected locality, including changes of the distant target. It concerns initial characteristics only, not existence of a process from arbitrary coefficient choices.

Suppose the physical boundary is analytic near p and the source colors are fixed there. Let G denote this embedded colored source germ. For a chart f extending conformally across 0, define

$$a(f) = |f'(0)| > 0, \qquad P(f) = \mathrm{Re}\,\frac{f''(0)}{f'(0)}. \tag{DR16}$$

If another physical marked domain has the same source germ but any distant target, let k be its inverse chordal chart with $k(0) = p$. The local map $h = k^{-1} \circ f$ is analytic across 0, takes the real boundary interval to a real interval, and has $h(0) = 0$, $d = h'(0) > 0$, $e = h''(0) \in \mathbb{R}$. The two initial physical traces agree by corrected locality. The local Loewner coordinate comparison is exactly (DR4), with the same derivation from pole cancellation and Itô's formula; a globally defined deletion map is unnecessary. Consequently $V(f) = V(k)$ depends only on G.

Proposition (proved). Under the semimartingale assumptions above, corrected locality at analytic source germs is equivalent at the level of initial characteristic covariance to the existence of two source-germ quantities $V(G)$ and $\beta(G)$ such that

$$\boxed{V(f) = V(G), \qquad B(f) = a(f)\,\beta(G) + \left(3 - \frac{V(G)}{2}\right) P(f).} \tag{DR17}$$

Here "equivalent at the level of covariance" means precisely that every covariant initial-characteristic assignment has this form, and every assignment of this form obeys the local chart covariance identity. It does not mean that arbitrary V, β yield a domain Markov exploration family.

Proof. Since $k = f \circ h^{-1}$, the chain rule gives

$$a(k) = \frac{a(f)}{d}, \qquad P(k) = \frac{P(f)}{d} - \frac{e}{d^2}.$$

Write $c = V(G)/2 - 3$. Combining these transformations with $B(k) = B(f)/d + ce/d^2$ yields

$$\frac{B(k) + cP(k)}{a(k)} = \frac{B(f) + cP(f)}{a(f)}.$$

This common value is the source-germ invariant $\beta(G)$, proving (DR17). Conversely direct substitution gives the required covariance. □

Physical translation invariance makes $V(G)$ and $\beta(G)$ invariant under translations of the colored germ. If dilation covariance is separately assumed, then $V(\lambda G) = V(G)$ and $\beta(\lambda G) = \lambda^{-1}\beta(G)$. For a straight source-boundary germ, which is unchanged up to translation by dilation, this would force $\beta(G) = 0$; the dimensionless variance can still depend on the boundary orientation before an affine metric has been identified.

The boundary pre-Schwarzian term in (DR17) explains the quantitative anomaly in the vanishing-slit test. Even when the physical source germ is unchanged, $P(f_\epsilon)$ may grow as ϵ^{-3} while $f_\epsilon \to f$ on every interior compact set. Thus source locality by itself gives a connection-type transformation law, rather than vanishing drift. The locally bounded bulk closure theorem removes exactly this uncontrolled boundary dependence. The missing global step is to understand how DMP constrains these local assignments at rough, dynamically created source germs, where the analytic boundary jet in (DR16) no longer exists.

3 A wrong affine metric cannot have tame driving drift in every bounded domain

Status. The conclusion here is stronger than the earlier wrong-coordinate consequence of drift boundedness. For a fixed non-similarity affine deformation of SLE_6, regular semimartingale driving with absolutely continuous right-continuous drift cannot hold in every bounded domain in the wrong Euclidean complex structure. No local boundedness of the initial drift functional is needed. The proof uses exact scale invariance only for the *known fixed-affine SLE_6 half-plane law*; it does not assume that arbitrary critical subsequential limits are scale invariant. All characteristic regularity assumptions remain explicit.

3.1 Exact Brownian scaling and a regular initial bracket

Scaling lemma (proved). Let W be a continuous real process on $[0,\infty)$ with $W_0 = 0$, whose law has exact Brownian scaling:

$$(c^{-1/2} W_{ct})_{t\geq 0} \overset{\text{law}}{=} (W_t)_{t\geq 0} \qquad (c > 0). \tag{SM1}$$

Suppose up to a strictly positive stopping time it is a continuous semimartingale

$$W_t = M_t + \int_0^t b_s \, ds, \tag{SM2}$$

where $M_0 = 0$ is a continuous local martingale and b is finite and right-continuous at 0. Suppose its bracket has a finite deterministic initial right derivative

$$\lim_{t\downarrow 0} \frac{\langle M\rangle_t}{t} = v \in [0,\infty) \quad \text{almost surely.} \tag{SM3}$$

Then the entire law of W is that of $\sqrt{v}B$. This includes the identically zero process if $v = 0$.

Proof. Fix $T < \infty$ and stop the local decomposition at a positive initial localizing time. As $c \downarrow 0$, with probability tending to one the whole interval $[0, cT]$ lies before that time. Right-continuity of b gives

$$\sup_{t\leq T} \left| c^{-1/2} \int_0^{ct} b_s \, ds \right| \leq \sqrt{c}\, T \sup_{s\leq cT} |b_s| \longrightarrow 0$$

almost surely on the localized event. Writing $C_t = \langle M\rangle_t$, (SM3) implies

$$\sup_{t\leq T} |C_{ct}/c - vt| \longrightarrow 0. \tag{SM4}$$

Indeed $|C_{ct}/c - vt| \leq t \sup_{0<s\leq cT} |C_s/s - v|$. The continuous-martingale functional central limit theorem now gives $c^{-1/2} M_{c\cdot} \Rightarrow \sqrt{v}B$ on $C[0,T]$. One may see this directly from Dambis–Dubins–Schwarz: extend its Brownian motion if necessary and write

$$c^{-1/2} M_{ct} = B^{(c)}_{C_{ct}/c}, \qquad B^{(c)}_u = c^{-1/2} \widetilde{B}_{cu}.$$

Each $B^{(c)}$ has standard Brownian law. Uniform bracket convergence to the deterministic clock and the Brownian modulus of continuity imply the stated limit; independence between $B^{(c)}$ and the bracket is unnecessary. Adding the negligible drift gives the same limit for W. By exact scaling, however, each rescaled process already has the law of W on $[0,T]$. That law is therefore Brownian with variance v. Arbitrariness of T proves the assertion. □

The regular initial bracket is not automatic for a general self-similar continuous process. In the affine-SLE application below it follows from locality and absolutely continuous drift in two domain charts, using the local bracket-regularization identity.

3.2 The half-plane affine-SLE obstruction

Let A be an invertible orientation-preserving real-linear map and let η be chordal SLE_6 in $A\mathbb{H}$ from 0 to ∞. Put $\gamma = A^{-1}\eta \subset \overline{\mathbb{H}}$ and let W be its half-plane-capacity driving function. Its continuous Loewner generation and infinite capacity lifetime are supplied by the affine-admissibility result already proved in the research record. Its law has exact Brownian scaling (SM1): ordinary SLE is dilation covariant, A commutes with positive scalar dilations, and capacity scales by the square of physical length.

Proposition (proved). If this W satisfies (SM2)–(SM3), then A is a Euclidean similarity.

Proof. The scaling lemma makes W ordinary Brownian driving of variance v. Thus the trace is ordinary chordal SLE_v in $\mathbb{H}$. We identify v without a Hausdorff-dimension theorem.

For $\kappa \le 4$ the nonzero real boundary points are never swallowed by SLE_κ: their Loewner distances divided by $\sqrt{\kappa}$ are Bessel processes of dimension $1+4/\kappa \ge 2$ (the case $\kappa = 0$ is immediate). An affine image of SLE_6 swallows boundary points, so $v > 4$. For any $\kappa > 4$, the boundary-race calculation gives

$$\mathbb{P}_\kappa(\tau_y < \tau_x) = F_\kappa\!\left(\frac{-x}{y-x}\right), \qquad F_\kappa(r) = \frac{\displaystyle\int_0^r [u(1-u)]^{-4/\kappa}\,du}{\displaystyle\int_0^1 [u(1-u)]^{-4/\kappa}\,du}, \quad x < 0 < y. \tag{SM5}$$

This includes $\kappa \ge 8$. To check it, the ratio $R = -X/(Y-X)$ has generator

$$\frac{\kappa}{2}\frac{d^2}{dr^2} + \frac{2(1-2r)}{r(1-r)}\frac{d}{dr}$$

after $ds = dt/(Y-X)^2$. Its scale density is $[r(1-r)]^{-4/\kappa}$, integrable at both ends exactly when $\kappa > 4$. The first swallowing is finite almost surely by the Bessel description; simultaneous swallowing is impossible since $Y - X$ is positive and nondecreasing. Bounded optional stopping gives (SM5).

Rotate $A\mathbb{H}$ conformally back to $\mathbb{H}$. The resulting map has matrix

$$T = \begin{pmatrix} \alpha & \beta \\ 0 & \delta \end{pmatrix}, \qquad \alpha, \delta > 0.$$

It acts on the real boundary by $x \mapsto \alpha x$, so it preserves the ratio $-x/(y-x)$ and the topological swallowing order. The law of γ therefore has boundary-race function F_6, while its Brownian driving gives F_v. Equality for every $r \in (0,1)$ forces $v = 6$: compare the derivatives in (SM5), proportional respectively to $[r(1-r)]^{-4/v}$ and $[r(1-r)]^{-2/3}$.

Finally $T^{-1}\mathrm{SLE}_6(\mathbb{H}; 0, \infty)$ is ordinary SLE_6. Half-plane affine identifiability now implies T is a similarity. For completeness, the strictly increasing SLE_6 left-passage function

$$L(u) = \frac{\int_{-\infty}^{u}(1+s^2)^{-2/3}\,ds}{\int_{-\infty}^{\infty}(1+s^2)^{-2/3}\,ds}$$

and preservation of topological side give $L(u) = L((\alpha u + \beta)/\delta)$ for all real u. Thus $\alpha = \delta$ and $\beta = 0$. Rotation does not affect whether A is a similarity, proving the proposition. □

3.3 Transfer from bounded-domain regularity to the half-plane germ

The preceding proposition is already a complete unbounded-domain statement. The following additional argument shows why requiring the tame drift structure in *every bounded domain* is enough to trigger it. We spell out the local coordinate transfer, rather than silently assume semimartingality is preserved by a general physical affine map.

Bounded-domain obstruction theorem (proved). Consider the canonical fixed-affine SLE_6 family

$$P_{D,p,q} = (A^{-1})_* \,\mathrm{Law}(\mathrm{SLE}_6(AD; Ap, Aq))$$

on bounded simply connected marked domains. Suppose in the original Euclidean complex structure every domain-chart driver is a continuous semimartingale whose finite-variation part is $\int_0^t b_s\,ds$, with finite right-continuous drift density and deterministic initial value $B(f)$, consistently under the usual conditional kernel interpretation. Then A is a Euclidean similarity. No boundedness or continuity of the map $f \mapsto B(f)$ is assumed.

Proof. Choose a bounded half-disk $D \subset \mathbb{H}$ with 0 in the interior of its real diameter, and a distinct target on that diameter away from 0. Choose the boundary colors so they agree near 0 with the half-plane problem $(\mathbb{H}, 0, \infty)$. Let $\phi : D \to \mathbb{H}$ be its chordal chart sending these two marks to $0, \infty$. It extends analytically across 0, with $\phi'(0) > 0$.

The local bracket-regularization lemma applies to this bounded-domain driver: delete one further vertical slit in its ϕ chart, away from its marks, with nonzero $h''(0)$. Both domains have tame absolutely continuous drift by hypothesis. The signed-measure identity

$$h_t''(W_t)(dC_t - 6\,dt) = 2\big[h_t'(W_t)^2 \widehat{b}_{u(t)} - h_t'(W_t) b_t\big]\,dt$$

therefore gives a finite right-continuous initial bracket density in D, with deterministic initial value v_D.

Corrected source-neighborhood locality of ordinary SLE_6, preserved by the fixed map A^{-1}, couples this bounded-domain curve to the canonical affine-SLE_6 half-plane curve so that their physical initial segments are identical until leaving a sufficiently small half-neighborhood of 0. This uses ordinary SLE_6 locality including distant target changes; it is not a new hypothesis about an unknown nonconformal family. On that common initial trace, the coordinate change is the analytic map ϕ at the source.

A local analytic coordinate change and its inverse preserve the tame semimartingale structure just obtained. More explicitly, if t is physical half-plane capacity and u is the bounded-chart capacity, the evolving local map satisfies

$$W_D(u(t)) = h_t(W_H(t)), \qquad \dot{u}(t) = h_t'(W_H(t))^2.$$

Initially $h_0 = \phi$, so $h_0'(0) > 0$ and all required local derivatives are finite. Before the trace leaves the common neighborhood, localize so the first derivative stays bounded away from zero and the derivatives stay bounded. Both capacity clocks are adapted functions of the same initial trace; their inverses are stopping-time clocks. The inverse field h_t^{-1} has finite variation in time, as follows by differentiating the Loewner composition. Applying the finite-variation Itô–Wentzell formula to the semimartingale W_D, and changing the capacity clock, produces a continuous-semimartingale decomposition of W_H on a positive initial interval. Its drift is absolutely continuous and right-continuous, and its initial bracket density is the same scalar v_D. These conclusions use only a *conformal local chart transition*, not the nonconformal map A.

For an explicit initial check, writing $d = \phi'(0)$, $e = \phi''(0)$, the transferred rates obey

$$V_H = v_D, \qquad B_H = dB_D - \left(\frac{v_D}{2} - 3\right)\frac{e}{d}, \tag{SM6}$$

so both are finite and deterministic. Thus the canonical half-plane driver satisfies (SM2)–(SM3). Its exact scaling and the half-plane proposition force A to be a similarity. □

Only a bounded domain flat at its source and one of its source-preserving slit deletions are needed for the bracket part of this contradiction. The full all-bounded-domain assumption is a convenient sufficient formulation. The obstruction does not identify the precise failure for a particular wrong affine metric: semimartingality, absolute continuity or right-continuity of the drift, or deterministic initial drift may fail. It does show that at least one must fail somewhere. It supplies no argument that an unknown exploration family admits a correct affine normalization with tame drift.

4 An affine metric is identifiable from one chordal law

For a bounded Jordan domain D, distinct $p, q \in \partial D$, and an orientation-preserving invertible real-linear map A, put

$$\nu^A_{D,p,q} = (A^{-1})_* \operatorname{Law}\big(\mathrm{SLE}_6(AD; Ap, Aq)\big). \tag{1}$$

Translations in an affine normalizer are immaterial. The tensor $A^{\mathsf{T}}A$ is its Euclidean conformal metric, determined only up to a positive scalar.

4.1 A boundary-swallowing race

Lemma 4.1 (Strict boundary-race probability)**.** *Let η be chordal SLE_6 in $\mathbb{H}$ from 0 to ∞. For $x < 0 < y$, let τ_x, τ_y denote their Loewner swallowing times. Then*

$$\mathbb{P}(\tau_y < \tau_x) = F\left(\frac{-x}{y-x}\right), \qquad F(r) = \frac{\int_0^r [u(1-u)]^{-2/3}\,du}{\int_0^1 [u(1-u)]^{-2/3}\,du}. \tag{2}$$

In particular F is strictly increasing on $(0,1)$. The event in (2) is invariant under orientation-preserving homeomorphisms of marked Jordan domains, when formulated as the order of separation from the terminal prime end by the filled exploration hull.

Proof. Use $\partial_t g_t(z) = 2/(g_t(z) - W_t)$ with $W_t = \sqrt{6}B_t$. Until the first swallowing time put

$$X_t = g_t(x) - W_t < 0, \quad Y_t = g_t(y) - W_t > 0, \quad T_t = Y_t - X_t, \quad R_t = -X_t/T_t \in (0,1).$$

Then

$$dX_t = \frac{2}{X_t}\,dt - \sqrt{6}\,dB_t, \qquad dY_t = \frac{2}{Y_t}\,dt - \sqrt{6}\,dB_t,$$
$$dT_t = \frac{2T_t}{-X_tY_t}\,dt > 0, \qquad dR_t = \frac{2(1-2R_t)}{R_t(1-R_t)T_t^2}\,dt + \frac{\sqrt{6}}{T_t}\,dB_t.$$

With $ds = dt/T_t^2$, the generator of R is

$$\mathcal{L} = 3\frac{d^2}{dr^2} + \frac{2(1-2r)}{r(1-r)}\frac{d}{dr}.$$

The increasing solution of $\mathcal{L}F = 0$, $F(0) = 0$, $F(1) = 1$ has $F'(r) = C[r(1-r)]^{-2/3}$, giving (2) by bounded optional stopping. To justify the terminal values, $|X|/\sqrt{6}$ and $Y/\sqrt{6}$ are Bessel processes of dimension $1 + 4/6 = 5/3$ up to their first zero; thus the first swallowing time is finite almost surely. Alternatively, the accessible endpoints of the displayed diffusion give the same stopping argument. There is no simultaneous swallowing, since $T_t \geq y - x > 0$, whereas both $X_t, Y_t \to 0$ would force $T_t \to 0$. At the first swallowing, $R = 1$ exactly when y is swallowed first. ODE solutions and the continuous driver stay finite up to that first time.

Swallowing means that the marked boundary end has been cut off from the target by the explored filled hull (including the boundary swallowed at first contact). This definition uses only the domain, the oriented path, and its order of traversal. A homeomorphism carries filled target components to filled target components. Capacity parametrization therefore has no role in the comparison event. □

4.2 Boundary rigidity and affine uniqueness

Lemma 4.2 (A single SLE law identifies its conformal boundary map)**.** *Let $h : \overline{D} \to \overline{D'}$ be an orientation-preserving homeomorphism between closed bounded Jordan domains, with $p' = h(p)$, $q' = h(q)$. If $h_*\mathrm{SLE}_6(D; p, q) = \mathrm{SLE}_6(D'; p', q')$, then $h|_{\partial D}$ agrees with the boundary extension of a conformal map $f : D \to D'$.*

Proof. Choose conformal maps $\phi : D \to \mathbb{H}$ and $\phi' : D' \to \mathbb{H}$ sending their marked endpoints to $0, \infty$. Their Jordan boundary extensions give an increasing homeomorphism

$$H = \phi' \circ h \circ \phi^{-1} : \overline{\mathbb{R}} \to \overline{\mathbb{R}}$$

fixing $0, \infty$. For every $x < 0 < y$, equality of the corresponding boundary-race probabilities and strictness in Lemma 4.1 give

$$\frac{-H(x)}{H(y) - H(x)} = \frac{-x}{y - x}, \quad \text{hence} \quad \frac{H(x)}{H(y)} = \frac{x}{y}.$$

Fixing one $x < 0$ and one $y > 0$ shows $H(s) = cs$ for all real s, with $c > 0$. Consequently $f = (\phi')^{-1} \circ (c\phi)$ has exactly the required boundary values. Equality of interior maps is not asserted for an arbitrary h. □

Theorem 4.3 (Single-domain affine identifiability)**.** *For even one bounded Jordan D and one pair of distinct boundary points p, q, the equality $\nu^A_{D,p,q} = \nu^B_{D,p,q}$ holds if and only if BA^{-1} is a Euclidean similarity. Equivalently,*

$$B^{\mathsf{T}}B = cA^{\mathsf{T}}A \qquad (c > 0).$$

The statement also allows orientation-reversing normalizers by composing one map with a reflection.

Proof. The equality of laws means $M = BA^{-1}$ carries the SLE law in AD to the SLE law in BD, at the corresponding endpoints. By Lemma 4.2, a conformal map $f : AD \to BD$ has boundary values equal to M. Both real coordinate functions of the real-linear map M are harmonic. The real and imaginary parts of f are also harmonic, continuous on the closure, with the same boundary values. The maximum principle on the bounded domain gives $f = M$ throughout AD. Thus M is complex-linear and invertible, hence a similarity. The converse is ordinary SLE similarity covariance. Reflection covariance reduces the remaining orientation case to this one. □

References

[1] G. F. Lawler, *Schramm–Loewner Evolution*, Park City lecture notes (2007), Lectures 3–4. https://math.uchicago.edu/~lawler/utah.pdf.

[2] Y. Yuan, *Topological characterisations of Loewner traces*, Indiana Univ. Math. J. **71** (2022), 1027–1046. https://arxiv.org/abs/2003.05535.

[3] *Research record: locality, terminal information, and planar exploration laws*, `progress.tex` in this repository, including the original-locality investigation and the corrected-locality continuation (26 October 2026).