Local exploration laws, initial germs, and affine metrics

Autonomous mathematical investigation, with independent critical checks

27 October 2026

Abstract

Independent-color resampling and RSW source shields imply an initial ordered trace-germ zero-one law, with quantitative decorrelation between small prefixes and the full exploration curve. This conclusion does not assume affine-SLE marginals or a limiting domain Markov theorem. Exact coalescence also gives total-variation prefix–suffix decoupling and exponential mixing of annular source-crossing statistics, with a centered strong law that does not require dilation invariance. A universal tensor decoder at analytic regular endpoints, exact suffix coalescence, and source splicing then show that a coherent family of affine-SLE_6 mixtures must use one deterministic common affine metric. The affine-mixture representation itself remains an unproved hypothesis. Separately, a conditional Loewner-driving classification permits arbitrary continuous finite-variation drift with a uniform first-order initial tangent in probability; tightness in law along mirrored slit tests can replace deterministic tangents. Locality and compatible strong DMP then force Brownian driving of variance six. A wrong-affine-coordinate obstruction holds under the weakened deterministic tangent assumptions. A separate signed conditional-bias criterion supplies semimartingality and the required initial tangents without postulating them; its discrete version passes to continuous driver limits without filtration convergence. The requisite bias cancellation and compatible limiting DMP remain unproved for general percolation. These results leave the corrected three-axiom classification and critical percolation universality open.

1 Current locality and scope

Two colored marked domains that agree near their common source must have identical initial exploration laws until exit from that neighborhood, even if their remote targets differ. This is the user's corrected locality axiom at commit `0e47018`. We fix one color orientation throughout. The terminal-germ construction for the earlier fixed-target axiom is excluded; its original proofs are retained in `archive/original-locality-results.tex` and the detailed research record.

The first theorem below needs only a deletion consequence of corrected locality. It does not claim that locality supplies regular driving drift. For rough restart domains we state the prime-end neighborhood or admissible test-deletion requirement explicitly. All driving clocks use $\partial_t g = 2/(g - W)$ and hence $\mathrm{hcap}(K_t) = 2t$. The standard conformal-map comparison for Loewner chains is derived below; ordinary SLE facts used for affine examples are described in [1, 2]. Continuous Loewner generation and infinite lifetime of affine SLE images were proved in the companion research record [4].

2 A near-source drift-boundedness criterion

Status. This is a new conditional rigidity result. It does not assume conformal invariance, Brownian driving, or local-martingale driving. Instead it shows that a locally bounded *initial driving*

drift, together with an absolutely continuous drift part and locality, already forces zero drift and variance six. No regularity of quadratic variation is assumed. Strong DMP then identifies the whole driving law. The boundedness and characteristic hypotheses are not derived from RSW or from the three project axioms. A finite-dimensional bulk-jet closure is given as one explicit sufficient hypothesis.

Only a particular deletion consequence of the corrected source-neighborhood locality is needed. Arbitrary target changes and the full force of corrected locality are not used in this theorem. Thus the result is compatible with, but does not exhaust, the corrected research objective.

2.1 Chart-dependent initial characteristics and their covariance

Let $f : \mathbb{H} \to D$ be an inverse chordal conformal chart, with source $f(0) = p$ and target $f(\infty) = q$, both interpreted as prime ends. For the physical ensemble in D, let W^f be its driving process in this chart, with half-plane capacity normalization

$$\partial_t g_t(z) = \frac{2}{g_t(z) - W_t^f}, \qquad W_0^f = 0.$$

Assume that it is a continuous semimartingale in its natural trace filtration, with decomposition

$$W_t^f = M_t^f + \int_0^t b_s^f \, ds, \qquad C_t^f = \langle M^f \rangle_t. \tag{DR1}$$

The drift-density paths b^f are assumed finite and right-continuous, after local stopping if necessary. No absolute continuity or regularity of C^f is assumed. At the initial time suppose

$$b_0^f = B(f) \in \mathbb{R} \tag{DR2}$$

is deterministic. It may depend on the entire physical marked domain and its conformal chart. In particular no Markov property of the single real driver is assumed. A finite deterministic initial variance rate $V(f)$ will be *derived* below from locality and (DR1)–(DR2).

Let $A \subset \overline{\mathbb{H}}$ be a half-plane hull disjoint from 0, let $h : \mathbb{H} \setminus A \to \mathbb{H}$ be hydrodynamically normalized, and set

$$\bar{h}(z) = h(z) - h(0), \qquad f_A = f \circ \bar{h}^{-1}. \tag{DR3}$$

The source is still the same physical prime end, since $\bar{h}(0) = 0$; the target is also unchanged. Put $d = h'(0) > 0$, $e = h''(0)$. Source locality, together with the usual deletion-chart calculation, will give a finite deterministic initial variance rate $V(f)$ and the covariance

$$\boxed{V(f_A) = V(f), \qquad B(f_A) = \frac{B(f)}{d} + \left(\frac{V(f)}{2} - 3\right)\frac{e}{d^2}.} \tag{DR4}$$

These are identities for *initial* characteristics and do not identify the two complete driving laws.

Here are the details without assuming a quadratic-variation density. If h_t is the evolving deletion map, then $du = h_t'(W_t)^2 dt$, and Itô's formula gives

$$d\widehat{W}_{u(t)} = h_t'(W_t)\, dM_t + h_t'(W_t) b_t \, dt + \frac{1}{2} h_t''(W_t)\, dC_t - 3h_t''(W_t)\, dt. \tag{DR5}$$

The -3 comes from

$$\partial_t h_t(z) = \frac{2h_t'(W_t)^2}{h_t(z) - h_t(W_t)} - \frac{2h_t'(z)}{z - W_t}, \qquad (\partial_t h_t)(W_t) = -3h_t''(W_t).$$

The comparison of filtrations is justified by geometric invertibility: from a deleted-domain initial trace one reconstructs its original trace and capacity, so the inverse capacity clock is adapted and its inverse is a stopping-time clock. Locality therefore compares the same trace filtration, not an unrelated enlargement. Adaptedness of an arbitrary increasing clock would not suffice for this step.

Local bracket-regularization lemma (proved). Suppose $h''(0) \neq 0$. Then on a positive initial interval, C_t is absolutely continuous and has a finite right-continuous density. In particular its initial right derivative $V(f)$ exists and is deterministic.

Proof. The deleted-domain driver also has an absolutely continuous drift, $\int \widehat{b}_u \, du$, by (DR1). Uniqueness of the continuous semimartingale decomposition applied to (DR5), after the justified capacity time change, gives the signed-measure identity

$$h_t''(W_t)(dC_t - 6\,dt) = 2\big[h_t'(W_t)^2 \widehat{b}_{u(t)} - h_t'(W_t) b_t\big] dt. \tag{DR5a}$$

The second derivative starts nonzero and remains so on a positive initial interval. On that interval the bracket is absolutely continuous, with

$$v_t = 6 + 2\frac{h_t'(W_t)^2 \widehat{b}_{u(t)} - h_t'(W_t) b_t}{h_t''(W_t)}. \tag{DR5b}$$

The derivative paths of h_t along the continuous driving point are continuous, while the two drift-density paths are right-continuous and finite. Thus v is right-continuous and finite there. In particular

$$V(f) = v_0 = 6 + 2\frac{d^2 B(f_A) - dB(f)}{e} \tag{DR5c}$$

is deterministic. This is an intrinsic initial rate of C, so it does not depend on which nonzero-e deletion was used to derive it. Applying the same time change to bracket gives $V(f_A) = V(f)$. Rearranging (DR5c) gives the drift identity (DR4). □

There is always an available nonzero-e test deletion, for example one vertical slit at any fixed real point different from 0. Right-continuity makes the initial drift and variance rates intrinsic, rather than arbitrary versions of densities at a single time. The bracket density has only been proved locally at the start; its global regularity is not being presumed.

Source-neighborhood convention. In half-plane coordinates these slit deletions preserve a full neighborhood of the source end. Under f this is a prime-end neighborhood. Thus (DR4) is justified on arbitrary rough domains if the corrected locality assumption is formulated for such neighborhoods, or if these particular deletions are explicitly admissible. For regular analytic source boundaries it is ordinary Euclidean neighborhood locality. If locality is instead stated only using Euclidean balls at non-singleton source impressions, the all-rough-domain extension must remain an explicit additional hypothesis.

2.2 Two vanishing slits force both characteristics

For $\epsilon > 0$ take the reflected vertical slits

$$A_\epsilon^{\pm} = \{\, \pm\epsilon^2 + iy : 0 \le y \le 2\epsilon \,\}. \tag{DR6}$$

Their hydrodynamic maps are

$$h_\epsilon^\pm(z) = \pm\epsilon^2 + \sqrt{(z \mp \epsilon^2)^2 + 4\epsilon^2}, \tag{DR7}$$

with the square-root branch asymptotic to $z \mp \epsilon^2$ at infinity. Let $f_\epsilon^\pm = f \circ (h_\epsilon^\pm - h_\epsilon^\pm(0))^{-1}$. Both sequences are conformal charts of bounded simply connected domains and

$$f_\epsilon^\pm \longrightarrow f \quad \text{locally uniformly in } \mathbb{H}. \tag{DR8}$$

Indeed the slits have diameter 2ϵ and approach 0; directly from (DR7), their maps and inverse maps converge to the identity on compact subsets of $\mathbb{H}$, while $h_\epsilon^\pm(0) \to 0$. Crucially their boundary derivatives do *not* converge to the identity jet:

$$d_\epsilon = (h_\epsilon^\pm)'(0) = \frac{\epsilon}{\sqrt{\epsilon^2+4}} \longrightarrow 0, \qquad e_\epsilon^\pm = (h_\epsilon^\pm)''(0) = \mp\frac{4}{\epsilon(\epsilon^2+4)^{3/2}}. \tag{DR9}$$

In particular

$$\frac{e_\epsilon^\pm}{d_\epsilon^2} = \mp\frac{4}{\epsilon^3\sqrt{\epsilon^2+4}}. \tag{DR10}$$

Thus compact convergence of inverse charts permits very large second-jet changes precisely at the source boundary.

Initial rigidity theorem (proved). Fix f and assume (DR1)–(DR4) for f and the charts $f_\epsilon^\pm$. If

$$\sup_{0<\epsilon<\epsilon_0} \max\{|B(f_\epsilon^+)|, |B(f_\epsilon^-)|\} < \infty \tag{DR11}$$

for some $\epsilon_0 > 0$, then

$$B(f) = 0, \qquad V(f) = 6. \tag{DR12}$$

No continuity of B is required.

Proof. Write $b = B(f)$, $v = V(f)$ and $b_\epsilon^\pm = B(f_\epsilon^\pm)$. Equation (DR4) gives

$$b_\epsilon^\pm = \frac{b}{d_\epsilon} + \left(\frac{v}{2} - 3\right)\frac{e_\epsilon^\pm}{d_\epsilon^2}.$$

Since $e_\epsilon^- = -e_\epsilon^+$,

$$b_\epsilon^+ + b_\epsilon^- = \frac{2b}{d_\epsilon}, \qquad b_\epsilon^+ - b_\epsilon^- = (v-6)\frac{e_\epsilon^+}{d_\epsilon^2}. \tag{DR13}$$

The first right side is bounded only if $b = 0$, since $d_\epsilon \to 0$. The second is bounded only if $v = 6$, since its multiplier has magnitude asymptotic to $2\epsilon^{-3}$. This proves (DR12). □

Exact instability and weaker rate conditions. For real numbers U, C one has $\max\{|U + C|, |U - C|\} = |U| + |C|$. Hence the two drift covariances give the exact identity

$$\boxed{\max\{|b_\epsilon^+|, |b_\epsilon^-|\} = \frac{|b|\sqrt{\epsilon^2+4}}{\epsilon} + \frac{4|v/2-3|}{\epsilon^3\sqrt{\epsilon^2+4}}.} \tag{DR14}$$

There is no possible cancellation between the two reflected tests. In particular, if the displayed maximum is $o(\epsilon^{-3})$, then $v = 6$. If it is $o(\epsilon^{-1})$, then $b = 0$ and $v = 6$. Boundedness is more than sufficient. Conversely, if $v \neq 6$, its leading blow-up is $|v-6|\epsilon^{-3}$; if $v = 6$ and $b \neq 0$, its leading blow-up is $2|b|\epsilon^{-1}$. This is a concrete, quantitative obstruction to any alternative semimartingale driving law with tame initial drift under source perturbations.

2.3 Local boundedness in marked-chart topology suffices

Equip inverse marked charts with locally uniform convergence on $\mathbb{H}$, retaining the fixed preimages $0, \infty$ of source and target. If B is locally bounded at every f in this topology, then (DR8) implies (DR11). In particular continuity of B is sufficient, but unnecessarily strong. This is a hypothesis about the driving characteristic itself; weak convergence of curve laws alone does not imply it.

Conditional classification theorem (proved). Suppose a Borel family of continuous Loewner-generating exploration laws on all bounded simply connected marked prime-end domains has strong DMP and source-neighborhood locality (including the test deletions (DR6) in each chart). Suppose its drivers satisfy (DR1)–(DR2), with finite right-continuous drift-density paths and the usual restart compatibility of characteristic kernels, with no initial assumption on quadratic variation. If the initial drift map B is locally bounded in marked inverse- chart topology, then every driving process is

$$W_t^f = \sqrt{6}\, B_t,$$

and the family is chordal SLE_6 in these conformal coordinates.

Proof. The initial rigidity theorem gives $B(f) = 0$ and the derived $V(f) = 6$ for every marked inverse chart. At any deterministic rational time r, condition on the initial exploration and use strong DMP in its remaining marked domain and current conformal chart. The centered future driver is exactly the restarted driver; its initial drift is b_r^f, by right-continuity. The restarted law has initial drift zero, so $b_r^f = 0$ almost surely. Countably many rational times and right-continuity give $b_t^f = 0$ throughout each localized lifetime. Therefore all domain chart drivers are continuous local martingales.

For completeness the no-density local-martingale locality argument then identifies their brackets. With both drifts zero, (DR5a) becomes $h_t''(W_t)(dC_t - 6\,dt) = 0$. A nonzero-$e$ vertical deletion yields $C_t = 6t$ on a positive initial interval. Let

$$T = \inf\{t \geq 0 : C_t - 6t \neq 0\}.$$

This is an entrance stopping time of a continuous adapted process into an open set, in the usual right-continuous filtration. The initial zero interval gives $T > 0$. If $T < \infty$, continuity gives $C_T = 6T$ and the first-deviation definition gives nonzero values arbitrarily soon after T. Strong DMP restarted at T gives another positive zero interval, a contradiction. Thus $C_t = 6t$ throughout. Lévy's characterization gives Brownian driving, and Loewner generation identifies SLE_6. □

The Borel-kernel condition means that the family admits the standard strong conditional-kernel interpretation on random slit domains, and that the characteristics can be chosen consistently on that family. It is not a claim that deterministic-domain convergence of discrete explorations proves this property. Translation invariance is not needed in this conditional theorem. If its hypotheses hold only after one fixed physical affine normalization, the conclusion is the corresponding fixed-affine SLE_6 family; an affine change of physical coordinates is not presumed to preserve semimartingale driving in an arbitrary preselected complex structure.

2.4 A finite-dimensional closure that is completely classified

For example, fix finitely many interior points $z_j \in \mathbb{H}$ and finitely many nonnegative derivative orders m_j. Let

$$J(f) = \big(f^{(k)}(z_j) : 1 \leq j \leq N, 0 \leq k \leq m_j\big).$$

Suppose the initial drift has a representation

$$B(f) = \mathcal{B}(J(f)), \tag{DR15}$$

where $\mathcal{B}$ is locally bounded at every realized jet vector. The bracket has no assumed density or finite-dimensional representation; its initial rate is supplied by the local regularization lemma. Locally uniform convergence of conformal maps implies convergence of all these interior jets by Cauchy's formula, so (DR15) implies local boundedness of B in marked-chart topology. The conditional classification theorem therefore applies: within (DR1)–(DR2), every source-local strong-DMP family with such a locally bounded finite-dimensional *bulk-jet drift closure* is SLE_6.

This statement does not cover a closure depending on unbounded source boundary derivatives such as $f'(0)$ or a boundary pre-Schwarzian. Those quantities can change without control under (DR8), and can carry exactly the singular terms in (DR13). Nor does it show that true percolation limits have semimartingale driving in a fixed physical complex structure. The new rigorous reduction is that any non-SLE candidate with regular characteristic rates must violate the displayed near-source drift bound, in the explicit ϵ^{-1} or ϵ^{-3} manner above.

2.5 The criterion also detects a wrong affine coordinate system

Corollary (proved). Let A be an orientation-preserving real-linear map which is not a Euclidean similarity, and let the physical ensemble be the fixed-affine pullback

$$P_{D,p,q} = (A^{-1})_* \,\mathrm{Law}(\mathrm{SLE}_6(AD; Ap, Aq)).$$

Although this family satisfies corrected source-neighborhood locality and strong DMP, in the original Euclidean complex structure it cannot satisfy both the regular semimartingale characteristic assumptions (DR1)–(DR2) in all marked domains and the locally bounded initial-drift condition above.

Proof. If both properties held, the conditional classification theorem would make this ensemble ordinary SLE_6 in the original complex structure. The single-domain affine identifiability theorem would then make A a similarity, a contradiction. □

Thus the drift criterion selects a conformal structure. It is not a coordinate-free consequence of coarse crossing bounds: bounded affine changes preserve such bounds, whereas a wrong affine structure necessarily violates at least one of the criterion's additional analytic hypotheses. The corollary does not decide whether the failure is semimartingality, right-continuity of the drift density, deterministic initial drift, or boundedness of the initial drift; distinguishing these failures is a separate question.

2.6 What corrected locality says at a regular analytic source germ

Here is a separate necessary local structure statement that uses the full corrected locality, including changes of the distant target. It concerns initial characteristics only, not existence of a process from arbitrary coefficient choices.

Suppose the physical boundary is analytic near p and the source colors are fixed there. Let G denote this embedded colored source germ. For a chart f extending conformally across 0, define

$$a(f) = |f'(0)| > 0, \qquad P(f) = \mathrm{Re}\,\frac{f''(0)}{f'(0)}. \tag{DR16}$$

If another physical marked domain has the same source germ but any distant target, let k be its inverse chordal chart with $k(0) = p$. The local map $h = k^{-1} \circ f$ is analytic across 0, takes the real boundary interval to a real interval, and has $h(0) = 0$, $d = h'(0) > 0$, $e = h''(0) \in \mathbb{R}$. The two initial physical traces agree by corrected locality. The local Loewner coordinate comparison is exactly (DR4), with the same derivation from pole cancellation and Itô's formula; a globally defined deletion map is unnecessary. Consequently $V(f) = V(k)$ depends only on G.

Proposition (proved). Under the semimartingale assumptions above, corrected locality at analytic source germs is equivalent at the level of initial characteristic covariance to the existence of two source-germ quantities $V(G)$ and $\beta(G)$ such that

$$\boxed{V(f) = V(G), \qquad B(f) = a(f)\,\beta(G) + \left(3 - \frac{V(G)}{2}\right) P(f).} \tag{DR17}$$

Here "equivalent at the level of covariance" means precisely that every covariant initial-characteristic assignment has this form, and every assignment of this form obeys the local chart covariance identity. It does not mean that arbitrary V, β yield a domain Markov exploration family.

Proof. Since $k = f \circ h^{-1}$, the chain rule gives

$$a(k) = \frac{a(f)}{d}, \qquad P(k) = \frac{P(f)}{d} - \frac{e}{d^2}.$$

Write $c = V(G)/2 - 3$. Combining these transformations with $B(k) = B(f)/d + ce/d^2$ yields

$$\frac{B(k) + cP(k)}{a(k)} = \frac{B(f) + cP(f)}{a(f)}.$$

This common value is the source-germ invariant $\beta(G)$, proving (DR17). Conversely direct substitution gives the required covariance. □

Physical translation invariance makes $V(G)$ and $\beta(G)$ invariant under translations of the colored germ. If dilation covariance is separately assumed, then $V(\lambda G) = V(G)$ and $\beta(\lambda G) = \lambda^{-1}\beta(G)$. For a straight source-boundary germ, which is unchanged up to translation by dilation, this would force $\beta(G) = 0$; the dimensionless variance can still depend on the boundary orientation before an affine metric has been identified.

The boundary pre-Schwarzian term in (DR17) explains the quantitative anomaly in the vanishing-slit test. Even when the physical source germ is unchanged, $P(f_\epsilon)$ may grow as ϵ^{-3} while $f_\epsilon \to f$ on every interior compact set. Thus source locality by itself gives a connection-type transformation law, rather than vanishing drift. The locally bounded bulk closure theorem removes exactly this uncontrolled boundary dependence. The missing global step is to understand how DMP constrains these local assignments at rough, dynamically created source germs, where the analytic boundary jet in (DR16) no longer exists.

3 Rigidity from an averaged initial finite-variation characteristic

Scope. This conditional theorem weakens the preceding driving-regularity premise. The canonical finite-variation characteristic need not be absolutely continuous and need not possess a right-continuous density. We assume only a deterministic first-order tangent in probability, uniformly on shrinking initial intervals. No such property is derived from RSW or the three axioms.

Let the driver in inverse chart f have canonical decomposition

$$W^f = M^f + A^f, \qquad W_0^f = M_0^f = A_0^f = 0, \tag{FV1}$$

where M^f is a continuous local martingale and A^f is continuous, adapted, and locally of finite variation. Write $C^f = \langle M^f \rangle$. Assume for every deterministic state and inherited chordal chart that a finite deterministic number $B(f)$ satisfies

$$\frac{1}{t} \sup_{0 \le s \le t} |A_s^f - B(f)s| \xrightarrow[t \downarrow 0]{\mathbb{P}} 0. \tag{FV2}$$

An almost-sure derivative $A_t^f/t \to B(f)$ implies (FV2), but (FV2) is weaker. No bound on $\mathrm{TV}(A^f; [0,t])/t$ is assumed.

Lemma 3.1 (A small finite-variation displacement in an analytic field)**.** *Let $W = M + A$ as in (FV1), let $C = \langle M \rangle$, and suppose $\sup_{s \le t} |A_s| = O_{\mathbb{P}}(t)$ and $A_t/t \to B$ in probability. Let $H_t(x)$ be a random field analytic in x on a fixed neighborhood of zero for a positive initial time interval. Suppose it has absolutely continuous finite variation in t, and, after localization, its first three spatial derivatives and the first spatial derivative of $\partial_t H_t$ are bounded there. Assume $H_t'(x) \to d$ as $(t,x) \to (0,0)$. Then*

$$\frac{1}{t} \int_0^t H_s'(W_s)\, dA_s \longrightarrow dB \quad \textit{in probability.} \tag{FV3}$$

Proof. Localize so M, A, W stay in the spatial neighborhood. Apply the finite-variation Itô–Wentzell formula to $F_t(x) = H_t(x + A_t) - H_t(x)$ at $x = M_t$. It gives

$$\begin{aligned} \int_0^t H_s'(W_s)\, dA_s = {} & H_t(M_t + A_t) - H_t(M_t) \\ & - \int_0^t [H_s'(M_s + A_s) - H_s'(M_s)]\, dM_s \\ & - \frac{1}{2} \int_0^t [H_s''(M_s + A_s) - H_s''(M_s)]\, dC_s \\ & - \int_0^t [(\partial_s H_s)(M_s + A_s) - (\partial_s H_s)(M_s)]\, ds. \end{aligned}$$

The first term divided by t converges to dB by the mean-value formula. The second term is a continuous local martingale N_t with bracket at most $K^2 \sup_{s \le t} |A_s|^2 C_t$. Thus $N_t/t \to 0$ in probability: stop when $|A_s| > Lt$ or $C_s > \eta$, apply the martingale maximal inequality, then let $t \downarrow 0$, $\eta \downarrow 0$, and $L \uparrow \infty$. Here $C_t \to 0$ almost surely and $\sup |A_s|/t$ is tight. The third term divided by t is bounded by $K \sup |A_s| C_t/t = o_{\mathbb{P}}(1)$; the fourth is bounded by $K \sup |A_s| = o_{\mathbb{P}}(1)$. No total-variation estimate was used. □

Proposition 3.2 (Initial covariance without a drift density)**.** *Suppose source locality allows one test deletion with hydrodynamic map h, $d = h'(0) > 0$, $e = h''(0) \neq 0$, and both original and deleted drivers obey (FV1)–(FV2), with their canonical characteristics in the same trace filtration. Then*

$$C_t^f/t \longrightarrow V(f) \in [0, \infty) \quad \textit{in probability,}$$

where $V(f)$ is deterministic, and

$$V(f_A) = V(f), \qquad B(f_A) = \frac{B(f)}{d} + \left(\frac{V(f)}{2} - 3 \right) \frac{e}{d^2}. \tag{FV4}$$

Proof. Use the same evolving deletion maps and stopping-time capacity clocks as in (DR5), now without assuming a drift density. Put $p_t = h_t'(W_t)$, $q_t = h_t''(W_t)$ and $u(t) = \int_0^t p_s^2 ds$. Uniqueness of the canonical semimartingale decomposition gives

$$\widehat{A}_{u(t)} = \int_0^t p_s \, dA_s + \frac{1}{2} \int_0^t q_s \, dC_s - 3 \int_0^t q_s \, ds. \tag{FV5}$$

The deletion field satisfies the analytic-field lemma after localization; its Loewner field derivative and spatial derivatives are bounded near the uncollided source. Therefore the first integral divided by t tends to $dB(f)$, without assuming C_t/t is tight. Also $u(t)/t \to d^2$. Condition (FV2) for the deleted state implies $\widehat{A}_{u(t)}/t \to d^2 B(f_A)$: on the event $u(t) \le Kt$, bound the error by $t^{-1} \sup_{s \le Kt} |\widehat{A}_s - B(f_A)s|$, and choose fixed $K > d^2$.

Since $q_t \to e \neq 0$, (FV5) first implies C_t/t is tight: with probability tending to one q_s has fixed sign and magnitude at least $|e|/2$ on $[0, t]$, so $|\int q_s dC_s| \ge |e| C_t/2$. Consequently

$$\frac{1}{t} \int_0^t q_s \, dC_s - e \frac{C_t}{t} \longrightarrow 0 \quad \text{in probability.}$$

Taking (FV5) divided by t now gives

$$V(f) = 6 + \frac{2\{d^2 B(f_A) - dB(f)\}}{e}.$$

This is intrinsic, nonnegative, and deterministic. The bracket relation $\widehat{C}_{u(t)} = \int_0^t p_s^2 dC_s$ gives $V(f_A) = V(f)$. For this last step the tangents of $C, \widehat{C}$ may be evaluated at asymptotically deterministic clocks by monotonic squeezing; convergence in probability of an arbitrary signed process would not justify that step. □

Lemma 3.3 (A finite-variation process with zero right tangent at every stopping time)**.** *Let $A_0 = 0$ be a continuous adapted locally finite-variation process. If for every bounded stopping time τ before lifetime*

$$\frac{A_{\tau+h} - A_\tau}{h} \longrightarrow 0 \quad \textit{in probability as } h \downarrow 0, \tag{FV6}$$

then A is indistinguishably zero.

Proof. Work on a compact localized interval. Pathwise put $V_t = \mathrm{TV}(A; [0, t])$, and use its generalized inverse $\theta(u) = \inf\{t : V_t \ge u\}$ for $0 < u < V_T$. The usual arclength parametrization of a continuous finite-variation path is $G(u) = A_{\theta(u)}$: it is 1-Lipschitz, satisfies $A_t = G(V_t)$, and has $|G'(u)| = 1$ for Lebesgue almost every u. Indeed the variation of G on every interval is its interval length. The monotone function θ has a finite derivative $\theta'(u) \in [0, \infty)$ for almost every u. Exclude also the countably many levels corresponding to nontrivial flat intervals of V.

At a level where these properties hold, let $s = \theta(u)$ and $v(h) = V_{s+h} - u$. Then $v(h) > 0$ and $v(h) \to 0$ as $h \downarrow 0$. The generalized-inverse bounds $\theta((u + v(h))-) \le s + h \le \theta((u + v(h))+)$ and differentiability of θ at u give $h/v(h) \to \theta'(u)$. Combining this with differentiability of G gives

$$\lim_{h \downarrow 0} \frac{A_{s+h} - A_s}{h} = \begin{cases} G'(u)/\theta'(u), & \theta'(u) > 0, \\ \mathrm{sgn}(G'(u)) \, \infty, & \theta'(u) = 0. \end{cases} \tag{FV7}$$

Thus the extended right derivative is nonzero at almost every variation level, including levels whose inverse belongs to a singular part of dV.

For deterministic $u > 0$ use the stopping times $\tau_u = \inf\{t : V_t \geq u\}$, truncated at a deterministic endpoint T. Fubini and the preceding pathwise statement show that for Lebesgue almost every deterministic u, on $\{\tau_u < T\}$ the quotient in (FV6) has a nonzero or infinite pathwise limit almost surely. But (FV6) gives convergence in probability to zero at this stopping time, which forbids such an event of positive probability. Thus $\mathbb{P}(\tau_u < T) = 0$ for Lebesgue almost every u, and $\mathbb{E}[V_T] = \int_0^\infty \mathbb{P}(\tau_u < T)du = 0$ after localizing V_T to be integrable. Hence $V_T = 0$ almost surely. Localization and a countable exhaustion of the lifetime prove the claim. □

Theorem 3.4 (Conditional classification from a finite-variation initial tangent)**.** *Suppose a Borel family on all relevant marked slit states has strong DMP and source locality, including the same mirrored test deletions (DR6). Suppose canonical decompositions (FV1), compatible with conditional restart in the inherited chordal charts, obey (FV2). If $B(f)$ is locally bounded in inverse-chart topology, or just satisfies the paired slit bound (DR11) at every state, the family is chordal SLE$_6$ in these coordinates.*

Proof. Proposition 3.2 gives exactly (DR4), hence the reflected-slit sum and difference argument gives $B(f) = 0$, $V(f) = 6$ at every deterministic state.

At a stopping time τ, the centered future driver is the driver of the remaining physical state in the inherited conformal chart. Its capacity parameter is exactly $s = t - \tau$ in the original half-plane coordinate: recentering by W_τ introduces no additional scale clock. Strong DMP and conditional-characteristic compatibility identify its canonical finite-variation part with $A_{\tau+s} - A_\tau$ under the regular conditional law. Applying (FV2) with $B = 0$, and integrating these conditional probabilities over the random state, gives (FV6). Lemma 3.3 therefore makes $A = 0$.

The same conditional restart argument gives $(C_{\tau+h} - C_\tau)/h \to 6$ in probability at every stopping time. Apply Lemma 3.3 to the continuous finite-variation process $C_t - 6t$: it is identically zero. Thus $W = M$ and $\langle M\rangle_t = 6t$ globally. Lévy's characterization and continuous Loewner generation identify SLE$_6$. □

Strict weakening and remaining limitation. Even an absolutely continuous drift with density $b_s = B + \sin(1/s)$, $s > 0$, has no right limit at zero, whereas $\int_0^t b_s ds = Bt + O(t^2)$ and therefore satisfies (FV2). The theorem also allows an a priori singular finite-variation drift; its exclusion is a conclusion, not an assumption. A claim that an actual exploration family has these initial characteristic tangents remains unproved. The theorem still depends on choosing one correct physical conformal structure and on tame behavior under the source slit tests.

3.1 The wrong-coordinate obstruction also only needs finite-variation tangents

Scaling lemma with probability tangents. The exact-scaling lemma (SM1)–(SM4) remains true if its drift assumption is replaced by continuous finite variation and (FV2), and its bracket assumption by $C_t/t \to v$ in probability for deterministic v. Indeed (FV2) makes $\sup_{s\leq T} |c^{-1/2}A_{cs}| = O_{\mathbb{P}}(\sqrt{c}) \to 0$. Since C is increasing, a finite deterministic mesh and monotonic squeezing show

$$\sup_{s\leq T} |C_{cs}/c - vs| \longrightarrow 0 \quad \text{in probability.}$$

The continuous-martingale functional central limit argument in the earlier scaling lemma then applies unchanged. Exact scaling makes the whole law Brownian with variance v.

Local transfer of the tangent. In Proposition 3.2, the analytic-field argument actually gives the uniform version

$$\frac{1}{t}\sup_{s\le t}\left|\int_0^s h_r'(W_r)\,dA_r - dB(f)s\right| \longrightarrow 0 \quad \text{in probability.}$$

Use the martingale maximal inequality in its proof, and the same uniform initial bound for A. Monotonic squeezing similarly makes $C_{ts}/t \to V(f)s$ uniformly on compact s intervals in probability. Equation (FV5) therefore transfers (FV2) through a local analytic chart change, once the initial bracket tangent has been obtained; it also transfers its inverse, since the first chart derivative is positive. No total-variation rate bound is introduced by this transfer.

Corollary 3.5 (Wrong-affine obstruction under finite-variation tangents)**.** *Consider the canonical fixed-affine SLE_6 family with matrix A on all bounded marked domains. If every Euclidean chart driver admits (FV1)–(FV2), then A is a Euclidean similarity. No local boundedness of $B(f)$ is needed.*

Proof. Choose the bounded flat-source domain and one test deletion used in the bounded-domain obstruction theorem. Proposition 3.2 gives its deterministic bracket tangent in probability. Couple its initial physical trace by known affine-SLE locality to the canonical half-plane affine-SLE trace. The local analytic chart transfer just proved gives a finite-variation decomposition and uniform deterministic drift tangent for the half-plane driver, and the same deterministic bracket tangent. Its exact Brownian scaling and the extended scaling lemma make its law ordinary Brownian driving. The boundary-swallowing-race and affine identification arguments in the half-plane obstruction theorem then give variance six and force A to be a similarity. □

3.2 A tightness-in-law variant without deterministic initial tangents

Proposition (proved). There is a useful alternative formulation of the preceding theorem. Allow (FV2) to have a finite random tangent β^f rather than a deterministic $B(f)$. Assume that for every fixed chart f the family of initial tangents in its two shrinking slit domains is uniformly tight:

$$\lim_{K\to\infty}\sup_{0<\epsilon<\epsilon_0}\max_{\sigma\in\{+,-\}}\mathbb{P}\{|\beta^{f_\epsilon^\sigma}|>K\}=0. \tag{FV8}$$

Then the same classification follows, with all other family and restart hypotheses unchanged.

The analytic-field lemma and initial covariance proof apply verbatim to finite random tangents, producing an intrinsic finite random bracket tangent V^f. For each fixed ϵ, source locality and the invertible local coordinate changes put the original and both deleted initial traces on one probability space. Only their initial germs are needed, so no simultaneous extension of this coupling past their exit times is required. On that space the covariances give

$$\beta^{f_\epsilon^+}+\beta^{f_\epsilon^-}=2\beta^f/d_\epsilon, \qquad \beta^{f_\epsilon^+}-\beta^{f_\epsilon^-}=(V^f-6)e_\epsilon^+/d_\epsilon^2.$$

Uniform tightness of the two marginals makes their sum and difference uniformly tight under any coupling. Since $d_\epsilon\to 0$ and $|e_\epsilon^+|/d_\epsilon^2\to\infty$, these identities force $\beta^f=0$ and $V^f=6$ almost surely. Formally, for $a>0$,

$$\mathbb{P}\{|\beta^f|>a\}\le\sum_{\sigma=\pm}\mathbb{P}\{|\beta^{f_\epsilon^\sigma}|>a/d_\epsilon\}\longrightarrow 0,$$

and the identical argument with threshold $a|e_\epsilon^+|/(2d_\epsilon^2)$ gives $V^f=6$. The stopping-time lemma then kills the finite-variation drift and identifies the bracket as before. Local boundedness of a deterministic drift tangent is one special case of (FV8); no deterministic initial-tangent premise is otherwise necessary for this version.

4 Rigidity from conditional first-moment bounds

Scope. The next criterion replaces the a priori semimartingale and initial-tangent premises by conditional first-moment bounds. It does not derive those bounds from RSW. They must hold in one fixed physical conformal structure, with the stated control along the specified shrinking slit tests. No quadratic-moment bound or initial drift derivative is assumed.

Lemma 4.1 (Conditional bias produces a bounded drift density)**.** *Let X be a bounded continuous adapted real process, with $X_0 = 0$, in a usual filtration. Suppose a finite deterministic K satisfies*

$$|\mathbb{E}[X_t - X_s \mid \mathcal{F}_s]| \le K(t-s) \quad \text{almost surely,} \qquad 0 \le s \le t. \tag{BI1}$$

Then $X = M + \int_0^{\cdot} b_r dr$, where M is a continuous local martingale and b is predictable with $|b| \le K$ almost everywhere.

Proof. On a fixed finite time interval, $X_t + Kt$ and $-X_t + Kt$ are class-D submartingales. Their Doob–Meyer decompositions have increasing predictable parts A^+ and A^-. Adding the decompositions and using uniqueness gives $A^+ + A^- = 2Kt$. Thus A^+ and A^- are absolutely continuous, with densities in $[0, 2K]$. The decomposition of X has drift density $b = dA^+/dt - K \in [-K, K]$. Since X and its drift are continuous, its martingale part is continuous. Uniqueness patches the decompositions over finite time intervals. □

Proposition 4.2 (Controlled rate bounds force the tangents to exist)**.** *Use the mirrored shrinking slits $A_\epsilon^\pm$ in (DR6), with initial derivatives d_ϵ and $e_\epsilon^\pm$ in (DR9). Assume the driver in a state f, and the drivers in all these deleted states, have canonical continuous-semimartingale decompositions with absolutely continuous drift. In each state the drift density is bounded on some strictly positive initial stopping interval. The interval may depend on the state and sample. Let $K_\epsilon < \infty$ be a deterministic bound common to the two deleted states. If $d_\epsilon K_\epsilon \to 0$ along a sequence tending to zero, then in the original state*

$$\frac{1}{t}\sup_{s \le t} |A_s| \longrightarrow 0, \qquad \frac{C_t}{t} \longrightarrow 6 \quad \text{almost surely as } t \downarrow 0. \tag{BI2}$$

Here A is canonical drift and C is the martingale bracket. No right continuity or initial tangent of the drift densities is presumed.

Proof. By locality, compare the original and each deleted driver using the same trace filtration and the stopping-time capacity clocks. With $p_t^\sigma = (h_t^\sigma)'(W_t)$ and $q_t^\sigma = (h_t^\sigma)''(W_t)$, uniqueness of canonical characteristics gives

$$\frac{1}{2} q_t^\sigma (dC_t - 6dt) = \big[(p_t^\sigma)^2 \widehat{b}^\sigma_{u_\sigma(t)} - p_t^\sigma b_t\big] dt. \tag{BI3}$$

One fixed nonzero-q_0 test makes C absolutely continuous on a positive initial interval; write $dC_t = v_t dt$ there. For each fixed ϵ, both mirrored identities hold almost everywhere on a positive initial interval, as a two-by-two system in b_t and $(v_t - 6)/2$. Its coefficients converge as $t \downarrow 0$ to

$$\begin{pmatrix} d_\epsilon & e_\epsilon^+ \\ d_\epsilon & -e_\epsilon^+ \end{pmatrix},$$

whose determinant is nonzero. The right sides are $(p_t^\sigma)^2 \widehat{b}^\sigma_{u_\sigma(t)}$, with each transformed density bounded by K_ϵ. Continuity of the coefficients and matrix inversion therefore give the pathwise bounds

$$\operatorname*{ess\,limsup}_{t \downarrow 0} |b_t| \le d_\epsilon K_\epsilon, \qquad \operatorname*{ess\,limsup}_{t \downarrow 0} |v_t - 6| \le \frac{2 d_\epsilon^2 K_\epsilon}{|e_\epsilon^+|}. \tag{BI4}$$

These assertions concern the same original densities. Intersect the probability-one sets for a countable sequence $\epsilon \downarrow 0$. By assumption $d_\epsilon K_\epsilon \to 0$; also $d_\epsilon/|e_\epsilon^+| \to 0$, so both right sides tend to zero. Integrating the resulting essential right limits proves (BI2). The positive comparison intervals may shrink with ϵ: the essential limsup is taken at fixed ϵ first. □

Theorem 4.3 (SLE from controlled conditional driving bias)**.** *Let a Borel exploration family have continuous Loewner drivers on all $t \geq 0$, source locality including the test deletions, and strong DMP. Require locality to identify the completed natural trace filtrations via the invertible capacity clocks. Under conditional restart in an inherited chart, require the canonical martingale, drift, and bracket to coincide with those of the restarted kernel. These are the same explicit filtration and characteristic compatibility premises as in the earlier conditional driving theorems. Suppose that, in every deterministic state f, there is an increasing sequence of stopping times $\rho_{f,m}$ increasing almost surely to infinity, with $\rho_{f,1} > 0$ almost surely, such that each $X_t^{f,m} = W^f_{t\wedge\rho_{f,m}}$ is bounded and satisfies (BI1), with a finite deterministic bound $K(f,m)$. Require*

$$\lim_{n\to\infty} \epsilon_n \max_{\sigma=\pm} K(f^\sigma_{\epsilon_n}, 1) = 0 \tag{BI5}$$

at every state, for some sequence $\epsilon_n \downarrow 0$. In particular uniformly bounded constants suffice; growth $o(\epsilon_n^{-1})$ is allowed. Then the family is chordal SLE_6 in these coordinates.

Proof. Lemma 4.1, applied to the localizing sequence, gives global continuous-semimartingale decompositions with locally bounded absolutely continuous drift. Uniqueness patches the decompositions on overlapping stopped intervals. Proposition 4.2 then gives the zero drift tangent and bracket tangent six in every state.

Use the conditional-kernel compatibility of canonical decompositions in the inherited natural trace filtration, as in the earlier conditional characteristic theorems. At every bounded stopping time τ, conditional DMP and (BI2) give

$$\frac{A_{\tau+h} - A_\tau}{h} \longrightarrow 0, \qquad \frac{C_{\tau+h} - C_\tau}{h} \longrightarrow 6 \quad \text{in probability.}$$

The zero-stopping-tangent lemma (Lemma 3.3) applied to A and $C - 6t$ gives $A = 0$ and $C_t = 6t$. Lévy's characterization identifies the driver as $\sqrt{6}B$, hence the trace as SLE_6. □

Rates in the slit test. Since $d_\epsilon \sim \epsilon/2$ and $|e_\epsilon^+| \sim 1/(2\epsilon)$, the two bounds in (BI4) are of orders ϵK_ϵ and $\epsilon^3 K_\epsilon$. The weaker condition $\epsilon^3 K_\epsilon \to 0$ alone forces the initial bracket tangent to be six. The full classification above also needs the stronger drift condition in (BI5).

What this reduction gains and what it leaves open. Conditional bias bounds concern first moments of observable driver increments, rather than unknown canonical decompositions. They produce semimartingality, bounded drift densities, and the missing initial tangents without assuming quadratic variation estimates. They are nevertheless substantial cancellation estimates: a bound on $\mathbb{E}|W_{t+h} - W_t|$ of Brownian order $\sqrt{h}$, or tightness from RSW, does not imply (BI1). The rate control in (BI5) is coordinate sensitive and is not justified by convergence of curve laws in the shrinking slit domains. Thus this is a conditional route toward the desired regularity, not a proof of percolation universality.

4.1 An observable discrete criterion that passes to curve limits

There is no need to identify the discrete or limiting canonical characteristics in order to test the preceding criterion. Signed conditional first moments on a discrete exploration clock suffice. The following elementary transfer also avoids convergence of filtrations. It does not establish the separate strong-DMP premise of the rigidity theorem.

Proposition 4.4 (Discrete conditional bias passes to the natural limit filtration)**.** *Fix $T > 0$. For each n, let $0 = t_0^n < t_1^n < \cdots < t_{N_n}^n = T$ be a deterministic partition with maximum spacing tending to zero. Let X^n be its continuous linear interpolation of grid values, all bounded by one deterministic constant L. Suppose the grid values are adapted to filtrations $(\mathcal{G}_j^n)_j$ and*

$$\left|\mathbb{E}\left[X^n_{t^n_{j+1}} - X^n_{t^n_j} \mid \mathcal{G}_j^n\right]\right| \le (K + \delta_n)(t_{j+1}^n - t_j^n), \qquad \delta_n \downarrow 0. \tag{DB1}$$

Suppose X^n converges in distribution in $C([0,T])$ to X. Then in the usual augmentation of the natural filtration of X,

$$|\mathbb{E}[X_t - X_s \mid \mathcal{F}_s]| \le K(t-s), \qquad 0 \le s \le t \le T. \tag{DB2}$$

Consequently X has a continuous local-martingale part and an absolutely continuous canonical drift with density bounded by K.

Proof. Summing the conditional inequalities and using the tower property shows that the grid processes $X^n_{t^n_j} \pm (K + \delta_n)t_j^n$ are respectively a submartingale and a supermartingale. In particular, for grid points $s_n \le t_n$ and every bounded nonnegative $\mathcal{G}^n_{j(s_n)}$-measurable H_n,

$$\left|\mathbb{E}[(X^n_{t_n} - X^n_{s_n})H_n]\right| \le (K + \delta_n)(t_n - s_n)\mathbb{E}H_n. \tag{DB3}$$

Choose grid points $s_n \to s$ and $t_n \to t$, with $s_n \le t_n$, and set $H_n = H(X^n_{\cdot\wedge s_n})$, where H is any bounded nonnegative continuous function on $C([0,T])$. The stopped interpolated path is measurable at the grid time s_n. Convergence in $C([0,T])$ and uniform continuity of each limiting path give joint convergence of the evaluations and stopped paths. The deterministic bound L permits passage of their expectations to the limit in (DB3).

Bounded continuous tests determine finite measures on the stopped-path space. Thus the resulting two inequalities extend to all bounded nonnegative tests measurable in the raw natural filtration at s. This proves (DB2) there. Equivalently $X_t + Kt$ and $-X_t + Kt$ are continuous class-D submartingales. These remain submartingales under the usual augmentation: for $u \downarrow s$, apply the inequality at $u < t$, then use reverse-martingale convergence and continuity of X to condition on $\bigcap_{u>s} \mathcal{F}_u^{\mathrm{raw}}$. Completion changes only null sets. Lemma 4.1 now supplies the asserted canonical decomposition. □

Exploration-time variant. An equivalent usable sufficient hypothesis allows a random discrete clock. Let the successive capacity times t_j^n and grid values X_j^n be adapted to the discrete revealed-color filtration, with uniform deterministic bounds on the drivers and clock horizons. Assume the times are nondecreasing, interpolation is well defined, and $J_n(s) = \inf\{j : t_j^n \ge s\} < \infty$ almost surely for each $s < T$. Suppose their largest spacing tends to zero in probability and

$$\left|\mathbb{E}[X_{j+1}^n - X_j^n \mid \mathcal{G}_j^n]\right| \le (K + \delta_n)\mathbb{E}[t_{j+1}^n - t_j^n \mid \mathcal{G}_j^n]. \tag{DB4}$$

After continuous interpolation, assume the same uniform-path convergence. Then (DB2) again follows. Indeed $X_j^n \pm (K + \delta_n)t_j^n$ are bounded discrete sub/supermartingales. Use optional sampling

at $J_n(s) = \inf\{j : t_j^n \geq s\}$ and $J_n(t)$, with a test of the path stopped at $t_{J_n(s)}^n$. The indices are stopping times because $\{J_n(s) \leq j\} = \{t_j^n \geq s\}$. The clock overshoots tend to zero in probability; tightness in the continuous-path topology makes the stopped paths and sampled values converge to their deterministic-time counterparts. Boundedness then passes the inequalities to the limit as above. At T, use continuity from times below T.

Fixed-interval block estimates also suffice. The per-reveal condition (DB4) is sufficient, not necessary for this transfer argument. It is enough that for each fixed $s < t$ and every bounded nonnegative continuous stopped-path test H, the tested inequalities (DB3), using approximations of s, t, hold up to an error $o(1)\|H\|_\infty$. The same weak-convergence proof then gives (DB2). Thus one may instead seek signed cancellation over mesoscopic blocks, allowing microscopic biases to cancel before a fixed capacity interval. No per-reveal estimate is asserted for lattice percolation.

Thus a possible lattice-level research target is the signed cancellation estimate (DB4), after suitable bounded stopping, in an unknown but fixed affine normalization. The constants must satisfy the shrinking-slit control in Theorem 4.3, and the limit must separately have its stated strong-DMP and conditional-characteristic compatibility. Neither a Brownian-size absolute increment estimate nor RSW alone is being substituted for this signed conditional estimate.

5 A wrong affine metric cannot have tame driving drift in every bounded domain

Status. The conclusion here is stronger than the earlier wrong-coordinate consequence of drift boundedness. For a fixed non-similarity affine deformation of SLE_6, regular semimartingale driving with absolutely continuous right-continuous drift cannot hold in every bounded domain in the wrong Euclidean complex structure. No local boundedness of the initial drift functional is needed. The proof uses exact scale invariance only for the *known fixed-affine SLE_6 half-plane law*; it does not assume that arbitrary critical subsequential limits are scale invariant. All characteristic regularity assumptions remain explicit.

5.1 Exact Brownian scaling and a regular initial bracket

Scaling lemma (proved). Let W be a continuous real process on $[0, \infty)$ with $W_0 = 0$, whose law has exact Brownian scaling:

$$(c^{-1/2} W_{ct})_{t \geq 0} \overset{\text{law}}{=} (W_t)_{t \geq 0} \qquad (c > 0). \tag{SM1}$$

Suppose up to a strictly positive stopping time it is a continuous semimartingale

$$W_t = M_t + \int_0^t b_s \, ds, \tag{SM2}$$

where $M_0 = 0$ is a continuous local martingale and b is finite and right-continuous at 0. Suppose its bracket has a finite deterministic initial right derivative

$$\lim_{t \downarrow 0} \frac{\langle M \rangle_t}{t} = v \in [0, \infty) \quad \text{almost surely.} \tag{SM3}$$

Then the entire law of W is that of $\sqrt{v} B$. This includes the identically zero process if $v = 0$.

Proof. Fix $T < \infty$ and stop the local decomposition at a positive initial localizing time. As $c \downarrow 0$, with probability tending to one the whole interval $[0, cT]$ lies before that time. Right-continuity of b gives

$$\sup_{t \le T} \left| c^{-1/2} \int_0^{ct} b_s \, ds \right| \le \sqrt{c}\, T \sup_{s \le cT} |b_s| \longrightarrow 0$$

almost surely on the localized event. Writing $C_t = \langle M \rangle_t$, (SM3) implies

$$\sup_{t \le T} |C_{ct}/c - vt| \longrightarrow 0. \tag{SM4}$$

Indeed $|C_{ct}/c - vt| \le t \sup_{0<s\le cT} |C_s/s - v|$. The continuous-martingale functional central limit theorem now gives $c^{-1/2} M_{c\cdot} \Rightarrow \sqrt{v} B$ on $C[0,T]$. One may see this directly from Dambis–Dubins–Schwarz: extend its Brownian motion if necessary and write

$$c^{-1/2} M_{ct} = B^{(c)}_{C_{ct}/c}, \qquad B^{(c)}_u = c^{-1/2} \widetilde{B}_{cu}.$$

Each $B^{(c)}$ has standard Brownian law. Uniform bracket convergence to the deterministic clock and the Brownian modulus of continuity imply the stated limit; independence between $B^{(c)}$ and the bracket is unnecessary. Adding the negligible drift gives the same limit for W. By exact scaling, however, each rescaled process already has the law of W on $[0,T]$. That law is therefore Brownian with variance v. Arbitrariness of T proves the assertion. □

The regular initial bracket is not automatic for a general self-similar continuous process. In the affine-SLE application below it follows from locality and absolutely continuous drift in two domain charts, using the local bracket-regularization identity.

5.2 The half-plane affine-SLE obstruction

Let A be an invertible orientation-preserving real-linear map and let η be chordal SLE_6 in $A\mathbb{H}$ from 0 to ∞. Put $\gamma = A^{-1}\eta \subset \overline{\mathbb{H}}$ and let W be its half-plane-capacity driving function. Its continuous Loewner generation and infinite capacity lifetime are supplied by the affine-admissibility result already proved in the research record. Its law has exact Brownian scaling (SM1): ordinary SLE is dilation covariant, A commutes with positive scalar dilations, and capacity scales by the square of physical length.

Proposition (proved). If this W satisfies (SM2)–(SM3), then A is a Euclidean similarity.

Proof. The scaling lemma makes W ordinary Brownian driving of variance v. Thus the trace is ordinary chordal SLE_v in $\mathbb{H}$. We identify v without a Hausdorff-dimension theorem.

For $\kappa \le 4$ the nonzero real boundary points are never swallowed by SLE_κ: their Loewner distances divided by $\sqrt{\kappa}$ are Bessel processes of dimension $1 + 4/\kappa \ge 2$ (the case $\kappa = 0$ is immediate). An affine image of SLE_6 swallows boundary points, so $v > 4$. For any $\kappa > 4$, the boundary-race calculation gives

$$\mathbb{P}_\kappa(\tau_y < \tau_x) = F_\kappa\!\left(\frac{-x}{y-x}\right), \qquad F_\kappa(r) = \frac{\displaystyle\int_0^r [u(1-u)]^{-4/\kappa}\, du}{\displaystyle\int_0^1 [u(1-u)]^{-4/\kappa}\, du}, \quad x < 0 < y. \tag{SM5}$$

This includes $\kappa \geq 8$. To check it, the ratio $R = -X/(Y - X)$ has generator

$$\frac{\kappa}{2}\frac{d^2}{dr^2} + \frac{2(1-2r)}{r(1-r)}\frac{d}{dr}$$

after $ds = dt/(Y - X)^2$. Its scale density is $[r(1-r)]^{-4/\kappa}$, integrable at both ends exactly when $\kappa > 4$. The first swallowing is finite almost surely by the Bessel description; simultaneous swallowing is impossible since $Y - X$ is positive and nondecreasing. Bounded optional stopping gives (SM5).

Rotate $A\mathbb{H}$ conformally back to $\mathbb{H}$. The resulting map has matrix

$$T = \begin{pmatrix} \alpha & \beta \\ 0 & \delta \end{pmatrix}, \qquad \alpha, \delta > 0.$$

It acts on the real boundary by $x \mapsto \alpha x$, so it preserves the ratio $-x/(y - x)$ and the topological swallowing order. The law of γ therefore has boundary-race function F_6, while its Brownian driving gives F_v. Equality for every $r \in (0, 1)$ forces $v = 6$: compare the derivatives in (SM5), proportional respectively to $[r(1-r)]^{-4/v}$ and $[r(1-r)]^{-2/3}$.

Finally $T^{-1}\mathrm{SLE}_6(\mathbb{H}; 0, \infty)$ is ordinary SLE_6. Half-plane affine identifiability now implies T is a similarity. For completeness, the strictly increasing SLE_6 left-passage function

$$L(u) = \frac{\int_{-\infty}^{u}(1+s^2)^{-2/3}\,ds}{\int_{-\infty}^{\infty}(1+s^2)^{-2/3}\,ds}$$

and preservation of topological side give $L(u) = L((\alpha u + \beta)/\delta)$ for all real u. Thus $\alpha = \delta$ and $\beta = 0$. Rotation does not affect whether A is a similarity, proving the proposition. □

5.3 Transfer from bounded-domain regularity to the half-plane germ

The preceding proposition is already a complete unbounded-domain statement. The following additional argument shows why requiring the tame drift structure in *every bounded domain* is enough to trigger it. We spell out the local coordinate transfer, rather than silently assume semimartingality is preserved by a general physical affine map.

Bounded-domain obstruction theorem (proved). Consider the canonical fixed-affine SLE_6 family

$$P_{D,p,q} = (A^{-1})_* \,\mathrm{Law}(\mathrm{SLE}_6(AD; Ap, Aq))$$

on bounded simply connected marked domains. Suppose in the original Euclidean complex structure every domain-chart driver is a continuous semimartingale whose finite-variation part is $\int_0^t b_s\,ds$, with finite right-continuous drift density and deterministic initial value $B(f)$, consistently under the usual conditional kernel interpretation. Then A is a Euclidean similarity. No boundedness or continuity of the map $f \mapsto B(f)$ is assumed.

Proof. Choose a bounded half-disk $D \subset \mathbb{H}$ with 0 in the interior of its real diameter, and a distinct target on that diameter away from 0. Choose the boundary colors so they agree near 0 with the half-plane problem $(\mathbb{H}, 0, \infty)$. Let $\phi : D \to \mathbb{H}$ be its chordal chart sending these two marks to $0, \infty$. It extends analytically across 0, with $\phi'(0) > 0$.

The local bracket-regularization lemma applies to this bounded-domain driver: delete one further vertical slit in its ϕ chart, away from its marks, with nonzero $h''(0)$. Both domains have tame absolutely continuous drift by hypothesis. The signed-measure identity

$$h_t''(W_t)(dC_t - 6\,dt) = 2\big[h_t'(W_t)^2\widehat{b}_{u(t)} - h_t'(W_t)b_t\big]dt$$

therefore gives a finite right-continuous initial bracket density in D, with deterministic initial value v_D.

Corrected source-neighborhood locality of ordinary SLE_6, preserved by the fixed map A^{-1}, couples this bounded-domain curve to the canonical affine-SLE_6 half-plane curve so that their physical initial segments are identical until leaving a sufficiently small half-neighborhood of 0. This uses ordinary SLE_6 locality including distant target changes; it is not a new hypothesis about an unknown nonconformal family. On that common initial trace, the coordinate change is the analytic map ϕ at the source.

A local analytic coordinate change and its inverse preserve the tame semimartingale structure just obtained. More explicitly, if t is physical half-plane capacity and u is the bounded-chart capacity, the evolving local map satisfies

$$W_D(u(t)) = h_t(W_H(t)), \qquad \dot{u}(t) = h_t'(W_H(t))^2.$$

Initially $h_0 = \phi$, so $h_0'(0) > 0$ and all required local derivatives are finite. Before the trace leaves the common neighborhood, localize so the first derivative stays bounded away from zero and the derivatives stay bounded. Both capacity clocks are adapted functions of the same initial trace; their inverses are stopping-time clocks. The inverse field h_t^{-1} has finite variation in time, as follows by differentiating the Loewner composition. Applying the finite-variation Itô–Wentzell formula to the semimartingale W_D, and changing the capacity clock, produces a continuous-semimartingale decomposition of W_H on a positive initial interval. Its drift is absolutely continuous and right-continuous, and its initial bracket density is the same scalar v_D. These conclusions use only a *conformal local chart transition*, not the nonconformal map A.

For an explicit initial check, writing $d = \phi'(0)$, $e = \phi''(0)$, the transferred rates obey

$$V_H = v_D, \qquad B_H = dB_D - \left(\frac{v_D}{2} - 3\right)\frac{e}{d}, \tag{SM6}$$

so both are finite and deterministic. Thus the canonical half-plane driver satisfies (SM2)–(SM3). Its exact scaling and the half-plane proposition force A to be a similarity. □

Only a bounded domain flat at its source and one of its source-preserving slit deletions are needed for the bracket part of this contradiction. The full all-bounded-domain assumption is a convenient sufficient formulation. The obstruction does not identify the precise failure for a particular wrong affine metric: semimartingality, absolute continuity or right-continuity of the drift, or deterministic initial drift may fail. It does show that at least one must fail somewhere. It supplies no argument that an unknown exploration family admits a correct affine normalization with tame drift.

6 An affine metric is identifiable from one chordal law

For a bounded Jordan domain D, distinct $p, q \in \partial D$, and an orientation-preserving invertible real-linear map A, put

$$\nu^A_{D,p,q} = (A^{-1})_* \,\mathrm{Law}\big(\mathrm{SLE}_6(AD; Ap, Aq)\big). \tag{1}$$

Translations in an affine normalizer are immaterial. The tensor $A^{\mathsf{T}}A$ is its Euclidean conformal metric, determined only up to a positive scalar.

6.1 A boundary-swallowing race

Lemma 6.1 (Strict boundary-race probability)**.** *Let η be chordal SLE_6 in $\mathbb{H}$ from 0 to ∞. For $x < 0 < y$, let τ_x, τ_y denote their Loewner swallowing times. Then*

$$\mathbb{P}(\tau_y < \tau_x) = F\left(\frac{-x}{y-x}\right), \qquad F(r) = \frac{\int_0^r [u(1-u)]^{-2/3}\,du}{\int_0^1 [u(1-u)]^{-2/3}\,du}. \tag{2}$$

In particular F is strictly increasing on $(0,1)$. The event in (2) is invariant under orientation-preserving homeomorphisms of marked Jordan domains, when formulated as the order of separation from the terminal prime end by the filled exploration hull.

Proof. Use $\partial_t g_t(z) = 2/(g_t(z) - W_t)$ with $W_t = \sqrt{6}B_t$. Until the first swallowing time put

$$X_t = g_t(x) - W_t < 0, \quad Y_t = g_t(y) - W_t > 0, \quad T_t = Y_t - X_t, \quad R_t = -X_t/T_t \in (0,1).$$

Then

$$dX_t = \frac{2}{X_t}\,dt - \sqrt{6}\,dB_t, \qquad dY_t = \frac{2}{Y_t}\,dt - \sqrt{6}\,dB_t,$$
$$dT_t = \frac{2T_t}{-X_tY_t}\,dt > 0, \qquad dR_t = \frac{2(1-2R_t)}{R_t(1-R_t)T_t^2}\,dt + \frac{\sqrt{6}}{T_t}\,dB_t.$$

With $ds = dt/T_t^2$, the generator of R is

$$\mathcal{L} = 3\frac{d^2}{dr^2} + \frac{2(1-2r)}{r(1-r)}\frac{d}{dr}.$$

The increasing solution of $\mathcal{L}F = 0$, $F(0) = 0$, $F(1) = 1$ has $F'(r) = C[r(1-r)]^{-2/3}$, giving (2) by bounded optional stopping. To justify the terminal values, $|X|/\sqrt{6}$ and $Y/\sqrt{6}$ are Bessel processes of dimension $1 + 4/6 = 5/3$ up to their first zero; thus the first swallowing time is finite almost surely. Alternatively, the accessible endpoints of the displayed diffusion give the same stopping argument. There is no simultaneous swallowing, since $T_t \geq y - x > 0$, whereas both $X_t, Y_t \to 0$ would force $T_t \to 0$. At the first swallowing, $R = 1$ exactly when y is swallowed first. ODE solutions and the continuous driver stay finite up to that first time.

Swallowing means that the marked boundary end has been cut off from the target by the explored filled hull (including the boundary swallowed at first contact). This definition uses only the domain, the oriented path, and its order of traversal. A homeomorphism carries filled target components to filled target components. Capacity parametrization therefore has no role in the comparison event. □

6.2 Boundary rigidity and affine uniqueness

Lemma 6.2 (A single SLE law identifies its conformal boundary map)**.** *Let $h : \overline{D} \to \overline{D'}$ be an orientation-preserving homeomorphism between closed bounded Jordan domains, with $p' = h(p)$, $q' = h(q)$. If $h_*\mathrm{SLE}_6(D; p, q) = \mathrm{SLE}_6(D'; p', q')$, then $h|_{\partial D}$ agrees with the boundary extension of a conformal map $f : D \to D'$.*

Proof. Choose conformal maps $\phi : D \to \mathbb{H}$ and $\phi' : D' \to \mathbb{H}$ sending their marked endpoints to $0, \infty$. Their Jordan boundary extensions give an increasing homeomorphism

$$H = \phi' \circ h \circ \phi^{-1} : \overline{\mathbb{R}} \to \overline{\mathbb{R}}$$

fixing $0, \infty$. For every $x < 0 < y$, equality of the corresponding boundary-race probabilities and strictness in Lemma 6.1 give

$$\frac{-H(x)}{H(y) - H(x)} = \frac{-x}{y - x}, \quad \text{hence} \quad \frac{H(x)}{H(y)} = \frac{x}{y}.$$

Fixing one $x < 0$ and one $y > 0$ shows $H(s) = cs$ for all real s, with $c > 0$. Consequently $f = (\phi')^{-1} \circ (c\phi)$ has exactly the required boundary values. Equality of interior maps is not asserted for an arbitrary h. □

Theorem 6.3 (Single-domain affine identifiability)**.** *For even one bounded Jordan D and one pair of distinct boundary points p, q, the equality $\nu^A_{D,p,q} = \nu^B_{D,p,q}$ holds if and only if BA^{-1} is a Euclidean similarity. Equivalently,*

$$B^\mathsf{T} B = cA^\mathsf{T} A \qquad (c > 0).$$

The statement also allows orientation-reversing normalizers by composing one map with a reflection.

Proof. The equality of laws means $M = BA^{-1}$ carries the SLE law in AD to the SLE law in BD, at the corresponding endpoints. By Lemma 6.2, a conformal map $f : AD \to BD$ has boundary values equal to M. Both real coordinate functions of the real-linear map M are harmonic. The real and imaginary parts of f are also harmonic, continuous on the closure, with the same boundary values. The maximum principle on the bounded domain gives $f = M$ throughout AD. Thus M is complex-linear and invertible, hence a similarity. The converse is ordinary SLE similarity covariance. Reflection covariance reduces the remaining orientation case to this one. □

7 An affine tensor is readable from a regular endpoint germ

Status. The argument below extends the initial half-plane trace-germ decoder to analytic regular source germs. By ordinary SLE_6 reversibility, it gives a decoder from the ordered terminal trace germ at any analytic regular target, even when the other endpoint and the rest of the domain are rough. It does not decode the unlabelled accessible frontier at a random tip. All almost-sure assertions are made separately under each fixed affine tensor; no uncountable intersection of full-probability events is used.

Let $\mathcal{X}$ be the usual Polish space of oriented unparameterized continuous finite curves, with the uniform Fréchet metric. Curves are identified under increasing reparameterization, including deletion of pauses. For a continuous path Y starting at 0 and exiting every disk, write $\tau_a(Y)$ for its first hit of $\{|z| = a\}$ and $\mathrm{Stop}_a Y = Y[0, \tau_a(Y)]$. The operations used here are Borel in standard continuous-curve coding; first hits can equivalently be defined by rational-time approximation. For a rough opposite prime end the full curve need not land continuously there. Only finite prefixes near the decoded regular endpoint are coded in $\mathcal{X}$; the unbounded half-plane curves exit every finite disk. This avoids any assumption of a singleton impression at the other mark.

Lemma 7.1 (Unit-radius stopping is almost surely continuous)**.** *Let Y have a law invariant under positive scalar spatial dilation and suppose it is continuous, starts at 0, and exits every disk. Then almost surely*

$$\tau_{1-\delta}(Y) \uparrow \tau_1(Y), \qquad \tau_{1+\delta}(Y) \downarrow \tau_1(Y) \quad (\delta \downarrow 0).$$

Consequently the diameter of $Y[\tau_{1-\delta}, \tau_{1+\delta}]$ tends to zero.

Proof. The first assertion on the left holds pathwise by continuity. The nondecreasing function $a \mapsto \tau_a(Y)$ has only countably many right jumps. Thus the set of radii with $\tau_{a+}(Y) > \tau_a(Y)$ has Lebesgue measure zero for every sample. Spatial dilation invariance implies that the probability of such a jump is the same at every fixed radius. Fubini's theorem on $[1, 2]$ makes this probability zero. Continuity of Y on a compact time interval containing these exit times then gives the diameter assertion. The argument is independent of the chosen continuous strictly increasing curve clock. □

For an orientation-preserving real linear map L carrying $\mathbb{H}$ onto itself, let μ_L be the law of $L\eta$, where η is chordal SLE_6 in $\mathbb{H}$ from 0 to ∞. Scalar multiples of L give the same law. Normalize

$$L = \begin{pmatrix} 1 & u \\ 0 & v \end{pmatrix}, \qquad (u, v) \in \Theta := \mathbb{R} \times (0, \infty). \tag{ED1}$$

The normalized metric represented by this law is $[(L^{-1})^{\mathsf{T}} L^{-1}]$.

Choose a countable bounded Lipschitz determining class $(f_m)_{m \geq 1}$ on $\mathcal{X}$, with values in $[0, 1]$ and Lipschitz constant at most one, allowing constant rescaling of an arbitrary countable bounded Lipschitz determining class. Such a class can be built from truncated distances to a countable dense set and finite rational combinations. Set

$$v_m(L) = \mathbb{E}_{\mu_L} f_m(\mathrm{Stop}_1 Y). \tag{ED2}$$

The vector $v(L) = (v_m(L))_m$ is Borel in L. Indeed, linear image and first-exit stopping are jointly Borel, so integration against the fixed ordinary SLE law preserves Borel measurability.

Brownian spatial scaling is mixing: under the Lamperti transform $Z_s = e^{-s/2} B_{e^s}$ it becomes a shift of a stationary Gaussian process with covariance $e^{-|s-u|/2}$. Its SLE trace factor, and each fixed linear image, are therefore dilation mixing. This gives the ergodicity used below without a separate path-space independence assumption.

The map $L \mapsto v(L)$ is injective. If two unit-radius stopped laws agreed, scaling invariance would identify their stopped laws at every radius. Those restrictions determine the entire ordered unbounded curve, contradicting the half-plane affine identifiability theorem unless the normalized matrices agree. Therefore the Lusin–Souslin theorem makes $v(\Theta)$ Borel and its inverse Borel.

Fix $\rho > 1$ and $r_n = \rho^{-n}$. For a curve γ starting at p define the local observable

$$X_n(\gamma, p) = r_n^{-1}(\gamma[0, \tau_{r_n}(\gamma - p)] - p). \tag{ED3}$$

If a finite curve fails to exit one of the large initial disks, assign a cemetery value to that term. It exits all sufficiently small disks, and finitely many cemetery values do not affect the limits below. Define

$$\ell_m(\gamma, p) = \lim_{N \to \infty} \frac{1}{N} \sum_{n=1}^{N} f_m(X_n(\gamma, p)), \tag{ED4}$$

using a cemetery vector if any limit fails to exist. Each limit is a function of every sufficiently small ordered initial trace prefix: deleting finitely many summands changes no Cesàro limit. In particular this is a Borel ordered-trace-germ statistic.

Lemma 7.2 (The decoder survives a quadratic local error)**.** *Let η be ordinary half-plane SLE_6 and let Φ be a local C^1 embedding at 0, defined on a half-neighborhood of 0, with*

$$\Phi(0) = 0, \qquad \Phi(z) = Lz + O(|z|^2), \tag{ED5}$$

where L is invertible. Assume after a fixed rotation that L maps $\mathbb{H}$ onto itself. Let γ agree with $\Phi(\eta)$ up to the first exit from this local chart. Then almost surely

$$\ell_m(\gamma, 0) = v_m(L) \quad \textit{for every } m. \tag{ED6}$$

Scalar normalization of L is understood in (ED2).

Proof. Put $Y_n = r_n^{-1} L\eta$. Spatial dilation of the trace includes the induced capacity reparameterization. Brownian scaling is mixing; its trace factor and every fixed affine image are therefore dilation ergodic, as proved in the initial half-plane decoder theorem. Hence

$$\frac{1}{N}\sum_{n=1}^{N} f_m(\mathrm{Stop}_1 Y_n) \longrightarrow v_m(L) \quad \text{almost surely.} \tag{ED7}$$

One must control the error in this average; pointwise stopping continuity at all the scaled random samples is not asserted.

For $0 < \delta < 1/4$ define

$$h_\delta(Y) = \min\{2, \operatorname{diam} Y[\tau_{1-\delta}(Y), \tau_{1+\delta}(Y)]\}\,. \tag{ED8}$$

The linearization error on a fixed bounded scaled chart is at most Cr_n, uniformly over the sample path until its linear image exits the disk of radius $1+\delta$. This uniformity is deterministic: the inverse linear map bounds the relevant preimage disk, and (ED5) bounds the error there. For all sufficiently large n, the first time that $r_n^{-1}\Phi(\eta)$ reaches radius one lies between the corresponding Y_n exit times at radii $1-\delta$ and $1+\delta$. Thus the two stopped curves consist of matched portions with pointwise error at most Cr_n, together with an unmatched terminal portion whose diameter is bounded by the diameter in (ED8). Increasing reparameterizations, with pauses allowed, give

$$|f_m(X_n(\gamma, 0)) - f_m(\mathrm{Stop}_1 Y_n)| \le Cr_n + h_\delta(Y_n). \tag{ED9}$$

The bounded range of f_m justifies capping the diameter by two.

Take a countable sequence $\delta_k \downarrow 0$. Birkhoff's theorem applied to every h_{δ_k} gives

$$\limsup_N \frac{1}{N}\sum_{n=1}^{N} |f_m(X_n(\gamma, 0)) - f_m(\mathrm{Stop}_1 Y_n)| \le \mathbb{E}_{\mu_L} h_{\delta_k}(Y) \tag{ED10}$$

simultaneously for all m, k. Lemma 7.1 and dominated convergence make the right side tend to zero as $k \to \infty$. Combine this with (ED7). Only the initial segment of η enters the argument, since all sufficiently small physical exits occur inside the fixed local chart. This proves (ED6). □

Theorem 7.3 (A universal analytic-endpoint tensor decoder)**.** *There is a Borel rule, depending only on the ordered initial trace germ, the endpoint, and its oriented tangent half-plane, that recovers the projective affine tensor from a single affine-SLE$_6$ sample in every bounded simply connected marked domain with analytic regular source. The rest of its boundary and its target may be rough prime-end data. There is also a terminal version of the rule for every domain with analytic regular target; the source may then be rough. The rule is the same for two domains agreeing near that regular endpoint.*

Proof. Let the law be $(A^{-1})_* \mathrm{SLE}_6(AD, Ap, Aq)$ and assume p is an analytic regular source. A conformal map $f : \mathbb{H} \to AD$ with marks $0, \infty$ extends conformally across 0 by analytic boundary

reflection, with $f(0) = Ap$ and $f'(0) \neq 0$. In a physical tangent frame Q carrying the interior tangent half-plane to $\mathbb{H}$, the local map is

$$\Phi(z) = Q(A^{-1}f(z) - p) = Lz + O(|z|^2), \qquad L = QA^{-1}M_{f'(0)}, \tag{ED11}$$

where M_c denotes multiplication by the nonzero complex number c. The linear map L carries $\mathbb{H}$ onto itself and has positive diagonal entries in the boundary frame. Apply Lemma 7.2. The Borel inverse of (ED2) recovers the scalar-normalized L, hence the projective tensor

$$[(L^{-1})^{\mathsf{T}}L^{-1}] = [QA^{\mathsf{T}}AQ^{\mathsf{T}}]. \tag{ED12}$$

Conjugating back by the known orthogonal frame Q recovers $[A^{\mathsf{T}}A]$. The derivative of the conformal map contributes only a positive scalar to this tensor. This rule uses the endpoint germ and tangent half-plane, not the rest of the conformal chart.

Ordinary SLE_6 reversibility [3] says the reversed curve has the same affine-SLE form with marks swapped. Apply the source decoder to that reversed curve at the regular target. Reversibility on prime-end domains follows by conformal transport from the half-plane theorem; only the regular target germ needs a Euclidean analytic boundary extension. □

Corollary 7.4 (Terminal singularity and identifiable mixtures)**.** *For one fixed marked domain with analytic regular target, affine-SLE_6 laws with distinct projective affine tensors are mutually singular on the ordered terminal trace-germ sigma-field. More strongly, if π is any Borel probability on normalized affine tensors and $P_\pi = \int P_T\,\pi(dT)$, then the terminal decoder has law π under P_π. Consequently $\pi \mapsto P_\pi$ is injective, and no nontrivial affine mixture can equal one pure affine-SLE law.*

Proof. The same Borel decoder equals T almost surely under P_T. Integrating that equality proves the distribution statement. Distinct pure laws concentrate on disjoint decoder fibers. □

Additional initial-germ obstruction for random mixtures. At an analytic regular source, the initial decoder likewise returns the latent tensor almost surely. Therefore a nontrivial mixture of affine metrics has a nontrivial initial trace-germ sigma-field. At every positive exploration time the metric posterior has already collapsed to a point, even though its prior was nontrivial. This rules out such mixtures under an additional initial-germ zero-one premise. It does not rule out a deterministic source-germ metric assignment; that mechanism still asks whether the current unlabelled frontier recovers the tensor.

Proposition 7.5 (DMP forces restarted mixing measures to be pure)**.** *Suppose an all-state family has the mixed-affine representation $P_S = \int \nu_S^T\,\pi_S(dT)$, with Borel mixing kernel π_S. Start it in a domain with analytic regular source and target. If it has strong DMP, then at each admissible stopping time $\tau > 0$ before the target is reached,*

$$\pi_{S_\tau} = \delta_{D_s(\gamma)} \qquad \textit{almost surely,} \tag{ED13}$$

where D_s is the initial tensor decoder. In particular a rule that uses a nondegenerate prior at every restarted state cannot satisfy DMP.

Proof. Sample the latent tensor T with law π_S and then sample the fixed-T affine SLE. The source decoder gives $D_s(\gamma) = T$ almost surely, and this variable is measurable from every positive prefix, including the prefix up to τ. Conditional on this prefix, ordinary fixed-T SLE DMP therefore gives the pure continuation law $\nu_{S_\tau}^{D_s(\gamma)}$. The family's DMP gives the mixed law with mixing measure π_{S_τ}. Since τ precedes reaching the target, the surviving target end has the same analytic regular germ: the compact prefix avoids a sufficiently small target neighborhood. Corollary 7.4 applied to this current state makes the two mixing measures equal, proving (ED13). □

8 A discrete source shield and stability of two exploration starts

Status and model. The deterministic shielding statement and the ensuing probabilistic estimate are proved here for independent site percolation on a triangular-lattice triangulation of a Jordan disk. Boundary vertices are wired black or white, with exactly two color changes, the marked source and target. The exploration is the nonbranching face-dual black–white interface between those changes. The same argument applies to a cellular model with the corresponding nonbranching primal/dual geometry and the stated white RSW bounds. It is an actual common-color coupling argument. It is *not* inferred for an arbitrary random-curve family from abstract locality or G2 alone.

Work at mesh δ. Two triangulated domains, their source positions, and their boundary colors may differ inside $B(p, \epsilon)$, but agree outside it. Their common target lies outside $B(p, R_0)$. Outside the source patch and inside this latter ball, assume the common geometry is either a lattice half-plane with straight boundary, or a canonical thin straight slit with its two boundary sides. The lattice approximations are chosen with free first rows adjacent to the wired boundary rows; slit width is $O(\delta)$. The black and white wired sides are the same for the two configurations outside the patch. Couple all free colors outside the patch identically. Those shared colors have the independent percolation law. The colors inside the patch may be coupled arbitrarily, with their correct marginal laws.

8.1 The deterministic one-edge crosscut

In a triangular face, either zero or two of its primal edges have endpoints of different colors. Hence the dual interface graph has degree zero or two at every interior dual vertex. In the Dobrushin disk it has a unique open simple path between the source and target; other components are loops.

Lemma 8.1 (A one-heterochromatic-edge shield)**.** *Let a simple primal crosscut c of a triangulated disk connect a white wired boundary vertex to a black wired boundary vertex. Suppose every edge of c is white–white except its final edge e, which joins a white* interior *vertex to the black boundary vertex. Suppose c separates the entire source patch from the target. Then the exploration must use the dual edge e^*, with the direction determined by its black and white sides. After using e^* it cannot reenter the source side of the crosscut.*

Consequently, if the same such crosscut and its colors occur in both configurations above, their explorations have identical ordered suffixes from e^* to the common target.

Proof. The source–target dual path must cross the primal crosscut. Primal and dual vertices are distinct, and dual edges cross primal edges only at their matched transverse intersections. No interface edge can cross a white–white primal edge. Thus the crossing must be e^*, and matching colors give the same direction in both configurations.

The dual interface path is simple, so it cannot traverse e^* twice. All other primal edges of c remain monochromatic, so they cannot be crossed later either. The path therefore stays on the target side after using e^*. In particular it never returns to the altered source patch. Outside the patch the two triangulations and all colors coincide. At each subsequent dual vertex the outgoing edge is uniquely determined by the local colors and the incoming edge. Starting at the same directed edge e^*, the two paths consequently have the same entire suffix. □

Using boundary *vertices* in this formulation avoids an ambiguity of cell-exterior pictures: e is an interior primal edge incident to a boundary vertex, so its dual edge is an ordinary interior edge. Neither a crossing through a dual/primal common vertex nor an unexplained passage around the end of a physical boundary edge is being used.

8.2 Uniform probability of a white arch

Assume a uniform white RSW lower bound for fixed-aspect rectangles of free vertices. No connection to a prescribed individual vertex is assumed. We give the geometry for a horizontal half-plane source, translated to $p = 0$, with black boundary on the negative real side and white boundary on the positive side. Let $r \gg \epsilon + \delta$.

Use a white vertical crossing in each of

$$[-3r, -2r] \times [0, 3r], \qquad [2r, 3r] \times [0, 3r],$$

and a white horizontal crossing of

$$[-3r, 3r] \times [r, 2r].$$

At the bottom of a stem, start the rectangle in the first free vertex row, not in the fixed wired row. Rounding by $O(\delta)$ changes the aspect ratios by a uniformly bounded amount. The crossings in the overlapping squares intersect by planarity. Thus their union contains a white path from a free vertex adjacent to the black boundary to a free vertex adjacent to the white boundary. Add a white–white edge at the white endpoint and choose an edge e to a black boundary vertex at the other endpoint. Trim the white path at its last white-wired-boundary contact before reaching the left endpoint, so that all its subsequent white vertices are interior; then erase loops. This gives a proper simple primal crosscut with just the one heterochromatic edge e.

All these vertices lie, allowing lattice rounding, in $A(0, r/2, 6r)$. The crosscut and the boundary arc through the source enclose the source patch. Its source-side component is inside $B(0, 6r)$: the crosscut and that local boundary arc form a Jordan polygon contained there, so their bounded interior is also contained there. The common half-annulus between the patch and crosscut shows that the entire altered domain portion of the source patch is on that side.

Each rectangle event is increasing in white colors of independent free vertices. RSW and FKG therefore give a constant $\vartheta > 0$, independent of r, δ and the contents of the source patch, for this white-arch event. Importantly, the left endpoint is *some* first-row white vertex adjacent to the black boundary, selected after the crossing occurs. Requiring connection to one predetermined microscopic vertex would not have this uniform lower bound.

Straight slit geometry. The same construction works for a finite-width lattice slit along the negative real axis, with black upper lip and white lower lip, by using five crossings around a square annulus: an upper-left vertical stem, a top horizontal crossing, a right vertical crossing, a bottom horizontal crossing, and a lower-left vertical stem. The stems occupy $[-3r, -2r] \times [0, 3r]$ and $[-3r, -2r] \times [-3r, 0]$ on their respective sides of the slit. The other three rectangles may be

$$[-3r, 3r] \times [2r, 3r], \quad [2r, 3r] \times [-3r, 3r], \quad [-3r, 3r] \times [-3r, -2r].$$

The first free upper row is adjacent to the black wired lip; the first free lower row is adjacent to the white wired lip. The resulting crosscut goes around the right side of the slit and separates its tip patch from the distant target. The boundary arc through the tip, including the two separate lattice lips, stays inside $B(0, 6r)$. The finite-width triangulated domain is Jordan; one need not identify its two physical boundary rows. RSW and FKG again give a uniform $\vartheta > 0$ after adjusting constants.

8.3 Polynomial source-patch stability

Theorem 8.2 (Common-color coalescence outside a source patch)**.** *Under the preceding geometric and percolation hypotheses, there are constants $C < \infty$ and $\alpha > 0$ depending only on the RSW*

bounds and the fixed geometry such that, for $0 < R < R_0/6$, the common-color coupling of the two full exploration curves satisfies

$$\mathbb{P}\{d_{\mathrm{F}}(\gamma_1, \gamma_2) > 2R\} \le C \left(\frac{\epsilon + \delta}{R}\right)^{\alpha}. \tag{SH1}$$

More concretely, outside an event of the displayed probability the curves coalesce at a directed dual edge inside $B(p, R)$; both prefixes before that edge stay inside this ball, and the entire suffixes are identical.

Proof. Put $r_0 = 16(\epsilon + \delta)$ and $r_j = 16^j r_0$. For every j with $6r_j \le R$, the corresponding white-arch event depends only on shared free vertices in $A(p, r_j/2, 6r_j)$. These vertex sets are disjoint for different j, up to an immaterial lattice enlargement absorbed by increasing r_0. The events are therefore independent and each has probability at least ϑ. If there are N available bands, the probability of no shield is at most $(1 - \vartheta)^N$.

On any shield event select a white crosscut by a deterministic finite-graph rule using only shared colors, for example a lexicographic simple path in the crossing cluster. The two configurations select the same crosscut and the same single heterochromatic edge. Lemma 8.1 gives a common suffix. Before crossing the shield, each path stays in its source-side component, contained in $B(p, 6r_j) \subset B(p, R)$. Synchronize the two prefixes by any increasing reparameterization and then align their common suffix. The prefixes are in a ball of diameter $2R$, so their oriented Fréchet distance is at most $2R$.

The number of bands is a logarithmic function of $R/(\epsilon + \delta)$. For instance with $\alpha = -\log(1 - \vartheta)/\log 16 > 0$, enlarging a fixed constant C gives $(1 - \vartheta)^N \le C((\epsilon + \delta)/R)^{\alpha}$. This also covers the case with no available band by making its right-hand side at least one. □

Passage to marginal curve limits. Suppose along a common mesh subsequence the two exploration laws converge in the usual Polish oriented uniform-Fréchet curve space. The coupled pairs are tight because their two marginals are tight. Choose a joint subsequential limit. The set $\{d_{\mathrm{F}} > 2R\}$ is open in the product curve space; the Portmanteau inequality transfers (SH1), with $\delta \downarrow 0$, to this limiting coupling. Thus the same quantitative source-patch stability holds for the two marginal curve limits. It requires neither convergence of random slit-domain kernels nor a limiting strong DMP theorem.

If source patches themselves shrink and the mesh is negligible relative to their radii, this gives continuity in law for these particular local source perturbations. One may first fix R and let $\epsilon \downarrow 0$, then let $R \downarrow 0$, or balance the coupling estimate to obtain an explicit Prokhorov rate.

What is not proved. This is not a consequence of abstract unforced-crossing estimates for one curve. It uses a common independent coloring and the planar nonbranching interface to force *coalescence at one edge*. It also is not the full marked inverse-conformal-chart stability condition used in the affine-ansatz support theorems: arbitrary converging charts may create forced thin fjords, noncommon microscopic boundaries, or source approaches outside this shared half-plane/slit geometry. Extending the shield coupling to such approximations requires additional work. No classification of arbitrary source-local exploration ensembles follows from (SH1) alone.

9 Exact suffix coalescence survives curve limits

The source-shield estimate contains more information than a small Fréchet distance. Its exact common suffix is stable under taking a joint curve limit, provided that the prefixes stay in a fixed

closed set away from the target. This allows the discontinuous terminal metric decoder to be used without asserting that it is a continuous observable.

Let $K \subset \mathbb{C}$ be closed and consider two oriented continuous curves. Define E_K to be the relation that there exist cut times s_1, s_2 in continuous representatives on $[0,1]$ such that

$$\gamma_i[0,s_i] \subset K \quad (i=1,2), \qquad \gamma_1[s_1,1] = \gamma_2[s_2,1] \quad \text{as ordered unparameterized curves.} \tag{CB1}$$

The relation is independent of the chosen representatives and allows pauses, as does the usual uniform Fréchet curve space. It implies a common endpoint, which may belong to K in this definition. All applications below have limiting common target $q \notin K$.

Lemma 9.1 (A closed relation of common ordered suffixes)**.** *E_K is closed in the product uniform-Fréchet curve space.*

Proof. Suppose $(\gamma_1^n, \gamma_2^n) \to (\gamma_1, \gamma_2)$ and the pairs belong to E_K. Choose representatives of each convergent sequence converging uniformly to fixed continuous representatives of its limit. This is possible by the definition of the Fréchet metric; reparameterize the cut times along with the curves. Pass to a subsequence on which both cut times s_i^n converge to $s_i \in [0,1]$.

Closedness of K and uniform convergence give $\gamma_i[0,s_i] \subset K$. Linear rescaling of $[s_i^n, 1]$ to $[0,1]$ makes the restricted curves converge uniformly to the restrictions on $[s_i, 1]$, by uniform continuity of γ_i (a cut at $s_i = 1$ gives a constant curve). The two restrictions at stage n have Fréchet distance zero. Their limits therefore also have distance zero and are the same ordered curve. Thus the limiting pair belongs to E_K. □

Proposition 9.2 (Coalescence transfer and comparison of metric priors)**.** *Suppose pairs of discrete curves with a common target converge marginally to laws P_1, P_2 on the oriented curve space, and their couplings satisfy*

$$\mathbb{P}(E_K) \geq 1 - e_n, \qquad \limsup_n e_n \leq e < 1, \tag{CB2}$$

where K is a fixed closed ball not containing the target. Each limiting domain is analytic and regular near that target, and the two colored target germs agree. Suppose

$$P_i = \int P_{i,T}\, \pi_i(dT), \tag{CB3}$$

where $P_{i,T}$ is the affine-SLE$_6$ law with normalized tensor T in domain i, and π_i are Borel probability measures on tensors. Then, with total variation defined as $\sup_B |\pi_1(B) - \pi_2(B)|$,

$$\|\pi_1 - \pi_2\|_{\mathrm{TV}} \leq e. \tag{CB4}$$

In particular if both marginals are pure affine-SLE laws, any positive uniform lower bound on the discrete coalescence probability forces their tensors to be equal.

Proof. Tightness of the marginals makes the coupled pairs tight. Take a joint subsequential limit. Lemma 9.1 and the closed-set Portmanteau inequality give $\mathbb{P}(E_K) \geq 1 - e$ in the limit. Because the target is outside K, the common suffix on E_K includes a nonconstant ordered terminal germ at that target. The universal terminal decoder of Theorem 7.3 is the same for the two domains, since their colored target germs agree. Let T_i be its value on curve i. The laws of these random variables are π_i, and on E_K they are equal. Hence

$$\sup_B |\mathbb{P}(T_1 \in B) - \mathbb{P}(T_2 \in B)| \leq \mathbb{P}(T_1 \neq T_2) \leq e.$$

For two distinct point masses the left-hand side equals one, proving the last assertion. The proof uses exact suffix equality, not continuity of the decoder and not just small curve distance. □

For the source-shield coupling, take $K = \overline{B}(p, R)$ with the regular target outside this ball. Common discrete targets may vary with mesh, provided $q_n \to q \notin K$; the closed-relation proof already allows this. Equation (SH1) then gives

$$\|\pi_1 - \pi_2\|_{\mathrm{TV}} \le C(\epsilon/R)^{\alpha} \tag{CB5}$$

for two source-patch perturbations with common flat or straight-slit annuli. This is a quantitative stability statement for mixing measures, and yields exact tensor equality for pure marginals once the right side is less than one. It is conditional on the affine-mixture marginal ansatz; no such ansatz is derived from RSW here.

9.1 Stopping an ordered unparameterized curve on a circle

Fix $p \in \mathbb{R}^2$ and $R_0 > 0$. Work in the usual metric space of oriented continuous curves modulo increasing reparameterization, with the uniform Fréchet topology. Curves start at p and end outside $\overline{B(p, R_0)}$. Small changes of the starting points are harmless; for the sequence version they need only lie inside every fixed smaller ball for sufficiently large indices. For a continuous representative $\gamma : [0, 1] \to \mathbb{R}^2$ and $0 < r < R_0$, put

$$\tau_\gamma(r) = \inf\{t : |\gamma(t) - p| \ge r\}, \qquad E_r\gamma = \gamma|_{[0,\tau_\gamma(r)]}. \tag{LT1}$$

The stopped path is regarded as an ordered unparameterized curve, so this definition is independent of pauses or changes of clock.

Lemma 9.3 (Almost every circle is a continuity circle)**.** *For each fixed continuous curve γ, the map $(\eta, s) \mapsto E_s\eta$ is continuous at (γ, r) for all but at most countably many radii $r \in (0, R_0)$. The stopping map is Borel. Consequently, for any probability law μ on these curves, E_r is μ-almost surely continuous for Lebesgue-almost every deterministic r.*

Proof. Choose any continuous representative of γ. The function $r \mapsto \tau_\gamma(r)$ is nondecreasing and hence has at most countably many discontinuities. Suppose it is continuous at r. If $(\gamma_n, r_n) \to (\gamma, r)$ in the product of Fréchet and ordinary metrics, choose increasing reparameterizations for which representatives $\widetilde{\gamma}_n$ converge uniformly to γ. Writing $\varepsilon_n = \|\widetilde{\gamma}_n - \gamma\|_\infty + |r_n - r|$, first-hit times obey, for every slightly larger positive error tending to zero,

$$\tau_\gamma(r - 2\varepsilon_n) \;\le\; \tau_{\widetilde{\gamma}_n}(r_n) \;\le\; \tau_\gamma(r + 2\varepsilon_n). \tag{LT2}$$

If $\varepsilon_n = 0$ the assertion is immediate; using a strictly larger error avoids any immaterial equality convention. Continuity of τ_γ at r gives convergence of the middle times to $\tau_\gamma(r)$. Uniform continuity of γ then makes the stopped representatives, linearly recoded onto $[0, 1]$, converge uniformly. This proves joint continuity at (γ, r).

The stopping map is understood on all curves reaching the selected radius, including already stopped curves reaching a larger radius. For parameterized curves the stopping time is Borel, since

$$\{\tau_\gamma(r) \le t\} = \left\{ \max_{0 \le u \le t} |\gamma(u) - p| \ge r \right\}.$$

Stopping and linear recoding are therefore Borel path operations. They commute with reparameterization and give the usual Borel stopping map on ordered unparameterized curves.

For completeness, the almost-every-radius assertion does not require a measurable choice of representatives. For a map between metric spaces its set of continuity points is a G_δ set, obtained from its oscillations on neighborhoods. Apply this to the joint stopping map. Its discontinuity set in the product of curve space and radius space is Borel, and every vertical section has Lebesgue measure zero by the countability argument above. Fubini's theorem now gives the assertion under μ. □

10 Independent color resampling, initial germs, and a common affine metric

Theorem 10.1 (Initial-germ zero-one law from independent colors and shielding)**.** *Let P be a subsequential exploration-curve limit of the independent color model in a fixed domain with flat source germ. Assume the Polish oriented Fréchet curve topology and the flat-source shield estimate of Theorem 8.2. Then the initial ordered trace-germ sigma-field is P-trivial. Precisely, if $\sigma_r\gamma$ denotes the ordered prefix stopped on the radius-r circle about the source, this field is $\bigcap_{r>0}\overline{\sigma(\sigma_r\gamma)}^{P}$. It is the usual ordered positive-prefix germ, with no clock chosen. No affine ansatz, terminal tensor decoder, limiting DMP theorem, or continuum configuration-measurability theorem is required.*

More quantitatively, for each sufficiently small r, every bounded observable F of the prefix stopped at radius r and every bounded Lipschitz observable G of the entire curve satisfy

$$|\mathbb{E}_P(FG) - \mathbb{E}_P F\,\mathbb{E}_P G| \le C\|F\|_\infty(\operatorname{Lip} G + \|G\|_\infty)r^{\alpha/(1+\alpha)}, \tag{EDG}$$

first at almost every stopping-continuity radius r. Constants and length units are fixed by the source geometry. The zero-one assertion itself holds on the full initial-germ sigma-field.

Proof. Fix a source-patch radius ϵ and couple two discrete configurations with independent colors inside $B(p,\epsilon)$ and shared independent colors outside. For any fixed $r < \epsilon/3$ and all small enough meshes, the first exploration stopped at radius r is determined entirely by its own interior colors. It is therefore independent of the *entire second exploration*, which uses only the shared outer colors and the second interior color set.

Take a joint subsequential curve limit. Both marginals equal P. Choose an almost-sure stopping-continuity radius $r < \epsilon/3$. The continuous mapping theorem preserves the product law, so the stopped prefix of γ_1 is independent of the full curve γ_2. Consequently any bounded initial-germ observable $F(\gamma_1)$ is independent of $G(\gamma_2)$ for every bounded Lipschitz G on the full curve space. This step concerns the limiting product law; it does not attempt to pass an arbitrary discontinuous germ statistic through a discrete limit.

The shield estimate supplies, in this same coupling, for a fixed small R and $\epsilon \ll R$,

$$\mathbb{P}\{d_{\mathrm{F}}(\gamma_1,\gamma_2) > 2R\} \le C(\epsilon/R)^\alpha.$$

Thus

$$\begin{aligned}|\mathbb{E}_P(FG) - \mathbb{E}_P F\,\mathbb{E}_P G| &= |\mathbb{E}F(\gamma_1)(G(\gamma_1) - G(\gamma_2))| \\ &\le \|F\|_\infty\left(2\operatorname{Lip}(G)R + 2C\|G\|_\infty(\epsilon/R)^\alpha\right).\end{aligned} \tag{EDG1}$$

The left side depends only on the marginal law P. Let $\epsilon \downarrow 0$ with R fixed and then $R \downarrow 0$. It vanishes. Bounded Lipschitz functions determine finite signed measures on a Polish space, so every bounded initial-germ observable is independent of the full curve sigma-field. Since it belongs to that sigma-field, it is almost surely constant. This proves initial trace-germ triviality, including its completion.

For the quantitative prefix assertion, start with a fixed good radius r, use $\epsilon = 4r$, and take $R = r^{\alpha/(1+\alpha)}$ after fixing the reference length unit. For sufficiently small r this is an admissible radius with $\epsilon \ll R$. Formula (EDG1) gives (EDG). □

Proposition 10.2 (Pure affine limit at a flat source)**.** *Let one independent site-percolation model satisfy the flat-source shield coupling of Theorem 8.2. Let D have a flat source germ. Suppose along*

one mesh subsequence its exploration converges to

$$P = \int \nu_{D,p,q}^{T}\, \pi(dT), \tag{ED14}$$

where π is a Borel probability on projective affine tensors. Assume the standard Polish oriented curve topology. Then π is a point mass. No limiting domain Markov theorem or continuum configuration-measurability theorem is needed.

Proof. The regular source endpoint decoder has distribution π under (ED14) and is measurable in the initial trace-germ sigma-field. This sigma-field is trivial by Theorem 10.1. Thus the decoder is almost surely one deterministic tensor, and π is a point mass. The target need only meet the curve-convergence assumptions here; analytic target regularity enters the source-comparison theorem below. □

Lemma 10.3 (Splicing a regular colored source germ)**.** *Fix a bounded reference domain D_0 with flat regular source $p = 0$, analytic regular target q, and one fixed boundary-color convention. Given any analytic regular colored source germ G at 0 with that convention, and any sufficiently small $\epsilon > 0$, there is a bounded marked domain $D_{G,\epsilon}$ agreeing with D_0 outside $B(0, \epsilon)$ and having exactly the germ G at 0. Its target germ at q agrees with D_0. The boundary can be chosen smooth except for any corners already present in D_0; only its two marked germs need to be analytic.*

Proof. For any analytic regular colored source germ G at p and any small ϵ, an orientation-preserving smooth diffeomorphism supported in $B(p, \epsilon)$ can map the source germ of one fixed flat-source reference domain to G, fixing p and the distant regular target. For an explicit local construction, use a radial twist to rotate a small core to the required tangent, followed by a cutoff graph shear to realize the analytic curvature in a still smaller core. The graph derivative can be made arbitrarily small, so the cutoff shear remains a diffeomorphism. Call the resulting domain $D_{G,\epsilon}$. It agrees with the reference outside the source patch and agrees with G in a still smaller neighborhood. The support avoids q, and ambient orientation preserves the boundary color order. Applying the diffeomorphism to D_0 proves the claim. □

Theorem 10.4 (One deterministic metric from a coherent affine mixture)**.** *Consider a translation-covariant source-local exploration family arising from one independent Bernoulli nonbranching cellular percolation model admitting the flat-source shield coupling of Theorem 8.2. Suppose each bounded-domain marginal with analytic regular marks is a Borel mixture of fixed affine-SLE_6 laws. Require the family to include the source splices of Lemma 10.3, and require coherent subsequential limits for these comparisons: for every such splice and the reference, compatible common-color discrete approximations along one common mesh subsequence realize the two specified marginal limits.*

Then one deterministic projective affine tensor is shared by every such marginal, and each marginal is the pure affine-SLE_6 law for that tensor. Strong DMP and marked inverse-chart continuity are not premises of this theorem.

Proof. The reference mixing measure π_0 is a point mass by Proposition 10.2. Fix another original domain with analytic regular source germ G. Translation covariance allows its source to be placed at 0 without changing its physical projective tensor. For every small ϵ, source locality identifies the original initial prefix law with that in $D_{G,\epsilon}$ through a common source neighborhood. The source tensor decoder is a function of this prefix and the common germ. Its distribution is the mixing measure in either state, so both have one common mixing measure π_G, independent of ϵ.

Coherence and the common-color shield coupling give a coupling of the two limiting curves in $D_{G,\epsilon}$ and D_0 such that their ordered terminal segments agree outside an event of probability at most $C(\epsilon/R)^\alpha$, for one fixed small R in the shared flat outer geometry. Their target germs coincide. Apply the two terminal decoders; the resulting tensor variables have respective laws π_G, π_0 and are equal on the coalescence event. The coupling inequality for total variation therefore gives

$$\|\pi_G - \pi_0\|_{\mathrm{TV}} \le C(\epsilon/R)^\alpha. \tag{ED16}$$

Here total variation is $\sup_B |\pi_G(B) - \pi_0(B)|$. Letting $\epsilon \downarrow 0$ gives $\pi_G = \pi_0$. Since π_0 is a point mass, the original marginal is pure and shares the same tensor. The argument applies separately to each original regular-source state; no simultaneous uncountable full-probability assertion is required. □

Thus, subject to the explicitly stated coherent independent-color scaling and shielding premises, the mixed-affine ansatz on regular domains already collapses to one deterministic common affine metric. This does not derive the mixed-affine ansatz for an arbitrary curve law, and does not classify arbitrary abstract three-axiom families.

Initial characteristic limits become deterministic. At any source to which the preceding zero-one argument applies, a finite initial limit in probability of an adapted characteristic A_t/t or C_t/t is deterministic. To see germ measurability, stop the characteristic at the first exit from an arbitrary source ball. That exit has strictly positive capacity time; the stopped and unstopped limits therefore agree. A limit in probability of variables measurable from the stopped prefix has a measurable version in its completed sigma-field. This holds for every small radius, so the limit belongs to the trivial initial germ. This removes the deterministic-value premise at these starts once existence of a tangent and natural-trace-filtration compatibility are known; neither tangent existence nor the required slit-uniform bound is obtained here.

11 Total-variation decoupling of an initial prefix and a distant suffix

The closed exact-coalescence relation gives a stronger consequence than bounded-Lipschitz covariance. It controls arbitrary measurable observables of the distant terminal suffix in total variation. This uses the actual independent-color source-shield coupling, not just abstract locality or an unforced-crossing bound.

Fix a bounded domain with source p, target q, flat source geometry, and a marginal curve scaling limit P satisfying the hypotheses of Theorem 10.1. Let $\sigma_r\gamma$ be the initial ordered prefix stopped on the radius-r circle about p. For $R < |q-p|$ define $\lambda_R\gamma$ to be the ordered terminal suffix beginning at the *last* visit to $\overline{B}(p,R)$. For any continuous representative on $[0,1]$ this means restriction from

$$t_R(\gamma) = \max\{t \in [0,1] : |\gamma(t) - p| \le R\}$$

to 1, followed by linear reparameterization. The resulting ordered curve does not depend on the representative or pauses.

Both stopping operations are Borel maps on the assumed Polish oriented Fréchet curve space. For completeness, on the continuous-parameter curve space, t_R is Borel because, for fixed a,

$$\{t_R \ge a\} = \left\{ \min_{a \le t \le 1} |\gamma(t) - p| \le R \right\}$$

is closed. Restriction and rescaling are jointly continuous in a curve and its cut parameter. Hence the terminal-suffix map to the quotient curve space is Borel on representatives and invariant under reparameterization. It descends to a Borel map on the Polish quotient: the preimages of Borel sets are saturated Borel sets, whose images and complements under the continuous quotient map are analytic; both are therefore Borel. First-exit stopping has the same Borel property.

Theorem 11.1 (Prefix–suffix decoupling)**.** *There exist $C < \infty$ and $\alpha > 0$, determined by the source-shield constants, such that for all sufficiently small R and all $0 < r < R/64$,*

$$\|\mathcal{L}_P(\sigma_r\gamma, \lambda_R\gamma) - \mathcal{L}_P(\sigma_r\gamma) \otimes \mathcal{L}_P(\lambda_R\gamma)\|_{\mathrm{TV}} \le C(r/R)^\alpha. \tag{SB1}$$

Here total variation is $\sup_A |\mu(A) - \nu(A)|$ on the product space. In particular this estimate holds at every *small radius r, not merely almost every stopping-continuity radius.*

Proof. Set $\epsilon = 8r$. At the discrete level take two configurations with independent colors in $B(p, \epsilon)$ and identical independent colors outside that ball. Both marginal explorations have the same scaling limit P. Their joint laws are tight. Choose a radius $\rho \in (r, 2r)$ at which first-exit stopping is P-almost surely continuous; such radii have full Lebesgue measure by Lemma 9.3. The first discrete exploration up to radius ρ uses only the first configuration's interior colors, once the mesh is sufficiently small. It is consequently independent of the entire second exploration. Continuous mapping along a joint subsequential limit gives

$$\sigma_\rho\gamma_1 \perp\!\!\!\perp \gamma_2.$$

Stopping the limiting first prefix again, now at radius r, is a Borel operation and requires no continuity assertion at r. Thus $\sigma_r\gamma_1$ is independent of the full curve γ_2.

Apply the source-shield estimate with shield radius $R/2$. Outside an event of probability at most $C(r/R)^\alpha$, both prefixes before coalescence stay in $K = \overline{B}(p, R/2)$ and the ordered suffixes are identical. Lemma 9.1 and the closed-set Portmanteau inequality preserve this statement in the same joint subsequential limit. Since $q \notin \overline{B}(p, R)$, each full curve's last visit to $\overline{B}(p, R)$ is on the common suffix. Therefore

$$\mathbb{P}\{\lambda_R\gamma_1 \ne \lambda_R\gamma_2\} \le C(r/R)^\alpha.$$

The pairs $(\sigma_r\gamma_1, \lambda_R\gamma_1)$ and $(\sigma_r\gamma_1, \lambda_R\gamma_2)$ form a coupling of the joint law in (SB1) with the displayed product law. The usual coupling bound for total variation proves (SB1). □

Corollary 11.2 (Expected conditional total-variation stability)**.** *Let $K_r(x, \cdot)$ be a regular conditional law of $\lambda_R\gamma$ given $\sigma_r\gamma = x$. Then*

$$\int \|K_r(x, \cdot) - \mathcal{L}_P(\lambda_R\gamma)\|_{\mathrm{TV}}\ \mathcal{L}_P(\sigma_r\gamma)(dx) \le C(r/R)^\alpha. \tag{SB2}$$

Consequently for bounded Borel functions F, G of the two respective variables,

$$|\operatorname{Cov}_P(F(\sigma_r\gamma), G(\lambda_R\gamma))| \le 2C\|F\|_\infty\|G\|_\infty(r/R)^\alpha. \tag{SB3}$$

Proof. On standard Borel spaces, total variation between a joint law and the product of its marginals is the integral of the total variations of the corresponding regular conditional laws. One may verify the identity using a countable generating algebra to compute the variation of each signed conditional measure. This gives (SB2). Integrating the bounded product observable against the difference of the two probability laws costs at most twice its supremum norm times the convention for total variation used in (SB1), proving (SB3). □

Variants and limitation. The same proof applies to any standard-Borel-valued observable Y_R that agrees whenever two curves belong to $E_{\overline{B}(p,R/2)}$. For example one may use the compact trace outside the open radius-R ball instead of the ordered distant suffix. Measurability of a given variant should be checked separately. Completed prefix sigma-fields are harmless, since versions may be chosen modulo their common marginal null sets.

This is quantitative spatial loss of memory at a deterministic regular source. It does not identify a conditional continuation kernel in an arbitrary random slit domain: the proof has no shield estimate uniform over rough explored frontiers. In particular it neither proves strong DMP in a curve limit nor the whole-slit stability premise of the driving-rigidity results. The whole-tube approach based on boundary three-arm estimates still needs the deterministic implication from ordered curve disagreement to an appropriate arm event.

12 Annular crossing statistics self-average at a regular source

The same source resampling argument applies to a macroscopic observable whenever the closed exact-coalescence relation forces its equality. It therefore controls annular crossings made *before the first outer exit*, even if the full curve later returns arbitrarily close to the source. No source nonreturn assumption is needed.

Proposition 12.1 (Decoupling for coalescence-invariant observations)**.** *Under the hypotheses of Theorem 11.1, let Y_R be a Borel map from the oriented curve space into a standard Borel space. Assume*

$$(\gamma,\widetilde{\gamma}) \in E_{\overline{B}(p,R/2)} \quad \Longrightarrow \quad Y_R(\gamma) = Y_R(\widetilde{\gamma}). \tag{AS1}$$

Then, for $0 < r < R/64$,

$$\big\|\mathcal{L}_P(\sigma_r\gamma, Y_R(\gamma)) - \mathcal{L}_P(\sigma_r\gamma)\otimes\mathcal{L}_P(Y_R(\gamma))\big\|_{\mathrm{TV}} \le C(r/R)^{\alpha}. \tag{AS2}$$

The same conclusion holds if the first variable is any Borel function of $\sigma_r\gamma$.

Proof. Use exactly the independent-inner/shared-outer coupling in the proof of Theorem 11.1, including a good stopping-continuity radius between r and $2r$. Its limiting curves obey $\sigma_r\gamma_1 \perp\!\!\!\perp \gamma_2$ and

$$\mathbb{P}\big(E_{\overline{B}(p,R/2)}\big) \ge 1 - C(r/R)^{\alpha}.$$

Assumption (AS1) implies $\mathbb{P}\{Y_R(\gamma_1) \ne Y_R(\gamma_2)\} \le C(r/R)^{\alpha}$. Couple the two pairs with common first variable $\sigma_r\gamma_1$, as in the earlier proof. The second pair has the product law by independence. This proves (AS2). Applying a Borel map to the first variable cannot increase total variation. □

Fix $a > 0$ sufficiently small that the source-shield construction applies up to scale a, and $|q-p| > a$. Put

$$r_n = a2^{-n}, \qquad n = 1,2,\dots.$$

For a continuous representative of γ define

$$\tau_n = \inf\{t : |\gamma(t)-p| = 2r_n\}, \qquad \eta_n = \max\{t \le \tau_n : |\gamma(t)-p| \le r_n\}. \tag{AS3}$$

Let $B_n(\gamma)$ be the ordered restriction to $[\eta_n,\tau_n]$, translated by $-p$ and dilated by r_n^{-1}. This is an oriented curve in $\{1 \le |z| \le 2\}$ from its inner to its outer boundary. It is the last inner-to-outer crossing before the first outer hit. Only this crossing is recorded; later returns of the full curve are immaterial.

The map B_n is Borel and invariant under changes of representative. Indeed τ_n is a Borel first-hitting operation. Restrict first to $[0,\tau_n]$, linearly reparameterized on $[0,1]$. On this stopped curve, η_n is a Borel last visit to the closed inner ball, by the last-visit argument in Section 11. Successive restriction, translation, dilation, and passage to the oriented quotient are Borel. In particular B_n is a Borel function of $\sigma_{2r_n}\gamma$.

There is a second property crucial to the proof: if $(\gamma,\widetilde{\gamma}) \in E_{\overline{B}(p,r_n/2)}$, then $B_n(\gamma) = B_n(\widetilde{\gamma})$. The prefixes before the common suffix remain strictly inside the inner circle, and the common suffix starts there. Both first outer hits thus occur on that suffix, and so do their last inner-ball visits before those hits. The corresponding ordered crossing segments agree exactly.

Theorem 12.2 (Exponential absolute regularity across dyadic source scales)**.** *Under the same source-shield and independent-color limit hypotheses, the annular crossing sequence $(B_n)_{n\ge1}$ satisfies*

$$\begin{aligned}\big\|\mathcal{L}_P\big((B_1,\dots,B_m),(B_{m+k},B_{m+k+1},\dots)\big)\\ -\mathcal{L}_P(B_1,\dots,B_m)\otimes\mathcal{L}_P(B_{m+k},B_{m+k+1},\dots)\big\|_{\mathrm{TV}} \le C_0 2^{-\alpha k}, \qquad m,k\ge1.\end{aligned} \tag{AS4}$$

The countable product on the fine-scale side is a standard Borel space. Constants are independent of m and k. No stationarity or dilation covariance of the marginal curve limit is assumed.

Proof. For $k \ge 8$, apply Proposition 12.1 with $R = r_m$ and $r = 2r_{m+k}$. The vector $Y_R = (B_1,\dots,B_m)$ satisfies (AS1): every one of its inner radii is at least r_m, so coalescence inside the radius-$r_m/2$ ball occurs before all the relevant inner and outer visits. The whole countable vector $(B_{m+k},B_{m+k+1},\dots)$ is a Borel function of $\sigma_{2r_{m+k}}\gamma$, because every corresponding outer radius is at most $2r_{m+k}$. Thus (AS2) gives the upper bound

$$C(2r_{m+k}/r_m)^\alpha = C2^\alpha 2^{-\alpha k}.$$

Enlarge C_0 to cover $1 \le k < 8$ by the trivial total-variation bound one. Interchanging the two coordinates of a joint law preserves total variation, giving (AS4) in the displayed order. □

Corollary 12.3 (A strong law for bounded annular observations)**.** *Let f_n be any deterministic sequence of Borel real functions of an annular oriented crossing, with $|f_n| \le 1$. Put $Z_n = f_n(B_n)$. Then*

$$\frac{1}{N}\sum_{n=1}^{N}\{Z_n - \mathbb{E}_P Z_n\} \longrightarrow 0 \qquad P\text{-almost surely.} \tag{AS5}$$

In particular, for a fixed bounded Borel f, if the deterministic Cesàro means

$$\frac{1}{N}\sum_{n=1}^{N}\mathbb{E}_P f(B_n)$$

converge to c_f, then the empirical means $N^{-1}\sum_{n\le N} f(B_n)$ converge almost surely to c_f. These assertions hold simultaneously for any prescribed countable collection of choices of the sequence (f_n).

Proof. The total-variation estimate (AS4) and $|Z_n| \le 1$ imply

$$|\operatorname{Cov}_P(Z_n,Z_j)| \le 2C_0 2^{-\alpha|n-j|} \quad (n \ne j).$$

Consequently, for $S_N = \sum_{n\le N}(Z_n - \mathbb{E}_P Z_n)$,

$$\mathbb{E}_P S_N^2 \le C_1 N.$$

Chebyshev's inequality and Borel–Cantelli give $S_{\ell^2}/\ell^2 \to 0$ almost surely, since for every $\zeta > 0$,

$$\sum_{\ell=1}^{\infty} \mathbb{P}\{|S_{\ell^2}| > \zeta \ell^2\} \le \frac{C_1}{\zeta^2} \sum_{\ell=1}^{\infty} \ell^{-2} < \infty.$$

For $\ell^2 \le N < (\ell+1)^2$, boundedness of the centered summands gives $|S_N - S_{\ell^2}| \le 2(2\ell+1)$. Dividing by N yields (AS5). Add the deterministic expectations for the second assertion, and intersect countably many probability-one events for the last one. □

Meaning and remaining gap. Annular crossing statistics of one limiting exploration self-average over logarithmically many source scales. The centering sequence is deterministic, even when the scaling limit itself has not been shown dilation invariant. Neither the convergence of those deterministic expectations nor their values is asserted. A source law that is invariant under dyadic dilation would make the expectations constant, but such invariance is an additional premise and is not inferred from precompactness of the collection of scaling limits. The result supplies a rigorous scale-statistics diagnostic without an affine-SLE ansatz; it does not classify those deterministic statistics or produce random slit-domain exploration kernels.

References

[1] G. F. Lawler, *Schramm–Loewner Evolution*, Park City lecture notes (2007), Lectures 3–4. https://math.uchicago.edu/~lawler/utah.pdf.

[2] Y. Yuan, *Topological characterisations of Loewner traces*, Indiana Univ. Math. J. **71** (2022), 1027–1046. https://arxiv.org/abs/2003.05535.

[3] J. Miller and S. Sheffield, *Imaginary geometry III: reversibility of SLE$_\kappa$ for $\kappa \in (4, 8)$*, Ann. of Math. **184** (2016), 455–486. https://annals.math.princeton.edu/2016/184-2/p03.

[4] *Research record: locality, terminal information, and planar exploration laws*, `progress.tex` in this repository, including the original-locality investigation and the corrected-locality continuation (27 October 2026).