

MATH 10A – Fall 2018
Quiz 10 – Prepared by John Yirong Zhen
Date: 10/30/2018

You are to finish this quiz in 10 minutes. You are allowed one single-sided letter-size cheat sheet. No calculators or other notes/books/devices are allowed.

Your cheatsheet must be handwritten by you, no photocopying or preprinted (unless you have written permission from the instructor). Try your best! Stay calm and good luck!

I. True/False (2 pts)

Circle T or F in the space provided in front of the statement to indicate whether it is true or false respectively. You get +1 for a correct answer, -1 for incorrect, and 0 for leaving it blank. (You should not guess if you don't know the answer.)

You do not need to justify your answers for T/F statements.

- ① F One equilibrium solution to the differential equations $\frac{dx}{dt} = 2x - 0.01xy$ and $\frac{dy}{dt} = -0.5y + 0.0001xy$ is $x = 5000$ and $y = 200$, and in this ecological system with two species, x is the prey and y is the predator.
- T ② If $a = \frac{7}{9}b$, the system $ax_1 + bx_2 = y_1$ and $7x_1 + 9x_2 = y_2$ will have a unique solution.

II. Written problems (10pts)

- You **MUST justify your answer** to undoubtably convince me that you solved and not guessed it. Partial credit will be given to good work and progress even if there is no final answer or the answer is incorrect. On the other hand, bogus justification for a correct answer will receive a 0.
- Keep your scratch work separate. Cross out writing you don't want to be graded and clearly label the parts you want to be graded.
- Points will be deducted for incorrect writings that you "forget to cross out."

See problem on back.

Consider the linear system of equations

$$a_{11}x_1 + a_{12}x_2 = b_1$$

$$a_{21}x_1 + a_{22}x_2 = b_2$$

Suppose that the matrix $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$ is nonsingular. Derive expressions for the solutions x_1 and x_2 .

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \frac{1}{a_{11}a_{22}-a_{21}a_{12}} \begin{bmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} = \frac{1}{a_{11}a_{22}-a_{21}a_{12}} \begin{bmatrix} b_1a_{22} - b_2a_{12} \\ -b_1a_{21} + b_2a_{11} \end{bmatrix}.$$

Therefore, $x_1 = \frac{b_1a_{22}-b_2a_{12}}{a_{11}a_{22}-a_{21}a_{12}}$ and $x_2 = \frac{-b_1a_{21}+b_2a_{11}}{a_{11}a_{22}-a_{21}a_{12}}$