



$$=5P \quad \text{교구} \quad M=:: \quad \text{H}::: \quad 0$$

$$>5 \quad =::: \quad Q$$

$$A5$$

|       |    |       |          |
|-------|----|-------|----------|
| TRj h | H  |       |          |
| =5P   | 교구 | M=: : | 3 HGC: 0 |
| >5GC: | Q  |       | HA::: 0  |
| A5    |    |       |          |

|              |        |   |                    |
|--------------|--------|---|--------------------|
| 5            | 3TRj h | 5 | m <sub>0</sub> 개게검 |
| 가원열만가원열만가원열만 |        | 5 |                    |

$$] E \mathbb{Q}$$
[illegible]

] E[ 46 한자서체외근의찬찬에5

] Eq

TRj h 5 % 고 간 %  
3 5 H  
각 5분과 15분 3개~결정간 공간 14mm

|       |   |
|-------|---|
| TRj h | H |
| =5    |   |
| >5    |   |
| A5    |   |
| B5    | 3 |

] E G a

|                   |       |   |   |
|-------------------|-------|---|---|
| m <sub>굴베게검</sub> | TRj h | 5 | 5 |
|-------------------|-------|---|---|

Figure 10: A 4x4 grid of scatter plots showing the relationship between the number of HTRj h and the number of HTRj h for different values of HTRj h (5, >5, A5, B5Rpi). The plots are arranged in a 4x4 grid, with the top row showing HTRj h = 5, the second row showing HTRj h > 5, the third row showing HTRj h = A5, and the bottom row showing HTRj h = B5Rpi. The x-axis for all plots is 'HTRj h' and the y-axis is 'HTRj h'. The plots show a positive correlation between the two variables, with the strength of the correlation increasing as HTRj h increases.