

EcoRI and NsiI RFLPs at a human PLA₂ gene on chromosome 1

Umesh Masharani, Roger T. Coleman, Lorin K. Johnson and Jeffrey J. Seilhamer

California Biotechnology Inc., 2450 Bayshore Parkway, Mountain View, CA 94043, USA

SOURCE/DESCRIPTION: pCB-04, a 0.7kb genomic fragment encoding the second exon of a phospholipase homologue (1,2) of human pancreatic phospholipase A₂ (3) was subcloned into the EcoRI site of pUC9.

POLYMORPHISMS: EcoRI (GAATTC) detects a single two allele polymorphism with fragments at either 8.4kb or 8.0kb.

NsiI (ATGCAT) detects a single two allele polymorphism with fragments at either 5.5kb or 5.0kb.

FREQUENCY: Studied in 105 unrelated North American caucasians

- | | | |
|------------|----------------|-------------|
| (i) EcoRI: | 8.4 kb allele: | 0.85 ± 0.03 |
| | 8.0 kb allele: | 0.15 ± 0.03 |
| (ii) NsiI: | 5.5 kb allele: | 0.09 ± 0.02 |
| | 5.0 kb allele: | 0.91 ± 0.02 |

CHROMOSOMAL LOCALIZATION: The genomic fragment encoding the second exon of a gene homologous to human pancreatic phospholipase A₂ has been assigned to the chromosome 1 by Southern filter hybridization of DNA from human-rodent hybrids (1,2).

MENDELIAN INHERITANCE: Codominant segregation of both RFLPs observed in two families (20 individuals).

PROBE AVAILABILITY: Requests for probe to J.J.S. at above address.

- REFERENCES:**
1. Seilhamer J J *et al.* 1988 J. Cell Biol. Suppl. 12E, 55
 2. Seilhamer J J *et al.* 1988 (manuscript in preparation).
 3. Seilhamer J J *et al.* 1986 DNA 5, 519-527.

