

Book Creator from Wikipedia

Contents

1	Selenium	1
1.1	Characteristics	1
1.1.1	Physical properties	1
1.1.2	Isotopes	2
1.2	Chemical compounds	2
1.2.1	Chalcogen compounds	2
1.2.2	Halogen compounds	2
1.2.3	Selenides	3
1.2.4	Other compounds	3
1.2.5	Organoselenium compounds	3
1.3	History	3
1.4	Occurrence	4
1.5	Production	4
1.6	Applications	4
1.6.1	Manganese electrolysis	4
1.6.2	Glass production	4
1.6.3	Alloys	5
1.6.4	Other uses	5
1.7	Biological role	5
1.7.1	Evolution in biology	6
1.7.2	Nutritional sources of selenium	6
1.7.3	Indicator plant species	6
1.7.4	Medical use	6
1.7.5	Detection in biological fluids	6
1.7.6	Toxicity	7
1.7.7	Deficiency	7
1.7.8	Controversial health effects	8
1.8	See also	8
1.9	References	8
1.10	External links	12
2	JScript	14
2.1	Comparison to JavaScript	14

2.2	Versions	14
2.2.1	JScript	14
2.2.2	JScript .NET	14
2.3	See also	15
2.4	Notes	15
2.5	References	15
2.6	External links	15
2.7	Text and image sources, contributors, and licenses	16
2.7.1	Text	16
2.7.2	Images	16
2.7.3	Content license	17

Chapter 1

Selenium


This article is about the chemical element. For the software testing framework, see [Selenium \(software\)](#).

Selenium is a chemical element with symbol **Se** and atomic number 34. It is a nonmetal with properties that are intermediate between the elements above and below in the periodic table, sulfur and tellurium. It rarely occurs in its elemental state or as pure ore compounds in the Earth's crust. Selenium (Greek *σελήνη* *selene* meaning "Moon") was discovered in 1817 by Jöns Jacob Berzelius, who noted the similarity of the new element to the previously discovered tellurium (named for the Earth).

Selenium is found in metal sulfide ores, where it partially replaces the sulfur. Commercially, selenium is produced as a byproduct in the refining of these ores, most often during production. Minerals that are pure selenide or selenate compounds are known but rare. The chief commercial uses for selenium today are glassmaking and pigments. Selenium is a semiconductor and is used in photocells. Applications in electronics, once important, have been mostly supplanted by silicon semiconductor devices. Selenium is still used in a few types of DC power surge protectors and one type of fluorescent quantum dot.

Selenium salts are toxic in large amounts, but trace amounts are necessary for cellular function in many organisms, including all animals. Selenium is an ingredient in many multivitamins and other dietary supplements, including infant formula. It is a component of the antioxidant enzymes glutathione peroxidase and thioredoxin reductase (which indirectly reduce certain oxidized molecules in animals and some plants). It is also found in three deiodinase enzymes, which convert one thyroid hormone to another. Selenium requirements in plants differ by species, with some plants requiring relatively large amounts and others apparently requiring none.^[4]

1.1 Characteristics



Structure of hexagonal (gray) selenium

1.1.1 Physical properties

Selenium forms several allotropes that interconvert with temperature changes, depending somewhat on the rate of temperature change. When prepared in chemical reactions, selenium is usually an amorphous, brick-red powder. When rapidly melted, it forms the black, vitreous form, usually sold commercially as beads.^[5] The structure of black selenium is irregular and complex and consists of polymeric rings with up to 1000 atoms per ring. Black Se is a brittle, lustrous solid that is slightly soluble in CS₂. Upon heating, it softens at 50 °C and converts to gray selenium at 180 °C; the transformation temperature is reduced by presence of halogens and amines.^[6]

The red α , β , and γ forms are produced from solutions of black selenium by varying the evaporation rate of the solvent (usually CS₂). They all have relatively low, monoclinic crystal symmetries and contain nearly identical puckered Se₈ rings with different arrangements, as in sulfur. The packing is most dense in the α form. In the Se₈ rings, the Se-Se distance is 233.5 pm and Se-Se-Se angle is 105.7°. Other selenium allotropes may contain Se₆ or Se₇ rings.^[6]

The most stable and dense form of selenium is gray and has a hexagonal crystal lattice consisting of helical polymeric chains, where the Se-Se distance is 237.3 pm and Se-Se-Se angle is 130.1°. The minimum distance be-

tween chains is 343.6 pm. Gray Se is formed by mild heating of other allotropes, by slow cooling of molten Se, or by condensing Se vapor just below the melting point. Whereas other Se forms are insulators, gray Se is a semiconductor showing appreciable **photoconductivity**. Unlike the other allotropes, it is insoluble in CS₂.^[6] It resists oxidation by air and is not attacked by nonoxidizing **acids**. With strong reducing agents, it forms polyselenides. Selenium does not exhibit the changes in viscosity that sulfur undergoes when gradually heated.^{[5][7]}

1.1.2 Isotopes

Main article: **Isotopes of selenium**

Selenium has six naturally occurring isotopes. Synthetic isotope ⁷⁹Se and 23 others have been identified:

⁸²Se is stable, for all practical purposes. See also *Selenium-79* for recent changes in the measured half-life of ⁷⁹Se, which are important for the dose calculations in the geological disposal of long-lived **radioactive waste**.^[10]



1.2 Chemical compounds

See also: **Category:Selenium compounds** and **organoselenium chemistry**

Selenium compounds commonly exist in the **oxidation states** −2, +2, +4, and +6.


1.2.1 Chalcogen compounds

Selenium forms two oxides: **selenium dioxide** (SeO₂) and **selenium trioxide** (SeO₃). Selenium dioxide is formed by the reaction of elemental selenium with oxygen:^[5]




Structure of the polymer SeO₂: The (pyramidal) Se atoms are yellow.

It is a polymeric solid that forms monomeric SeO₂ molecules in the gas phase. It dissolves in water to form **selenous acid**, H₂SeO₃. Selenous acid can also be made directly by oxidizing elemental selenium with **nitric acid**.^[11]




Unlike sulfur, which forms a stable trioxide, selenium trioxide is thermodynamically unstable and decomposes to the dioxide above 185 °C:^{[5][11]}



Selenium trioxide is produced in the laboratory by the reaction of **anhydrous potassium selenate** (K₂SeO₄) and sulfur trioxide (SO₃).^[12]


Salts of selenous acid are called selenites. These include **silver selenite** (Ag₂SeO₃) and **sodium selenite** (Na₂SeO₃).

Hydrogen sulfide reacts with aqueous selenous acid to produce **selenium disulfide**:



Selenium disulfide consists of 8-membered rings. It has an approximate composition of SeS₂, with individual rings varying in composition, such as Se₄S₄ and Se₂S₆. Selenium disulfide has been used in shampoo as an **antidandruff** agent, an inhibitor in polymer chemistry, a glass dye, and a reducing agent in **fireworks**.^[11]


Selenium trioxide may be synthesized by dehydrating **selenic acid**, H₂SeO₄, which is itself produced by the oxidation of selenium dioxide with **hydrogen peroxide**.^[13]



Hot, concentrated selenic acid can react with gold to form gold(III) selenate.^[14]

1.2.2 Halogen compounds


Iodides of selenium are not well known. The only stable chloride is **selenium monochloride** (Se₂Cl₂), which might be better known as selenium(I) chloride; the corresponding **bromide** is also known. These species are structurally analogous to the corresponding **disulfur dichloride**. Selenium dichloride is an important reagent in the preparation of selenium compounds (e.g. the preparation of Se₇). It is prepared by treating selenium with **sulfuryl chloride** (SO₂Cl₂).^[15] Selenium reacts with **fluorine** to form **selenium hexafluoride**:



In comparison with its sulfur counterpart (**sulfur hexafluoride**), **selenium hexafluoride** (SeF₆) is more reactive and is a toxic **pulmonary irritant**.^[16] Some of the selenium oxyhalides, such as **selenium oxyfluoride** (SeOF₂) and **selenium oxychloride** (SeOCl₂) have been used as specialty solvents.^[5]

1.2.3 Selenides

Analogous to the behavior of other chalcogens, selenium forms **hydrogen selenide**, H_2Se . It is a strongly **odiferous**, toxic, and colorless gas. It is more acidic than H_2S . In solution it ionizes to HSe^- . The selenide dianion Se^{2-} forms a variety of compounds, including the minerals from which selenium is obtained commercially. Illustrative selenides include **mercury selenide** (HgSe), **lead selenide** (PbSe), **zinc selenide** (ZnSe), and **copper indium gallium diselenide** ($\text{Cu}(\text{Ga},\text{In})\text{Se}_2$). These materials are **semiconductors**. With highly electropositive metals, such as **aluminium**, these selenides are prone to hydrolysis:^[5]




Alkali metal selenides react with selenium to form polyselenides, Se_2^{n-} , which exist as chains.

1.2.4 Other compounds

Tetraselenium tetranitride, Se_4N_4 , is an explosive orange compound analogous to **tetrasulfur tetranitride** (S_4N_4).^{[5][17][18]} It can be synthesized by the reaction of **selenium tetrachloride** (SeCl_4) with $[(\text{CH}_3)_3\text{Si}]_2\text{N}_2$.^[19]

Selenium reacts with **cyanides** to yield selenocyanates:^[5]



1.2.5 Organoselenium compounds

Main article: **Organoselenium chemistry**

Selenium, especially in the II oxidation state, forms stable bonds to **carbon**, which are structurally analogous to the corresponding **organosulfur compounds**. Especially common are selenides (R_2Se , analogues of **thioethers**), diselenides (R_2Se_2 , analogues of **disulfides**), and **selenols** (RSeH , analogues of **thiols**). Representatives of selenides, diselenides, and selenols include respectively **selenomethionine**, **diphenyldiselenide**, and **benzeneselenol**. The sulfoxide in sulfur chemistry is represented in selenium chemistry by the selenoxides (formula $\text{RSe}(\text{O})\text{R}$), which are intermediates in organic synthesis, as illustrated by the **selenoxide elimination reaction**. Consistent with trends indicated by the **double bond rule**, selenoketones, $\text{R}(\text{C}=\text{Se})\text{R}$, and selenaldehydes, $\text{R}(\text{C}=\text{Se})\text{H}$, are rarely observed.^[20]

1.3 History

Selenium (Greek *σελήνη* *selene* meaning “Moon”) was discovered in 1817 by Jöns Jakob Berzelius and Johan Gottlieb Gahn.^[21] Both chemists owned a chemistry plant near Gripsholm, Sweden, producing **sulfuric acid** by the **lead chamber process**. The pyrite from the **Falun mine** created a red precipitate in the lead chambers which was presumed to be an arsenic compound, so the pyrite’s use to make acid was discontinued. Berzelius and Gahn wanted to use the pyrite and they also observed that the red precipitate gave off a smell like **horseradish** when burned. This smell was not typical of arsenic, but a similar odor was known from **tellurium** compounds. Hence, Berzelius’s first letter to **Alexander Marcet** stated that this was a tellurium compound. However, the lack of tellurium compounds in the **Falun mine** minerals eventually led Berzelius to reanalyze the red precipitate, and in 1818 he wrote a second letter to Marcet describing a newly found element similar to **sulfur** and tellurium. Because of its similarity to tellurium, named for the Earth, Berzelius named the new element after the **Moon**.^{[22][23]}

In 1873, **Willoughby Smith** found that the electrical resistance of grey selenium was dependent on the ambient light. This led to its use as a cell for sensing light. The first commercial products using selenium were developed by **Werner Siemens** in the mid-1870s. The selenium cell was used in the **photophone** developed by **Alexander Graham Bell** in 1879. Selenium transmits an electric current proportional to the amount of light falling on its surface. This phenomenon was used in the design of **light meters** and similar devices. Selenium’s semiconductor properties found numerous other applications in electronics.^{[24][25][26]} The development of **selenium rectifiers** began during the early 1930s, and these replaced **copper oxide** rectifiers because they were more efficient.^{[27][28][29]} These lasted in commercial applications until the 1970s, following which they were replaced with less expensive and even more efficient **silicon rectifiers**.

Selenium came to medical notice later because of its toxicity to **human beings** working in industries. Selenium was also recognized as an important veterinary toxin, which is seen in animals that have eaten high-selenium plants. In 1954, the first hints of specific biological functions of selenium were discovered in **microorganisms**.^{[30][31]} It was discovered to be essential for mammalian life in 1957.^{[32][33]} In the 1970s, it was shown to be present in two independent sets of **enzymes**. This was followed by the discovery of **selenocysteine** in proteins. During the 1980s, selenocysteine was shown to be encoded by the **codon UGA**. The recoding mechanism was worked out first in **bacteria** and then in **mammals** (see **SECIS element**).^[34]

1.4 Occurrence



Native selenium in sandstone, from a uranium mine near Grants, New Mexico

See also: [Category:Selenide minerals](#).

Native (i.e., elemental) selenium is a rare mineral, which does not usually form good crystals, but, when it does, they are steep rhombohedra or tiny acicular (hair-like) crystals.^[35] Isolation of selenium is often complicated by the presence of other compounds and elements.

Selenium occurs naturally in a number of inorganic forms, including selenide, selenate, and selenite, but these minerals are rare. The common mineral selenite is not a selenium mineral, and contains no selenite ion, but is rather a type of gypsum (calcium sulfate hydrate) named like selenium for the moon well before the discovery of selenium. Selenium is most commonly found as an impurity, replacing a small part of the sulfur in sulfide ores of many metals.^{[36][37]}

In living systems, selenium is found in the amino acids selenomethionine, selenocysteine, and methylselenocysteine. In these compounds, selenium plays a role analogous to that of sulfur. Another naturally occurring organoselenium compound is dimethyl selenide.^{[38][39]}

Certain solids are selenium-rich, and selenium can be bioconcentrated by some plants. In soils, selenium most often occurs in soluble forms such as selenate (analogous to sulfate), which are leached into rivers very easily by runoff.^{[36][37]} Ocean water contains significant amounts of selenium.^{[40][41]}

Anthropogenic sources of selenium include coal burning, and the mining and smelting of sulfide ores.^[42]

1.5 Production

Selenium is most commonly produced from selenide in many sulfide ores, such as those of copper, nickel, or lead.

Electrolytic metal refining is particularly productive of selenium as a byproduct, obtained from the anode mud of copper refineries. Another source was the mud from the lead chambers of sulfuric acid plants, a process that is no longer used. Selenium can be refined from these muds by a number of methods. However, most elemental selenium comes as a byproduct of refining copper or producing sulfuric acid.^{[43][44]} Since its invention, solvent extraction and electrowinning (SX/EW) production of copper produces an increasing share of the worldwide copper supply.^[45] This changes the availability of selenium because only a comparably small part of the selenium in the ore is leached with the copper.^[46]

Industrial production of selenium usually involves the extraction of selenium dioxide from residues obtained during the purification of copper. Common production from the residue then begins by oxidation with sodium carbonate to produce selenium dioxide, which is mixed with water and acidified to form selenous acid (oxidation step). Selenous acid is bubbled with sulfur dioxide (reduction step) to give elemental selenium.^{[47][48]}

About 2,000 tonnes of selenium were produced in 2011 worldwide, mostly in Germany (650 t), Japan (630 t), Belgium (200 t), and Russia (140 t), and the total reserves were estimated at 93,000 tonnes. These data exclude two major producers, the United States and China. A previous sharp increase was observed in 2004 from 4–5 to \$27/lb. The price was relatively stable during 2004–2010 at about US\$30 per pound (in 100-pound lots) but increased to \$65 /lb in 2011. The consumption in 2010 was divided as follows: metallurgy – 30%, glass manufacturing – 30%, agriculture – 10%, chemicals and pigments – 10%, and electronics – 10%. China is the dominant consumer of selenium at 1,500–2,000 tonnes/year.^[49]

1.6 Applications

1.6.1 Manganese electrolysis

During the electro winning of manganese, the addition of selenium dioxide decreases the power necessary to operate the electrolysis cells. China is the largest consumer of selenium dioxide for this purpose. For every tonne of manganese, an average 2 kg selenium oxide is used.^{[49][50]}

1.6.2 Glass production

The largest commercial use of Se, accounting for about 50% of consumption, is for the production of glass. Se compounds confer a red color to glass. This color cancels out the green or yellow tints that arise from iron impurities typical for most glass. For this purpose, various selenite and selenate salts are added. For other applications, a red color may be desired, produced by mixtures of CdSe and CdS.^[51]

1.6.3 Alloys

Selenium is used with **bismuth** in **brasses** to replace more toxic **lead**. The regulation of lead in drinking water applications with the **Safe Drinking Water Act** of 1974 made a reduction of lead in brass necessary. The new brass is marketed under the name **EnviroBrass**.^[52] Like lead and sulfur, selenium improves the machinability of steel at concentrations around 0.15%.^{[53][54]} Selenium produces the same machinability improvement in copper alloys.^[55]

Solar cells

Copper indium gallium selenide is a material used in solar cells.^[56]

1.6.4 Other uses

Small amounts of organoselenium compounds are used to modify the **vulcanization** catalysts for the production of rubber.^[46]

The demand for selenium by the electronics industry is declining, despite a number of continuing applications.^[49] Its **photovoltaic** and **photoconductive** properties are still useful in **photocopying**,^{[57][58][59][60]} **photocells**, **light meters** and **solar cells**. Its use as a photoconductor in plain-paper copiers once was a leading application, but in the 1980s, the photoconductor application declined (although it was still a large end-use) as more and more copiers switched to organic photoconductors. Though once widely used, **selenium rectifiers** have mostly been replaced (or are being replaced) by silicon-based devices. The most notable exception is in power **DC surge protection**, where the superior energy capabilities of selenium suppressors make them more desirable than **metal oxide varistors**.

Zinc selenide was the first material for blue **LEDs**, but gallium nitride is dominating the market now.^[61] **Cadmium selenide** was an important component in **quantum dots**. Sheets of amorphous selenium convert **X-ray** images to patterns of charge in **xeroradiography** and in solid-state, flat-panel X-ray cameras.^[62] Ionized selenium (Se+24) is one of the active mediums used in X-ray lasers.^[63]

Selenium is a catalyst in some chemical reactions, but it is not widely used because of issues with toxicity. In **X-ray crystallography**, incorporation of one or more selenium atoms in place of sulfur helps with multiple-wavelength anomalous dispersion and **single wavelength anomalous dispersion phasing**.^[64]

Selenium is used in the **toning of photographic prints**, and it is sold as a toner by numerous photographic manufacturers. Selenium intensifies and extends the tonal range of black-and-white photographic images and improves the permanence of prints.^{[65][66][67]}

⁷⁵Se is used as a gamma source in industrial radiography.^[68]

1.7 Biological role

Main article: **Selenium in biology**

Although it is toxic in large doses, selenium is an essential **micronutrient** for animals. In plants, it occurs as a bystander mineral, sometimes in toxic proportions in **forage** (some plants may accumulate selenium as a defense against being eaten by animals, but other plants, such as **locoweeds**, require selenium, and their growth indicates the presence of selenium in soil).^[4] See more on plant nutrition below.

Selenium is a component of the unusual **amino acids** **selenocysteine** and **selenomethionine**. In humans, selenium is a **trace element** nutrient that functions as **cofactor** for **reduction of antioxidant enzymes**, such as **glutathione peroxidases**^[69] and certain forms of **thioredoxin reductase** found in animals and some plants (this enzyme occurs in all living organisms, but not all forms of it in plants require selenium).

The **glutathione peroxidase** family of enzymes (GSH-Px) catalyze certain reactions that remove reactive oxygen species such as **hydrogen peroxide** and organic **hydroperoxides**:



The thyroid gland and every cell that uses thyroid hormone use selenium, which is a cofactor for the three of the four known types of **thyroid hormone deiodinases**, which activate and then deactivate various thyroid hormones and their metabolites; the **iodothyronine deiodinases** are the subfamily of deiodinase enzymes that use selenium as the otherwise rare amino acid selenocysteine. (Only the deiodinase, **iodotyrosine deiodinase**, which works on the last breakdown products of thyroid hormone, does not use selenium.)^[70]

Selenium may inhibit **Hashimoto's disease**, in which the body's own thyroid cells are attacked as alien. A reduction of 21% on TPO antibodies is reported with the dietary intake of 0.2 mg of selenium.^[71]

Increased dietary selenium reduces the effects of mercury toxicity,^{[72][73][74]} although it is effective only at low to modest doses of mercury.^[75] Evidence suggests that the molecular mechanisms of mercury toxicity includes the irreversible inhibition of selenoenzymes that are required to prevent and reverse oxidative damage in brain and endocrine tissues.^{[76][77]}

1.7.1 Evolution in biology

Main article: [Evolution of dietary antioxidants](#)

From about three billion years ago, prokaryotic selenoprotein families drive the evolution of selenocysteine, an amino acid. Selenium is incorporated into several prokaryotic selenoprotein families in bacteria, archaea, and eukaryotes as selenocysteine,^[78] where selenoprotein peroxiredoxins protect bacterial and eukaryotic cells against oxidative damage. Selenoprotein families of GSH-Px and the deiodinases of eukaryotic cells seem to have a bacterial phylogenetic origin. The selenocysteine-containing form occurs in species as diverse as green algae, diatoms, sea urchin, fish, and chicken. Selenium enzymes are involved in the small reducing molecules [glutathione](#) and [thioredoxin](#). One family of selenium-bearing molecules (the [glutathione peroxidases](#)) destroys peroxide and repairs damaged peroxidized cell membranes, using glutathione. Another selenium-bearing enzyme in some plants and in animals ([thioredoxin reductase](#)) generates reduced thioredoxin, a dithiol that serves as an electron source for peroxidases and also the important reducing enzyme [ribonucleotide reductase](#) that makes DNA precursors from RNA precursors.^[79]

Trace elements involved in GSH-Px and superoxide dismutase enzymes activities, i.e. selenium, [vanadium](#), [magnesium](#), [copper](#), and [zinc](#), may have been lacking in some terrestrial mineral-deficient areas.^[78] Marine organisms retained and sometimes expanded their selenoproteomes, whereas the selenoproteomes of some terrestrial organisms were reduced or completely lost. These findings suggest that, with the exception of [vertebrates](#), aquatic life supports selenium use, whereas terrestrial habitats lead to reduced use of this trace element.^[80] Marine fishes and vertebrate thyroid glands have the highest concentration of selenium and iodine. From about 500 million years ago, freshwater and terrestrial plants slowly optimized the production of “new” endogenous antioxidants such as [ascorbic acid](#) (vitamin C), [polyphenols](#) (including flavonoids), [tocopherols](#), etc. A few of these appeared more recently, in the last 50–200 million years, in fruits and flowers of angiosperm plants. In fact, the angiosperms (the dominant type of plant today) and most of their antioxidant pigments evolved during the late [Jurassic](#) period.

The deiodinase [isoenzymes](#) constitute another family of eukaryotic selenoproteins with identified enzyme function. Deiodinases are able to extract electrons from iodides, and iodides from iodothyronines. They are, thus, involved in thyroid-hormone regulation, participating in the protection of [thyrocytes](#) from damage by H_2O_2 produced for thyroid-hormone biosynthesis.^[81] About 200 million years ago, new selenoproteins were developed as mammalian GSH-Px enzymes.^{[82][83][84][85]}

1.7.2 Nutritional sources of selenium

Dietary selenium comes from nuts, cereals and mushrooms. [Brazil nuts](#) are the richest dietary source (though this is soil-dependent, since the Brazil nut does not require high levels of the element for its own needs).^{[86][87]}

Recommended Dietary Allowance ~ 55 $\mu\text{g/day}$. Selenium as a dietary supplement is available in many forms, including multi-vitamins/mineral supplements - typically 20 $\mu\text{g/day}$. Selenium-specific supplements may have ~200 $\mu\text{g/day}$.

In June 2015 the U.S. [Food and Drug Administration](#) (FDA) published its final rule establishing the requirement of minimum and maximum levels of selenium in [infant formula](#).^[88]

The selenium content in the human body is believed to be in the 13–20 milligram range.^[89]

1.7.3 Indicator plant species

Certain species of plants are considered indicators of high selenium content of the soil because they require high levels of selenium to thrive. The main selenium indicator plants are *Astragalus* species (including some [locoweeds](#)), prince's plume (*Stanleya* sp.), woody asters (*Xylorhiza* sp.), and false goldenweed (*Oenopsis* sp.)^[90]

1.7.4 Medical use

The substance loosely called [selenium sulfide](#) (approximate formula, SeS_2) is the active ingredient in some anti-dandruff shampoos.^[91] The selenium compound kills the scalp fungus *Malassezia*, which causes shedding of dry skin fragments. The ingredient is also used in body lotions to treat [tinea versicolor](#) due to infection by a different species of *Malassezia* fungus.^[92]

1.7.5 Detection in biological fluids

Selenium may be measured in blood, plasma, serum, or urine to monitor excessive environmental or occupational exposure, to confirm a diagnosis of poisoning in hospitalized victims, or investigate a suspected case of fatal overdose. Some analytical techniques are capable of distinguishing organic from inorganic forms of the element. Both organic and inorganic forms of selenium are largely converted to monosaccharide conjugates (selenosugars) in the body prior elimination in the urine. Cancer patients receiving daily oral doses of selenothionine may achieve very high plasma and urine selenium concentrations.^[93]

1.7.6 Toxicity


Although selenium is an essential **trace element**, it is toxic if taken in excess. Exceeding the **Tolerable Upper Intake Level** of 400 micrograms per day can lead to selenosis.^[94] This 400 μg Tolerable Upper Intake Level is based primarily on a 1986 study of five Chinese patients who exhibited overt signs of selenosis and a follow up study on the same five people in 1992.^[95] The 1992 study actually found the maximum safe dietary Se intake to be approximately 800 micrograms per day (15 micrograms per kilogram body weight), but suggested 400 micrograms per day to avoid creating an imbalance of nutrients in the diet and to accord with data from other countries.^[96] In China, people who ingested corn grown in extremely selenium-rich stony coal (carbonaceous **shale**) have suffered from selenium toxicity. This coal was shown to have selenium content as high as 9.1%, the highest concentration in coal ever recorded.^[97]

Signs and symptoms of selenosis include a garlic odor on the breath, gastrointestinal disorders, hair loss, **sloughing** of nails, fatigue, irritability, and neurological damage. Extreme cases of selenosis can exhibit **cirrhosis** of the liver, **pulmonary edema**, or death.^[98] Elemental selenium and most metallic **selenides** have relatively low toxicities because of low **bioavailability**. By contrast, **selenates** and **selenites** have an oxidant mode of action similar to that of arsenic trioxide and are very toxic. The chronic toxic dose of selenite for humans is about 2400 to 3000 micrograms of selenium per day.^[99] **Hydrogen selenide** is an extremely toxic, corrosive gas.^[100] Selenium also occurs in organic compounds, such as dimethyl selenide, **selenomethionine**, **selenocysteine** and **methylselenocysteine**, all of which have high **bioavailability** and are toxic in large doses.

On 19 April 2009, 21 **polo** ponies died shortly before a match in the United States Polo Open. Three days later, a pharmacy released a statement explaining that the horses had received an incorrect dose of one of the ingredients used in a vitamin/mineral supplement compound that had been incorrectly compounded by a **compounding pharmacy**. Analysis of blood levels of **inorganic compounds** in the supplement indicated the selenium concentrations were ten to fifteen times higher than normal in the **blood samples**, and 15 to 20 times higher than normal in the liver samples. Selenium was later confirmed to be the toxic factor.^[101]

Selenium poisoning of water systems may result whenever new agricultural runoff courses through normally dry, undeveloped lands. This process leaches natural soluble selenium compounds (such as selenates) into the water, which may then be concentrated in new “wetlands” as the water evaporates. Selenium pollution of waterways also occurs when selenium is leached from coal flue ash, mining and metal smelting, crude oil processing, and landfill.^[102] The resultant high selenium levels in waterways were found to cause congenital disor-

ders in oviparous species, including wetland birds^[103] and fish.^[104] Elevated dietary methylmercury levels can amplify the harm of selenium toxicity in oviparous species.^{[105][106]}



Relationship between survival of juvenile salmon and concentration of selenium in their tissues after 90 days (Chinook salmon^[107]) or 45 days (Atlantic salmon^[108]) exposure to dietary selenium. The 10% lethality level ($\text{LC}_{10} = 1.84 \mu\text{g/g}$) was derived by applying the biphasic model of Brain and Cousens^[109] to only the Chinook salmon data. The Chinook salmon data comprise two series of dietary treatments, combined here because the effects on survival are indistinguishable.

In fish and other wildlife, selenium is necessary for life, but toxic in high doses. For salmon, the optimal concentration of selenium is about 1 microgram selenium per gram of whole body weight. Much below that level, young salmon die from deficiency;^[108] much above, they die from toxic excess.^[107]

The Occupational Safety and Health Administration (OSHA) has set the legal limit (Permissible exposure limit) for selenium in the workplace at 0.2 mg/m^3 over an 8-hour workday. The National Institute for Occupational Safety and Health (NIOSH) has set a Recommended exposure limit (REL) of 0.2 mg/m^3 over an 8-hour workday. At levels of 1 mg/m^3 , selenium is immediately dangerous to life and health.^[110]

1.7.7 Deficiency

Main article: **Selenium deficiency**

Selenium deficiency is rare in otherwise healthy, well-nourished individuals. It can occur in patients with severely compromised **intestinal function**, those undergoing **total parenteral nutrition**, and^[111] in those of advanced age (over 90). Also, people dependent on food grown from selenium-deficient soil are at risk. Although New Zealand soil has low levels of selenium, adverse health effects have not been detected in the residents.^[112]

Selenium deficiency, defined by low (<60% of normal) selenoenzyme activity levels in brain and endocrine tissues, occurs only when a low selenium level is linked

with an additional stress, such as high exposures to mercury^[113] or increased oxidant stress from vitamin E deficiency.^[114]

Selenium interacts with other nutrients, such as iodine and vitamin E. The effect of selenium deficiency on health remains uncertain, particularly in relation to Kashin-Beck disease.^[115] Also, selenium interacts with other minerals, such as zinc and copper. High doses of Se supplements in pregnant animals might disturb the Zn:Cu ratio and lead to Zn reduction; in such treatment cases, Zn levels should be monitored. Further studies are needed to confirm these interactions.^[116]

In the regions (e.g. various regions within North America) where low selenium soil levels lead to low concentrations in the plants, some animal species may be deficient unless selenium is supplemented with diet or injection.^[117] Ruminants are particularly susceptible. In general, absorption of dietary selenium is lower in ruminants than other animals, and is lower from forages than from grain.^[118] Ruminants grazing certain forages, e.g., some white clover varieties containing cyanogenic glycosides, may have higher selenium requirements,^[118] presumably because cyanide is released from the aglycone by glucosidase activity in the rumen^[119] and glutathione peroxidases is deactivated by the cyanide acting on the glutathione moiety.^[120] Neonate ruminants at risk of WMD (white muscle disease) may be administered both selenium and vitamin E by injection; some of the WMD myopathies respond only to selenium, some only to vitamin E, and some to either.^[121]

1.7.8 Controversial health effects

Main article: [Selenium in biology](#)

A number of correlative epidemiological studies have implicated selenium deficiency (measured by blood levels) in a number of serious or chronic diseases, such as cancer,^{[122][123]} diabetes,^[122] HIV/AIDS,^[124] and tuberculosis. In addition, selenium supplementation has been found to be a chemopreventive for some types of cancer in some types of rodents. However, in randomized, blinded, controlled prospective trials in humans, selenium supplementation has not succeeded in reducing the incidence of any disease, nor has a meta-analysis of such selenium supplementation studies detected a decrease in overall mortality.^[125]

1.8 See also

- [ACES \(nutritional supplement\)](#)
- [Abundance of elements in Earth's crust](#)
- [Selenium yeast](#)

- [Selene](#)

1.9 References

- [1] Standard Atomic Weights 2013. Commission on Isotopic Abundances and Atomic Weights
- [2] Greenwood, Norman N.; Earnshaw, Alan (1997). *Chemistry of the Elements* (2nd ed.). Butterworth-Heinemann. ISBN 0-08-037941-9.
- [3] Magnetic susceptibility of the elements and inorganic compounds, in Lide, D. R., ed. (2005). *CRC Handbook of Chemistry and Physics* (86th ed.). Boca Raton (FL): CRC Press. ISBN 0-8493-0486-5.
- [4] Ruyle, George. "Poisonous Plants on Arizona Rangelands" (PDF). The University of Arizona. Retrieved 2009-01-05.
- [5] House, James E. (2008). *Inorganic chemistry*. Academic Press. p. 524. ISBN 0-12-356786-6.
- [6] Greenwood, Norman N.; Earnshaw, Alan (1997). *Chemistry of the Elements* (2nd ed.). Butterworth-Heinemann. pp. 751–752. ISBN 0-08-037941-9.
- [7] Video of selenium heating on YouTube
- [8] "The half-life of ⁷⁹Se". Physikalisch-Technische Bundesanstalt. 2010-09-23. Retrieved 2012-05-29.
- [9] Jörg, Gerhard; Bühnenmann, Rolf; Hollas, Simon; et al. (2010). "Preparation of radiochemically pure ⁷⁹Se and highly precise determination of its half-life". *Applied Radiation and Isotopes*. **68** (12): 2339–2351. doi:10.1016/j.apradiso.2010.05.006. PMID 20627600.
- [10] Audi, Georges; Bersillon, O.; Blachot, J.; Wapstra, A.H. (2003). "The NUBASE Evaluation of Nuclear and Decay Properties". *Nuclear Physics A. Atomic Mass Data Center*. **729**: 3–128. Bibcode:2003NuPhA.729....3A. doi:10.1016/j.nuclphysa.2003.11.001.
- [11] Wiberg, Egon; Wiberg, Nils; Holleman, Arnold Frederick (2001). *Inorganic chemistry*. San Diego: Academic Press. p. 583. ISBN 0-12-352651-5.
- [12] Greenwood, Norman N.; Earnshaw, Alan (1997). *Chemistry of the Elements* (2nd ed.). Butterworth-Heinemann. p. 780. ISBN 0-08-037941-9.
- [13] Seppelt, K.; Desmarteau, Darryl D. (1980). "Selenoyl difluoride". *Inorganic Syntheses*. Inorganic Syntheses. **20**: 36–38. doi:10.1002/9780470132517.ch9. ISBN 0-471-07715-1. The report describes the synthesis of selenic acid.
- [14] Lenher, V. (April 1902). "Action of selenic acid on gold". *Journal of the American Chemical Society*. **24** (4): 354–355. doi:10.1021/ja02018a005.
- [15] Xu, Zhengtao (2007). Devillanova, Francesco A., ed. *Handbook of chalcogen chemistry: new perspectives in sulfur, selenium and tellurium*. Royal Society of Chemistry. p. 460. ISBN 0-85404-366-7.

- [16] Proctor, Nick H.; Hathaway, Gloria J. (2004). James P. Hughes, ed. *Proctor and Hughes' chemical hazards of the workplace* (5th ed.). Wiley-IEEE. p. 625. ISBN 0-471-26883-6.
- [17] Woollins, Derek; Kelly, Paul F. (1993). "The Reactivity of Se_4N_4 in Liquid Ammonia". *Polyhedron*. **12** (10): 1129–1133. doi:10.1016/S0277-5387(00)88201-7.
- [18] Kelly, P. F.; Slawin, A. M. Z.; Soriano-Rama, A. (1997). "Use of Se_4N_4 and $\text{Se}(\text{NSO})_2$ in the preparation of palladium adducts of diselenium dinitride, Se_2N_2 ; crystal structure of $[\text{PPh}_4][\text{Pd}(\text{Se}_2\text{N}_2)_2]$ ". *Dalton Transactions* (4): 559–562. doi:10.1039/a606311j.
- [19] Siivari, Jari; Chivers, Tristram; Laitinen, Risto S. (1993). "A simple, efficient synthesis of tetraselenium tetranitride". *Inorganic Chemistry*. **32** (8): 1519–1520. doi:10.1021/ic00060a031.
- [20] Erker, G.; Hock, R.; Krüger, C.; Werner, S.; Klärner, F. G.; Artschwager-Perl, U. (1990). "Synthesis and Cycloadditions of Monomeric Selenobenzophenone". *Angewandte Chemie International Edition in English*. **29** (9): 1067–1068. doi:10.1002/anie.199010671.
- [21] Berzelius, J. J. (1818) "Lettre de M. Berzelius à M. Berthollet sur deux métaux nouveaux" (Letter from Mr. Berzelius to Mr. Berthollet on two new metals), *Annales de chimie et de physique*, series 2, vol. 7, pp. 199–206.
- [22] Weeks, Mary Elvira (1932). "The discovery of the elements. VI. Tellurium and selenium". *Journal of Chemical Education*. **9** (3): 474. Bibcode:1932JChEd...9..474W. doi:10.1021/ed009p474.
- [23] Trofast, Jan (2011). "Berzelius' Discovery of Selenium". *Chemistry International*. **33** (5): 16–19. PDF
- [24] "Action of light on selenium". *Popular Science*. **10** (1): 116. 1876.
- [25] Levinshtein, M. E; Simin, G. S (1992-12-01). *Earliest semiconductor device*. Getting to Know Semiconductors. pp. 77–79. ISBN 978-981-02-3516-1.
- [26] Winston, Brian (1998-05-29). *Media Technology and Society: A History: From the Telegraph to the Internet*. p. 89. ISBN 978-0-415-14229-8.
- [27] Morris, Peter Robin (1990). *A History of the World Semiconductor Industry*. p. 18. ISBN 978-0-86341-227-1.
- [28] Bergmann, Ludwig (1931). "Über eine neue Selen-Sperrschicht-Photozelle". *Physikalische Zeitschrift*. **32**: 286–288.
- [29] Waitkins, G. R.; Bearse, A. E.; Shutt, R. (1942). "Industrial Utilization of Selenium and Tellurium". *Industrial & Engineering Chemistry*. **34** (8): 899–910. doi:10.1021/ie50392a002.
- [30] Pinsent, Jane (1954). "The need for selenite and molybdate in the formation of formic dehydrogenase by members of the Coli-aerogenes group of bacteria". *Biochem J*. **57** (1): 10–16. PMC 1269698. PMID 13159942.
- [31] Stadtman, Thressa C. (2002). "Some Functions of the Essential Trace Element, Selenium". *Trace Elements in Man and Animals* 10: 831–836. doi:10.1007/0-306-47466-2_267. ISBN 0-306-46378-4.
- [32] Schwarz, Klaus; Foltz, Calvin M. (1957). "Selenium as an Integral Part of Factor 3 Against Dietary Necrotic Liver Degeneration". *Journal of the American Chemical Society*. **79** (12): 3292–3293. doi:10.1021/ja01569a087.
- [33] Oldfield, James E. (2006). "Selenium: A historical perspective". *Selenium*: 1–6. doi:10.1007/0-387-33827-6_1. ISBN 978-0-387-33826-2.
- [34] Hatfield, D. L.; Gladyshev, V. N. (2002). "How Selenium Has Altered Our Understanding of the Genetic Code". *Molecular and Cellular Biology*. **22** (11): 3565–3576. doi:10.1128/MCB.22.11.3565-3576.2002. PMC 133838. PMID 11997494.
- [35] "Native Selenium". Webminerals. Retrieved 2009-06-06.
- [36] Kabata-Pendias, A. (1998). "Geochemistry of selenium". *Journal of environmental pathology, toxicology and oncology: official organ of the International Society for Environmental Toxicology and Cancer*. **17** (3–4): 173–177. PMID 9726787.
- [37] Fordyce, Fiona (2007). "Selenium Geochemistry and Health". *AMBIO: A Journal of the Human Environment*. **36**: 94–97. doi:10.1579/0044-7447(2007)36[94:SGAH]2.0.CO;2.
- [38] Wessjohann, Ludger A.; Schneider, Alex; Abbas, Muhammad; Brandt, Wolfgang (2007). "Selenium in chemistry and biochemistry in comparison to sulfur". *Biological Chemistry*. **388** (10): 997–1006. doi:10.1515/BC.2007.138. PMID 17937613.
- [39] Birringer, Marc; Pilawa, Sandra; Flohé, Leopold (2002). "Trends in selenium biochemistry". *Natural Product Reports*. **19** (6): 693–718. doi:10.1039/B205802M. PMID 12521265.
- [40] Amouroux, David; Liss, Peter S; Tessier, Emmanuel; et al. (2001). "Role of oceans as biogenic sources of selenium". *Earth and Planetary Science Letters*. **189** (3–4): 277–283. Bibcode:2001E&PSL.189..277A. doi:10.1016/S0012-821X(01)00370-3.
- [41] Haug, Anna; Graham, Robin D.; Christophersen, Olav A.; Lyons, Graham H. (2007). "How to use the world's scarce selenium resources efficiently to increase the selenium concentration in food". *Microbial Ecology in Health and Disease*. **19** (4): 209–228. doi:10.1080/08910600701698986. PMC 2556185. PMID 18833333.
- [42] "Public Health Statement: Selenium" (PDF). Agency for Toxic Substances and Disease Registry. Retrieved 2009-01-05.

- [43] “Public Health Statement: Selenium – Production, Import/Export, Use, and Disposal” (PDF). Agency for Toxic Substances and Disease Registry. Retrieved 2009-01-05.
- [44] “Chemistry: Periodic Table: selenium: key information”. webelements. Retrieved 2009-01-06.
- [45] Bartos, P.J. (2002). “SX-EW copper and the technology cycle”. *Resources Policy*. **28** (3–4): 85–94. doi:10.1016/S0301-4207(03)00025-4.
- [46] Naumov, A. V. (2010). “Selenium and tellurium: State of the markets, the crisis, and its consequences”. *Metallurgist*. **54** (3–4): 197–200. doi:10.1007/s11015-010-9280-7.
- [47] Hoffmann, James E. (1989). “Recovering selenium and tellurium from copper refinery slimes”. *JOM*. **41** (7): 33–38. Bibcode:1989JOM....41g..33H. doi:10.1007/BF03220269.
- [48] Hyvärinen, Olli; Lindroos, Leo; Yllö, Erkki (1989). “Recovering selenium from copper refinery slimes”. *JOM*. **41** (7): 42–43. Bibcode:1989JOM....41g..42H. doi:10.1007/BF03220271.
- [49] “Selenium and Tellurium: Statistics and Information”. United States Geological Survey. Retrieved 2012-05-30.
- [50] Sun, Yan; Tian, Xike; He, Binbin; et al. (2011). “Studies of the reduction mechanism of selenium dioxide and its impact on the microstructure of manganese electrodeposit”. *Electrochimica Acta*. **56** (24): 8305–8310. doi:10.1016/j.electacta.2011.06.111.
- [51] Bernd E. Langner “Selenium and Selenium Compounds” in Ullmann’s Encyclopedia of Industrial Chemistry, 2005, Wiley-VCH, Weinheim. doi:10.1002/14356007.a23_525.
- [52] Davis, Joseph R (2001). *Copper and Copper Alloys*. ASM Int. p. 91. ISBN 978-0-87170-726-0.
- [53] Isakov, Edmund (2008-10-31). *Cutting Data for Turning of Steel*. p. 67. ISBN 978-0-8311-3314-6.
- [54] Gol’Dshtein, Ya. E.; Mushtakova, T. L.; Komissarova, T. A. (1979). “Effect of selenium on the structure and properties of structural steel”. *Metal Science and Heat Treatment*. **21** (10): 741–746. doi:10.1007/BF00708374.
- [55] Davis, Joseph R. (2001). *Copper and Copper Alloys*. ASM International. p. 278. ISBN 978-0-87170-726-0.
- [56] Deutsche Gesellschaft für Sonnenenergie (2008). “Copper indium diselenide (CIS) cell”. *Planning and Installing Photovoltaic Systems: A Guide for Installers, Architects and Engineers*. Earthscan. pp. 43–44. ISBN 978-1-84407-442-6.
- [57] Springett, B. E. (1988). “Application of Selenium-Tellurium Photoconductors to the Xerographic Copying and Printing Processes”. *Phosphorus and Sulfur and the Related Elements*. **38** (3–4): 341–350. doi:10.1080/03086648808079729.
- [58] Williams, Rob (2006). *Computer Systems Architecture: A Networking Approach*. Prentice Hall. pp. 547–548. ISBN 978-0-321-34079-5.
- [59] Diels, Jean-Claude; Arissian, Ladan (2011). “The Laser Printer”. *Lasers*. Wiley-VCH. pp. 81–83. ISBN 978-3-527-64005-8.
- [60] Meller, Gregor & Grasser, Tibor (2009). *Organic Electronics*. Springer. pp. 3–5. ISBN 978-3-642-04537-0.
- [61] Normile, Dennis (2000). “The birth of the Blues”. *Popular Science*. p. 57.
- [62] Kasap, Safa; Frey, Joel B.; Belev, George; et al. (2009). “Amorphous selenium and its alloys from early xeroradiography to high resolution X-ray image detectors and ultrasensitive imaging tubes”. *Physica status solidi (b)*. **246** (8): 1794–1805. Bibcode:2009PSSBR.246.1794K. doi:10.1002/pssb.200982007.
- [63] Svelto, Orazio (1998). *Principles of LASERs fourth ed.* Plenum. p. 457. ISBN 0-306-45748-2.
- [64] Hai-Fu, F.; Woolfson, M. M.; Jia-Xing, Y. (1993). “New Techniques of Applying Multi-Wavelength Anomalous Scattering Data”. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*. **442** (1914): 13–32. Bibcode:1993RSPSA.442...13H. doi:10.1098/rspa.1993.0087.
- [65] MacLean, Marion E. (1937). “A project for general chemistry students: Color toning of photographic prints”. *Journal of Chemical Education*. **14**: 31. Bibcode:1937JChEd..14...31M. doi:10.1021/ed014p31.
- [66] Penichon, Sylvie (1999). “Differences in Image Tonality Produced by Different Toning Protocols for Matte Collodion Photographs”. *Journal of the American Institute for Conservation*. **38** (2): 124–143. doi:10.2307/3180042. JSTOR 3180042.
- [67] McKenzie, Joy (2003). *Exploring Basic Black & White Photography*. Delmar. p. 176. ISBN 978-1-4018-1556-1.
- [68] Hayward, Peter; Currie, Dean. “Radiography of Welds Using Selenium 75, Ir 192 and X-rays” (PDF).
- [69] Linus Pauling Institute at Oregon State University lpi.oregonstate.edu
- [70] “Selenium”. Linus Pauling Institute at Oregon State University. Retrieved 2009-01-05.
- [71] Mazokopakis, EE; Papadakis, JA; Papadomanolaki, MG; et al. (2007). “Effects of 12 months treatment with L-selenomethionine on serum anti-TPO Levels in Patients with Hashimoto’s thyroiditis”. *Thyroid*. **17** (7): 609–612. doi:10.1089/thy.2007.0040. PMID 17696828.
- [72] Ralston, NV; Ralston, CR; Blackwell, JL III; Raymond, LJ (2008). “Dietary and tissue selenium in relation to methylmercury toxicity” (PDF). *Neurotoxicology*. **29** (5): 802–811. doi:10.1016/j.neuro.2008.07.007. PMID 18761370.
- [73] Penglase, S; Hamre, K; Ellingsen, S (2014). “Selenium prevents downregulation of antioxidant selenoprotein genes by methylmercury”. *Free Radical Biology and Medicine*. **75**: 95–104. doi:10.1016/j.freeradbiomed.2014.07.019. PMID 25064324.

- [74] Usuki, F; Yamashita, A; Fujimura, M (2011). "Post-transcriptional defects of antioxidant selenoenzymes cause oxidative stress under methylmercury exposure". *The Journal of Biological Chemistry*. **286** (8): 6641–6649. doi:10.1074/jbc.M110.168872.
- [75] Ohi, G; Seki, H; Maeda, H; Yagyu, H (1975). "Protective effect of selenite against methylmercury toxicity: observations concerning time, dose and route factors in the development of selenium attenuation". *Industrial Health*. **13**: 93–99. doi:10.2486/indhealth.13.93.
- [76] Ralston, NVC; Raymond, LJ (2010). "Dietary selenium's protective effects against methylmercury toxicity". *Toxicology*. **278** (1): 112–123. doi:10.1016/j.tox.2010.06.004. PMID 20561558.
- [77] Carvalho, CML; Chew, Hashemy SI; Hashemy, J; et al. (2008). "Inhibition of the human thioredoxin system: A molecular mechanism of mercury toxicity". *Journal of Biological Chemistry*. **283** (18): 11913–11923. doi:10.1074/jbc.M710133200. PMID 18321861.
- [78] Gladyshev, Vadim N.; Hatfield, Dolph L. (1999). "Selenocysteine-containing proteins in mammals". *Journal of Biomedical Science*. **6** (3): 151–160. doi:10.1007/BF02255899. PMID 10343164.
- [79] Stadtman, TC (1996). "Selenocysteine". *Annual Review of Biochemistry*. **65**: 83–100. doi:10.1146/annurev.bi.65.070196.000503. PMID 8811175.
- [80] Lobanov, Alexey V; Fomenko, Dmitri E; Zhang, Yan; et al. (2007). "Evolutionary dynamics of eukaryotic selenoproteomes: large selenoproteomes may associate with aquatic life and small with terrestrial life". *Genome Biology*. **8** (9): R198. doi:10.1186/gb-2007-8-9-r198. PMC 2375036. PMID 17880704.
- [81] Venturi, Sebastiano; Venturi, Mattia (2007), "Evolution of Dietary Antioxidant Defences" (PDF), *European Epi-Marker*, **11** (3), pp. 1–11
- [82] Castellano, Sergi; Novoselov, Sergey V; Kryukov, Gregory V; et al. (2004). "Reconsidering the evolution of eukaryotic selenoproteins: a novel nonmammalian family with scattered phylogenetic distribution". *EMBO Reports*. **5** (1): 71–7. doi:10.1038/sj.embor.7400036. PMC 1298953. PMID 14710190.
- [83] Kryukov, Gregory V; Gladyshev, Vadim N (2004). "The prokaryotic selenoproteome". *EMBO Reports*. **5** (5): 538–43. doi:10.1038/sj.embor.7400126. PMC 1299047. PMID 15105824.
- [84] Wilting, R.; Schorling, S.; Persson, B.C.; Böck, A. (1997). "Selenoprotein synthesis in archaea: identification of an mRNA element of *Methanococcus jannaschii* probably directing selenocysteine insertion". *Journal of Molecular Biology*. **266** (4): 637–41. doi:10.1006/jmbi.1996.0812. PMID 9102456.
- [85] Zhang, Yan; Fomenko, Dmitri E; Gladyshev, Vadim N (2005). "The microbial selenoproteome of the Sargasso Sea". *Genome Biology*. **6** (4): R37. doi:10.1186/gb-2005-6-4-r37. PMC 1088965. PMID 15833124.
- [86] Barclay, Margaret N. I.; MacPherson, Allan; Dixon, James (1995). "Selenium content of a range of UK food". *Journal of food composition and analysis*. **8** (4): 307–318. doi:10.1006/jfca.1995.1025.
- [87] A list of selenium-rich foods can be found on The Office of Dietary Supplements Selenium Fact Sheet.
- [88] "FDA Issues Final Rule to Add Selenium to List of Required Nutrients for Infant Formula". *www.fda.gov*. Retrieved 2015-09-10.
- [89] A common reference for this is Schroeder, HA; Frost, DV; Balassa, JJ (1970). "Essential trace metals in man: Selenium". *Journal of chronic diseases*. **23** (4): 227–43. doi:10.1016/0021-9681(70)90003-2. PMID 4926392.
- [90] Zane Davis, T. (2008-03-27). "Selenium in Plants" (PDF). p. 8. Retrieved 2008-12-05.
- [91] "Selenium(IV)_sulfide". Pharmacy Codes. Retrieved 2009-01-06.
- [92] "Selenium sulfide". DermNet NZ. Retrieved 2009-01-06.
- [93] Baselt, R. (2008). *Disposition of Toxic Drugs and Chemicals in Man* (8th ed.). Foster City, CA: Biomedical Publications. pp. 1416–1420. ISBN 978-0-9626523-5-6.
- [94] "Dietary Supplement Fact Sheet: Selenium". National Institutes of Health; Office of Dietary Supplements. Retrieved 2009-01-05.
- [95] Panel on Dietary Antioxidants and Related Compounds, Subcommittees on Upper Reference Levels of Nutrients and Interpretation and Uses of DRIs, Standing Committee on the Scientific Evaluation of Dietary Reference Intakes, Food and Nutrition Board, Institute of Medicine (August 15, 2000). *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids*. Institute of Medicine. pp. 314–315. ISBN 0-309-06949-1.
- [96] Yang, G.; Zhou, R. (1994). "Further Observations on the Human Maximum Safe Dietary Selenium Intake in a Seleniferous Area of China". *Journal of trace elements and electrolytes in health and disease*. **8** (3–4): 159–165. PMID 7599506.
- [97] Yang, Guang-Qi; Xia, Yi-Ming (1995). "Studies on Human Dietary Requirements and Safe Range of Dietary Intakes of Selenium in China and Their Application in the Prevention of Related Endemic Diseases". *Biomedical and Environmental Sciences*. **8** (3): 187–201. PMID 8561918.
- [98] "Public Health Statement: Health Effects" (PDF). Agency for Toxic Substances and Disease Registry. Retrieved 2009-01-05.
- [99] Wilber, C. G. (1980). "Toxicology of selenium". *Clinical Toxicology*. **17** (2): 171–230. doi:10.3109/15563658008985076. PMID 6998645.
- [100] Olson, O.E. (1986). "Selenium Toxicity in Animals with Emphasis on Man". *International Journal of Toxicology*. **5**: 45–70. doi:10.3109/10915818609140736.

- [101] “Polo pony selenium levels up to 20 times higher than normal”. 2009-05-06. Retrieved 2009-05-05.
- [102] Lemly, D (2004). “Aquatic selenium pollution is a global environmental safety issue”. *Ecotoxicology and Environmental Safety*. **59**: 44–56. doi:10.1016/S0147-6513(03)00095-2.
- [103] Ohlendorf, H. M. (2003). *Ecotoxicology of selenium*. Handbook of ecotoxicology. Boca Raton: Lewis Publishers. pp. 466–491. ISBN 978-1-56670-546-2.
- [104] Lemly, A.D (1997). “A teratogenic deformity index for evaluating impacts of selenium on fish populations”. *Ecotoxicology and Environmental Safety*. **37** (3): 259–266. doi:10.1006/eesa.1997.1554.
- [105] Penglase, S; Hamre, K; Ellingsen, S (2014). “Selenium and mercury have a synergistic negative effect on fish reproduction”. *Aquatic Toxicology*. **149**: 16–24. doi:10.1016/j.aquatox.2014.01.020.
- [106] Heinz, GH; Hoffman, DJ (1998). “Methylmercury chloride and selenomethionine interactions on health and reproduction in mallards”. *Environmental Toxicology and Chemistry*. **17** (2): 139–145. doi:10.1002/etc.5620170202.
- [107] Hamilton, Steven J.; Buhl, Kevin J.; Faerber, Neil L.; et al. (1990). “Toxicity of organic selenium in the diet to chinook salmon”. *Environ. Toxicol. Chem.* **9** (3): 347–358. doi:10.1002/etc.5620090310.
- [108] Poston, H. A.; Combs Jr., GF; Leibovitz, L (1976). “Vitamin E and selenium interrelations in the diet of Atlantic salmon (*Salmo salar*): gross, histological and biochemical signs”. *Journal of Nutrition*. **106** (7): 892–904. PMID 932827.
- [109] Brain, P.; Cousens, R. (1989). “Weed Research”. *Weed Research*. **29** (2): 93–96. doi:10.1111/j.1365-3180.1989.tb00845.x.
- [110] “CDC - NIOSH Pocket Guide to Chemical Hazards - Selenium”. www.cdc.gov. Retrieved 2015-11-21.
- [111] Ravaglia, G; Forti, P; Maioli, F; et al. (2000). “Effect of micronutrient status on natural killer cell immune function in healthy free-living subjects aged ≥ 90 y1”. *American Journal of Clinical Nutrition*. **71** (2): 590–598. PMID 10648276.
- [112] MedSafe Editorial Team. “Selenium”. *Prescriber Update Articles*. New Zealand Medicines and Medical Devices Safety Authority. Retrieved 2009-07-13.
- [113] Ralston, N.V.C.; Raymond, L.J. (2010). “Dietary selenium’s protective effects against methylmercury toxicity”. *Toxicology*. **278** (1): 112–123. doi:10.1016/j.tox.2010.06.004. PMID 20561558.
- [114] Mann, Jim; Truswell, A. Stewart (2002). *Essentials of Human Nutrition* (2nd ed.). Oxford University Press. ISBN 978-0-19-262756-8.
- [115] Moreno-Reyes, R; Mathieu, F; Boelaert, M; et al. (2003). “Selenium and iodine supplementation of rural Tibetan children affected by Kashin-Beck osteoarthropathy”. *American Journal of Clinical Nutrition*. **78** (1): 137–144. PMID 12816783.
- [116] Kachuee, R; Moeini, M; Suori, M (2013). “The effect of dietary organic and inorganic selenium supplementation on serum Se, Cu, Fe and Zn status during the late pregnancy in Merghoz goats and their kids”. *Small Ruminant Research*. **110** (1): 20–27. doi:10.1016/j.smallrumres.2012.08.010.
- [117] National Research Council, Subcommittee on Sheep Nutrition (1985). *Nutrient requirements of sheep*. 6th ed., National Academy Press, Washington, ISBN 0309035961.
- [118] National Research Council, Committee on Nutrient Requirements of Small Ruminants (2007). *Nutrient requirements of small ruminants*. National Academies Press, Washington, ISBN 0-309-10213-8.
- [119] Coop, I. E.; Blakely, R. L. (1949). “The metabolism and toxicity of cyanides and cyanogenic glycosides in sheep”. *N. Z. J. Sci. Technol.* **30**: 277–291.
- [120] Kraus, R. J.; Prohaska, J. R.; Ganther, H. E. (1980). “Oxidized forms of ovine erythrocyte glutathione peroxidase. Cyanide inhibition of 4-glutathione: 4-selenoenzyme”. *Biochem. Biophys. Acta*. **615** (1): 19–26. doi:10.1016/0005-2744(80)90004-2. PMID 7426660.
- [121] Kahn, C. M. (ed.) (2005). *Merck veterinary manual*. 9th ed. Merck & Co., Inc., Whitehouse Station, ISBN 0911910506.
- [122] Ip, C (1998). “Lessons from basic research in selenium and cancer prevention” (PDF). *The Journal of Nutrition*. **128** (11): 1845–54. PMID 9808633.
- [123] Amaral, A. F. S.; Cantor, K. P.; Silverman, D. T.; Malats, N. (31 August 2010). “Selenium and Bladder Cancer Risk: a Meta-analysis”. *Cancer Epidemiology Biomarkers & Prevention*. **19** (9): 2407–2415. doi:10.1158/1055-9965.EPI-10-0544.
- [124] Rayman, Margaret P. (2000). “The importance of selenium to human health”. *The Lancet*. **356** (9225): 233–41. doi:10.1016/S0140-6736(00)02490-9. PMID 10963212.
- [125] Bjelakovic, G; Nikolova, D; Gluud, LL; et al. (2012). Bjelakovic, Goran, ed. “Antioxidant supplements for prevention of mortality in healthy participants and patients with various diseases”. *Cochrane database of systematic reviews (Online)*. **3**: CD007176. doi:10.1002/14651858.CD007176.pub2. PMID 22419320.

1.10 External links

- Selenium at *The Periodic Table of Videos* (University of Nottingham)

- [National Institutes of Health page on Selenium](#)
- [Assay](#)
- [ATSDR – Toxicological Profile: Selenium](#)
- [CDC - NIOSH Pocket Guide to Chemical Hazards](#)
- [Peter van der Krogt elements site](#)

Chapter 2

JScript

Not to be confused with **JavaScript**.

JScript is Microsoft's dialect of the **ECMAScript** standard^[2] that is used in **Microsoft's Internet Explorer**.

JScript is implemented as an **Active Scripting** engine. This means that it can be “plugged in” to **OLE Automation** applications that support **Active Scripting**, such as **Internet Explorer**, **Active Server Pages**, and **Windows Script Host**.^[3] It also means such applications can use multiple **Active Scripting** languages, e.g., **JScript**, **VBScript** or **PerlScript**.

JScript was first supported in the **Internet Explorer 3.0** browser released in August 1996. Its most recent version is **JScript 9.0**, included in **Internet Explorer 9**.

JScript 10.0^[4] is a separate dialect, also known as **JScript .NET**, which adds several new features from the abandoned fourth edition of the **ECMAScript** standard. It must be compiled for **.NET Framework** version 2 or version 4, but static type annotations are optional.

2.1 Comparison to JavaScript

As explained by **JavaScript** guru **Douglas Crockford** in his talk titled *The JavaScript Programming Language* on **YUI Theater**,

[Microsoft] did not want to deal with **Sun Microsystems** about the trademark issue, and so they called their implementation **JScript**. A lot of people think that **JScript** and **JavaScript** are different but similar languages. That's not the case. They are just different names for the same language, and the reason the names are different was to get around trademark issues.^[5]

However, **JScript** supports **conditional compilation**, which allows a programmer to selectively execute code within **block comments**. This is an extension to the **ECMAScript** standard that is not supported in other **JavaScript** implementations, thus making the above statement not completely true.

2.2 Versions

2.2.1 JScript

The original **JScript** is an **Active Scripting** engine. Like other **Active Scripting** languages, it is built on the **COM/OLE Automation** platform and provides scripting capabilities to host applications.

This is the version used when hosting **JScript** inside a **Web** page displayed by **Internet Explorer**, in an **HTML** application, in **classic ASP**, in **Windows Script Host** scripts and several other **Automation** environments.

JScript is sometimes referred to as “classic **JScript**” or “**Active Scripting JScript**” to differentiate it from newer **.NET**-based versions.

Some versions of **JScript** are available for multiple versions of **Internet Explorer** and **Windows**. For example, **JScript 5.7** was introduced with **Internet Explorer 7.0** and is also installed for **Internet Explorer 6.0** with **Windows XP Service Pack 3**, while **JScript 5.8** was introduced with **Internet Explorer 8.0** and is also installed with **Internet Explorer 6.0** on **Windows Mobile 6.5**.

Microsoft's implementation of **ECMAScript 5th Edition** in **Windows 8 Consumer Preview** is called *JavaScript* and the corresponding **Visual Studio 11 Express Beta** includes a “completely new”, full-featured **JavaScript** editor with **IntelliSense** enhancements for **HTML5** and **ECMAScript 5** syntax, “**VSDOC**” annotations for multiple overloads, simplified **DOM** configuration, brace matching, collapsible outlining and “go to definition”.^[6]

JScript is also available on **Windows CE** (included in **Windows Mobile**, optional in **Windows Embedded CE**). The **Windows CE** version lacks **Active Debugging**.

2.2.2 JScript .NET

Main article: **JScript .NET**

JScript .NET is a **Microsoft .NET** implementation of **JScript**. It is a **CLS** language and thus inherits very powerful features, but lacks many features of the origi-

nal JScript language, making it inappropriate for many scripting scenarios. JScript .NET can be used for ASP.NET pages and for complete .NET applications, but the lack of support for this language in Microsoft Visual Studio places it more as an upgrade path for classic ASP using classic JScript than as a new first-class language.

JScript .NET is not supported in the .NET Compact Framework.

Note: JScript .NET versions are not related to classic JScript versions. JScript .NET is a separate product. Even though JScript .NET is not supported within the Visual Studio IDE, its versions are in sync with other .NET languages versions (C#, VB.NET, VC++) that follow their corresponding Visual Studio versions.

.NET Framework 3.0 and 3.5 are built on top of 2.0 and do not include the newer JScript.NET release (version 10.0 for .NET Framework 4.0).

(Source: file version of jsc.exe JScript.NET compiler and Microsoft.JScript.dll installed with .NET Framework)

2.3 See also

- JScript.Encode
- Windows Script File
- Windows Script Host

2.4 Notes

- [1] JScript supports various features not specified in the ECMA standard,^[8] as does JavaScript.
- [2] Microsoft said JScript 3.0 was “the first scripting language to fully conform to the ECMA-262 standard”.^[9]
- [3] JScript 5.7 includes an implementation of the ECMAScript Compact Profile (ECMA-327) which turns off features not required by the ES-CP when using the “JScript.Compact” ProgID.
- [4] JScript .NET is “being developed in conjunction with ECMAScript Edition 4”.^[10]

2.5 References

- [1] “Types of Script Files”. Msdn.microsoft.com. Retrieved 2012-08-17.
- [2] “JScript (ECMAScript3)”. Msdn.microsoft.com. Retrieved 2012-08-17.
- [3] “What Is WSH?”. Msdn.microsoft.com. Retrieved 2012-08-17.
- [4] What is JScript 10.0?

- [5] Douglas Crockford, *The JavaScript Programming Language*
- [6] “What’s New in ASP.NET 4.5 and Visual Web Developer 11 Beta: The Official Microsoft ASP.NET Site”. “What’s New for ASP.NET 4.5 and Web Development in Visual Studio 11 Beta”.
- [7] *Version Information (Windows Scripting - JScript)*, Microsoft, retrieved 2010-05-31
- [8] *Microsoft JScript Features - Non-ECMA (Windows Scripting - JScript)*, Microsoft, retrieved 2010-05-31
- [9] *Microsoft Embraces ECMA Internet Scripting Standard; Delivers Industry’s First ECMA-Compliant Scripting Language, JScript 3.0, In Key Microsoft Products*, Microsoft, 1997-06-30
- [10] *What Is JScript .NET?*, Microsoft

2.6 External links

- JScript documentation in the MSDN Library
- JScript 5.7 Release Notes
- JScript .NET documentation in the MSDN Library
- JScript blog
- JavaScript - JScript - ECMAScript version history
- jPaq - A Fully Customizable JScript Library

2.7 Text and image sources, contributors, and licenses

2.7.1 Text

- Selenium** *Source:* <https://en.wikipedia.org/wiki/Selenium?oldid=745309467> *Contributors:* Kpjas, CYD, Vicki Rosenzweig, Mav, Bryan Derksen, Malcolm Farmer, JeLuF, Heron, Ewen, Stevertigo, DennisDaniels, Dwmyers, RTC, JohnOwens, Erik Zachte, Zashaw, Tannin, Iluvcapra, Minesweeper, Looxix~enwiki, Ahoerstemeier, Mgimpel~enwiki, Andres, Schneelocke, Ideyal, Emperorbma, Crusadeonilliteracy, Nohat, Stone, David Latapie, Wik, Zoicon5, Peregrine981, Svante~enwiki, Grendelkhan, Topbanana, Elwoz, Wetman, Donarreiskoffer, Robbot, Sharkey, Ee00224, Peak, Nurg, Romanm, Naddy, Modulatum, Mervyn, Martin Hampl~enwiki, Lupo, Carnildo, Giftlite, DocWatson42, Bork, Muke, Curps, Niteowlneils, Yekrats, Luigi30, Darrien, Bobblewik, Delta G, Ragib, Mmm~enwiki, Pgan002, Semprini, Yath, Antandrus, Doc Taxon, Elroch, Icairns, Tooki, Adashiel, Timelord, Imroy, DanielCD, Discospinster, Rich Farmbrough, Vsmith, Mani1, Bender235, Sunborn, Mashford, CanisRufus, BenjBot, El C, Joanjoc~enwiki, Kwamikagami, Worldtraveller, Remember, Femto, Davidruben, Arcadian, Eddideigel, A2Kafir, Jumbuck, Larry Grossman, Alansohn, Paleorthid, Benjah-bmm27, SI, Walkerma, Phyllis1753, TenOfAllTrades, Etherwings, Rentastrawberry, Gene Nygaard, Tainter, Corpet~enwiki, Megan1967, Syamil, David Haslam, Seansheep, Benbest, Sengkang, GregorB, CharlesC, Graham87, V8rik, FreplySpang, DePiep, Grammarbot, Saperaud~enwiki, Rjwilmsi, Jake Wartenberg, Ttaring, FlaBot, Wegsjac, Ground Zero, Nihiltres, Kerowyn, Lonfa, Kolbasz, Swtpc6800, Aspro, Colenso, Chobot, Jaraalbe, Moocha, Bgwhite, WriterHound, YurikBot, Wavelength, JWB, Chris Capoccia, Hellbus, Stephenb, Yyy, Cryptic, The Hokkaido Crow, Badagnani, Joel7687, Dogcow, Pyrotec, Biolinker, Rallette, Silverhill, Tetracube, Lt-wiki-bot, Pb30, Geoffrey.landis, Anclation~enwiki, JLaTondre, JDSpeeder1, Elijahmeeks, GrinBot~enwiki, Groyolo, Itub, SmackBot, Oxford Comma, M dorothy, Gigs, David.Mestel, Giraldufaber, Bomac, Edgar181, Gilliam, Skizzik, Mirokado, Chris the speller, Thumperward, Pylori, MalafayaBot, Uthbrian, DHN-bot~enwiki, Sbharis, Firetrap9254, KojieroSaske, Tscabot, Kemperb, CaptainCarrot, Vladislav, Rrburke, VMS Mosaic, SundarBot, Grover cleveland, Cybercobra, Bigturtle, Nakon, Pwjb, Smokefoot, Drphilharmonic, Jitterro, PeterJeremy, SashatoBot, Archimerged, DO11.10, Khazar, John, Coredesat, Chiragsdoshi, Rhelmich, IronGargoyle, 16@r, Beetstra, Mr Stephen, Jose77, Fan-1967, JoeBot, Andreas Rejbrand, W0lfie, GreyMage668, Igoldste, Thricecube, Nmadhubala, Poolkris, Vices, SeL, Stifynsemons, DanielRigal, WeggeBot, Neelix, Marqueeed, A876, Rhode Island Red, Rifleman 82, Travelbird, Difluoroethene, Roberta F., Narayanese, JGabbard, Malleus Fatuorum, Thijs!bot, Epb123, Headbomb, Escarbot, AntiVandalBot, Shirt58, Dr. Blofeld, Smartse, Tillman, JAnDbot, Deflective, Josephmarty, Plantsurfer, Sheitan, IanOsgood, .anacondabot, Acroterion, Currenz, Magioladitis, WolfmanSF, Karlhahn, Bongwarrior, VoABot II, Kajasudhakarakababu, Brewhaha@edmc.net, Baccyak4H, Dmaddison, WhatamIdoing, Cgingold, Lethaniol, TheBoy765, WLU, MartinBot, NReitzel, Cosumel, Nono64, Leyo, Watch37264, Lithui, Kimse, Nbauman, Rod57, Kevspencer, Jeepday, Mikael Häggström, Coppertwig, Ltioe, Mary Moor, Sapphirine, Thesis4Eva, Julianclouton, Treisijs, Kvdveer, Xpansion, Chatvisual, 28bytes, VolkovBot, Testbenchdude, DrMicro, TheMindsEye, Zotamedu, TXiKiBoT, Cloudswrest, Una Smith, Psyche825, Taylork1992, Eubulides, CMatts, Jswiftrap, WatermelonPotion, Onceonthisisland, AlleborgoBot, EmxBot, PedroDaGr8, Fanatix, SieBot, PlanetStar, Sebastiano venturi, Cwkmall, Xenophon777, Flyer22 Reborn, Alexbrn, Cablehorn, Arjen Dijkman, Oxymoron83, Hak-ká-ngin, Double Vigie, Nergaal, Msjayhawk, ClueBot, Trojancowboy, PipepBot, The Thing That Should Not Be, ImperfectlyInformed, Pras, DragonBot, Shinkolobwe, Razorflame, HelenRidley3, Plasmic Physics, DumZiBoT, Fiquei, Cheese munkay, Golddragon24, XLinkBot, Vanished 45kd09la13, Alexius08, SkyLined, Mr0t1633, DOI bot, Viper9411, Ironholds, CarsracBot, 5 albert square, Alchemist-hp, Tassedethe, Numbo3-bot, Erutuo, Tide rolls, Jarble, Lileensorchick, Luckas-bot, TheSuave, Yobot, Lethalgeek, Pbtotgourou, II MusLiM HyBRiD II, Tempodivalse, Ectuohey, AnomieBOT, Rubinbot, Jim1138, Jo3saml, Flewis, MaterialsScientist, Citation bot, DynamoDegsy, ArthurBot, LilHelpa, MauritsBot, Xqbot, Schmutz MDPI, Gigemag76, YBG, Wordclock, Srich32977, Nedim Ardoğa, Doulos Christos, Josh 2369, LucienBOT, Riventree, VHacker, Jeandry, Eldin raigmore, Atyrstan, Finalius, Robo37, Citation bot 1, Pinethicket, HRoestBot, Rameshngbot, Tom.Reding, Tincopper2, Serols, Rousseu, Mikespedia, Jututacular, Xeworlebi, Kibi78704, Jauhienij, Titsnitsmaliits, Double sharp, TobeBot, Trappist the monk, Attinio, Jynto, DARTH SIDIOUS 2, Andrea105, EmausBot, WikitanvirBot, Curly daus, 478jjjz, Rbaselt, Racexr11, GoingBatty, Wikipelli, Peterindelft, Kazem.f.a, Fæ, StringTheory11, Chemicalinterest, SporkBot, Ohnodoodle, Donner60, Era2, Blazingice30672, DASHBotAV, ClueBot NG, Horoporo, Azminch, This lousy T-shirt, DPG1993, Edgar1024132, Denis Zhilin, Skoot13, Frietjes, Sampenglase, Rezabot, Widr, Helpful Pixie Bot, Кайдзэн, Bibcode Bot, Missclark, BG19bot, TinyEdit, ElphiBot, NVCRalston, Pengdu, Venturis1947, BattyBot, WBeckon, Sfarney, Schafhirt, TylerDurden8823, JYBot, Dexbot, Hmainsbot1, Stas1995, JZNIOSH, Ventur-sebastiano, TwoTwoHello, James12345, T42N24T, Reatlas, WikiUser44, Sawerchessread, EdGMnt, Ramendocor, Anrnusna, Davejones7, Grand'mere Eugene, ZYjacklin, Medgirl131, Dr.maishi, Sajawal Ahmed Ali Sial, Banclark80, Emily Temple-Wood (NIOSH), LL221W, KasparBot, Smikail, TerpeneOtto, Lulztech and Anonymous: 436
- JScript** *Source:* <https://en.wikipedia.org/wiki/JScript?oldid=730364424> *Contributors:* General Wesc, William Avery, Maury Markowitz, Frecklefoot, Voidvector, LittleDan, Julesd, Guaka, Dysprosia, Grendelkhan, Bevo, Chealer, Ancheta Wis, Robbrown, AlistairMcMillan, Abu badali, Gschizas, Asqueella, Porges, Hydrox, Rowaasr13, Danakil, John Vandenberg, Minghong, Wayfarer, Andrew Gray, Paul1337, Oleg Alexandrov, WadeSimMiser, Isnow, Crpietschmann, Toussaint, Jbarta, Rjwilmsi, Tizio, Koavf, Amire80, Aegicen~enwiki, FlaBot, Latka, Margosbot~enwiki, Ysangkok, RexNL, YurikBot, FrenchIsAwesome, Me and, Pi Delport, Hydrargyrum, Mipadi, Razorox, Five-strokes, Gadget850, Vlad, Black Falcon, Ms2ger, Tyomitch, SmackBot, Maian, Myrdred, Renku, Bluebot, Thumperward, Gyrobo, Frap, Jacob Poon, TimJackson, Ohconfucius, Soumyasch, Arbruijn, Phuzion, Sander Sæde, Grayczyk, FatalError, SkyWalker, FleetCommand, Cydebot, Thijs!bot, Hervegirod, Mdsy, Magioladitis, Tanitall, Danpetitpas, Joshua Issac, Josh Tumath, Ajfweb, VolkovBot, TXiKiBoT, MrFirewall, Cminor9, Synthebot, Digita, S.Örvarr.S, Brenont, PhMajerus, Sfan00 IMG, Alksentrs, Mattspac, John Nevard, El bot de la dieta, TimotheusMax, Addbot, GhettoBlaster, Mabdu, Ricvelozo, Lucas-bot, Yobot, AnomieBOT, Mlb1992, Jaselg, Sass24, Krinkle, FrescoBot, Sae1962, Machine Elf 1735, MastiBot, JeepdaySock, Black Yoshi, ZéroBot, DonToto, M93A, Solde9, ClueBot NG, Helpful Pixie Bot, Damaca, Chmarkine, Jfd34, Codename Lisa, Eatingbaconsince2000, Editorfun and Anonymous: 90

2.7.2 Images

- File:Commons-logo.svg** *Source:* <https://upload.wikimedia.org/wikipedia/en/4/4a/Commons-logo.svg> *License:* CC-BY-SA-3.0 *Contributors:* ? *Original artist:* ?
- File:Crystal_Clear_app_browser.png** *Source:* https://upload.wikimedia.org/wikipedia/commons/f/fe/Crystal_Clear_app_browser.png *License:* LGPL *Contributors:* All Crystal icons were posted by the author as LGPL on kde-look *Original artist:* Everaldo Coelho and YellowIcon
- File:Folder_Hexagonal_Icon.svg** *Source:* https://upload.wikimedia.org/wikipedia/en/4/48/Folder_Hexagonal_Icon.svg *License:* Cc-by-sa-3.0 *Contributors:* ? *Original artist:* ?

- **File:Free-to-read_lock_75.svg** *Source:* https://upload.wikimedia.org/wikipedia/commons/8/80/Free-to-read_lock_75.svg *License:* CC0 *Contributors:* Adapted from 9pxlOpen_Access_logo_PLoS_white_green.svg *Original artist:* This version:Trappist_the_monk (talk) (Uploads)
- **File:Hexagonal.svg** *Source:* <https://upload.wikimedia.org/wikipedia/commons/c/c2/Hexagonal.svg> *License:* BSD *Contributors:* Transferred from en.wikipedia to Commons. *Original artist:* The original uploader was Danieljamesscott at English Wikipedia
- **File:Internet_Explorer_10_logo.svg** *Source:* https://upload.wikimedia.org/wikipedia/commons/2/2f/Internet_Explorer_10_logo.svg *License:* Public domain *Contributors:* Extracted from this color and this shape. *Original artist:* Microsoft
- **File:Jscript_icon.gif** *Source:* https://upload.wikimedia.org/wikipedia/en/e/e8/Jscript_icon.gif *License:* Fair use *Contributors:* **Original publication:** unsure

Immediate source: <http://www.newobjects.com/default.asp?category=78> *Original artist:* Microsoft

- **File:M_box.svg** *Source:* https://upload.wikimedia.org/wikipedia/commons/9/94/M_box.svg *License:* Public domain *Contributors:* Own work based on: File:Microsoft.svg *Original artist:* Ariesk47 (talk)
- **File:Se_dose-response_curve_for_juvenile_salmon_mortality_-_percent_scale.jpg** *Source:* https://upload.wikimedia.org/wikipedia/commons/0/02/Se_dose-response_curve_for_juvenile_salmon_mortality_-_percent_scale.jpg *License:* CC BY-SA 3.0 *Contributors:* Own work *Original artist:* William N. Beckon
- **File:Selenium-dioxide-chain-3D-balls.png** *Source:* <https://upload.wikimedia.org/wikipedia/commons/8/8e/Selenium-dioxide-chain-3D-balls.png> *License:* Public domain *Contributors:* Own work *Original artist:* Ben Mills
- **File:Selenium_in_sandstone_Westwater_Canyon_Section_23_Mine_Grants,_New_Mexico.jpg** *Source:* https://upload.wikimedia.org/wikipedia/commons/e/e6/Selenium_in_sandstone_Westwater_Canyon_Section_23_Mine_Grants%2C_New_Mexico.jpg *License:* CC BY 2.0 *Contributors:* Selenium in sandstone (Westwater Canyon Member, Morrison Formation, Upper Jurassic; Section 23 Mine, north of Grants, New Mexico, USA) 1 *Original artist:* James St. John
- **File:Selenium_trigonal.jpg** *Source:* https://upload.wikimedia.org/wikipedia/en/d/df/Selenium_trigonal.jpg *License:* CC-BY-SA-3.0 *Contributors:*
I created this work entirely by myself.
Original artist:
Materialscentist (talk)
- **File:Symbol_list_class.svg** *Source:* https://upload.wikimedia.org/wikipedia/en/d/db/Symbol_list_class.svg *License:* Public domain *Contributors:* ? *Original artist:* ?
- **File:Transparent.gif** *Source:* <https://upload.wikimedia.org/wikipedia/commons/c/ce/Transparent.gif> *License:* Public domain *Contributors:* Own work *Original artist:* Edokter
- **File:Wikibooks-logo-en-noslogan.svg** *Source:* <https://upload.wikimedia.org/wikipedia/commons/d/df/Wikibooks-logo-en-noslogan.svg> *License:* CC BY-SA 3.0 *Contributors:* Own work *Original artist:* User:Bastique, User:Ramac et al.
- **File:Wiktionary-logo-v2.svg** *Source:* <https://upload.wikimedia.org/wikipedia/commons/0/06/Wiktionary-logo-v2.svg> *License:* CC BY-SA 4.0 *Contributors:* Own work *Original artist:* Dan Polansky based on work currently attributed to Wikimedia Foundation but originally created by Smurrayinchester
- **File:Windows_logo_-_2012_(dark_blue).svg** *Source:* https://upload.wikimedia.org/wikipedia/commons/e/ee/Windows_logo_%E2%80%93_2012_%28dark_blue%29.svg *License:* Public domain *Contributors:* Windows 10 *Original artist:* File:Windows 10 Logo.svg: Multiple editors; see image description page

2.7.3 Content license

- Creative Commons Attribution-Share Alike 3.0