

Statistical Thinking: Monte Carlo Simulation

by Prof. Seungchul Lee
iSystems Design Lab
<http://isystems.unist.ac.kr/>
UNIST

Table of Contents

- I. 1. Example: Probability of having head with a fair coin
- II. 2. Example: the expected number of trials upto the first hitting H
- III. 3. Example: Integration
- IV. 4. Example: compute π statistically

1. Example: Probability of having head with a fair coin

Head 1 and Tail 0

In [1]:

```
import numpy as np
import matplotlib.pyplot as plt

%matplotlib inline
```

In [2]:

```
n_trials = 100
n_H = 0

for i in range(n_trials):
    flip = np.random.randint(2)
    if flip == 1:
        n_H += 1

print(n_H/n_trials)
```

0.49

2. Example: the expected number of trials upto the first hitting H

$$\text{coin} \begin{cases} H : \frac{1}{2} \\ T : \frac{1}{2} \end{cases}$$

Question: the expected number of trials upto the first hitting H ?

$$\begin{array}{lll} 1 & H & \frac{1}{2} \\ 2 & TH & \frac{1}{2} \frac{1}{2} \\ 3 & TTH & \frac{1}{2} \frac{1}{2} \frac{1}{2} \\ & \vdots & \end{array} \quad \sum_{n=1}^{\infty} n \left(\frac{1}{2} \right)^n$$

In [3]:

```
val = 0

for n in range(1,20):
    val += n*(1/2)**n

print(val)
```

1.999959945678711

In [4]:

```
n_trials = 1000

NUM = []

for i in range(n_trials):
    num = 1
    while np.random.randint(2) != 0:
        num += 1
    NUM.append(num)

print(np.mean(NUM))
```

1.956

Remark:

how to compute $\sum_{n=1}^{\infty} n \left(\frac{1}{2}\right)^n = 1 \cdot \frac{1}{2} + 2 \left(\frac{1}{2}\right)^2 + 3 \left(\frac{1}{2}\right)^3 + \dots$

$$\begin{aligned} \frac{d}{dx} \sum_{n=1}^{\infty} (1-x)^{n+1} &= \frac{d}{dx} \frac{(1-x)^2}{1-(1-x)} = \frac{d}{dx} \frac{(1-x)^2}{x} \\ \sum_{n=1}^{\infty} (n+1)(1-x)^n &= \sum_{n=1}^{\infty} n(1-x)^n + \sum_{n=1}^{\infty} (1-x)^n \\ &= \frac{(1-x)^2 + 2(1-x)x}{x^2} \\ x = \frac{1}{2} &\Rightarrow \sum_{n=1}^{\infty} n \left(\frac{1}{2}\right)^n + \frac{1 - \frac{1}{2}}{\frac{1}{2}} = \frac{\frac{1}{4} + 2 \cdot \frac{1}{2} \cdot \frac{1}{2}}{\frac{1}{4}} \\ &\Rightarrow \sum_{n=1}^{\infty} n \left(\frac{1}{2}\right)^n = 2 \end{aligned}$$


or

$$\begin{aligned} y &= 1 \cdot \frac{1}{2} + 2 \left(\frac{1}{2}\right)^2 + 3 \left(\frac{1}{2}\right)^3 + \dots \\ 2y &= 1 + 2 \left(\frac{1}{2}\right) + 3 \left(\frac{1}{2}\right)^2 + \dots \\ \Rightarrow y &= 1 + \left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^2 + \dots = \frac{1}{1 - \frac{1}{2}} = 2 \end{aligned}$$

3. Example: Integration

$$\int_0^1 x^2 dx = \frac{1}{2} x^3 \Big|_0^1 = \frac{1}{2}$$

Question : how to solve integration with computers ?



$$\Delta A = f(x) \Delta x$$

$$A \approx \sum \Delta A = \sum f(x_k) \Delta x$$

In [5]:

```
# by summing up

dx = 0.001
x = np.arange(0,1,dx)

A = 0
for xk in x:
    A += (xk**2)*dx

print(A)
```

0.3328335

In [6]:



```
# shortened version

A = np.sum(x**2)*dx
print(A)
```


0.3328335

Question: another method ? (use randomness)

$x = \text{rand}(n, 1)$
 $y = \text{rand}(n, 1)$ \implies $\text{plot}(x, y)$



$$\frac{\# \text{ under } y = x^2}{\# \text{ total}}$$



$$\frac{\text{area under } y = x^2}{\text{total area}}$$

It is known as **Monte Carlo simulation**

\implies extremely powerful

\implies can apply to many, many, many (engineering) problems

In [7]:

```
# the number of points below curve out of the total number is a fraction of area

n = 10000

# generate n random numbers x and y

x = np.random.rand(n,1)
y = np.random.rand(n,1)

count = 0
for i in range(n):
    # compute y to f(x)
    if y[i,0] < x[i,0]**2:
        count += 1

# result normalized by total #
print(count/n)
```

0.3349

In [8]:

```
# shortened version

A = np.sum(y < x**2)/n
print(A)
```

0.3349

4. Example: compute π statistically

In [9]:

```
n = 5000

y = np.random.rand(n,1)
x = np.random.rand(n,1)

idx = np.empty((n,1))
count = 0

for i in range(n):
    if np.sqrt(x[i]**2 + y[i]**2) < 1:
        count += 1
        idx[i] = 1
    else:
        idx[i] = 0

print((count/n)*4)
```

3.1288