

MAT292—ODE

Jonah Chen

September 13, 2021

Contents

1	Newton's Law of Cooling	1
---	-------------------------	---

1 Newton's Law of Cooling

Newtons law of cooling states that

$$\frac{du}{dt} = -k(u - T_0) \quad (1.1)$$

Note that there is one trivial solution, the equilibrium solution is $u(t) = T_0$. The meaning of this solution is the temperature of an object doesn't change when it is already at the equilibrium temperature.

$$\frac{\frac{du}{dt}}{u - T_0} = -k \quad (1.2)$$

$$\frac{d}{dt} \log(u - T_0) = -k \quad (1.3)$$

$$\log(u - T_0) = -kt + c_1 \quad (1.4)$$

$$u = T_0 + \exp(c_1) \exp(-kt) = T_0 + c_2 \exp(-kt) \quad (1.5)$$

Note that $c_2 = \exp(c_1) > 0$. However, this is not a complete solution as it cannot describe the solutions with $u < T_0$.

Warning: note that the integral of $\frac{1}{x}$ is $\log|x|$, **NOT** $\log(x)$. This is what caused the solution to be incomplete.

Hence, $c_2 = \pm \exp(c_1)$. Note that $c_1 = \pm\infty$ is allowed thus so c_2 can take any value.

Below are the integral curves.