


# MAT347 Abstract Algebra

Jonah Chen

## 1 GROUPS

Groups are generally associated with symmetries. Consider the equilateral triangle:



We know that there are six symmetries of the triangle:

- Identity transformation (do nothing) denoted as  $\text{id}$  or  $e$
- Two rotations ( $A \rightarrow B \rightarrow C \rightarrow A$  and  $A \rightarrow C \rightarrow B \rightarrow A$ )
- Three reflections  $A \leftrightarrow B$ ,  $A \leftrightarrow C$ ,  $B \leftrightarrow C$

Note that these symmetries preserve the structure of the triangle, hence the composition of two symmetries must also be a symmetry. Let

- $\rho$  be the rotation  $A \rightarrow B \rightarrow C \rightarrow A$
- $\sigma$  be the reflections  $B \leftrightarrow C$

Note that  $\rho\sigma$  is the  $A \leftrightarrow C$  reflection and  $\sigma\rho$  is the  $A \leftrightarrow B$  reflection. Hence they may not be commutative.

We also know that all symmetries can be reversed.  $\alpha$  has an inverse  $\alpha^{-1}$  such that  $\alpha\alpha^{-1} = \alpha^{-1}\alpha = e$ . These inspires the following definition:

**Definition:** A **group** is a set  $G$  with a composition

$$G \times G \rightarrow G \tag{1}$$

$$(g, h) \mapsto g \cdot h \tag{2}$$

Satisfying:

- Associativity:  $(g \cdot h) \cdot k = g \cdot (h \cdot k)$

- Identity:  $\exists e \in G$  such that  $g \cdot e = e \cdot g = g$  for all  $g \in G$
- Inverse:  $\forall g \in G, \exists g^{-1} \in G$  such that  $g \cdot g^{-1} = g^{-1} \cdot g = e$

Examples:

- $\mathbb{Z}$  with  $+$  is a group. It is associative,  $e = 0$  and  $g^{-1} = -g$ .
- $\mathbb{Z}/n\mathbb{Z}$  with addition modulo  $n$ .
- If  $F$  is a field, it implicitly has two group structures:
  - Additive group:  $(F, +)$  is a group. It is associative,  $e = 0$  and  $g^{-1} = -g$ .
  - Multiplicative group:  $(F \setminus \{0\}, \times)$  is a group. It is associative,  $e = 1$  and  $g^{-1} = 1/g$ .
- $GL(n, F)$  – “general linear group” contains all invertible  $n \times n$  matrices.
- $SL(n, F)$  – “special linear group” contains all invertible  $n \times n$  matrices with determinant 1.
- $SO(n, F)$  – “special orthogonal group”  $= \{A \in SL(n, F) | A^t = A^{-1}\}$ .


## 1.1 CYCLIC GROUPS

One of the simplest groups is  $\mathbb{Z}/n\mathbb{Z}$  for  $n \in \mathbb{N}$  with the operation addition modulo  $n$ . This is known as the “cyclic group of order  $n$ ” or  $C_n$ . i.e. for  $n = 8$ ,  $5 + 7 = 4 \pmod{8}$ , which we denote  $\bar{5} + \bar{7} = \bar{4}$ .

We know the inverse  $\bar{k}^{-1} = \overline{n - k}$  for nonzero  $k$  or  $\bar{0}^{-1} = \bar{0}$ .

Another way to express the cyclic group is  $\bar{k} \leftrightarrow e^{2\pi i k/n}$  with multiplication operation. Then,

$$\overline{k+n} = e^{2\pi i(k+n)/n} = e^{2\pi i k/n} e^{2\pi i n/n} = e^{2\pi i k/n} = \bar{k}. \quad (3)$$



**Definition:** [Order] The **order** of a group  $G$  is its cardinality denoted  $\text{ord}(G)$  or  $|G|$ . It could be a finite or infinite ordinal. In particular,  $|C_n| = n$ .

## 1.2 QUATERNION GROUP

The quaternion group  $\mathbb{H} = \{\pm 1, \pm i, \pm j, \pm k\}$  is a group of order 8 with the multiplication operation. It has

**Definition:** [Subgroup] A **subgroup** of a group  $G$  is a subset  $H \subseteq G$  such that  $H$  is a group.

**Definition:** [Coset] If  $G$  is a group and  $H \leq G$ , consider sets of the form

$$Hg = \{hg | h \in H\} \quad (4)$$

This is a **right coset** of  $H$ .

**Theorem:** [Partitioning with Cosets] Consider  $Hg$  and  $Hg'$  for  $g, g' \in G$ . There are two cases:

- They might be disjoint:  $Hg \cap Hg' = \emptyset$ .
- They might intersect. Suppose  $hg = h'g'$  for some  $h, h' \in H$

$$h^{-1}hg = h^{-1}h'g' \quad (5)$$

$$g = h^{-1}h'g' \in Hg' \quad (6)$$

Similarly,  $g' \in Hg$ . Consider an arbitrary element of  $Hg$  with  $k \in H$ . Then,  $kg = kh^{-1}h'g' \in Hg'$  i.e.  $Hg \leq Hg'$ . Similarly,  $Hg' \leq Hg$ . Thus,  $Hg = Hg'$ .

The right cosets of  $H$  partition  $G$ . In particular,

$$G = \bigsqcup Hg_i \quad (7)$$

For fixed  $g$ , if  $hg = h'g$  for  $h, h' \in H$  then  $hgg^{-1} = h'gg^{-1}$  so  $h = h'$ . So in  $Hg$ , every element can be matched with an element of  $H$ . So,  $|Hg| = |H|$ .

**Theorem:** [Lagrange] If  $|G| < \infty$  and  $H \leq G$ , then  $|H| \mid |G|$

**Definition:** [Index] For  $H \leq G$ , the **index** of  $H$  in  $G$  is  $[G : H] = |G|/|H|$ .

If  $|G| = 13$ , the only subgroups of  $G$  are  $\{e\}, G$ .

If  $G = \mathbb{Z}$  and  $H = 2\mathbb{Z}$  (even numbers). Then  $H + 0 = H$  is one coset, and  $H + 1 =$  the odd integers is another coset. So,  $\mathbb{Z} = (2\mathbb{Z}) \sqcup (2\mathbb{Z} + 1)$ .

Same for Left Cosets Interaction of left and right cosets?

Consider the triangle group with rotations  $e, \rho, \rho^2$  and reflections  $\sigma_A, \sigma_B, \sigma_C$ . Consider the subgroup  $H = \{e, \sigma_A\}$ .

$$He = \{e, \sigma_A\} \quad (8)$$

$$H\rho = \{\rho, \sigma_B\} \quad (9)$$

$$H\rho^2 = \{\rho^2, \sigma_C\} \quad (10)$$

$$eH = \{e, \sigma_A\} \quad (11)$$

$$\rho H = \{\rho, \sigma_C\} \quad (12)$$

$$\rho^2 H = \{\rho^2, \sigma_B\} \quad (13)$$

Note that the left and right cosets are different. They are the same if the group is commutative.

**Definition:** [Action] An **action** of a group  $G$  on a set  $X$  is a map

$$G \times X \rightarrow X \quad (14)$$

$$(g, x) \mapsto gx \quad (15)$$

such that

$$(gh)x = g(hx) \quad (16)$$

$$ex = x \quad (17)$$

If  $G$  is a group, it acts on itself. This is called a “left translation” or “left regular action”.

How about the right action  $(g, x) \mapsto xg$ . The second condition may not be true

$$(gh, x) = xgh \quad (18)$$

$$(g, (hx)) = (g, xh) = xhg \quad (19)$$

which is not true. Instead, let  $(g, x) = xg^{-1}$ . Then,

$$(gh, x) = x(gh)^{-1} = xh^{-1}g^{-1} \quad (20)$$

$$(g, (hx)) = (g, xh^{-1}) = xh^{-1}g^{-1} \quad (21)$$

This is the definition of the right action.

There is a third action of  $G$  on itself by  $(g, x) = xgx^{-1}$ . This action is called conjugation.

Take the following example: Let  $G = SO(3)$  and let  $X = S^2$ .  $G$  acts on  $X$  by rotation. Let  $H = \left\{ \begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \right\}$  be the subgroup of rotations that fixes the  $z$ -axis.

$H$  also acts on  $X$  ??

**Definition:** [Orbit] If  $G$  acts on  $X$ , the **orbit** of  $x \in X$  is the set  $Gx = \{gx | g \in G\}$ . i.e. the set of all points  $x$  is taken to by elements of  $G$ .

The orbits of  $H \approx SO(2)$  on the sphere are the lines of latitude (and the north and south poles).

$H$  fixes the north pole, thus every coset  $gH$  takes the north pole to a point. Suppose  $gH$  and  $g'H$  are cosets such that  $gHN = g'HN \implies gN = g'N \implies (g')^{-1}gN = N \implies (g')^{-1}g \in H \implies gH \dots$  so the points ofn the sphere are in 1-1 correspondence with the left cosets of  $H$ .

**Definition:** [Stabilizer] If  $G$  acts on  $X$  and  $x \in X$ , the “stabilizer” of  $x$  in  $G$  is  $\{g \in G | gx = x\}$

**Definition:** [Centralizer] If  $A \subset G$ , the **centralizer** of  $A$  in  $G$  is  $C_G(A) = \{g \in G | ga = ag \forall a \in A\}$

- If  $G$  is abelian, then  $C_G(A) = G$  for any  $A$ .
- In the triangle group,  $C_G(\{\rho\}) = \{e, \rho, \rho^2\}$

**Definition:** [Center] The **center** of  $G$  is  $Z(G) = \{g \in G | gg' = g'g \forall g' \in G\} = C_G(G)$

**Proposition:** For any  $A \subset G$ ,  $C_G(A) \leq Z(G)$  (is a subgroup).

Consider the regular  $n$ -gon ( $n \geq 3$ ), what are its rigid motion symmetries?

- There are always  $n$  rotations by  $\frac{2\pi}{n}$  about the origin.
- When  $n$  is even, there are  $n/2$  reflections in each pair of edges, and each pair of vertices. When  $n$  is odd, there are  $n$  reflections in each pair of (edge, vertex). There are always  $n$  reflections.
- Write  $\rho$  for clockwise rotation by  $\frac{2\pi}{n}$ . Fix one vertex and let  $\sigma$  be the reflection that fixes that vertex.
- Note that  $\rho\sigma = \sigma\rho^{-1}$ . To show this, it suffices to find where two of the vertices gets mapped.

**Proposition:** The symmetries are  $e, \rho, \rho^2, \dots, \rho^{n-1}, \sigma, \sigma\rho, \sigma\rho^2, \dots, \sigma\rho^{n-1}$

**Definition:** [dihedral group] The group of symmetries of the regular  $n$ -gon is  $D_{2n}$ , the **dihedral group** of order  $2n$ .

Given  $H \leq G$  we write  $G/H$  as the set of left cosets

$$G/H = \{gH | g \in G\} \quad (22)$$

$$H \backslash G = \{Hg | g \in G\} \quad (23)$$

Both of these are called “ $G \bmod H$ ”. In general, the two are different.

Now we want to ask, is  $H \backslash G$  a group?

- The most naive idea is to reuse multiplication in  $G$ , i.e.  $Hg \cdot Hg' = Hgg'$ , but it only sometimes works.
- This formula means:  $hg \cdot h'g' = h''gg'$ . For any  $h, h' \in H, \exists h''$  s.t. this holds.
- Trick:  $hg \cdot h'g' = hgh'e'g' = hgh'(g^{-1}g)g' = h(ghg^{-1})gg'$ . Now we can ask if  $ghg^{-1} \in H$  (for every  $h' \in H$ )

**Definition:** [Normal Subgroup] A subgroup  $H \leq G$  is **normal** if  $ghg^{-1} \in H \forall g \in G, h \in H$ , which is abbreviated as  $gHg^{-1} = H$ .  $H \trianglelefteq G$  means  $H$  is a normal subgroup of  $G$

- Notice that if  $gHg^{-1} = H$  then  $gH = Hg$ . So  $H$  is normal, the left and right cosets must be the same.

**Definition:** [Quotient Group] If  $H \trianglelefteq G$ , then  $G/H$  is called the quotient group.

### 1.3 HOMOMORPHISMS

**Definition:** [Homomorphism] If  $G, K$  are groups, a **homomorphism** is a map  $\varphi : G \rightarrow K$  such that  $\varphi(gg') = \varphi(g)\varphi(g') \forall g, g' \in G$ .

Observations: IF  $\varphi : G \rightarrow K$  is a homomorphism and  $g \in G$ , then

1.  $\varphi(g) = \varphi(eg) = \varphi(e)\varphi(g)$ , so  $\varphi(e) = e$  (the identity element of  $K$ )
2.  $e = \varphi(e) = \varphi(gg^{-1}) = \varphi(g)\varphi(g^{-1})$ , so  $\varphi(g^{-1}) = \varphi(g)^{-1}$

#### Examples

- $G = \mathbb{Z}$  and  $\varphi : \mathbb{Z} \rightarrow \mathbb{Z}, \varphi(n) = 2n$  is a homomorphism, as  $\varphi(n+m) = 2(n+m) = 2n+2m = \varphi(n) + \varphi(m)$
- $G = \mathbb{Z}, K = \mathbb{R}$  and  $\varphi : \mathbb{Z} \rightarrow \mathbb{R}, \varphi(n) = n$ . This mapping is called an **inclusion** as  $\mathbb{Z} \subset \mathbb{R}$ .
- If  $G$  is a group and  $g_0 \in G$ , then  $C_{g_0} : G \rightarrow G, g \mapsto g_0 g g_0^{-1}$  is a homomorphism.
- A linear transformation  $T : V \rightarrow W$  if  $V, W$  are vector spaces (the additive group).
- Note that  $\varphi : g \mapsto g^{-1}$  is **only** a homomorphism if  $G$  is abelian.

**Definition:** [Kernel/Image] If  $\varphi : G \rightarrow G'$  is a homomorphism, then the **kernel** of  $\varphi$  is

$$\ker(\varphi) = \{g \in G \mid \varphi(g) = e\}. \quad (24)$$

The **image** of  $\varphi$  is

$$\text{im}(\varphi) = \{\varphi(g) \mid g \in G\} \subseteq G' \quad (25)$$

**Theorem:**  $\ker(\varphi) \leq G$  and  $\text{im}(\varphi) \leq G'$   $\ker(\varphi) \trianglelefteq G$

*Proof.* Since  $\varphi(e) = e$ ,  $e \in \ker(\varphi)$ , and  $e \in \text{im}(\varphi)$ . So both are nonempty. Suppose  $g, h \in \ker(\varphi)$ ,  $e = \varphi(e) = \varphi(hh^{-1}) = \varphi(h)\varphi(h^{-1}) \dots$  □

- Suppose  $N \trianglelefteq G$  and then define  $G \rightarrow G/N, g \mapsto Ng$ . We claim this is a homomorphism. Proof is simple  $\varphi(gg') = Ng g' = NgNg' = NgN(g^{-1}gg') = N(gNg^{-1})gg' = NNgg' = Ng g'$
- This map is called the (natural) **projection** of  $G$  onto  $G/N$ . Sometimes written  $\Pi_{G/N}$  or  $\text{proj}_{G/N}$ .
- $\text{im}(\Pi_{G/N}) = G/N$  and  $\ker(\Pi_{G/N}) = N$ .
- Any homomorphism is related to this one, so this could be considered as the “generic homomorphism”.

**Definition:** [Isomorphism] If  $\varphi : G \rightarrow H$  is a homomorphism, and  $\ker(\varphi) = \{e\}$  then  $\varphi$  is injective. If  $\varphi(G) = H$  then  $\varphi$  is surjective. Thinking of  $G$  and  $H$  as sets, there is an inverse  $\varphi^{-1} : H \rightarrow G$  such that  $\varphi^{-1} \circ \varphi = 1_G$  and  $\varphi \circ \varphi^{-1} = 1_H$ . It is easy to check that  $\varphi^{-1}$  is also a homomorphism. In this case,  $\varphi$  is an **isomorphism**

- Suppose we have an injective homomorphism  $\varphi : G \rightarrow H$  where  $\ker(\varphi) = \{e\}$ . Then, we can consider  $\varphi : G \rightarrow \text{im}(\varphi) \leq H$ . Sometimes we say  $\varphi : G \rightarrow H$  is an **isomorphism into**  $H$ , as opposed to an isomorphism **onto**  $H$  or between  $G$  and  $H$ .

**Definition:** [Automorphism] If  $G$  is a group, an **automorphism** of  $G$  is an isomorphism  $\varphi : G \rightarrow G$ .

#### Examples:

- If  $G = \mathbb{Z}$ ,  $n \mapsto -n$  is the only automorphism apart from the identity.
- If  $G$  is abelian,  $g \mapsto g^{-1}$  is an automorphism.
- If  $F$  is a field, and  $G = GL(n, F)$  then  $g \mapsto (g^t)^{-1}$  (transposed inverse) is an automorphism.
- If we fix  $g_0 \in G$  then the conjugation  $C_{g_0} : G \rightarrow G$  where  $C_{g_0}(g) = g_0 g g_0^{-1}$  is an automorphism.

**Definition:** [Automorphism Group]  $\text{Alt}(G)$  is the **group** of automorphisms of  $G$ .

**Definition:** [Inner/Outer Automorphisms] The **inner automorphisms** of  $G$  are

$$\text{Inn}(G) = \{\varphi \in \text{Alt}(G) \mid \varphi = C_{g_0} \text{ for some } g_0 \in G\}. \quad (26)$$

If an element of  $\text{Alt}(G)$  that is not inner is **outer**.

- It is easy to show that  $\text{Inn}(G) \leq \text{Alt}(G)$ .
- Observe that if  $G$  is abelian, then  $\text{Inn}(G) = \{\text{id}\}$
- In general,  $\{\text{id}\} \leq \text{Inn}(G) \leq \text{Alt}(G)$ .
- The map

$$G \rightarrow \text{Alt}(G) \quad (27)$$

$$g \mapsto C_g \quad (28)$$

is a homomorphism. Its image is  $\text{Inn}(G)$  and its kernel is  $Z_G$  (the center).

**Definition:** [Fiber] If  $p$  is a projection, then  $p^{-1}(x)$  is the **fiber** over  $x$

- If  $N \triangleleft G$ , the projection  $\pi : G \rightarrow G/N$  is a homomorphism. The fibers of  $\pi$  is the cosets  $gN = Ng$ , and they are all the same size.
- Suppose  $\varphi : G \rightarrow H$  is a homomorphism, and  $N = \ker(\varphi) \trianglelefteq G$ . The fibers of  $\varphi$  is the cosets of  $G/N$ .
- We have  $\varphi : G \rightarrow H$  and  $\pi : G \rightarrow G/N$ . Wouldn't it be nice if  $G/N \rightarrow H$  "induced by  $\varphi$ " were a homomorphism? Well, it is.

**Theorem:** [(First) Isomorphism] If  $\varphi : G \rightarrow H$  is a homomorphism, and  $N = \ker(\varphi)$ , then there is a homomorphism  $\bar{\varphi} : G/N \rightarrow H$  such that  $\bar{\varphi} \circ \pi = \varphi$ . Moreover,  $\ker(\bar{\varphi}) = \{eN\}$ , the trivial subgroup of  $G/N$ , so  $\bar{\varphi}$  is injective. So,  $\bar{\varphi} : G/N \rightarrow \text{im}(\varphi)$  is an **isomorphism**.

- This theorem suggests that you can construct an isomorphism from an arbitrary homomorphism. First,  $\varphi$  factors through  $G/N$ , then we can include it into  $H$ .

$$G \xrightarrow{\pi} G/N \xrightarrow{\bar{\varphi}} \text{im}(\varphi) \xrightarrow{\text{inclusion}} H \quad (29)$$

**Theorem:** [(Third) Isomorphism]  $N \trianglelefteq G$  and  $H \leq G$ , then  $N \leq H \implies N \trianglelefteq G$ .

*Proof.* ????

□

**Theorem:**

$$G/H \cong G/N \big/ H/N \quad (30)$$

*Proof.* Define  $\varphi : G \rightarrow G/N \big/ H/N$  by

$$\varphi(g) = (gN)H/N \quad (31)$$

We need to show  $\varphi$  is a homomorphism. Let

$$\varphi(gg') = gg'N H/N \quad (32)$$

$$= gNg'N H/N \quad (33)$$

$$= gN H/N \cdot g'N H/N \quad (34)$$

$$= \varphi(g)\varphi(g') \quad (35)$$

$$(36)$$

□

We will then ask what is  $\ker(\varphi)$ . Suppose  $\varphi(g) = H/N$ , so  $gN H/N = H/N$ . But  $g$  is a representation for  $gN$ , so  $gH/N$  for this to be in  $H/N$  we want  $g \in H$  so  $\ker(\varphi) = H$ . An arbitrary element of  $G/N \big/ H/N$  is  $gN H/N$  for some  $g \in G$ , so  $\text{im}(\varphi) = G/N \big/ H/N$ .

- $G = \mathbb{Z}, H = 3\mathbb{Z}, K = 4\mathbb{Z}$ . By the second isomorphism theorem,  $\mathbb{Z}/3\mathbb{Z} \cong 4\mathbb{Z}/12\mathbb{Z}$ , and also  $\mathbb{Z}/4\mathbb{Z} \cong 3\mathbb{Z}/12\mathbb{Z}$ .