

Sensitivity Analysis Report

(Supplement to Section 4.4 and Table 3)

1. Sample, variables and conventions

All models are binary logistic regressions estimated by maximum likelihood on the analytical sample of $n = 320$ physicians (Section 3.4 of the manuscript). Two outcomes are used. **Own use** is 1 where the respondent reported having used AI without formal approval ($n = 95$) and 0 otherwise. **Exposure** is 1 where the respondent reported own use or observation of colleagues ($n = 161$) and 0 where no case was known ($n = 159$). The exposure outcome is a superset of the own use outcome.

Predictors. **AI provision by employer**, **AI-specific training** and **Guidelines on AI use** are binary; responses of 'do not know' or 'cannot say' are coded 0 (Section 4.3). The **governance index** is their unweighted sum (0 to 3) and replaces the three components in the specifications marked accordingly. **Male** is coded against female. **Age** enters either as one ordinal predictor across the six bands of the questionnaire, with the youngest band (20 to 29 years) coded 0 so that the odds ratio refers to a one category increase, or as five dummies with 40 to 49 years as the reference. **Setting** enters either as one dummy for hospital or medical care centre against ambulatory practice, or as two separate dummies for hospital and for medical care centre against practice.

Specifications S01 to S16 cover every combination of outcome, age coding, setting coding and governance coding. S14 and S13 are the primary and secondary models of Table 3. A1 and A2 re-estimate the Table 3 models with 'do not know' responses on tool provision and guidelines treated as missing rather than as absent, which reduces the sample to $n = 261$. Variance inflation factors for the Table 3 predictor set are given in Section 6. All figures are produced by 04_regression_main.py, 05_regression_sensitivity.py and 07_sensitivity_report.py in the code directory of the repository.

2. Overview of specifications

Spec.	Outcome	Age	Setting	Governance	N	LLR p	Pseudo R ²	AIC	BIC
S01	Exposure	categorical	separate	components	320	<.001	0.137	407.03	452.25
S02	Own use	categorical	separate	components	320	<.001	0.092	377.37	422.59
S03	Exposure	categorical	separate	index	320	<.001	0.121	409.79	447.47
S04	Own use	categorical	separate	index	320	<.001	0.089	374.56	412.25
S05	Exposure	categorical	bundled	components	320	<.001	0.136	405.21	446.66
S06	Own use	categorical	bundled	components	320	<.001	0.092	375.62	417.07
S07	Exposure	categorical	bundled	index	320	<.001	0.121	407.82	441.74
S08	Own use	categorical	bundled	index	320	<.001	0.089	372.69	406.60
S09	Exposure	ordinal	separate	components	320	<.001	0.127	403.37	433.52
S10	Own use	ordinal	separate	components	320	<.001	0.072	377.22	407.36
S11	Exposure	ordinal	separate	index	320	<.001	0.112	405.87	428.48
S12	Own use	ordinal	separate	index	320	<.001	0.068	374.59	397.20
S13 *	Exposure	ordinal	bundled	components	320	<.001	0.127	401.42	427.80
S14 *	Own use	ordinal	bundled	components	320	<.001	0.071	375.50	401.88
S15	Exposure	ordinal	bundled	index	320	<.001	0.112	403.88	422.72
S16	Own use	ordinal	bundled	index	320	<.001	0.068	372.77	391.61
A1	Own use	ordinal	bundled	components, DK missing	261	0.003	0.059	325.37	350.33
A2	Exposure	ordinal	bundled	components, DK missing	261	<.001	0.145	322.54	347.49

* Table 3 of the manuscript: S14 primary model (own use), S13 secondary model (exposure). DK missing: 'do not know' responses treated as missing.

3. Table 3 models

S14. Table 3, primary model (own use)

Outcome: Own use. Age: ordinal (youngest band coded 0). Setting: hospital and medical care centre bundled (reference: practice). Governance: three binary components.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-0.279	0.374	-0.75	0.455	0.76	[0.36, 1.57]
AI provision by employer	0.426	0.286	1.49	0.136	1.53	[0.87, 2.68]
Guidelines on AI use	-0.250	0.509	-0.49	0.623	0.78	[0.29, 2.11]
AI-specific training	0.386	0.356	1.08	0.279	1.47	[0.73, 2.96]
Male	0.562	0.269	2.09	0.037	1.75	[1.03, 2.97]
Age (ordinal, one category increase)	-0.494	0.118	-4.20	<.001	0.61	[0.48, 0.77]
Hospital or medical care centre	-0.005	0.291	-0.02	0.987	1.00	[0.56, 1.76]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.071$. AIC = 375.50. BIC = 401.88.

S13. Table 3, secondary model (exposure)

Outcome: Exposure. Age: ordinal (youngest band coded 0). Setting: hospital and medical care centre bundled (reference: practice). Governance: three binary components.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	0.694	0.346	2.00	0.045	2.00	[1.02, 3.95]
AI provision by employer	0.606	0.286	2.12	0.034	1.83	[1.05, 3.21]
Guidelines on AI use	-0.887	0.496	-1.79	0.074	0.41	[0.16, 1.09]
AI-specific training	0.323	0.358	0.90	0.367	1.38	[0.68, 2.78]
Male	0.102	0.256	0.40	0.691	1.11	[0.67, 1.83]
Age (ordinal, one category increase)	-0.527	0.107	-4.92	<.001	0.59	[0.48, 0.73]
Hospital or medical care centre	0.566	0.264	2.15	0.032	1.76	[1.05, 2.95]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.127$. AIC = 401.42. BIC = 427.80.

4. Specification grid

S01. Exposure, age categorical, setting separate, governance components

Outcome: Exposure. Age: categorical (five dummies, reference 40 to 49 years). Setting: hospital and medical care centre entered separately (reference: practice). Governance: three binary components.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-0.552	0.349	-1.58	0.114	0.58	[0.29, 1.14]
Age 20 to 29 (ref. 40 to 49)	2.201	0.814	2.70	0.007	9.03	[1.83, 44.58]
Age 30 to 39 (ref. 40 to 49)	0.650	0.380	1.71	0.087	1.91	[0.91, 4.03]
Age 50 to 59 (ref. 40 to 49)	-0.193	0.375	-0.52	0.606	0.82	[0.40, 1.72]
Age 60 to 69 (ref. 40 to 49)	-1.085	0.440	-2.47	0.014	0.34	[0.14, 0.80]
Age 70 and over (ref. 40 to 49)	-0.800	0.673	-1.19	0.235	0.45	[0.12, 1.68]
AI provision by employer	0.608	0.289	2.10	0.035	1.84	[1.04, 3.24]
Guidelines on AI use	-0.948	0.507	-1.87	0.061	0.39	[0.14, 1.05]
AI-specific training	0.380	0.362	1.05	0.293	1.46	[0.72, 2.97]
Male	0.077	0.260	0.30	0.767	1.08	[0.65, 1.80]
Hospital	0.526	0.297	1.77	0.077	1.69	[0.95, 3.03]
Medical care centre	0.726	0.446	1.63	0.104	2.07	[0.86, 4.96]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.137$. AIC = 407.03. BIC = 452.25.

S02. Own use, age categorical, setting separate, governance components

Outcome: Own use. Age: categorical (five dummies, reference 40 to 49 years). Setting: hospital and medical care centre entered separately (reference: practice). Governance: three binary components.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-1.208	0.379	-3.19	0.001	0.30	[0.14, 0.63]
Age 20 to 29 (ref. 40 to 49)	1.458	0.573	2.54	0.011	4.30	[1.40, 13.21]

Predictor	Coef.	SE	z	p	OR	95% CI
Age 30 to 39 (ref. 40 to 49)	0.142	0.398	0.36	0.721	1.15	[0.53, 2.52]
Age 50 to 59 (ref. 40 to 49)	-0.338	0.402	-0.84	0.401	0.71	[0.32, 1.57]
Age 60 to 69 (ref. 40 to 49)	-1.784	0.570	-3.13	0.002	0.17	[0.05, 0.51]
Age 70 and over (ref. 40 to 49)	-0.793	0.734	-1.08	0.280	0.45	[0.11, 1.91]
AI provision by employer	0.380	0.291	1.31	0.192	1.46	[0.83, 2.59]
Guidelines on AI use	-0.275	0.526	-0.52	0.601	0.76	[0.27, 2.13]
AI-specific training	0.431	0.364	1.18	0.236	1.54	[0.75, 3.14]
Male	0.589	0.275	2.14	0.032	1.80	[1.05, 3.09]
Hospital	0.023	0.328	0.07	0.943	1.02	[0.54, 1.95]
Medical care centre	0.278	0.488	0.57	0.569	1.32	[0.51, 3.44]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.092$. AIC = 377.37. BIC = 422.59.

S03. Exposure, age categorical, setting separate, governance index

Outcome: Exposure. Age: categorical (five dummies, reference 40 to 49 years). Setting: hospital and medical care centre entered separately (reference: practice). Governance: composite index, 0 to 3.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-0.386	0.339	-1.14	0.255	0.68	[0.35, 1.32]
Age 20 to 29 (ref. 40 to 49)	2.119	0.809	2.62	0.009	8.33	[1.70, 40.69]
Age 30 to 39 (ref. 40 to 49)	0.548	0.373	1.47	0.142	1.73	[0.83, 3.60]
Age 50 to 59 (ref. 40 to 49)	-0.275	0.369	-0.75	0.455	0.76	[0.37, 1.56]
Age 60 to 69 (ref. 40 to 49)	-1.174	0.434	-2.70	0.007	0.31	[0.13, 0.72]
Age 70 and over (ref. 40 to 49)	-0.932	0.666	-1.40	0.162	0.39	[0.11, 1.45]
Governance index (0 to 3)	0.186	0.148	1.26	0.208	1.20	[0.90, 1.61]
Male	0.079	0.257	0.31	0.758	1.08	[0.65, 1.79]
Hospital	0.503	0.293	1.72	0.086	1.65	[0.93, 2.94]
Medical care centre	0.591	0.440	1.34	0.179	1.81	[0.76, 4.28]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.121$. AIC = 409.79. BIC = 447.47.

S04. Own use, age categorical, setting separate, governance index

Outcome: Own use. Age: categorical (five dummies, reference 40 to 49 years). Setting: hospital and medical care centre entered separately (reference: practice). Governance: composite index, 0 to 3.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-1.132	0.370	-3.06	0.002	0.32	[0.16, 0.67]
Age 20 to 29 (ref. 40 to 49)	1.413	0.570	2.48	0.013	4.11	[1.35, 12.55]
Age 30 to 39 (ref. 40 to 49)	0.099	0.394	0.25	0.801	1.10	[0.51, 2.39]
Age 50 to 59 (ref. 40 to 49)	-0.377	0.399	-0.94	0.345	0.69	[0.31, 1.50]
Age 60 to 69 (ref. 40 to 49)	-1.842	0.568	-3.24	0.001	0.16	[0.05, 0.48]
Age 70 and over (ref. 40 to 49)	-0.871	0.729	-1.19	0.233	0.42	[0.10, 1.75]
Governance index (0 to 3)	0.259	0.156	1.66	0.097	1.30	[0.95, 1.76]
Male	0.585	0.274	2.13	0.033	1.79	[1.05, 3.07]
Hospital	0.020	0.325	0.06	0.950	1.02	[0.54, 1.93]
Medical care centre	0.203	0.486	0.42	0.677	1.22	[0.47, 3.17]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.089$. AIC = 374.56. BIC = 412.25.

S05. Exposure, age categorical, setting bundled, governance components

Outcome: Exposure. Age: categorical (five dummies, reference 40 to 49 years). Setting: hospital and medical care centre bundled (reference: practice). Governance: three binary components.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-0.549	0.349	-1.57	0.116	0.58	[0.29, 1.14]
Age 20 to 29 (ref. 40 to 49)	2.157	0.807	2.67	0.008	8.64	[1.78, 42.06]
Age 30 to 39 (ref. 40 to 49)	0.609	0.367	1.66	0.097	1.84	[0.90, 3.78]
Age 50 to 59 (ref. 40 to 49)	-0.202	0.374	-0.54	0.589	0.82	[0.39, 1.70]
Age 60 to 69 (ref. 40 to 49)	-1.078	0.439	-2.46	0.014	0.34	[0.14, 0.80]

Predictor	Coef.	SE	z	p	OR	95% CI
Age 70 and over (ref. 40 to 49)	-0.816	0.672	-1.21	0.225	0.44	[0.12, 1.65]
AI provision by employer	0.606	0.289	2.10	0.036	1.83	[1.04, 3.23]
Guidelines on AI use	-0.924	0.502	-1.84	0.066	0.40	[0.15, 1.06]
AI-specific training	0.383	0.361	1.06	0.289	1.47	[0.72, 2.98]
Male	0.088	0.259	0.34	0.734	1.09	[0.66, 1.81]
Hospital or medical care centre	0.576	0.272	2.12	0.034	1.78	[1.04, 3.03]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.136$. AIC = 405.21. BIC = 446.66.

S06. Own use, age categorical, setting bundled, governance components

Outcome: Own use. Age: categorical (five dummies, reference 40 to 49 years). Setting: hospital and medical care centre bundled (reference: practice). Governance: three binary components.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-1.201	0.379	-3.17	0.002	0.30	[0.14, 0.63]
Age 20 to 29 (ref. 40 to 49)	1.399	0.560	2.50	0.012	4.05	[1.35, 12.14]
Age 30 to 39 (ref. 40 to 49)	0.088	0.383	0.23	0.817	1.09	[0.52, 2.31]
Age 50 to 59 (ref. 40 to 49)	-0.351	0.401	-0.88	0.381	0.70	[0.32, 1.54]
Age 60 to 69 (ref. 40 to 49)	-1.777	0.569	-3.12	0.002	0.17	[0.06, 0.52]
Age 70 and over (ref. 40 to 49)	-0.815	0.732	-1.11	0.266	0.44	[0.11, 1.86]
AI provision by employer	0.375	0.291	1.29	0.198	1.46	[0.82, 2.57]
Guidelines on AI use	-0.237	0.518	-0.46	0.648	0.79	[0.29, 2.18]
AI-specific training	0.435	0.364	1.19	0.233	1.54	[0.76, 3.15]
Male	0.601	0.274	2.20	0.028	1.82	[1.07, 3.12]
Hospital or medical care centre	0.085	0.302	0.28	0.778	1.09	[0.60, 1.97]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.092$. AIC = 375.62. BIC = 417.07.

S07. Exposure, age categorical, setting bundled, governance index

Outcome: Exposure. Age: categorical (five dummies, reference 40 to 49 years). Setting: hospital and medical care centre bundled (reference: practice). Governance: composite index, 0 to 3.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-0.385	0.338	-1.14	0.255	0.68	[0.35, 1.32]
Age 20 to 29 (ref. 40 to 49)	2.100	0.803	2.62	0.009	8.17	[1.69, 39.43]
Age 30 to 39 (ref. 40 to 49)	0.531	0.362	1.47	0.142	1.70	[0.84, 3.45]
Age 50 to 59 (ref. 40 to 49)	-0.279	0.368	-0.76	0.449	0.76	[0.37, 1.56]
Age 60 to 69 (ref. 40 to 49)	-1.171	0.434	-2.70	0.007	0.31	[0.13, 0.73]
Age 70 and over (ref. 40 to 49)	-0.938	0.665	-1.41	0.159	0.39	[0.11, 1.44]
Governance index (0 to 3)	0.188	0.147	1.28	0.202	1.21	[0.90, 1.61]
Male	0.083	0.256	0.33	0.745	1.09	[0.66, 1.80]
Hospital or medical care centre	0.525	0.268	1.96	0.050	1.69	[1.00, 2.86]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.121$. AIC = 407.82. BIC = 441.74.

S08. Own use, age categorical, setting bundled, governance index

Outcome: Own use. Age: categorical (five dummies, reference 40 to 49 years). Setting: hospital and medical care centre bundled (reference: practice). Governance: composite index, 0 to 3.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-1.130	0.370	-3.05	0.002	0.32	[0.16, 0.67]
Age 20 to 29 (ref. 40 to 49)	1.373	0.557	2.46	0.014	3.95	[1.32, 11.76]
Age 30 to 39 (ref. 40 to 49)	0.062	0.380	0.16	0.870	1.06	[0.51, 2.24]
Age 50 to 59 (ref. 40 to 49)	-0.386	0.398	-0.97	0.333	0.68	[0.31, 1.48]
Age 60 to 69 (ref. 40 to 49)	-1.837	0.567	-3.24	0.001	0.16	[0.05, 0.48]
Age 70 and over (ref. 40 to 49)	-0.885	0.728	-1.21	0.224	0.41	[0.10, 1.72]
Governance index (0 to 3)	0.264	0.155	1.70	0.089	1.30	[0.96, 1.77]
Male	0.594	0.273	2.18	0.030	1.81	[1.06, 3.09]
Hospital or medical care centre	0.064	0.300	0.21	0.830	1.07	[0.59, 1.92]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.089$. AIC = 372.69. BIC = 406.60.

S09. Exposure, age ordinal, setting separate, governance components

Outcome: Exposure. Age: ordinal (youngest band coded 0). Setting: hospital and medical care centre entered separately (reference: practice). Governance: three binary components.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	0.715	0.358	1.99	0.046	2.04	[1.01, 4.12]
AI provision by employer	0.606	0.286	2.12	0.034	1.83	[1.05, 3.21]
Guidelines on AI use	-0.898	0.500	-1.80	0.072	0.41	[0.15, 1.08]
AI-specific training	0.320	0.358	0.89	0.372	1.38	[0.68, 2.78]
Male	0.097	0.257	0.38	0.706	1.10	[0.67, 1.82]
Age (ordinal, one category increase)	-0.533	0.111	-4.82	<.001	0.59	[0.47, 0.73]
Hospital	0.541	0.286	1.89	0.059	1.72	[0.98, 3.01]
Medical care centre	0.645	0.437	1.48	0.140	1.91	[0.81, 4.49]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.127$. AIC = 403.37. BIC = 433.52.

S10. Own use, age ordinal, setting separate, governance components

Outcome: Own use. Age: ordinal (youngest band coded 0). Setting: hospital and medical care centre entered separately (reference: practice). Governance: three binary components.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-0.225	0.387	-0.58	0.561	0.80	[0.37, 1.71]
AI provision by employer	0.427	0.286	1.49	0.136	1.53	[0.87, 2.68]
Guidelines on AI use	-0.282	0.514	-0.55	0.583	0.75	[0.28, 2.07]
AI-specific training	0.380	0.356	1.07	0.287	1.46	[0.73, 2.94]
Male	0.551	0.270	2.04	0.042	1.73	[1.02, 2.95]
Age (ordinal, one category increase)	-0.510	0.122	-4.18	<.001	0.60	[0.47, 0.76]
Hospital	-0.067	0.314	-0.21	0.832	0.94	[0.51, 1.73]
Medical care centre	0.194	0.470	0.41	0.679	1.21	[0.48, 3.05]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.072$. AIC = 377.22. BIC = 407.36.

S11. Exposure, age ordinal, setting separate, governance index

Outcome: Exposure. Age: ordinal (youngest band coded 0). Setting: hospital and medical care centre entered separately (reference: practice). Governance: composite index, 0 to 3.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	0.821	0.354	2.32	0.020	2.27	[1.14, 4.54]
Governance index (0 to 3)	0.178	0.145	1.23	0.220	1.19	[0.90, 1.59]
Male	0.086	0.254	0.34	0.736	1.09	[0.66, 1.79]
Age (ordinal, one category increase)	-0.537	0.110	-4.90	<.001	0.58	[0.47, 0.72]
Hospital	0.505	0.282	1.79	0.073	1.66	[0.95, 2.88]
Medical care centre	0.531	0.431	1.23	0.218	1.70	[0.73, 3.95]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.112$. AIC = 405.87. BIC = 428.48.

S12. Own use, age ordinal, setting separate, governance index

Outcome: Own use. Age: ordinal (youngest band coded 0). Setting: hospital and medical care centre entered separately (reference: practice). Governance: composite index, 0 to 3.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-0.170	0.383	-0.45	0.656	0.84	[0.40, 1.78]
Governance index (0 to 3)	0.260	0.152	1.71	0.087	1.30	[0.96, 1.75]
Male	0.539	0.270	2.00	0.045	1.72	[1.01, 2.91]
Age (ordinal, one category increase)	-0.514	0.122	-4.23	<.001	0.60	[0.47, 0.76]
Hospital	-0.076	0.312	-0.24	0.808	0.93	[0.50, 1.71]
Medical care centre	0.131	0.469	0.28	0.780	1.14	[0.45, 2.86]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.068$. AIC = 374.59. BIC = 397.20.

S15. Exposure, age ordinal, setting bundled, governance index

Outcome: Exposure. Age: ordinal (youngest band coded 0). Setting: hospital and medical care centre bundled (reference: practice). Governance: composite index, 0 to 3.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	0.816	0.342	2.39	0.017	2.26	[1.16, 4.42]
Governance index (0 to 3)	0.178	0.144	1.24	0.216	1.20	[0.90, 1.59]
Male	0.087	0.254	0.34	0.732	1.09	[0.66, 1.79]
Age (ordinal, one category increase)	-0.536	0.107	-5.03	<.001	0.59	[0.47, 0.72]
Hospital or medical care centre	0.511	0.260	1.97	0.049	1.67	[1.00, 2.78]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.112$. AIC = 403.88. BIC = 422.72.

S16. Own use, age ordinal, setting bundled, governance index

Outcome: Own use. Age: ordinal (youngest band coded 0). Setting: hospital and medical care centre bundled (reference: practice). Governance: composite index, 0 to 3.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-0.214	0.368	-0.58	0.560	0.81	[0.39, 1.66]
Governance index (0 to 3)	0.267	0.151	1.77	0.077	1.31	[0.97, 1.76]
Male	0.550	0.269	2.05	0.041	1.73	[1.02, 2.93]
Age (ordinal, one category increase)	-0.501	0.118	-4.27	<.001	0.61	[0.48, 0.76]
Hospital or medical care centre	-0.027	0.289	-0.09	0.926	0.97	[0.55, 1.72]

N = 320. LLR test $p < .001$. McFadden pseudo $R^2 = 0.068$. AIC = 372.77. BIC = 391.61.

5. Additional specifications: 'do not know' treated as missing

Respondents answering 'do not know' or 'cannot say' on tool provision ($n = 43$) or 'do not know' on guidelines ($n = 27$) are excluded (59 respondents, some answering both), leaving $n = 261$. Predictor set and coding otherwise as in Table 3.

A1. Own use, Table 3 specification, $n = 261$

Table 3 specification; 'do not know' responses on tool provision and guidelines treated as missing.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	-0.160	0.393	-0.41	0.684	0.85	[0.39, 1.84]
AI provision by employer	0.366	0.297	1.23	0.218	1.44	[0.81, 2.58]
Guidelines on AI use	-0.315	0.518	-0.61	0.543	0.73	[0.26, 2.01]
AI-specific training	0.334	0.369	0.91	0.365	1.40	[0.68, 2.88]
Male	0.304	0.284	1.07	0.285	1.35	[0.78, 2.36]
Age (ordinal, one category increase)	-0.424	0.122	-3.48	<.001	0.65	[0.52, 0.83]
Hospital or medical care centre	0.104	0.305	0.34	0.732	1.11	[0.61, 2.02]

N = 261. LLR test $p = 0.003$. McFadden pseudo $R^2 = 0.059$. AIC = 325.37. BIC = 350.33.

A2. Exposure, Table 3 specification, $n = 261$

Table 3 specification; 'do not know' responses on tool provision and guidelines treated as missing.

Predictor	Coef.	SE	z	p	OR	95% CI
Intercept	0.936	0.386	2.43	0.015	2.55	[1.20, 5.44]
AI provision by employer	0.546	0.309	1.77	0.077	1.73	[0.94, 3.17]
Guidelines on AI use	-1.005	0.521	-1.93	0.054	0.37	[0.13, 1.02]
AI-specific training	0.286	0.386	0.74	0.460	1.33	[0.62, 2.84]
Male	-0.030	0.284	-0.11	0.915	0.97	[0.56, 1.69]
Age (ordinal, one category increase)	-0.540	0.118	-4.56	<.001	0.58	[0.46, 0.74]
Hospital or medical care centre	0.718	0.292	2.45	0.014	2.05	[1.16, 3.63]

N = 261. LLR test $p < .001$. McFadden pseudo $R^2 = 0.145$. AIC = 322.54. BIC = 347.49.

6. Variance inflation factors, Table 3 predictor set

Predictor	VIF
AI provision by employer	1.13
Guidelines on AI use	1.22
AI-specific training	1.27
Male	1.10
Age (ordinal, one category increase)	1.32
Hospital or medical care centre	1.23

Maximum VIF = 1.32. The predictors are identical in both Table 3 models, so the factors apply to both.