

*Basic combinatory objects, uniform preorders and partial
combinatory algebras*

Jonas Frey

Category Theory Octoberfest

29 October 2022

Dedicated to the Memory of Pieter Hofstra

Remembering Pieter



Overview

Pieter Hofstra's Basic Combinatory objects

Basic Combinatory Objects

In his 2006 paper “All realizability is relative”¹, Pieter Hofstra introduced the notion of *basic combinatory object* (building on work with van Oosten on *ordered combinatory algebras*).

Definition

A **basic combinatory object (BCO)** is a set A equipped with a partial order \leq and a set \mathcal{F}_A of ‘computable’ partial endofunctions with down-closed domain, such that

1. $\exists i \in \mathcal{F} \forall a \in A. i(a) \leq a$
2. $\forall f, g \in \mathcal{F} \exists h \in \mathcal{F} \forall a \in \text{dom}(g \circ f). h(a) \leq g(f(a))$

BCOs form a locally ordered category **BCO** which admits a full embedding

$$\text{fam} : \mathbf{BCO} \hookrightarrow \mathbf{IOrd}$$

into the locally ordered category $\mathbf{IOrd} = [\mathbf{Set}^{\text{op}}, \mathbf{Ord}]$ of **Set-indexed preorders**, given by $\text{fam}(A)(J) = (A^J, \leq)$ where

$$\varphi \leq \psi \quad :\Leftrightarrow \quad \exists f \in \mathcal{F}_A \forall j \in J. f(\varphi(j)) \leq \psi(j)$$

for $\varphi, \psi : J \rightarrow A$.

¹Hofstra, Pieter JW. “All realizability is relative.” *Mathematical Proceedings of the Cambridge Philosophical Society*. Vol. 141. No. 2. Cambridge University Press, 2006.

Basic Combinatory objects II – finite meets

BCOs are closed under products in $[\mathbf{Set}^{\text{op}}, \mathbf{Ord}]$, thus $\mathbf{fam}(A)$ is an indexed meet-semilattice iff

$$A \rightarrow A \times A \quad \text{and} \quad A \rightarrow 1$$

have right adjoints

$$(- \wedge -) : A \rightarrow A \times A \quad \text{and} \quad \top : A \rightarrow 1$$

in **BCO**. We call such BCOs **cartesian**.

Basic Combinatory objects III – existential quantification

- Say that an indexed preorder $P : \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Ord}$ admits **existential quantification**, if the reindexing maps $f^* : P(I) \rightarrow P(J)$ have left adjoints $\exists_f : P(J) \rightarrow P(I)$ for all $f : J \rightarrow I$, subject to the **Beck–Chevalley condition**.
- Denote by $\exists\text{-}\mathbf{IOrd}$ the subcategory of \mathbf{IOrd} on indexed preorders admitting \exists and indexed monotone maps preserving \exists .
- Pieter Hofstra showed that
 1. the forgetful functor $\exists\text{-}\mathbf{IOrd} \rightarrow \mathbf{IOrd}$ is 2-monadic, and
 2. the induced (lax idempotent) 2-monad $D : \mathbf{IOrd} \rightarrow \mathbf{IOrd}$ restricts to \mathbf{BCO} .

$$\begin{array}{ccc} \mathbf{BCO} & \overset{D}{\dashrightarrow} & \mathbf{BCO} \\ \text{fam} \downarrow & & \downarrow \text{fam} \\ \mathbf{IOrd} & \xrightarrow{D} & \mathbf{IOrd} \end{array}$$

For a BCO A , the carrier of $D(A)$ is the set of **downsets**.

3. Furthermore, D plays well with finite meets: if P has finite meets then $D(P)$ has finite meets and moreover it satisfies the **Frobenius condition**.

Examples: BCOs from posets and (O)PCAs

- Every poset can be viewed as BCO where only the identity function is computable, which gives a full embedding

$$\mathbf{Pos} \rightarrow \mathbf{BCO}$$

- Every **PCA** \mathcal{A} can be viewed as a *cartesian* BCO where the ordering is trivial and

$$\mathcal{F}_{\mathcal{A}} = \{e \cdot (-) : \mathcal{A} \rightarrow \mathcal{A} \mid e \in \mathcal{A}\}.$$

- More generally, **filtered ordered PCAs** \mathcal{A} can be viewed as cartesian BCO with

$$\mathcal{F}_{\mathcal{A}} = \{e \cdot (-) : \mathcal{A} \rightarrow \mathcal{A} \mid e \in \Phi_{\mathcal{A}}\}.$$

- Pieter observed that in both cases the associated **realizability tripos** $\mathbf{rt}(\mathcal{A}) : \mathbf{Set}^{\mathbf{op}} \rightarrow \mathbf{Ord}$ is given by

$$\mathbf{rt}(\mathcal{A}) = D(\mathbf{fam}(\mathcal{A})) = \mathbf{fam}(D(\mathcal{A}))$$

- In particular this means that **realizability triposes are freely generated under existential quantification!**
- This is related to the fact that realizability toposes are ex/lex-completions.

Characterizing filtered OPCAs among BCOs

Theorem (Hofstra)

TFAE for a cartesian BCO A :

1. A is (induced by) a filtered OPCA.
2. $\mathbf{fam}(D(A))$ is a tripos.
3. The fibers of $D\mathbf{fam}(A)$ are Heyting algebras.

Proof.

The implications $1 \Rightarrow 2 \Rightarrow 3$ are clear.

For $3 \Rightarrow 1$ let $\iota \in \mathbf{fam}(D(A))(A)$ be the function sending every a to its principal downset, and let $\varepsilon \in \mathcal{F}_A$ be a witness of the inequality

$$(\pi_1(\iota) \Rightarrow \pi_2(\iota)) \wedge \pi_1(\iota) \leq \pi_2(\iota)$$

in $\mathbf{fam}(D(A))(A \times A)$. Then the application operation of the OPCA is given by $\varepsilon \circ \wedge$.

The filter Φ_A is given by the **designated truth values**, i.e. the $a \in A$ that are equivalent to \top in $\mathbf{fam}(A)(1)$.



Two variations : DCOs and uniform preorders

Overview

- Pieter's paper formed the starting point for my PhD thesis in which I gave characterizations of realizability triposes and toposes over PCAs.
- In hindsight, the only missing piece in the BCO-approach is that the image of $\mathbf{BCO} \hookrightarrow \mathbf{IOrd}$ does not have an easy characterization — if we could characterize (O)PCAs among BCOs and BCOs among indexed posets then we could characterize (O)PCAs among indexed preorders.
- In the following I introduce a sub- and a super-category of \mathbf{BCO} , which *do* have simple characterizations as subcategories of \mathbf{IOrd} , and explain how to adapt Pieter's techniques.

Discrete combinatory objects

Definition

A **discrete combinatory object (DCO)** is simply a BCO whose partial order structure is trivial. We write $\mathbf{DCO} \subseteq \mathbf{BCO}$ for the full subcategory of DCOs.

Definition

Given an indexed preorder $P : \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Ord}$, we call $\delta \in P(J)$ **discrete**, if it is right orthogonal to all cartesian maps over surjections in the total category $\int P$ of P .

Lemma

An indexed preorder $P : \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Ord}$ is equivalent to one of the form $\text{fam}(A)$ for a DCO A , iff it has a discrete generic predicate.

Proof.

Given a discrete predicate $\delta \in \mathcal{A}$, define DCO structure on A by taking as computable those partial functions $f : A \rightarrow A$ satisfying $\iota|_{\text{dom}(f)} \leq f^*(\iota)$ in $P(\text{dom}(f))$. □

Characterizing $\mathbf{fam}(\mathcal{A})$

We immediately get the following.

Lemma

An indexed meet-semilattice $\mathcal{H} : \mathbf{Set}^{\mathbf{op}} \rightarrow \mathbf{Ord}$ is equivalent to one of the form $\mathbf{fam}(\mathcal{A})$ for a filtered PCA \mathcal{A} iff it has a discrete generic predicate and $D(\mathcal{H})$ is a tripos. The filter is trivial iff $\mathcal{H}(1) \simeq 1$.

- Filtered PCAs are better known as **inclusions of PCAs**, their realizability toposes are called **relative realizability toposes**.
- To be able to characterize realizability *triposes*, we have to reconstruct \mathcal{H} from $D(\mathcal{H})$. This is what we do next.

\exists -prime predicates

Motivation/Analogy:

- Given a poset P , the set of $D(P)$ of **downsets** in P is a complete lattice under inclusion, and is in fact the *free sup-lattice* on P .
- The **principal downsets** $\downarrow x = \{y \in P \mid y \leq x\}$ can be characterized as **completely join-prime elements** in D – an element x of a sup-lattice L is called completely join-prime if we have

$$x \leq \bigvee_{j \in J} y_j \quad \Rightarrow \quad \exists j \in J. x \leq y_j$$

for all families $(y_j)_{j \in J}$ of elements.

- L is a free join-completion iff it has **enough** completely join-prime elements, i.e. if every element is a join of completely-join-primes.

\exists -prime predicates

Definition

Given an indexed preorder \mathcal{H} which admits existential quantification, a predicate $\pi \in \mathcal{H}(I)$ is called **\exists -prime** if for all functions $I \xleftarrow{u} J \xleftarrow{v} K$ and predicates $\theta \in \mathcal{H}(K)$ such that $u^*\pi \leq \exists_v \theta$, there exists a section s of v such that $u^*\pi \leq s^*\theta$.

Lemma

An indexed preorder \mathcal{H} is an \exists -completion iff it has enough \exists -prime predicates, i.e. if for every predicate $\varphi \in \mathcal{H}(I)$ there exists a function $u : J \rightarrow I$ and an \exists -prime predicate $\pi \in \mathcal{H}(J)$ with $\varphi \cong \exists_u \pi$.

In this case, we have $\mathcal{H} \simeq D(\mathcal{P})$ where $\mathcal{P} \subseteq \mathcal{H}$ is the indexed sub-preorder on \exists -prime predicates.

With this we can characterize (relative) realizability triposes!

Characterizing realizability triposes

Theorem

A tripos $\mathcal{H} : \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Ord}$ is a relative realizability tripos over an inclusion of PCAs, iff

1. \mathcal{H} has enough \exists -prime predicates, and
2. the full indexed sup-preorder $\mathcal{P} \subseteq \mathcal{H}$ on \exists -prime predicates is closed under finite meets and has a discrete generic predicate δ .

- The discreteness condition on δ can be stated in \mathcal{H} rather than \mathcal{P} , which is a slight strengthening.
- We get ordinary (non-relative) realizability if the tripos is **2-valued**, i.e. $\mathcal{H}(1) \simeq \mathbf{Bool}$.