

A refinement of Gabriel-Ulmer duality

Jonas Frey
CMU Pittsburgh

Cartmell’s generalized algebraic theories [Car86] – which extend algebra by type dependency – and Freyd’s essentially algebraic theories [Fre72] – which permit a controlled form of partiality – are commonly recognized as being equally expressive, and are both subsumed by the less syntactic and more abstract *finite-limit theories*, which are simply small finite-limit categories.

More specifically, for each generalized/essentially algebraic theory \mathcal{T} there exists a small finite-limit category \mathbb{C} such that

$$\mathrm{Mod}(\mathcal{T}) \simeq \mathbf{Lex}(\mathbb{C}, \mathbf{Set}),$$

i.e. the category of models of \mathcal{T} is equivalent to the category of finite-limit-preserving (‘lex’) functors from \mathbb{C} to the category of sets.

Classical Gabriel-Ulmer duality [GU71] states that the categories of models of such theories are precisely the *locally finitely presentable categories*, giving rise to a contravariant biequivalence

$$\mathbf{Lex} \simeq \mathbf{LFP}^{\mathrm{op}}$$

between the 2-categories of finite-limit categories and locally finitely presentable categories.

We propose a refinement of this duality based on the notion of *clan*, which was introduced by Taylor under the name ‘category with display maps’ [Tay87, 4.3.2], and later renamed by Joyal [Joy17, 1.1.1].

Definition 1 *A clan is a small category \mathcal{T} with terminal object 1 equipped with a class \mathcal{D} of display maps such that pullbacks of display maps along arbitrary maps exist and are again display maps, and display maps contain isomorphisms and terminal projections and are closed under composition.*

A model of a clan \mathcal{T} is a functor $A : \mathcal{T} \rightarrow \mathbf{Set}$ which preserves 1 and pullbacks of display maps. We denote the full subcategory of $[\mathcal{T}, \mathbf{Set}]$ on models by $\mathrm{Mod}(\mathcal{T})$.

Since corepresentable functors $\mathcal{T}(\Gamma, -)$ preserve all limits, the dual Yoneda embedding co-restricts to a fully faithful functor

$$Z : \mathcal{T}^{\mathrm{op}} \rightarrow \mathrm{Mod}(\mathcal{T}).$$

Clans refine finite-limit theories in that the ‘same’ finite-limit theory can be represented by different clans, thus we cannot expect to reconstruct a clan from its category of models alone.

To recover a duality we equip the category $\mathrm{Mod}(\mathcal{T})$ of models of \mathcal{T} with additional information in form of a weak factorization system $(\mathcal{E}, \mathcal{F})$ which is cofibrantly generated by the set

$$\mathcal{E}_0 = \{Z(f) \mid f \in \mathcal{D}\},$$

of morphisms¹. We call maps in \mathcal{F} *full maps*, maps in \mathcal{E} *extensions*, and models A such that $(0 \rightarrow A) \in \mathcal{E}$ *0-extensions*.

Using an instance of the *fat small object argument* [MRV14] we show that (the opposite of) \mathcal{T} can be recovered up to Cauchy completion from $\mathrm{Mod}(\mathcal{T})$ and $(\mathcal{E}, \mathcal{F})$ as the full subcategory of $\mathrm{Mod}(\mathcal{T})$ on *finitely presentable 0-extensions*, i.e. finitely presentable models A such that $(0 \rightarrow A) \in \mathcal{E}$.

Moreover, we give a characterization of those locally finitely presentable categories equipped with weak factorization systems that arise as categories of models of clans:

Proposition 1 *A lfp category \mathfrak{A} with wfs $(\mathcal{E}, \mathcal{F})$ is clan-algebraic, if*

¹The same w.f.s. construction was independently proposed by Simon Henry in a HoTTEST seminar.

1. the full subcategory \mathbb{C} on $\{A \in \mathfrak{A} \mid A \text{ finitely presented } 0\text{-extension}\}$ is dense in \mathfrak{A} ,
2. $(\mathcal{E}, \mathcal{F})$ is cofibrantly generated by $\mathcal{E} \cap \mathbf{mor}(\mathbb{C})$, and
3. for $C \in \mathbb{C}$ and $A \in \mathbf{Mod}(\mathbb{C}^{\mathrm{op}})$, the functor $\mathfrak{A}(C, -) : \mathfrak{A} \rightarrow \mathbf{Set}$ preserves $\mathrm{colim}(\int A \rightarrow \mathbb{C} \rightarrow \mathfrak{A})$.

We conjecture a simplification of condition 3 which refines the characterization of algebraic categories in [ARV10, Corollary 18.4]:

Conjecture 1 *Condition 3 of the theorem is implied by \mathfrak{A} having full and effective quotients of component-wise full equivalence relations.*

Finally we showcase a major advantage of clans over finite-limit theories, which is that different clanic representations of the same finite-limit theory when interpreted in higher types give rise to different ‘homotopy-coherent generalizations’ of the set-level algebraic structure.

For example, the finite-limit theory of categories admits (at least) 4 clanic refinements, whose models in higher types are respectively *Segal spaces*, *Segal categories*, *pre-categories* (in the sense of [Uni13]), and ‘discrete’ *1-categories*. This can be seen as a refinement of Cescavicius and Scholze’s concept of *animated* model of an algebraic theory [CS19].

Acknowledgement

The presented work is supported by the Air Force Office of Scientific Research under award number FA9550-20-1-0305, and by the U. S. Army Research Office under grant number W911NF-21-1-0121.

References

- [ARV10] J. Adámek, J. Rosický, and E.M. Vitale, *Algebraic theories: a categorical introduction to general algebra*, vol. 184, Cambridge University Press, 2010.
- [Car86] J. Cartmell, *Generalised algebraic theories and contextual categories*, Annals of pure and applied logic **32** (1986), 209–243.
- [CS19] K. Cescavicius and P. Scholze, *Purity for flat cohomology*, arXiv preprint arXiv:1912.10932 (2019).
- [Fre72] P. Freyd, *Aspects of topoi*, Bulletin of the Australian Mathematical Society **7** (1972), no. 1, 1–76.
- [GU71] P. Gabriel and F. Ulmer, *Lokal präsentierbare kategorien*, Lecture Notes in Math., vol. 221, Springer-Verlag, 1971.
- [Joy17] A. Joyal, *Notes on clans and tribes*, arXiv preprint arXiv:1710.10238 (2017).
- [MRV14] M. Makkai, J. Rosicky, and L. Vokrinek, *On a fat small object argument*, Advances in Mathematics **254** (2014), 49–68.
- [Tay87] P. Taylor, *Recursive domains, indexed category theory and polymorphism*, Ph.D. thesis, University of Cambridge, 1987.
- [Uni13] The Univalent Foundations Program, *Homotopy type theory: Univalent foundations of mathematics*, <https://homotopytypetheory.org/book>, Institute for Advanced Study, 2013.