



Universität Augsburg
Mathematisch-Naturwissenschaftlich-
Technische Fakultät

Classical networks for the Hubbard model with a tilted potential

- Master Thesis Colloquium -

Jonas Kell
Chair for theoretical Physics III

12th of March 2025

2025-03-12

Master Colloquium Presentation

1. Thank you for coming
2. Title: Classical networks for the Hubbard model with a tilted potential
3. Quickly what this will be about

Outline

- 1 Introduction of the Physical Problem
- 2 Time Evolution
- 3 Sampling & Simplifications
- 4 Variational Classical Networks
- 5 Implementation & Experiments
- 6 Conclusion & Outlook



2025-03-12

Master Colloquium Presentation

Outline

1.
 - Introduction to the physical theory
 - In depth: Time evolution - why and how
 - Quickly: Sampling Strategy and simplifications
 - To give the title justice: Variational classical networks - what is all about
 - End (if there is time or if questions lead into that direction): Implementation and experiments
 - Outlook

Outline

- 1 Introduction of the Physical Problem
- 2 Time Evolution
- 3 Sampling & Simplifications
- 4 Variational Classical Networks
- 5 Implementation & Experiments
- 6 Conclusion & Outlook

Particle Types

Introduction of the Physical Problem

- Bosonic operators
 - Commutation relations

$$\begin{aligned}\left[\hat{b}_l, \hat{b}_m\right] &= 0 \\ \left[\hat{b}_l^\dagger, \hat{b}_m^\dagger\right] &= 0 \\ \left[\hat{b}_l, \hat{b}_m^\dagger\right] &= \delta_{l,m}\end{aligned}$$



2025-03-12

Master Colloquium Presentation

- └ Introduction of the Physical Problem
 - └ Particle Types

1. Wanna discuss, because during the writing of the thesis quite new insight in what in fact the differences of behavior are in between the particle types
2. First of all wanna point out how we have lattice operators, so fixed sites where there can be particles
3. Occupation number from 0 to infinity
4. Nice to handle because they commute

Particle Types

Introduction of the Physical Problem

- Bosonic operators
- Commutation relations

$$\begin{aligned}\left[\hat{b}_l, \hat{b}_m\right] &= 0 \\ \left[\hat{b}_l^\dagger, \hat{b}_m^\dagger\right] &= 0 \\ \left[\hat{b}_l, \hat{b}_m^\dagger\right] &= \delta_{l,m}\end{aligned}$$

Particle Types

Introduction of the Physical Problem

- Bosonic operators
 - Commutation relations
- Fermionic operators
 - Anti-commutation relations

$$\{\hat{c}_l, \hat{c}_m\} = 0$$

$$\{\hat{c}_l^\dagger, \hat{c}_m^\dagger\} = 0$$

$$\{\hat{c}_l, \hat{c}_m^\dagger\} = \delta_{l,m}$$



2025-03-12

Master Colloquium Presentation

└ Introduction of the Physical Problem

└ Particle Types

1. Occupation number only 0 and one: nice for computational handling
2. Anti-commute: not so nice to handle in the preparation

Particle Types

Introduction of the Physical Problem

- Bosonic operators
- Commutation relations
- Fermionic operators
- Anti-commutation relations

$$\{\hat{c}_l, \hat{c}_m\} = 0$$
$$\{\hat{c}_l^\dagger, \hat{c}_m^\dagger\} = 0$$
$$\{\hat{c}_l, \hat{c}_m^\dagger\} = \delta_{l,m}$$

Particle Types

Introduction of the Physical Problem

- Bosonic operators
 - Commutation relations
- Fermionic operators
 - Anti-commutation relations
- Hard-core bosonic operators
 - Commutation relations
 - Occupation number limited to 0 and 1
 - Number operator is idempotent

$$[\hat{h}_l, \hat{h}_m] = 0$$

$$[\hat{h}_l^\dagger, \hat{h}_m^\dagger] = 0$$

$$[\hat{h}_l, \hat{h}_m^\dagger] = \left(1 - 2 \cdot \hat{h}_m^\dagger \hat{h}_m\right) \cdot \delta_{l,m}$$

$$\left(\hat{h}_l^\dagger \hat{h}_l\right)^n = \hat{h}_l^\dagger \hat{h}_l \quad \forall n \in \mathbb{N}$$



2025-03-12

Master Colloquium Presentation

└ Introduction of the Physical Problem

└ Particle Types

1. Hard-core bosons: best fo both worlds
2. Like the fermions nice because only occupation number 0 and 1
3. but COMMUTATION relations, because of that we will be able to generate an important mapping
4. Finally: number operator is *idempotent* (power = potenz in english) which we need for one mathematical step later

Particle Types

Introduction of the Physical Problem

- Bosonic operators
 - Commutation relations
- Fermionic operators
 - Anti-commutation relations
- Hard-core bosonic operators
 - Commutation relations
 - Occupation number limited to 0 and 1
 - Number operator is idempotent

$$[\hat{h}_l, \hat{h}_m] = 0$$

$$[\hat{h}_l^\dagger, \hat{h}_m^\dagger] = 0$$

$$[\hat{h}_l, \hat{h}_m^\dagger] = \left(1 - 2 \cdot \hat{h}_m^\dagger \hat{h}_m\right) \cdot \delta_{l,m}$$

$$\left(\hat{h}_l^\dagger \hat{h}_l\right)^n = \hat{h}_l^\dagger \hat{h}_l \quad \forall n \in \mathbb{N}$$

Particle Types

Introduction of the Physical Problem

- Bosonic operators
 - Commutation relations
- Fermionic operators
 - Anti-commutation relations
- Hard-core bosonic operators
 - Commutation relations
 - Occupation number limited to 0 and 1
 - Number operator is idempotent
- Spins (Pauli operators/matrices)
 - Particles from different sites commute
 - Different spin directions on the same site anti-commute
 - They obey angular momentum commutation relations

$$\hat{\sigma}^0 = \mathbb{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\hat{\sigma}^2 = \hat{\sigma}^y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$

$$\hat{\sigma}^1 = \hat{\sigma}^x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$\hat{\sigma}^3 = \hat{\sigma}^z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\left\{ \hat{\sigma}_l^\alpha, \hat{\sigma}_l^\beta \right\} = 2 \cdot \delta_{\alpha, \beta}$$

$$\left[\hat{\sigma}_l^\alpha, \hat{\sigma}_m^\beta \right] = 2 \cdot i \cdot \varepsilon_{\alpha, \beta, \gamma} \cdot \delta_{l, m} \cdot \hat{\sigma}_l^\gamma$$



2025-03-12

Master Colloquium Presentation

└ Introduction of the Physical Problem

└ Particle Types

1. Final particle type: spins - watch the 1/2 if it is a pauli or spin operator
2. Has also mostly commutative properties (also mainly *angular momentum* relations), but also one anti-commutation relation because of the properties of the Pauli matrices
3. Will be necessary for one mathematical re-formulation

Particle Types	
Introduction of the Physical Problem	
■ Bosonic operators	$\hat{\sigma}^0 = \mathbb{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$
■ Commutation relations	$\hat{\sigma}^2 = \hat{\sigma}^y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$
■ Fermionic operators	
■ Anti-commutation relations	$\hat{\sigma}^1 = \hat{\sigma}^x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$
■ Hard-core bosonic operators	$\hat{\sigma}^3 = \hat{\sigma}^z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$
■ Commutation relations	
■ Occupation number limited to 0 and 1	
■ Number operator is idempotent	
■ Spins (Pauli operators/matrices)	
■ Particles from different sites commute	$\left\{ \hat{\sigma}_l^\alpha, \hat{\sigma}_l^\beta \right\} = 2 \cdot \delta_{\alpha, \beta}$
■ Different spin directions on the same site anti-commute	$\left[\hat{\sigma}_l^\alpha, \hat{\sigma}_m^\beta \right] = 2 \cdot i \cdot \varepsilon_{\alpha, \beta, \gamma} \cdot \delta_{l, m} \cdot \hat{\sigma}_l^\gamma$
■ They obey angular momentum commutation relations	

Particle Types - Mappings

Introduction of the Physical Problem

■ Properties of Pauli matrices

- Base of the complex 2x2 matrices
- Can be directly mapped to hard-core bosons

$$\hat{\sigma}_l^x = \hat{h}_l + \hat{h}_l^\dagger$$

$$\hat{\sigma}_l^y = i \cdot (\hat{h}_l - \hat{h}_l^\dagger)$$

$$\hat{\sigma}_l^z = 2 \cdot \hat{h}_l^\dagger \hat{h}_l - 1$$

$$|\downarrow\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} = |0\rangle$$

$$\hat{\sigma}^+ |\downarrow\rangle = |\uparrow\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} = |1\rangle = \hat{h}^\dagger |0\rangle$$



2025-03-12

Master Colloquium Presentation

└ Introduction of the Physical Problem

└ Particle Types - Mappings

1. Pauli matrices are a base for the complex 2x2 matrices
2. Kronecker product of two pauli matrices therefore base for complex 4x4 matrices
3. Simple transformation that can be verified for its commutation relations or derived from two *Jordan-Wigner Transformations*
4. Important to watch the order of base states and keep convention or -1 will be introduced unexpectedly

Particle Types - Mappings
Introduction of the Physical Problem

- Properties of Pauli matrices
 - Base of the complex 2x2 matrices
 - Can be directly mapped to hard-core bosons

$$\begin{aligned}\hat{\sigma}_l^x &= \hat{h}_l + \hat{h}_l^\dagger \\ \hat{\sigma}_l^y &= i \cdot (\hat{h}_l - \hat{h}_l^\dagger) \\ \hat{\sigma}_l^z &= 2 \cdot \hat{h}_l^\dagger \hat{h}_l - 1 \\ |1\rangle &= \begin{pmatrix} 0 \\ 1 \end{pmatrix} = |\downarrow\rangle \\ \hat{\sigma}^+ |1\rangle = |\uparrow\rangle &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} = |1\rangle = \hat{h}^\dagger |0\rangle\end{aligned}$$

Hamiltonian of the system

Introduction of the Physical Problem

- Hubbard model Hamiltonian
- Constant external electric field
- Two particle types (spin degrees)
 - Different *spins* than for Pauli spin operators

$$\mathcal{H} = \mathcal{H}_0 + \hat{V}$$

$$\mathcal{H}_0 = U \cdot \sum_l \hat{n}_{l,\uparrow} \hat{n}_{l,\downarrow} + \sum_{l,\sigma} \underbrace{\left(\vec{E} \cdot \vec{r}_l \right)}_{\varepsilon_l} \hat{n}_{l,\sigma}$$

$$\hat{V} = -J \cdot \sum_{\langle l,m \rangle, \sigma} \left(\hat{h}_{l,\sigma}^\dagger \hat{h}_{m,\sigma} + \hat{h}_{m,\sigma}^\dagger \hat{h}_{l,\sigma} \right)$$



2025-03-12

Master Colloquium Presentation

└ Introduction of the Physical Problem

└ Hamiltonian of the system

1. Hubbard model:

- Double occupation term
- Two particle degrees of freedom (= spin but not same spin as before: so also written as h and d)
- Here: electrical field that acts symmetrical on both spin directions
- Hopping term that is here treated as a perturbation

Hamiltonian of the system Introduction of the Physical Problem


- Hubbard model Hamiltonian
- Constant external electric field
- Two particle types (spin degrees)
 - Different spins than for Pauli spin operators

$$\mathcal{H} = \mathcal{H}_0 + \hat{V}$$
$$\mathcal{H}_0 = U \cdot \sum_l \hat{n}_l \hat{n}_l + \sum_{l,\sigma} \left(\vec{E} \cdot \vec{r}_l \right) \hat{n}_{l,\sigma}$$
$$\hat{V} = -J \cdot \sum_{\langle l,m \rangle, \sigma} \left(\hat{h}_{l,\sigma}^\dagger \hat{h}_{m,\sigma} + \hat{h}_{m,\sigma}^\dagger \hat{h}_{l,\sigma} \right)$$

Geometry of the system

Introduction of the Physical Problem

- 2-dimensional geometry
- Square lattice arrangement
- Computation for general size and field angle




2025-03-12

Master Colloquium Presentation

- └ Introduction of the Physical Problem
- └ Geometry of the system

1. two-dimensional
2. square
3. Electrical field and we want to calculate generally for angle, field strength and number of sites

Geometry of the system
Introduction of the Physical Problem



Goal of the Calculation

Time Evolution

- Evaluate the time-evolution of the system for various observables
 - Uses calculation in the *Interaction Picture*
 - Introduce operator-less *effective Hamiltonian*

$$|\Psi^S(t)\rangle = \sum_N e^{\mathcal{H}_{\text{eff}}(N,t)} \Psi_N |N\rangle$$



2025-03-12

Master Colloquium Presentation

Time Evolution

Goal of the Calculation

1. Goal: evaluate the time-evolution for different observables (effectively)
2. Brute-force calculation is at least NP-hard, definitely exponential and one of the hardest problems to solve computationally generally
3. Strategy: perform a controlled expansion in the perturbation V and re-express the formulas as a dependency on a operator-free "effective Hamiltonian"
 - Can be re-ordered without concern as no operators
 - For the re-formulation we will use the interaction picture
4. Particle-type dependency is stored in this formula in the BASE STATES. Because in this re-formulation, the $\mathcal{H}_{\text{eff}}$ is a pure number (no operator) and the initial state is a number (not operator dependent at all). But the base-state contains different info for fermions than for bosons, as the operators for hc-bosons commute and so the base-states are uniquely defined. But there are multiple possibilities for teh fermions, because swaps introduce minus signs. (State is not just 0s and 1s, but vacuum that is pre-set with operators and they have an order!!!!)

Goal of the Calculation

Time Evolution

- Evaluate the time-evolution of the system for various observables
 - Uses calculation in the *Interaction Picture*
 - Introduce operator-less *effective Hamiltonian*

$$|\Psi^S(t)\rangle = \sum_N e^{\mathcal{H}_{\text{eff}}(N,t)} \Psi_N |N\rangle$$

The effective Hamiltonian

Time Evolution

$$\mathcal{H}_{\text{eff}}(N, t) = -iE_0(N)t + \mathcal{H}_N(t)$$

- Constructed from the contributions of the base energy and the perturbation
- Base energy contribution:

$$\begin{aligned} E_0(N) &= \frac{\langle N | \mathcal{H}_0 | N \rangle}{\langle N | N \rangle} \\ &= \langle N | U \cdot \sum_l \hat{n}_{l,\uparrow} \hat{n}_{l,\downarrow} | N \rangle + \langle N | \sum_{l,\sigma} \varepsilon_l \hat{n}_{l,\sigma} | N \rangle \\ &= U \cdot \sum_l n_{l,\uparrow} n_{l,\downarrow} + \sum_{l,\sigma} \varepsilon_l n_{l,\sigma} \end{aligned}$$



2025-03-12

Master Colloquium Presentation

Time Evolution

The effective Hamiltonian

1. Comprised of two parts:

- Simple, quickly stated term that depends on the base energy
- Complicated one that will be expanded in the perturbation

The effective Hamiltonian

Time Evolution

$$\mathcal{H}_{\text{eff}}(N, t) = -iE_0(N)t + \mathcal{H}_N(t)$$

- Constructed from the contributions of the base energy and the perturbation

- Base energy contribution:

$$\begin{aligned} E_0(N) &= \frac{\langle N | \mathcal{H}_0 | N \rangle}{\langle N | N \rangle} \\ &= \langle N | U \cdot \sum_l \hat{n}_{l,\uparrow} \hat{n}_{l,\downarrow} | N \rangle + \langle N | \sum_{l,\sigma} \varepsilon_l \hat{n}_{l,\sigma} | N \rangle \\ &= U \cdot \sum_l n_{l,\uparrow} n_{l,\downarrow} + \sum_{l,\sigma} \varepsilon_l n_{l,\sigma} \end{aligned}$$

The effective Hamiltonian

Time Evolution

- To evaluate the time-evolution of the perturbation

$$\hat{V} = -J \cdot \sum_{\langle l,m \rangle} \left(\hat{h}_l^\dagger \hat{h}_m + \hat{h}_m^\dagger \hat{h}_l + \hat{d}_l^\dagger \hat{d}_m + \hat{d}_m^\dagger \hat{d}_l \right)$$

- Solve the equation of motion for the ladder operators

$$\begin{aligned} \hat{h}_m^{\dagger I}(t) &= e^{i \cdot \varepsilon_m \cdot t} \left(1 + (e^{iU \cdot t} - 1) \hat{d}_m^{\dagger S} \hat{d}_m^S \right) \hat{h}_m^{\dagger S} \\ \hat{h}_m^I(t) &= e^{-i \cdot \varepsilon_m \cdot t} \left(1 + (e^{-iU \cdot t} - 1) \hat{d}_m^{\dagger S} \hat{d}_m^S \right) \hat{h}_m^S \\ \hat{d}_m^{\dagger I}(t) &= e^{i \cdot \varepsilon_m \cdot t} \left(1 + (e^{iU \cdot t} - 1) \hat{h}_m^{\dagger S} \hat{h}_m^S \right) \hat{d}_m^{\dagger S} \\ \hat{d}_m^I(t) &= e^{-i \cdot \varepsilon_m \cdot t} \left(1 + (e^{-iU \cdot t} - 1) \hat{h}_m^{\dagger S} \hat{h}_m^S \right) \hat{d}_m^S \end{aligned}$$



2025-03-12

Master Colloquium Presentation

Time Evolution

The effective Hamiltonian

1. Need: V-Operator in the Interaction picture
2. For that: first solve the equations of motion for the ladder operators and the plug in
3. Solving the equations of motions in the interaction picture requires their number operator being *idempotent*

The effective Hamiltonian

Time Evolution

- To evaluate the time-evolution of the perturbation

$$\hat{V} = -J \cdot \sum_{\langle l,m \rangle} \left(\hat{h}_l^\dagger \hat{h}_m + \hat{h}_m^\dagger \hat{h}_l + \hat{d}_l^\dagger \hat{d}_m + \hat{d}_m^\dagger \hat{d}_l \right)$$

- Solve the equation of motion for the ladder operators

$$\hat{h}_m^{\dagger I}(t) = e^{i \cdot \varepsilon_m \cdot t} \left(1 + (e^{iU \cdot t} - 1) \hat{d}_m^{\dagger S} \hat{d}_m^S \right) \hat{h}_m^{\dagger S}$$

$$\hat{h}_m^I(t) = e^{-i \cdot \varepsilon_m \cdot t} \left(1 + (e^{-iU \cdot t} - 1) \hat{d}_m^{\dagger S} \hat{d}_m^S \right) \hat{h}_m^S$$

$$\hat{d}_m^{\dagger I}(t) = e^{i \cdot \varepsilon_m \cdot t} \left(1 + (e^{iU \cdot t} - 1) \hat{h}_m^{\dagger S} \hat{h}_m^S \right) \hat{d}_m^{\dagger S}$$

$$\hat{d}_m^I(t) = e^{-i \cdot \varepsilon_m \cdot t} \left(1 + (e^{-iU \cdot t} - 1) \hat{h}_m^{\dagger S} \hat{h}_m^S \right) \hat{d}_m^S$$

The effective Hamiltonian

Time Evolution

- Insert and reorder the operators

$$\begin{aligned}
 \hat{V}^I(t) &= \left\{ \hat{V}^S \right\}^I(t) = -J \cdot \sum_{[l,m]} \left\{ \left(\hat{h}_l^{\dagger S} \hat{h}_m^S + \hat{d}_l^{\dagger S} \hat{d}_m^S \right) \right\}^I(t) \\
 &= -J \cdot \sum_{[l,m]} \left(\hat{h}_l^{\dagger I}(t) \hat{h}_m^I(t) + \hat{d}_l^{\dagger I}(t) \hat{d}_m^I(t) \right) \\
 &= -J \cdot \sum_{[l,m]} \left[\Lambda_A(l, m, t) \cdot \hat{F}_A(l, m) + \Lambda_B(l, m, t) \cdot \hat{F}_B(l, m) + \Lambda_C(l, m, t) \cdot \hat{F}_C(l, m) \right]
 \end{aligned}$$

$$\Lambda_A(l, m, t) = e^{i(\varepsilon_l - \varepsilon_m) \cdot t} \quad \hat{F}_A(l, m) = \sum_{\sigma \in \{\uparrow, \downarrow\}} \hat{h}_{l, \sigma}^{\dagger S} \hat{h}_{m, \sigma}^S \left(1 + 2 \cdot \hat{n}_{l, \bar{\sigma}}^S \hat{n}_{m, \bar{\sigma}}^S - \hat{n}_{l, \bar{\sigma}}^S - \hat{n}_{m, \bar{\sigma}}^S \right)$$

$$\Lambda_B(l, m, t) = e^{i(\varepsilon_l - \varepsilon_m + U) \cdot t} \quad \hat{F}_B(l, m) = \sum_{\sigma \in \{\uparrow, \downarrow\}} \hat{h}_{l, \sigma}^{\dagger S} \hat{h}_{m, \sigma}^S \left(\hat{n}_{l, \bar{\sigma}}^S - \hat{n}_{l, \bar{\sigma}}^S \hat{n}_{m, \bar{\sigma}}^S \right)$$

$$\Lambda_C(l, m, t) = e^{i(\varepsilon_l - \varepsilon_m - U) \cdot t} \quad \hat{F}_C(l, m) = \sum_{\sigma \in \{\uparrow, \downarrow\}} \hat{h}_{l, \sigma}^{\dagger S} \hat{h}_{m, \sigma}^S \left(\hat{n}_{m, \bar{\sigma}}^S - \hat{n}_{m, \bar{\sigma}}^S \hat{n}_{l, \bar{\sigma}}^S \right)$$



2025-03-12

Master Colloquium Presentation

Time Evolution

The effective Hamiltonian

1. Inserting yields TWO kinds of terms

- time-dependent analytical pre-factors (or their integral as one sees later)
- dressed hopping terms, basically only depending on the occupation of the state N

The effective Hamiltonian

Time Evolution

- Insert and reorder the operators

$$\begin{aligned}
 \hat{V}^I(t) &= \left\{ \hat{V}^S \right\}^I(t) = -J \cdot \sum_{[l,m]} \left\{ \left(\hat{h}_l^{\dagger S} \hat{h}_m^S + \hat{d}_l^{\dagger S} \hat{d}_m^S \right) \right\}^I(t) \\
 &= -J \cdot \sum_{[l,m]} \left(\hat{h}_l^{\dagger I}(t) \hat{h}_m^I(t) + \hat{d}_l^{\dagger I}(t) \hat{d}_m^I(t) \right) \\
 &= -J \cdot \sum_{[l,m]} \left[\Lambda_A(l, m, t) \cdot \hat{F}_A(l, m) + \Lambda_B(l, m, t) \cdot \hat{F}_B(l, m) + \Lambda_C(l, m, t) \cdot \hat{F}_C(l, m) \right] \\
 \Lambda_A(l, m, t) &= e^{i(\varepsilon_l - \varepsilon_m) \cdot t} & \hat{F}_A(l, m) &= \sum_{\sigma \in \{\uparrow, \downarrow\}} \hat{h}_{l, \sigma}^{\dagger S} \hat{h}_{m, \sigma}^S \left(1 + 2 \cdot \hat{n}_{l, \bar{\sigma}}^S \hat{n}_{m, \bar{\sigma}}^S - \hat{n}_{l, \bar{\sigma}}^S - \hat{n}_{m, \bar{\sigma}}^S \right) \\
 \Lambda_B(l, m, t) &= e^{i(\varepsilon_l - \varepsilon_m + U) \cdot t} & \hat{F}_B(l, m) &= \sum_{\sigma \in \{\uparrow, \downarrow\}} \hat{h}_{l, \sigma}^{\dagger S} \hat{h}_{m, \sigma}^S \left(\hat{n}_{l, \bar{\sigma}}^S - \hat{n}_{l, \bar{\sigma}}^S \hat{n}_{m, \bar{\sigma}}^S \right) \\
 \Lambda_C(l, m, t) &= e^{i(\varepsilon_l - \varepsilon_m - U) \cdot t} & \hat{F}_C(l, m) &= \sum_{\sigma \in \{\uparrow, \downarrow\}} \hat{h}_{l, \sigma}^{\dagger S} \hat{h}_{m, \sigma}^S \left(\hat{n}_{m, \bar{\sigma}}^S - \hat{n}_{m, \bar{\sigma}}^S \hat{n}_{l, \bar{\sigma}}^S \right)
 \end{aligned}$$

The effective Hamiltonian

Time Evolution

- Evaluate the contribution to the effective Hamiltonian
 - Requires previously calculated value of the V-operator in the Interaction Picture
 - Controllable *cumulant expansion*

$$\begin{aligned}
 \mathcal{H}_N(t) &= \sum_{v=1}^{\infty} \frac{(-i)^v}{v!} \int_0^t dt_1 \int_0^t dt_2 \cdots \int_0^t dt_v \langle \mathbb{T} \hat{V}^I(t_1) \hat{V}^I(t_2) \cdots \hat{V}^I(t_v) \rangle_{c(N)} \\
 &= -i \int_0^t dt_1 \frac{\langle N | \hat{V}^I(t_1) | \Psi^S \rangle}{\langle N | \Psi^S \rangle} \\
 &\quad - \frac{1}{2} \int_0^t dt_1 \int_0^t dt_2 \left(\frac{\langle N | \mathbb{T} \hat{V}^I(t_1) \hat{V}^I(t_2) | \Psi^S \rangle}{\langle N | \Psi^S \rangle} - \frac{\langle N | \hat{V}^I(t_1) | \Psi^S \rangle \cdot \langle N | \hat{V}^I(t_2) | \Psi^S \rangle}{\langle N | \Psi^S \rangle^2} \right) \\
 &\quad + \cdots
 \end{aligned}$$



2025-03-12

Master Colloquium Presentation

Time Evolution

The effective Hamiltonian

1. Generally any expansion possible, but here we expand the *argument of the exponential* - cumulant
2. first order easily derivable with the info from the previous slide (integrate)
3. second and higher orders very complicated occupation terms and the pre-factors require quite a lot of edge-cases to integrate with the time-ordering operator
 - Hint onto the thesis for this exact derivation

The effective Hamiltonian

Time Evolution

- Evaluate the contribution to the effective Hamiltonian

- Requires previously calculated value of the V-operator in the Interaction Picture
- Controllable cumulant expansion

$$\begin{aligned}
 H_E(t) &= \sum_{v=1}^{\infty} \frac{(-i)^v}{v!} \int_0^t dt_1 \int_0^t dt_2 \cdots \int_0^t dt_v \langle \mathbb{T} \hat{V}^I(t_1) \hat{V}^I(t_2) \cdots \hat{V}^I(t_v) \rangle_{c(N)} \\
 &= -i \int_0^t dt_1 \frac{\langle N | \hat{V}^I(t_1) | \Psi^S \rangle}{\langle N | \Psi^S \rangle} \\
 &\quad - \frac{1}{2} \int_0^t dt_1 \int_0^t dt_2 \left(\frac{\langle N | \mathbb{T} \hat{V}^I(t_1) \hat{V}^I(t_2) | \Psi^S \rangle}{\langle N | \Psi^S \rangle} - \frac{\langle N | \hat{V}^I(t_1) | \Psi^S \rangle \cdot \langle N | \hat{V}^I(t_2) | \Psi^S \rangle}{\langle N | \Psi^S \rangle^2} \right) \\
 &\quad + \cdots
 \end{aligned}$$

Handling of Observables

Time Evolution

- This allows for general evaluation of expectation values
 - Requires a probability for the state
 - Requires a local observable

$$\frac{\langle \Psi^S(t) | \hat{\mathcal{O}} | \Psi^S(t) \rangle}{\langle \Psi^S(t) | \Psi^S(t) \rangle} =$$

$$\sum_N P(N, t) \underbrace{\sum_K \langle N | \hat{\mathcal{O}} | K \rangle e^{\mathcal{H}_{\text{eff}}(K, t) - \mathcal{H}_{\text{eff}}(N, t)} \frac{\Psi_K}{\Psi_N}}_{\hat{\mathcal{O}}_{\text{loc}}(N, t)} = \sum_N P(N, t) \hat{\mathcal{O}}_{\text{loc}}(N, t)$$



2025-03-12

Master Colloquium Presentation

Time Evolution

Handling of Observables

1. Why? General evaluation of an expectation value for an observable.
 - Probability for state
 - Local observable (can be swapped for many different ones)
2. Does ONLY require difference of effective Hamiltonian for evaluation (notice for later)

Handling of Observables

Time Evolution

- This allows for general evaluation of expectation values
 - Requires a probability for the state
 - Requires a local observable

$$\frac{\langle \Psi^S(t) | \hat{\mathcal{O}} | \Psi^S(t) \rangle}{\langle \Psi^S(t) | \Psi^S(t) \rangle} = \sum_N P(N, t) \underbrace{\sum_K \langle N | \hat{\mathcal{O}} | K \rangle e^{\mathcal{H}_{\text{eff}}(K, t) - \mathcal{H}_{\text{eff}}(N, t)} \frac{\Psi_K}{\Psi_N}}_{\hat{\mathcal{O}}_{\text{loc}}(N, t)} = \sum_N P(N, t) \hat{\mathcal{O}}_{\text{loc}}(N, t)$$

Simple Observables

Time Evolution

- Local observable for double occupation measurement
 - Observables generally are very sparse matrices
 - Reduces to pure occupation measurement

$$\hat{\mathcal{O}}_{\text{loc}}(N, t) = \sum_K \langle N | \hat{\mathcal{O}}_{\text{do}}(l) | K \rangle e^{\mathcal{H}_{\text{eff}}(K, t) - \mathcal{H}_{\text{eff}}(N, t)} \frac{\Psi_K}{\Psi_N} = n_{l, \uparrow} \cdot n_{l, \downarrow}$$

- Local observable for particle current measurement
 - Requires evaluation of the difference of two effective Hamiltonians

$$\hat{\mathcal{O}}_{\text{loc}}(N, t) = i \cdot \left[n_{m, \sigma} \cdot (1 - n_{l, \sigma}) - n_{l, \sigma} \cdot (1 - n_{m, \sigma}) \right] \cdot \frac{\Psi_{\tilde{N}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{N}, t) - \mathcal{H}_{\text{eff}}(N, t)}$$



2025-03-12

Master Colloquium Presentation

Time Evolution

Simple Observables

1. Complex valued observable, make sure this cancels in the case of approximations
2. *SHOULD MENTION*: Energy and variance (most expensive ones, are assumed to keep constant, can be used to gauge the quality of the expansion)

Simple Observables

Time Evolution

- Local observable for double occupation measurement
 - Observables generally are very sparse matrices
 - Reduces to pure occupation measurement

$$\hat{\mathcal{O}}_{\text{do}}(N, t) = \sum_K \langle N | \hat{\mathcal{O}}_{\text{do}}(l) | K \rangle e^{\mathcal{H}_{\text{eff}}(K, t) - \mathcal{H}_{\text{eff}}(N, t)} \frac{\Psi_K}{\Psi_N} = n_{l, \uparrow} \cdot n_{l, \downarrow}$$

- Local observable for particle current measurement
 - Requires evaluation of the difference of two effective Hamiltonians

$$\hat{\mathcal{O}}_{\text{loc}}(N, t) = i \cdot \left[n_{m, \sigma} \cdot (1 - n_{l, \sigma}) - n_{l, \sigma} \cdot (1 - n_{m, \sigma}) \right] \cdot \frac{\Psi_{\tilde{N}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{N}, t) - \mathcal{H}_{\text{eff}}(N, t)}$$

Access to Density-Matrices

Time Evolution

- Non-classical (*quantum*) measurements depend on the (reduced) density matrix
 - Access to *purity*, *concurrence* and other entanglement measures/monotones
- Direct calculation not possible

$$\rho_A = \text{Tr}_B(\rho_{AB}) = \sum_{k=0}^M (\mathbb{1}_A \otimes \langle \chi_k |) \rho_{AB} (\mathbb{1}_A \otimes | \chi_k \rangle)$$

- Use Pauli matrices to expand the complex 4x4 matrix

$$\rho_A(t) = \rho_{l,\sigma,m,\mu}(t) = \frac{1}{4} \sum_{\alpha,\beta \in \{0,x,y,z\}} \langle \hat{\sigma}_{l,\sigma}^\alpha \hat{\sigma}_{m,\mu}^\beta \rangle_{(t)} (\hat{\sigma}^\alpha \otimes \hat{\sigma}^\beta)$$



2025-03-12

Master Colloquium Presentation

Time Evolution

Access to Density-Matrices

1. Quite interesting, because of this extra mentioned the Pauli operators
2. No access to the full density matrix, can not trace out, because too many states to trace out
3. BUT: as the complex 4x4 matrix can be written in the base of the kronecker product of two pauli matrices, can be expressed (because the pauli operators can be translated into hard-core bosonic ones again and for these we have the general formula for expectation values)

Access to Density-Matrices

Time Evolution

- Non-classical (*quantum*) measurements depend on the (reduced) density matrix
 - Access to *purity*, *concurrence* and other entanglement measures/monotones
- Direct calculation not possible

$$\rho_A = \text{Tr}_B(\rho_{AB}) = \sum_{k=0}^M (\mathbb{1}_A \otimes \langle \chi_k |) \rho_{AB} (\mathbb{1}_A \otimes | \chi_k \rangle)$$

- Use Pauli matrices to expand the complex 4x4 matrix

$$\rho_A(t) = \rho_{l,\sigma,m,\mu}(t) = \frac{1}{4} \sum_{\alpha,\beta \in \{0,x,y,z\}} \langle \hat{\sigma}_{l,\sigma}^\alpha \hat{\sigma}_{m,\mu}^\beta \rangle_{(t)} (\hat{\sigma}^\alpha \otimes \hat{\sigma}^\beta)$$

Access to Density-Matrices

Time Evolution

$$\begin{aligned}
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^x \hat{\sigma}_{m,\mu}^0 : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{N}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{N},t) - \mathcal{H}_{\text{eff}}(N,t)} \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^y \hat{\sigma}_{m,\mu}^0 : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{N}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{N},t) - \mathcal{H}_{\text{eff}}(N,t)} \cdot i \cdot (1 - 2 \cdot n_{l,\sigma}) \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^z \hat{\sigma}_{m,\mu}^0 : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{N}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{N},t) - \mathcal{H}_{\text{eff}}(N,t)} \cdot (2 \cdot n_{l,\sigma} - 1) \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^x \hat{\sigma}_{m,\mu}^x : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{\tilde{N}}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{\tilde{N}},t) - \mathcal{H}_{\text{eff}}(N,t)} \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^y \hat{\sigma}_{m,\mu}^y : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{\tilde{N}}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{\tilde{N}},t) - \mathcal{H}_{\text{eff}}(N,t)} \cdot (2 \cdot n_{l,\sigma} - 1) \cdot (1 - 2 \cdot n_{m,\mu}) \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^x \hat{\sigma}_{m,\mu}^y : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{\tilde{N}}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{\tilde{N}},t) - \mathcal{H}_{\text{eff}}(N,t)} \cdot i \cdot (1 - 2 \cdot n_{m,\mu}) \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^y \hat{\sigma}_{m,\mu}^x : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{\tilde{N}}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{\tilde{N}},t) - \mathcal{H}_{\text{eff}}(N,t)} \cdot i \cdot (1 - 2 \cdot n_{l,\sigma})
 \end{aligned}$$



2025-03-12

Master Colloquium Presentation

Time Evolution

Access to Density-Matrices

1. Sixteen terms (with missing cases trivially derived from the ones that are given)
2. Again all depend on the difference of effective hamiltonians between two states ith local modifications

Access to Density-Matrices

Time Evolution

$$\begin{aligned}
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^x \hat{\sigma}_{m,\mu}^0 : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{N}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{N},t) - \mathcal{H}_{\text{eff}}(N,t)} \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^y \hat{\sigma}_{m,\mu}^0 : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{N}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{N},t) - \mathcal{H}_{\text{eff}}(N,t)} \cdot i \cdot (1 - 2 \cdot n_{l,\sigma}) \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^z \hat{\sigma}_{m,\mu}^0 : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{N}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{N},t) - \mathcal{H}_{\text{eff}}(N,t)} \cdot (2 \cdot n_{l,\sigma} - 1) \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^x \hat{\sigma}_{m,\mu}^x : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{\tilde{N}}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{\tilde{N}},t) - \mathcal{H}_{\text{eff}}(N,t)} \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^y \hat{\sigma}_{m,\mu}^y : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{\tilde{N}}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{\tilde{N}},t) - \mathcal{H}_{\text{eff}}(N,t)} \cdot (2 \cdot n_{l,\sigma} - 1) \cdot (1 - 2 \cdot n_{m,\mu}) \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^x \hat{\sigma}_{m,\mu}^y : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{\tilde{N}}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{\tilde{N}},t) - \mathcal{H}_{\text{eff}}(N,t)} \cdot i \cdot (1 - 2 \cdot n_{m,\mu}) \\
 \hat{\mathcal{O}} = \hat{\sigma}_{l,\sigma}^y \hat{\sigma}_{m,\mu}^x : \quad \hat{\mathcal{O}}_{\text{loc}}(N,t) &= \frac{\Psi_{\tilde{\tilde{N}}}}{\Psi_N} \cdot e^{\mathcal{H}_{\text{eff}}(\tilde{\tilde{N}},t) - \mathcal{H}_{\text{eff}}(N,t)} \cdot i \cdot (1 - 2 \cdot n_{l,\sigma})
 \end{aligned}$$

Probability Calculations

Sampling & Simplifications

- Full probability still requires normalization
- Switch to the Metropolis *Hastings* algorithm
 - Non-exponential number of samples
 - Depends only on a transition probability
 - Does not need normalization because it cancels

$$\begin{aligned} \alpha &= \frac{P(\tilde{N}, t)}{P(N, t)} = \frac{f(\tilde{N}, t)}{f(N, t)} = \frac{|e^{\mathcal{H}_{\text{eff}}(\tilde{N}, t)}|^2 |\Psi_{\tilde{N}}|^2}{|e^{\mathcal{H}_{\text{eff}}(N, t)}|^2 |\Psi_N|^2} \\ &= \frac{|\Psi_{\tilde{N}}|^2 e^{\Re(\mathcal{H}_{\text{eff}}(\tilde{N}, t)) + i\Im(\mathcal{H}_{\text{eff}}(\tilde{N}, t)) + \Re(\mathcal{H}_{\text{eff}}(\tilde{N}, t)) - i\Im(\mathcal{H}_{\text{eff}}(\tilde{N}, t))}}{|\Psi_N|^2 e^{\Re(\mathcal{H}_{\text{eff}}(N, t)) + i\Im(\mathcal{H}_{\text{eff}}(N, t)) + \Re(\mathcal{H}_{\text{eff}}(N, t)) - i\Im(\mathcal{H}_{\text{eff}}(N, t))}} \\ &= \frac{|\Psi_{\tilde{N}}|^2}{|\Psi_N|^2} e^{2\Re(\mathcal{H}_{\text{eff}}(\tilde{N}, t)) - 2\Re(\mathcal{H}_{\text{eff}}(N, t))} \\ &= \frac{|\Psi_{\tilde{N}}|^2}{|\Psi_N|^2} e^{2\Re(\mathcal{H}_{\text{eff}}(\tilde{N}, t) - \mathcal{H}_{\text{eff}}(N, t))} \end{aligned}$$

$$\frac{\langle \Psi^S(t) | \hat{\mathcal{O}} | \Psi^S(t) \rangle}{\langle \Psi^S(t) | \Psi^S(t) \rangle} = \sum_N P(N, t) \cdot \hat{\mathcal{O}}_{\text{loc}}(N, t) \approx \frac{1}{|\{N\}_{\text{MC}}|} \sum_{\{N\}_{\text{MC}}} \hat{\mathcal{O}}_{\text{loc}}(N, t)$$



2025-03-12

Master Colloquium Presentation

Sampling & Simplifications

Probability Calculations

1. Normalization requires knowing exponential amount of states and we need to sample an exponential amount to get all
2. switch to METROPOLIS HASTINGS algorithm
 - only fixed number of samples required (depends on the desired precision)
 - No normalization needed, because it cancels in the division

Probability Calculations
Sampling & Simplifications

- Full probability still requires normalization
- Switch to the Metropolis *Hastings* algorithm
 - Non-exponential number of samples
 - Depends only on a transition probability
 - Does not need normalization because it cancels

$$\alpha = \frac{P(\tilde{N}, t)}{P(N, t)} = \frac{f(\tilde{N}, t)}{f(N, t)} = \frac{|e^{\mathcal{H}_{\text{eff}}(\tilde{N}, t)}|^2 |\Psi_{\tilde{N}}|^2}{|e^{\mathcal{H}_{\text{eff}}(N, t)}|^2 |\Psi_N|^2}$$

$$= \frac{|\Psi_{\tilde{N}}|^2 e^{\Re(\mathcal{H}_{\text{eff}}(\tilde{N}, t)) + i\Im(\mathcal{H}_{\text{eff}}(\tilde{N}, t)) + \Re(\mathcal{H}_{\text{eff}}(\tilde{N}, t)) - i\Im(\mathcal{H}_{\text{eff}}(\tilde{N}, t))}}{|\Psi_N|^2 e^{\Re(\mathcal{H}_{\text{eff}}(N, t)) + i\Im(\mathcal{H}_{\text{eff}}(N, t)) + \Re(\mathcal{H}_{\text{eff}}(N, t)) - i\Im(\mathcal{H}_{\text{eff}}(N, t))}}$$

$$= \frac{|\Psi_{\tilde{N}}|^2}{|\Psi_N|^2} e^{2\Re(\mathcal{H}_{\text{eff}}(\tilde{N}, t)) - 2\Re(\mathcal{H}_{\text{eff}}(N, t))}$$

$$= \frac{|\Psi_{\tilde{N}}|^2}{|\Psi_N|^2} e^{2\Re(\mathcal{H}_{\text{eff}}(\tilde{N}, t) - \mathcal{H}_{\text{eff}}(N, t))}$$

$$\frac{\langle \Psi^S(t) | \hat{\mathcal{O}} | \Psi^S(t) \rangle}{\langle \Psi^S(t) | \Psi^S(t) \rangle} = \sum_N P(N, t) \cdot \hat{\mathcal{O}}_{\text{loc}}(N, t) \approx \frac{1}{|\{N\}_{\text{MC}}|} \sum_{\{N\}_{\text{MC}}} \hat{\mathcal{O}}_{\text{loc}}(N, t)$$

Analytical Simplifications

Sampling & Simplifications

- Choose a helpful initial state

$$\Psi_N = \frac{1}{\sqrt{\#(\text{states})}} = \frac{1}{\sqrt{2^{\#(\text{sites}) \cdot 2}}} = \frac{1}{2^{\#(\text{sites})}}$$

$$|\Psi^S(t=0)\rangle = \bigotimes_{l=1}^{\#(\text{states})} \frac{1}{2} \left(1 + \hat{h}_{l,\uparrow}^{\dagger S} + \hat{h}_{l,\downarrow}^{\dagger S} + \hat{h}_{l,\uparrow}^{\dagger S} \hat{h}_{l,\downarrow}^{\dagger S} \right) |0\rangle$$

- Now *almost all* terms cancel in *differences* of the effective Hamiltonian

- Exemplary for single flip on base energy term

$$\begin{aligned} E_0(N) - E_0(\tilde{N}) &= U \sum_l n_{l,\downarrow} n_{l,\uparrow} - U \sum_l \tilde{n}_{l,\downarrow} \tilde{n}_{l,\uparrow} + \sum_{l,\sigma} \varepsilon_l n_{l,\sigma} - \sum_{l,\sigma} \varepsilon_l \tilde{n}_{l,\sigma} \\ &= \varepsilon_i (2n_{i,\sigma_i} - 1) + U \cdot \begin{cases} \sigma_i = \uparrow : & n_{i,\downarrow} (2n_{i,\uparrow} - 1) \\ \sigma_i = \downarrow : & n_{i,\uparrow} (2n_{i,\downarrow} - 1) \end{cases} \end{aligned}$$



2025-03-12

Master Colloquium Presentation

Sampling & Simplifications

Analytical Simplifications

1. Choose the helpful *initial state* with all base states having the same probability
2. because of this the divisions of Psi-N over Psi-K everywhere give 1 and many terms cancel (outside of the local interaction range of the modification)

Analytical Simplifications

Sampling & Simplifications

- Choose a helpful initial state

$$\Psi_N = \frac{1}{\sqrt{\#(\text{states})}} = \frac{1}{\sqrt{2^{\#(\text{sites}) \cdot 2}}} = \frac{1}{2^{\#(\text{sites})}}$$

$$|\Psi^S(t=0)\rangle = \bigotimes_{l=1}^{\#(\text{states})} \frac{1}{2} \left(1 + \hat{h}_{l,\uparrow}^{\dagger S} + \hat{h}_{l,\downarrow}^{\dagger S} + \hat{h}_{l,\uparrow}^{\dagger S} \hat{h}_{l,\downarrow}^{\dagger S} \right) |0\rangle$$

- Now almost all terms cancel in differences of the effective Hamiltonian

$$\begin{aligned} E_0(N) - E_0(\tilde{N}) &\approx U \sum_l n_{l,\downarrow} n_{l,\uparrow} - U \sum_l \tilde{n}_{l,\downarrow} \tilde{n}_{l,\uparrow} + \sum_{l,\sigma} \varepsilon_l n_{l,\sigma} - \sum_{l,\sigma} \varepsilon_l \tilde{n}_{l,\sigma} \\ &= \varepsilon_i (2n_{i,\sigma_i} - 1) + U \cdot \begin{cases} \sigma_i = \uparrow : & n_{i,\downarrow} (2n_{i,\uparrow} - 1) \\ \sigma_i = \downarrow : & n_{i,\uparrow} (2n_{i,\downarrow} - 1) \end{cases} \end{aligned}$$

Why and How should Parameters be variational?

Variational Classical Networks

- Theoretically: always better approximation through higher order terms
 - Hard to calculate analytically and expensive to evaluate
- Hope: Better behavior if parameters are *learned/optimized*, starting from the analytical result

$$|\Psi_{\vec{\eta}+\delta\cdot\dot{\vec{\eta}}}(t+\delta)\rangle \leftrightarrow e^{-iH\delta} |\Psi_{\vec{\eta}}(t)\rangle$$

$$\vec{\eta}(t+\delta) = \vec{\eta}(t) + \delta \cdot \dot{\vec{\eta}}(t)$$

$$O_k(N) = \frac{\partial \ln \Psi_{N,\vec{\eta}}(t)}{\partial \eta_k} = \frac{\partial \ln}{\partial \eta_k} e^{\mathcal{H}_{\text{eff}}(N,\vec{\eta},t) + \ln \Psi_N}$$

$$E_{\text{loc}}(N) = \frac{\langle N | \mathcal{H} | \Psi_{\vec{\eta}}(t) \rangle}{\langle N | \Psi_{\vec{\eta}}(t) \rangle}$$



2025-03-12

Master Colloquium Presentation

└ Variational Classical Networks

└ Why and How should Parameters be variational?

1. Argument:

- Could go on performing the controlled expansion for all perturbation-strengths and get nice result
- But higher orders complicated and expensive
- Try to get more from the lower orders, by using Variational parameters (optimized by the problem, we hope they can incorporate influence from higher order terms without them being present)

2. Time dependent variational principle (TDVP)

- Find a parametrized *wavefunction*
- Update the parameters so that a tiny nudge into the direction of the *time DERIVATIVE* of the parameter vector has the same effect as a time evolution with the full hamiltonian
- The first order numerical integration (could do higher orders, but for the concept here)

3. O: Variational derivative E: local energy (we had already earlier)

Why and How should Parameters be variational?

Variational Classical Networks

- Theoretically: always better approximation through higher order terms
- Hard to calculate analytically and expensive to evaluate
- Hope: Better behavior if parameters are *learned/optimized*, starting from the analytical result

$$|\Psi_{\vec{\eta}+\delta\cdot\dot{\vec{\eta}}}(t+\delta)\rangle \leftrightarrow e^{-iH\delta} |\Psi_{\vec{\eta}}(t)\rangle$$
$$\vec{\eta}(t+\delta) = \vec{\eta}(t) + \delta \cdot \dot{\vec{\eta}}(t)$$
$$O_k(N) = \frac{\partial \ln \Psi_N(\vec{\eta})}{\partial \eta_k} = \frac{\partial \ln}{\partial \eta_k} e^{\mathcal{H}_{\text{eff}}(N,\vec{\eta}) + \ln \Psi_N}$$
$$E_{\text{loc}}(N) = \frac{\langle N | \mathcal{H} | \Psi_{\vec{\eta}}(t) \rangle}{\langle N | \Psi_{\vec{\eta}}(t) \rangle}$$

Why and How should Parameters be variational?

Variational Classical Networks

- Theoretically: always better approximation through higher order terms
 - Hard to calculate analytically and expensive to evaluate
- Hope: Better behavior if parameters are *learned/optimized*, starting from the analytical result

$$|\Psi_{\vec{\eta}+\delta\cdot\dot{\vec{\eta}}}(t+\delta)\rangle \leftrightarrow e^{-iH\delta} |\Psi_{\vec{\eta}}(t)\rangle$$

$$\vec{\eta}(t+\delta) = \vec{\eta}(t) + \delta \cdot \dot{\vec{\eta}}(t)$$

$$\vec{S}_{k,k'} = \langle O_k^* O_{k'} \rangle_{(\vec{\eta})} - \langle O_k^* \rangle_{(\vec{\eta})} \cdot \langle O_{k'} \rangle_{(\vec{\eta})}$$

$$\vec{F}_k = \langle E_{\text{loc}} O_k^* \rangle_{(\vec{\eta})} - \langle E_{\text{loc}} \rangle_{(\vec{\eta})} \cdot \langle O_k^* \rangle_{(\vec{\eta})}$$

$$\sum_{k'} \vec{S}_{k,k'} \dot{\vec{\eta}}_{k'} = -i \vec{F}_k$$

$$\dot{\vec{\eta}} = -i \overleftrightarrow{S}^{-1} \vec{F}$$



2025-03-12

Master Colloquium Presentation

└ Variational Classical Networks

└ Why and How should Parameters be variational?

1. Pseudo inversion
2. S: covariance matrix
3. F: TDVP force
4. *time DERIVATIVE*

Why and How should Parameters be variational?

Variational Classical Networks

- Theoretically: always better approximation through higher order terms
- Hard to calculate analytically and expensive to evaluate
- Hope: Better behavior if parameters are *learned/optimized*, starting from the analytical result

$$|\Psi_{\vec{\eta}+\delta\cdot\dot{\vec{\eta}}}(t+\delta)\rangle \leftrightarrow e^{-iH\delta} |\Psi_{\vec{\eta}}(t)\rangle$$

$$\vec{\eta}(t+\delta) = \vec{\eta}(t) + \delta \cdot \dot{\vec{\eta}}(t)$$

$$\vec{S}_{k,k'} = \langle O_k^* O_{k'} \rangle_{(\vec{\eta})} - \langle O_k^* \rangle_{(\vec{\eta})} \cdot \langle O_{k'} \rangle_{(\vec{\eta})}$$

$$\vec{F}_k = \langle E_{\text{loc}} O_k^* \rangle_{(\vec{\eta})} - \langle E_{\text{loc}} \rangle_{(\vec{\eta})} \cdot \langle O_k^* \rangle_{(\vec{\eta})}$$

$$\sum_{k'} \vec{S}_{k,k'} \dot{\vec{\eta}}_{k'} = -i \vec{F}_k$$

$$\dot{\vec{\eta}} = -i \overleftrightarrow{S}^{-1} \vec{F}$$

First Try: Cumulant Expansion Prefactors

Variational Classical Networks

- Idea: generate a variational Hamiltonian by replacing the analytical prefactors

$$\mathcal{H}_N(t) = \sum_i C_i(t) \cdot \Phi_i(N) \quad \leftrightarrow \quad \mathcal{H}_{\text{VCN}}(N, \vec{\eta}) = \sum_i \eta_i \cdot \Phi_i(N)$$

$C_1(t) = \Pi_A(0, 1, t)$	$\Phi_1(N) = J \sum_l \sum_{\langle l < m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N \hat{F}_A(l, m) K \rangle$	$C_5(t) = \Pi_B(1, 0, t)$	$\Phi_5(N) = J \sum_l \sum_{\langle l > m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N \hat{F}_B(l, m) K \rangle$
$C_2(t) = \Pi_B(0, 1, t)$	$\Phi_2(N) = J \sum_l \sum_{\langle l < m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N \hat{F}_B(l, m) K \rangle$	$C_6(t) = \Pi_C(1, 0, t)$	$\Phi_6(N) = J \sum_l \sum_{\langle l > m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N \hat{F}_C(l, m) K \rangle$
$C_3(t) = \Pi_C(0, 1, t)$	$\Phi_3(N) = J \sum_l \sum_{\langle l < m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N \hat{F}_C(l, m) K \rangle$	$C_7(t) = \Pi_A(0, M, t)$	$\Phi_7(N) = J \sum_l \sum_{\langle l < m \rangle^{+(y)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N \hat{F}_A(l, m) K \rangle$
$C_4(t) = \Pi_A(1, 0, t)$	$\Phi_4(N) = J \sum_l \sum_{\langle l > m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N \hat{F}_A(l, m) K \rangle$	...	...



2025-03-12

Master Colloquium Presentation

Variational Classical Networks

First Try: Cumulant Expansion Prefactors

- Idea: replace prefactors that depend on the time and keep the shape of the terms that depend on the occupation of the state
- Verification: as the TDVP is also derived from an *action principle* that must mean as per Noether's theorem that the energy MUST be conserved by a correct TDVP step

First Try: Cumulant Expansion Prefactors
Variational Classical Networks

■ Idea: generate a variational Hamiltonian by replacing the analytical prefactors

$$\mathcal{H}_N(t) = \sum_i C_i(t) \cdot \Phi_i(N) \quad \leftrightarrow \quad \mathcal{H}_{\text{VCN}}(N, \vec{\eta}) = \sum_i \eta_i \cdot \Phi_i(N)$$

$$C_1(t) = \Pi_A(0, 1, t) \quad \Phi_1(N) = J \sum_l \sum_{\langle l < m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N | \hat{F}_A(l, m) | K \rangle \quad C_5(t) = \Pi_B(1, 0, t) \quad \Phi_5(N) = J \sum_l \sum_{\langle l > m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N | \hat{F}_B(l, m) | K \rangle$$

$$C_2(t) = \Pi_B(0, 1, t) \quad \Phi_2(N) = J \sum_l \sum_{\langle l < m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N | \hat{F}_B(l, m) | K \rangle \quad C_6(t) = \Pi_C(1, 0, t) \quad \Phi_6(N) = J \sum_l \sum_{\langle l > m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N | \hat{F}_C(l, m) | K \rangle$$

$$C_3(t) = \Pi_C(0, 1, t) \quad \Phi_3(N) = J \sum_l \sum_{\langle l < m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N | \hat{F}_C(l, m) | K \rangle \quad C_7(t) = \Pi_A(0, M, t) \quad \Phi_7(N) = J \sum_l \sum_{\langle l < m \rangle^{+(y)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N | \hat{F}_A(l, m) | K \rangle$$

$$C_4(t) = \Pi_A(1, 0, t) \quad \Phi_4(N) = J \sum_l \sum_{\langle l > m \rangle^{+(x)}} \sum_K \frac{\Psi_K}{\Psi_N} \langle N | \hat{F}_A(l, m) | K \rangle \quad \dots \quad \dots$$

Correction: Watch Explicit Time-Dependency

Variational Classical Networks

- First strategy is not suitable
 - Energy- and variance is not conserved
- Problem: explicit time-dependency of base energy

$$\frac{\partial \mathcal{H}_{\text{eff}}(N, \vec{\eta}, t)}{\partial \vec{\eta}_k} = \frac{\partial (-iE_0(N)t + \mathcal{H}_{\text{VCN}}(N, \vec{\eta}))}{\partial \vec{\eta}_k}$$

- Solution: Replace base energy factors with variational parameters

$$E_0(N) = U \cdot \sum_m n_{m,\uparrow} n_{m,\downarrow} + \sum_{l,\sigma} \varepsilon_l n_{l,\sigma}$$

$$\eta_l^{\text{b.e.}} = \begin{cases} l=0 : & -i \cdot U \cdot t \\ l>0 : & -i \cdot \varepsilon_l \cdot t \end{cases}$$

$$\Phi_l^{\text{b.e.}}(N) = \begin{cases} l=0 : & \sum_m n_{m,\uparrow} n_{m,\downarrow} \\ l>0 : & \sum_{\sigma} n_{l,\sigma} \end{cases}$$

$$\mathcal{H}_{\text{eff}}^{\text{VCN}}(N, \vec{\eta}) = \sum_{n=1}^{\#(\text{var.})} \eta_n^{\text{var.}} \cdot \Phi_n^{\text{var.}}(N) + \sum_{l=0}^L \eta_l^{\text{b.e.}} \cdot \Phi_l^{\text{b.e.}}(N) = \sum_l \eta_l \cdot \Phi_l(N)$$

$$\vec{\eta} = (\eta_1^{\text{var.}}, \eta_2^{\text{var.}}, \dots, \eta_{\#(\text{var.})}^{\text{var.}}, \eta_0^{\text{b.e.}}, \eta_1^{\text{b.e.}}, \dots, \eta_L^{\text{b.e.}})^T$$



2025-03-12

Master Colloquium Presentation

Variational Classical Networks

Correction: Watch Explicit Time-Dependency

1. Fails because not taken the explicit time-dependency into account as it was *differentiated away* wrongly
2. As a solution: introduce variational base-energy factors - which is a successful strategy

Correction: Watch Explicit Time-Dependency

Variational Classical Networks

- First strategy is not suitable
 - Energy- and variance is not conserved
- Problem: explicit time-dependency of base energy

$$\frac{\partial \mathcal{H}_{\text{eff}}(N, \vec{\eta}, t)}{\partial \vec{\eta}_k} = \frac{\partial (-iE_0(N)t + \mathcal{H}_{\text{VCN}}(N, \vec{\eta}))}{\partial \vec{\eta}_k}$$
- Solution: Replace base energy factors with variational parameters

$$\mathcal{H}_{\text{eff}}(N) = U \cdot \sum_m n_{m,\uparrow} n_{m,\downarrow} + \sum_{l,\sigma} \varepsilon_l n_{l,\sigma}$$

$$\eta_l^{\text{b.e.}} = \begin{cases} l=0 : & -i \cdot U \cdot t \\ l>0 : & -i \cdot \varepsilon_l \cdot t \end{cases} \quad \mathcal{H}_{\text{eff}}^{\text{VCN}}(N, \vec{\eta}) = \sum_{n=1}^{\#(\text{var.})} \eta_n^{\text{var.}} \cdot \Phi_n^{\text{var.}}(N) + \sum_{l=0}^L \eta_l^{\text{b.e.}} \cdot \Phi_l^{\text{b.e.}}(N) = \sum_l \eta_l \cdot \Phi_l(N)$$

$$\vec{\eta} = (\eta_1^{\text{var.}}, \eta_2^{\text{var.}}, \dots, \eta_{\#(\text{var.})}^{\text{var.}}, \eta_0^{\text{b.e.}}, \eta_1^{\text{b.e.}}, \dots, \eta_L^{\text{b.e.}})^T$$

$$\Phi_l^{\text{b.e.}}(N) = \begin{cases} l=0 : & \sum_m n_{m,\uparrow} n_{m,\downarrow} \\ l>0 : & \sum_{\sigma} n_{l,\sigma} \end{cases}$$

Implementation

Implementation & Experiments

- Too much code to properly show
 - Look at repository for overview
- Testing: validation code to check the correct implementation
 - Independent implementations for same functionality
 - Sanity checks for state of data and program
- Optimization: runtime comparison for different code styles
 - For generated code fast execution over readability



2025-03-12

Master Colloquium Presentation

- └ Implementation & Experiments
 - └ Implementation

1. While nothing is looked at in detail maybe interesting for some
2. Testing: validation and assertion based programming
3. Optimization: code style runtime experiments


Implementation
Implementation & Experiments

- Too much code to properly show
 - Look at repository for overview
- Testing: validation code to check the correct implementation
 - Independent implementations for same functionality
 - Sanity checks for state of data and program
- Optimization: runtime comparison for different code styles
 - For generated code fast execution over readability

Numerical Experiments

Implementation & Experiments

- Sadly no presentation possible in the small time allocated for the presentation
- The plots are in the presentation in the supplementary material



2025-03-12


Master Colloquium Presentation
└ Implementation & Experiments
└ Numerical Experiments

1. See supplementary material for more plots and stuff

Numerical Experiments

Implementation & Experiments

- Sadly no presentation possible in the small time allocated for the presentation
- The plots are in the presentation in the supplementary material



Conclusion & Outlook

Conclusion & Outlook

Conclusion:

- Successfully transferred and validated the theory
- Working implementation
 - Extensively tested
 - Configurable and extensible
- Started validating expectations with measurements on the compute cluster

Outlook:

- Re-write in lower level language to gain performance over Python
- Start more real experiments now that the theory is validated to work



2025-03-12

Master Colloquium Presentation

└ Conclusion & Outlook

└ Conclusion & Outlook

1. Conclusion:

- Managed to Solve all problems
- Provide heavily *tested* and *extensible* library of code AND theoretical summary
- Started to validate expectations with computations on the cluster

2. Future plans:

- re-write in performant language after testing is successful
- Go through the parameters to start investigating the behavior of the systems

Conclusion & Outlook

Conclusion & Outlook

Conclusion:

- Successfully transferred and validated the theory
- Working implementation
 - Extensively tested
 - Configurable and extensible
- Started validating expectations with measurements on the compute cluster

Outlook:

- Re-write in lower level language to gain performance over Python
- Start more real experiments now that the theory is validated to work

Thank you for your kind attention



1. Thank you for your kind attention
2. All tools and other resources are referenced in the presentation
3. You can also find everything on my Github



References I

Summary & Conclusion

- [1] J. Kell, *Latex sources for this presentation and the main thesis*, (2025)
<https://github.com/jonas-kell/master-thesis-documents>.
- [2] J. Kell, *Thesis code implementation and reference*, (2025)
<https://github.com/jonas-kell/master-thesis-code>.
- [3] J. Kell, *Https://github.com/jonas-kell/beamer-theme-unia-mntf*, (2024)
<https://github.com/jonas-kell/math-manipulator>.



Extra Plots: Overview

Extra slides

- Exact comparison density matrix
- First orders: energy and variance comparison
- J parameter sweep
- Monte Carlo sampling convergence
- Runtime validation
- VCN: parametrization failure
- VCN: energy conservation can be controlled
- VCN: minimal square reference system
- VCN: larger square reference system in direct comparison
- VCN: system size dependency



2025-03-12

Master Colloquium Presentation

└ Extra slides

└ Extra Plots: Overview


Extra Plots: Overview

Extra slides

- Exact comparison density matrix
- First orders: energy and variance comparison
- J parameter sweep
- Monte Carlo sampling convergence
- Runtime validation
- VCN: parametrization failure
- VCN: energy conservation can be controlled
- VCN: minimal square reference system
- VCN: larger square reference system in direct comparison
- VCN: system size dependency

Exact Comparison: Concurrence

Extra slides




2025-03-12

Master Colloquium Presentation

Extra slides


Exact Comparison: Concurrence



1. Concurrence comparison of small system with exact calculations and comparison thereof
2. See extreme boost of second order
3. NOTICE: how the External calculation is done COMPLETELY on density matrices and diagonalization, so proves we can sample the reduced density matrix out

Exact Comparison: Purity

Extra slides



UNIA

2025-03-12


Master Colloquium Presentation

Extra slides

Exact Comparison: Purity

Exact Comparison: Purity


Extra slides



1. Purity measurements for the graph of the measurement from one page before

Exact Comparison: Pauli Measurements



Extra slides



1. Portrait mode format, because extracted from the thesis.
2. You can see what a problem the z-contributions are

First Orders: Energy Comparison

Extra slides



2025-03-12

Master Colloquium Presentation


└ Extra slides

└ First Orders: Energy Comparison

1. Second picture zoomed in



First Orders: Energy Comparison

Extra slides



First Orders: Variance Comparison

Extra slides




2025-03-12

Master Colloquium Presentation

└ Extra slides


└ First Orders: Variance Comparison

1. Second picture zoomed in



J-Sweep: Current Border

Extra slides



UJA


2025-03-12

Master Colloquium Presentation

Extra slides


J-Sweep: Current Border

1. Logarithmic plot
2. Measurements show the trend of the convergence of the relative error for small perturbations
3. Depends on the observable though, still



J-Sweep: Current Center

Extra slides




Uia

2025-03-12

Master Colloquium Presentation


Extra slides

J-Sweep: Current Center



J-Sweep: Single Occupation Center

Extra slides




Uia

2025-03-12

Master Colloquium Presentation


Extra slides

J-Sweep: Single Occupation Center



J-Sweep: Double Occupation Border

Extra slides




2025-03-12

Master Colloquium Presentation



Extra slides

J-Sweep: Double Occupation Border




Monte Carlo Convergence: Occupation

Extra slides




1. Well as we can see, the appriximation converges as the variance gos to zero



Monte Carlo Convergence: Occupation

Extra slides




2025-03-12

Master Colloquium Presentation


└ Extra slides

└ Monte Carlo Convergence: Occupation



Runtime Validation

Extra slides




2025-03-12

Master Colloquium Presentation

└ Extra slides



└ Runtime Validation

Runtime Validation
Extra slides




VCN Failure: Non-Explicit Parametrization

Extra slides




1. Complex part now properly vanishes (was long a problem)
2. See how the vcn goes like the base energy curve only in the beginning (suspiciously)



VCN Failure: Non-Explicit Parametrization

Extra slides



2025-03-12

Master Colloquium Presentation


└ Extra slides

└ VCN Failure: Non-Explicit Parametrization

1. See complex part jetting off to infinity because it doesn't know how to compensate

VCN Validation: Energy Conservation Controllable

Extra slides




2025-03-12

Master Colloquium Presentation

└ Extra slides



└ VCN Validation: Energy Conservation Controllable

1. Still no perfect conservation for infinite steps, but this is probably due to first order integration only




VCN: Small Square System

Extra slides





1. Notice how VCN now is MUCH better




VCN: Small Square System

Extra slides




1. Step size dependent version of previous slide



VCN: Bigger Square System



Extra slides



1. Now not a guarantee for immediately better results, but very promising

VCN: Bigger Square System

Extra slides




2025-03-12

Master Colloquium Presentation

└ Extra slides



└ VCN: Bigger Square System

1. Here one can see very quickly more steps doesn't only make it better



VCN End to End Test: System Size Dependency

Extra slides




2025-03-12

Master Colloquium Presentation

└ Extra slides



└ VCN End to End Test: System Size Dependency

1. This is the energy for validation



VCN End to End Test: System Size Dependency

Extra slides




2025-03-12

Master Colloquium Presentation

└ Extra slides



└ VCN End to End Test: System Size Dependency

1. This is the purity
2. Does not seem to have a huge variation



VCN End to End Test: System Size Dependency



Extra slides



1. This is the concurrence
2. Poorly resolved, this should be looked into possibly in the future

VCN End to End Test: System Size Dependency

Extra slides




2025-03-12

Master Colloquium Presentation

└ Extra slides

└ VCN End to End Test: System Size Dependency



1. This is the current
2. Most obvious effect, but still extremely stable over a order of magnitude of sites
3. See assymmetric behavior, because electrical field is set to not be symmetric