

## Foundations of Artificial Intelligence

Prof. Dr. J. Boedecker, Prof. Dr. W. Burgard, Prof. Dr. F. Hutter, Prof. Dr. B. Nebel,  
Dr. rer. nat. M. Tangemann  
T. Schulte, M. Krawez, R. Rajan, S. Adriaensen, K. Sirohi  
Summer Term 2020

University of Freiburg  
Department of Computer Science

### Exercise Sheet 5 — Solutions

#### Exercise 5.1 (Modeling, Proofs)

Consider the following knowledge base:

*If the unicorn is mythical, then it is immortal, but if it is not mythical, then it is a mortal mammal. If the unicorn is immortal or a mammal, then it is horned. The unicorn is magical if it is horned.*

Using this knowledge base, can you prove that the unicorn is (a) mythical, (b) magical or (c) horned? First, formalize the knowledge base with propositional logic. If a statement is valid or unsatisfiable, use resolution for prove. Else, write down one satisfying and one unsatisfying interpretation.

#### Solution:

The above statements can be formalized using the atomic propositions *mythical*, *mortal*, *mammal*, *magical* and *horned*:

$$\text{mythical} \rightarrow \neg \text{mortal} \quad (\text{i})$$

$$\neg \text{mythical} \rightarrow \text{mortal} \wedge \text{mammal} \quad (\text{ii})$$

$$\neg \text{mortal} \vee \text{mammal} \rightarrow \text{horned} \quad (\text{iii})$$

$$\text{horned} \rightarrow \text{magical} \quad (\text{iv})$$

Let  $KB = (i) \wedge (ii) \wedge (iii) \wedge (iv)$  be the set of those four propositions, in CNF:

$$\{\neg \text{mythical}, \neg \text{mortal}\} \quad (1)$$

$$\{\text{mythical}, \text{mortal}\} \quad (2)$$

$$\{\text{mythical}, \text{mammal}\} \quad (3)$$

$$\{\text{mortal}, \text{horned}\} \quad (4)$$

$$\{\neg \text{mammal}, \text{horned}\} \quad (5)$$

$$\{\neg \text{horned}, \text{magical}\} \quad (6)$$

- (a) We cannot tell from  $KB$  whether the unicorn is mythical or not, since there exist two models  $I_{\text{mythical}}$  and  $I_{\neg \text{mythical}}$  of  $KB$  with  $I_{\text{mythical}} \models \text{mythical}$  and  $I_{\neg \text{mythical}} \models \neg \text{mythical}$ .

More specific:

$$I_{\text{mythical}} = \{\text{mythical} \mapsto 1, \text{mortal} \mapsto 0, \text{mammal} \mapsto 0, \text{magical} \mapsto 1, \text{horned} \mapsto 1\}$$

and

$$I_{\neg \text{mythical}} = \{\text{mythical} \mapsto 0, \text{mortal} \mapsto 1, \text{mammal} \mapsto 1, \text{magical} \mapsto 1, \text{horned} \mapsto 1\}.$$

Thus,  $KB \not\models \neg \text{mythical}$  and  $KB \not\models \text{mythical}$ , since neither *mythical* nor  $\neg \text{mythical}$  are true in *all* models of  $KB$ . In other words, the unicorn could be mythical or not, depending on the choice of model for  $KB$ .

- (b)  $KB \models \text{horned}$  holds, since  $KB \cup \{\neg \text{horned}\}$  is unsatisfiable. Proof:

Let

$$\{\neg \text{horned}\} \tag{7a}$$

then

$$(5) + (7a) : \quad \{\neg \text{mammal}\} \tag{8a}$$

$$(4) + (7a) : \quad \{\text{mortal}\} \tag{9a}$$

$$(3) + (8a) : \quad \{\text{mythical}\} \tag{10a}$$

$$(1) + (10a) : \quad \{\neg \text{mortal}\} \tag{11a}$$

$$(11a) + (9a) : \quad \emptyset$$

- (c)  $KB \models \text{magical}$  holds, since  $KB \cup \{\neg \text{magical}\}$  is unsatisfiable. Proof:

Let

$$\{\neg \text{magical}\} \tag{7b}$$

then

$$(6) + (7b) : \quad \{\neg \text{horned}\} \tag{8b}$$

$$(5) + (8b) : \quad \{\neg \text{mammal}\} \tag{9b}$$

$$(4) + (8b) : \quad \{\text{mortal}\} \tag{10b}$$

$$(3) + (9b) : \quad \{\text{mythical}\} \tag{11b}$$

$$(1) + (11b) : \quad \{\neg \text{mortal}\} \tag{12b}$$

$$(12b) + (10b) : \quad \emptyset$$

### Exercise 5.2 (DPLL)

Use the Davis-Putnam-Logemann-Loveland (DPLL) procedure to check whether the given formulae  $\phi_1$  and  $\phi_2$  are satisfiable or not. Write down all steps carried out by the algorithm during the process. If you have to apply a splitting rule, split on variables in alphabetical order, trying *true* first, then *false*. Should the formula be satisfiable, please indicate the satisfying assignment.

(a)

$$\phi_1 = (D \vee C) \wedge (\neg A \vee \neg D \vee B \vee \neg C) \wedge (A \vee C) \wedge (\neg C \vee \neg B) \wedge (\neg C \vee B \vee \neg A \vee D) \wedge (C \vee \neg D \vee B) \wedge (\neg D \vee \neg B \vee \neg A)$$

**Solution:**

$$\begin{aligned} \text{Splitting } A \rightarrow 1 & \quad (D \vee C) \wedge (\neg D \vee B \vee \neg C) \wedge (\neg C \vee \neg B) \wedge (\neg C \vee B \vee D) \wedge \\ & (C \vee \neg D \vee B) \wedge (\neg D \vee \neg B) \\ \text{Splitting } B \rightarrow 1 & \quad (D \vee C) \wedge \neg C \wedge \neg D \\ \text{Unit-propagation } C \rightarrow 0 & \quad D \wedge \neg D \\ \text{Unit-propagation } D \rightarrow 1 & \quad \perp \\ \text{Backtracking } B \rightarrow 0 & \quad (D \vee C) \wedge (\neg D \vee \neg C) \wedge (\neg C \vee D) \wedge (C \vee \neg D) \\ \text{Splitting } C \rightarrow 1 & \quad \neg D \wedge D \\ \text{Unit-propagation } D \rightarrow 1 & \quad \perp \\ \text{Backtracking } C \rightarrow 0 & \quad D \wedge \neg D \\ \text{Unit-propagation } D \rightarrow 1 & \quad \perp \\ \text{Backtracking } A \rightarrow 0 & \quad (D \vee C) \wedge C \wedge (\neg C \vee \neg B) \wedge (C \vee \neg D \vee B) \\ \text{Unit-propagation } C \rightarrow 1 & \quad \neg B \\ \text{Unit-propagation } B \rightarrow 0 & \quad \top \end{aligned}$$

Satisfying assignment:  $A \rightarrow 0; B \rightarrow 0; C \rightarrow 1; D \rightarrow 1$  or  $0$ ;

(b)

$$\phi_2 = (D \vee \neg A \vee B) \wedge (\neg B \vee \neg C \vee A \vee D) \wedge (\neg B \vee \neg A) \wedge (B \vee \neg D) \wedge (A \vee C \vee \neg B) \wedge (\neg D \vee \neg C) \wedge (B \vee D)$$

**Solution:**

$$\begin{aligned} \text{Splitting } A \rightarrow 1 & \quad (D \vee B) \wedge \neg B \wedge (B \vee \neg D) \wedge (\neg D \vee \neg C) \wedge (B \vee D) \\ \text{Unit-propagation } B \rightarrow 0 & \quad D \wedge \neg D \wedge (\neg D \vee \neg C) \wedge D \\ \text{Unit-propagation } D \rightarrow 1 & \quad \perp \\ \text{Backtracking } A \rightarrow 0 & \quad (\neg B \vee \neg C \vee D) \wedge (B \vee \neg D) \wedge (C \vee \neg B) \wedge (\neg D \vee \neg C) \wedge (B \vee D) \\ \text{Splitting } B \rightarrow 1 & \quad (\neg C \vee D) \wedge C \wedge (\neg D \vee \neg C) \\ \text{Unit-propagation } C \rightarrow 1 & \quad D \wedge \neg D \\ \text{Unit-propagation } D \rightarrow 1 & \quad \perp \\ \text{Backtracking } B \rightarrow 0 & \quad \neg D \wedge (\neg D \vee \neg C) \wedge D \\ \text{Unit-propagation } D \rightarrow 1 & \quad \perp \end{aligned}$$

The formula is unsatisfiable.