

## Foundations of Artificial Intelligence

Prof. Dr. J. Boedecker, Prof. Dr. W. Burgard, Prof. Dr. F. Hutter, Prof. Dr. B. Nebel,  
Dr. rer. nat. M. Tangermann  
T. Schulte, M. Krawez, R. Rajan, S. Adriaensen, K. Sirohi  
Summer Term 2020

University of Freiburg  
Department of Computer Science

### Exercise Sheet 7 — Solutions

#### Exercise 7.1 (Planning)

Consider the following STRIPS-Task  $\Pi = \langle \mathcal{S}, O, I, G \rangle$ :

- $\mathcal{S}$ :  $\{X, Y, Z, G\}$
- $O$ :  $\{A, B, C, D, E, F\}$  where
$$\begin{array}{ll} A : pre(A) = \{X\}, & eff(A) = \{Y, Z\} \\ B : pre(B) = \{X\}, & eff(B) = \{\neg X, Z\} \\ C : pre(C) = \{\neg Y\}, & eff(C) = \{Z\} \\ D : pre(D) = \{\neg Z\}, & eff(D) = \{Y\} \\ E : pre(E) = \{\neg X, Y\}, & eff(E) = \{\neg Y, G\} \\ F : pre(F) = \{Z\}, & eff(F) = \{\neg Z, G\} \end{array}$$
- $I$ :  $\{X, Y\}$
- $G$ :  $\{G\}$

- (a) State for each operator from  $O$  if it is applicable in  $I$  or not. For each applicable operator also give the resulting state after applying that operator in  $I$ .

**Solution:**

Operator	Applicable?	Resulting State
$A$	Yes	$\{X, Y, Z\}$
$B$	Yes	$\{Y, Z\}$
$C$	No	-
$D$	Yes	$\{X, Y\}$
$E$	No	-
$F$	No	-

- (b) Give an applicable plan  $\pi$  that leads from  $I$  to  $G$ .

**Solution:**

$$\pi = \langle B, E \rangle, \langle A, F \rangle, \langle B, F \rangle, \dots$$

**Exercise 7.2** (Bayes' Rule)

In Freiburg 80% of all cars are red. You see a car at night that does *not* appear red to you. You know that you can correctly identify a red car only in 70% of the cases when the given car is red. And you can identify a non-red car correctly in 90% of the cases when the given car is non-red.

- (a) List all conditional and non-conditional probabilities that you can determine directly from the task description. Note: Differentiate between the statement that a car *is red* and the statement that you have *seen a red car*.
- (b) Compute the probability that the car is actually red, when you perceive a car as red in Freiburg at night.

**Solution:**

$P(R)$ : car is red

$P(PR)$ : car is perceived red

$P(R) = 0.8, P(\neg R) = 0.2$

$P(PR|R) = 0.7$

$P(\neg PR|\neg R) = 0.9$

$$\begin{aligned}
 P(R|PR) &= \frac{P(PR|R) \cdot P(R)}{P(PR)} \\
 &= \frac{P(PR|R) \cdot P(R)}{P(PR|R) \cdot P(R) + P(PR|\neg R) \cdot P(\neg R)} \\
 &= \frac{0.7 \cdot 0.8}{0.7 \cdot 0.8 + 0.1 \cdot 0.2} \\
 &= \frac{0.56}{0.56 + 0.02} = \frac{0.56}{0.58} = \frac{28}{29}
 \end{aligned}$$

**Exercise 7.3** (Independence and Joint and Conditional Probabilities)

- (a) A 6-sided die is rolled once. Which of the following events are independent? Show the probability values and reasoning.
  - $E$  : An even number is rolled
  - $O$  : An odd number is rolled
  - $T$  : A number  $\geq 3$  is rolled

**Solution:**

We know there are 6 possible outcomes for the roll of the die.

$$P(E) = 0.5$$

3 out of 6 possibilities are covered under the event

$$P(O) = 0.5$$

3 out of 6 possibilities are covered under the event

$$P(T) = \frac{2}{3}$$

4 out of 6 possibilities are covered under the event

$$P(E \cap O) = 0 \neq P(E) * P(O) = 0.25 \quad \text{E and O are disjoint events}$$

$$P(E \cap T) = \frac{1}{3} = P(E) * P(T) = \frac{1}{3} \quad \text{E and T cover 2 out of 6 possibilities}$$

$$P(O \cap T) = \frac{1}{3} = P(O) * P(T) = \frac{1}{3} \quad \text{O and T cover 2 out of 6 possibilities}$$

By definition of independence, E and T are independent and O and T are independent.

- (b) Make the joint probability distribution table for the events E and T.

**Solution:**

	$E = False$	$E = True$
$T = False$	0.167	0.167
$T = True$	0.333	0.333

- (c) Calculate the conditional probability  $P(\neg e | t)$ .

**Solution:**

$$P(\neg e | t) = P(\neg e \wedge t) / P(t) = 0.333 / 0.666 = 0.5$$