

# Numerical Optimal Control

## Lecture 8: Indirect Optimal Control

Sébastien Gros

ITK, NTNU

NTNU PhD course

# Outline

- 1 Introduction to the Pontryagin Maximum Principle (PMP)
- 2 Interpretation of  $H_u$
- 3 Input bounds in Indirect Optimal Control
- 4 Singular Optimal Control problems
- 5 General constraints in Indirect Optimal Control

## Overview - let's take a step back...

	Continuous Equations	Discrete Equations
Global	Hamilton-Jacobi-Bellman (HJB)	Dynamic Programming (DP)
Local	Pontryagin (PMP)	Direct Optimal Control (DOC)

**Continuous problem:**

$$\min_{\mathbf{u} \in \mathcal{U}} \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}),$$

$$\mathbf{x}(t_0) = \mathbf{x}_0$$

## Overview - let's take a step back...

	Continuous Equations	Discrete Equations
Global	Hamilton-Jacobi-Bellman (HJB)	Dynamic Programming (DP)
Local	Pontryagin (PMP)	Direct Optimal Control (DOC)

**Continuous problem:**

$$\min_{\mathbf{u} \in \mathcal{U}} \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}),$$

$$\mathbf{x}(t_0) = \mathbf{x}_0$$

**PMP:** define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$

Get the optimal input  $\mathbf{u}(\mathbf{x}, \boldsymbol{\lambda}) = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$

Use it in the state-costate integration:

$$\text{States : } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\text{Costates : } \dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}), \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$$

## Overview - let's take a step back...

	Continuous Equations	Discrete Equations
Global	Hamilton-Jacobi-Bellman (HJB)	Dynamic Programming (DP)
Local	Pontryagin (PMP)	Direct Optimal Control (DOC)

### Continuous problem:

$$\begin{aligned} \min_{\mathbf{u} \in \mathcal{U}} \quad & \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt \\ \text{s.t.} \quad & \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \\ & \mathbf{x}(t_0) = \mathbf{x}_0 \end{aligned}$$

- PMP equations provide an " $\infty$ "-dimensional input profile  $\mathbf{u}(\cdot)$
- State constraints hard to handle

PMP: define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$

Get the optimal input  $\mathbf{u}(\mathbf{x}, \boldsymbol{\lambda}) = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$

Use it in the state-costate integration:

$$\text{States : } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\text{Costates : } \dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}), \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$$

## Overview - let's take a step back...

	Continuous Equations	Discrete Equations
Global	Hamilton-Jacobi-Bellman (HJB)	Dynamic Programming (DP)
Local	Pontryagin (PMP)	Direct Optimal Control (DOC)

**Continuous problem:**

$$\min_{\mathbf{u} \in \mathcal{U}} \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}),$$

$$\mathbf{x}(t_0) = \mathbf{x}_0$$

**D.O.C.** describes the solution as a finite set of variables  $\mathbf{w}$  transform the problem into a discrete one

Solve the resulting Nonlinear Program (NLP):

$$\min_{\mathbf{w}} \Phi(\mathbf{w})$$

$$\begin{aligned} \text{s.t. } & \mathbf{g}(\mathbf{w}) = 0, \\ & \mathbf{h}(\mathbf{w}) \leq 0 \end{aligned}$$

## Overview - let's take a step back...

	Continuous Equations	Discrete Equations
Global	Hamilton-Jacobi-Bellman (HJB)	Dynamic Programming (DP)
Local	Pontryagin (PMP)	Direct Optimal Control (DOC)

### Continuous problem:

$$\min_{\mathbf{u} \in \mathcal{U}} \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \\ \mathbf{x}(t_0) = \mathbf{x}_0$$

- Input profile restricted to a finite-dimensional space (e.g. piecewise-constant)
- Easy to treat all types of constraints

**D.O.C.** describes the solution as a finite set of variables  $\mathbf{w}$  transform the problem into a discrete one

Solve the resulting Nonlinear Program (NLP):

$$\begin{aligned} & \min_{\mathbf{w}} \Phi(\mathbf{w}) \\ & \text{s.t. } \mathbf{g}(\mathbf{w}) = 0, \\ & \quad \mathbf{h}(\mathbf{w}) \leq 0 \end{aligned}$$

## Overview - let's take a step back...

	Continuous Equations	Discrete Equations
Global	Hamilton-Jacobi-Bellman (HJB)	Dynamic Programming (DP)
Local	Pontryagin (PMP)	Direct Optimal Control (DOC)

**The big distinction in Optimal Control...**

## Overview - let's take a step back...

	Continuous Equations	Discrete Equations
Global	Hamilton-Jacobi-Bellman (HJB)	Dynamic Programming (DP)
Local	Pontryagin (PMP)	Direct Optimal Control (DOC)

**The big distinction in Optimal Control...**

**"First optimise then discretize" (HJB & PMP)**

- First write the **continuous equations** describing the solution to the problem
- ... then **discretize the equations** & solve

Note: the PMP "family" is referred to as *Indirect* optimal control here

## Overview - let's take a step back...

	Continuous Equations	Discrete Equations
Global	Hamilton-Jacobi-Bellman (HJB)	Dynamic Programming (DP)
Local	Pontryagin (PMP)	Direct Optimal Control (DOC)

**The big distinction in Optimal Control...**

**"First optimise then discretize" (HJB & PMP)**

- First write the **continuous equations** describing the solution to the problem
- ... then **discretize the equations** & solve

Note: the PMP "family" is referred to as *Indirect* optimal control here  
vs.

**"First discretize then optimise" (DP & DOC)**

- **First discretize** the continuous OCP into a discrete one...
- ... then write the **discrete equations** describing the solution & solve

# Pontryagin Maximum Principle

**Simple continuous problem:**

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}),$$

$$\mathbf{x}(t_0) = \mathbf{x}_0$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$

# Pontryagin Maximum Principle

**Simple continuous problem:**

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}),$$

$$\mathbf{x}(t_0) = \mathbf{x}_0$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$

Get the optimal control solution from  $\mathbf{u} = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$

# Pontryagin Maximum Principle

## Simple continuous problem:


$$\min_{\mathbf{x}, \mathbf{u}} \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}),$$

$$\mathbf{x}(t_0) = \mathbf{x}_0$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$



Get the optimal control solution from  $\mathbf{u} = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$  with:

States :  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}), \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

# Pontryagin Maximum Principle

## Simple continuous problem:


$$\min_{\mathbf{x}, \mathbf{u}} \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}),$$

$$\mathbf{x}(t_0) = \mathbf{x}_0$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$



Get the optimal control solution from  $\mathbf{u} = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$  with:

States :  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}), \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

# Pontryagin Maximum Principle

## Simple continuous problem:


$$\min_{\mathbf{x}, \mathbf{u}} \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}),$$

$$\mathbf{x}(t_0) = \mathbf{x}_0$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$



Get the optimal control solution from  $\mathbf{u} = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$  with:

$$\text{States : } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\text{Costates : } \dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}), \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$$

## Two Points Boundary Value Problem

- Integrate forward ? We have  $\mathbf{x}(t_0) = \mathbf{x}_0$ , but we don't have  $\boldsymbol{\lambda}(t_0)$ ...

# Pontryagin Maximum Principle

## Simple continuous problem:


$$\min_{\mathbf{x}, \mathbf{u}} \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}),$$

$$\mathbf{x}(t_0) = \mathbf{x}_0$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$



Get the optimal control solution from  $\mathbf{u} = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$  with:

States :  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}), \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

## Two Points Boundary Value Problem

- Integrate forward ? We have  $\mathbf{x}(t_0) = \mathbf{x}_0$ , but we don't have  $\boldsymbol{\lambda}(t_0)$ ...
- Integrate forward-backward ? Integrate state forward from  $\mathbf{x}(t_0) = \mathbf{x}_0$  then backward from  $\boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$ ... but we don't know  $\mathbf{u}$ ...

# Pontryagin Maximum Principle

Simple continuous problem:


$$\min_{\mathbf{x}, \mathbf{u}} \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}),$$

$$\mathbf{x}(t_0) = \mathbf{x}_0$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$



Get the optimal control solution from  $\mathbf{u} = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$  with:

States :  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}), \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

## Two Points Boundary Value Problem

Note that the **entire** solution is "described by"  $\boldsymbol{\lambda}(t_0)$

# Solving the PMP equations / TPBVP



**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :



$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

**TPBVP**



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$



$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

**TPBVP**



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:



$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$

**TPBVP**



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:

$$\dot{x} = -\lambda - \sin(x)$$



$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$

**TPBVP**



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:

$$\dot{x} = -\lambda - \sin(x)$$


$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$

Guess  $\lambda_0 = 0.41$  / Current  $\lambda_0 = 0.41$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:

$$\dot{x} = -\lambda - \sin(x)$$


$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$

Guess  $\lambda_0 = 0.41$  / Current  $\lambda_0 = 1.6486$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:

$$\dot{x} = -\lambda - \sin(x)$$


$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$

Guess  $\lambda_0 = 0.41$  / Current  $\lambda_0 = 0.57211$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:

$$\dot{x} = -\lambda - \sin(x)$$


$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$

Guess  $\lambda_0 = 0.41$  / Current  $\lambda_0 = 0.49677$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:

$$\dot{x} = -\lambda - \sin(x)$$


$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$

Guess  $\lambda_0 = 0.41$  / Current  $\lambda_0 = 0.38536$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:

$$\dot{x} = -\lambda - \sin(x)$$


$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$

Guess  $\lambda_0 = 0.41$  / Current  $\lambda_0 = 0.53059$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:

$$\dot{x} = -\lambda - \sin(x)$$


$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$

Guess  $\lambda_0 = 0.41$  / Current  $\lambda_0 = 0.43026$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



# Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

---

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$

What is going on ?!?

Let's try to understand the relationship  
 $\lambda_0 \rightarrow \lambda(t)$

## Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

---

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



## Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

---

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



## Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

---

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



## Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

---

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



## Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

---

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



## Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

---

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



## Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

---

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



## Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

---

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



## Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

---

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



## Solving the PMP equations / TPBVP

**Input:** Initial conditions  $x_0$ , guess  $\lambda_0$

while  $\|r\| > \text{tol}$  do

Integrate with  $u = \arg \min_u H(x, \lambda, u)$ :

$$\dot{x} = F(x, u), \quad x(t_0) = x_0$$

$$\dot{\lambda} = -\nabla_x H(x, \lambda, u), \quad \lambda(t_0) = \lambda_0$$

Compute:

$$r = \lambda(t_f) - \nabla_x \phi(x(t_f)) \quad \text{and} \quad \frac{\partial r}{\partial \lambda_0}$$

$$\text{Newton step: } \lambda_0 \leftarrow \lambda_0 - \frac{\partial r}{\partial \lambda_0}^{-1} r$$

---

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

is minimised by  $u = -\lambda$ . Dynamics read as:


$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(4) = 0$  !!

**Example:**

$$\begin{aligned} \min_{x, u} \quad & \frac{1}{2} \int_0^4 (x^2 + u^2) dt \\ \dot{x} = u - \sin(x), \quad & x(0) = 1 \end{aligned}$$



## Conservation of volumes in the $x, \lambda$ space

Consider a compact domain  $D(t)$  in the  $x(t), \lambda(t)$  space, with boundary  $\partial D(t)$ . The volume of  $D(t)$  say  $\rho(D(t)) \in \mathbb{R}$  is given by:

$$\rho(D(t)) = \int_{D(t)} dD$$

and evolves with time as the integration of  $x(t), \lambda(t)$  proceeds.

## Conservation of volumes in the $x, \lambda$ space

Consider a compact domain  $D(t)$  in the  $x(t), \lambda(t)$  space, with boundary  $\partial D(t)$ . The volume of  $D(t)$  say  $\rho(D(t)) \in \mathbb{R}$  is given by:

$$\rho(D(t)) = \int_{D(t)} dD$$

and evolves with time as the integration of  $x(t), \lambda(t)$  proceeds. Its time-derivative reads as:

$$\frac{d}{dt} \rho(D(t)) = \int_{\partial D(t)} \left\langle \begin{bmatrix} \dot{x} \\ \dot{\lambda} \end{bmatrix}, \mathbf{n}_{\partial D} \right\rangle d\partial D$$

where is  $\mathbf{n}_{\partial D}$  the normal to the surface  $\partial D$

## Conservation of volumes in the $x, \lambda$ space

Consider a compact domain  $D(t)$  in the  $x(t), \lambda(t)$  space, with boundary  $\partial D(t)$ . The volume of  $D(t)$  say  $\rho(D(t)) \in \mathbb{R}$  is given by:

$$\rho(D(t)) = \int_{D(t)} dD$$

and evolves with time as the integration of  $x(t), \lambda(t)$  proceeds. Its time-derivative reads as:

$$\frac{d}{dt} \rho(D(t)) = \int_{\partial D(t)} \left\langle \begin{bmatrix} \dot{x} \\ \dot{\lambda} \end{bmatrix}, \mathbf{n}_{\partial D} \right\rangle d\partial D \stackrel{\text{Green's th.}}{\overbrace{=}} \int_{D(t)} \operatorname{div} \left( \begin{bmatrix} \dot{x} \\ \dot{\lambda} \end{bmatrix} \right) dD$$

where  $\mathbf{n}_{\partial D}$  the normal to the surface  $\partial D$ , and  $\operatorname{div}$  the divergence operator.

## Conservation of volumes in the $\mathbf{x}, \boldsymbol{\lambda}$ space

Consider a compact domain  $D(t)$  in the  $\mathbf{x}(t), \boldsymbol{\lambda}(t)$  space, with boundary  $\partial D(t)$ . The volume of  $D(t)$  say  $\rho(D(t)) \in \mathbb{R}$  is given by:

$$\rho(D(t)) = \int_{D(t)} dD$$

and evolves with time as the integration of  $\mathbf{x}(t), \boldsymbol{\lambda}(t)$  proceeds. Its time-derivative reads as:

$$\frac{d}{dt} \rho(D(t)) = \int_{\partial D(t)} \left\langle \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix}, \mathbf{n}_{\partial D} \right\rangle d\partial D \stackrel{\text{Green's th.}}{\cong} \int_{D(t)} \operatorname{div} \left( \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix} \right) dD$$

where is  $\mathbf{n}_{\partial D}$  the normal to the surface  $\partial D$ , and  $\operatorname{div}$  the divergence operator. Here it yields:

$$\operatorname{div} \left( \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix} \right) = \operatorname{Tr} \left( \frac{\partial \dot{\mathbf{x}}}{\partial \mathbf{x}} + \frac{\partial \dot{\boldsymbol{\lambda}}}{\partial \boldsymbol{\lambda}} \right) = \operatorname{Tr} \left( \frac{\partial \mathbf{F}}{\partial \mathbf{x}} - \frac{\partial}{\partial \boldsymbol{\lambda}} \frac{\partial H}{\partial \mathbf{x}} \right) = 0$$

## Conservation of volumes in the $\mathbf{x}, \boldsymbol{\lambda}$ space

Consider a compact domain  $D(t)$  in the  $\mathbf{x}(t), \boldsymbol{\lambda}(t)$  space, with boundary  $\partial D(t)$ . The volume of  $D(t)$  say  $\rho(D(t)) \in \mathbb{R}$  is given by:

$$\rho(D(t)) = \int_{D(t)} dD$$

and evolves with time as the integration of  $\mathbf{x}(t), \boldsymbol{\lambda}(t)$  proceeds. Its time-derivative reads as:

$$\frac{d}{dt} \rho(D(t)) = \int_{\partial D(t)} \left\langle \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix}, \mathbf{n}_{\partial D} \right\rangle d\partial D \stackrel{\text{Green's th.}}{\overbrace{=}} \int_{D(t)} \operatorname{div} \left( \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix} \right) dD$$

where  $\mathbf{n}_{\partial D}$  the normal to the surface  $\partial D$ , and  $\operatorname{div}$  the divergence operator. Here it yields:

$$\operatorname{div} \left( \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix} \right) = \operatorname{Tr} \left( \frac{\partial \dot{\mathbf{x}}}{\partial \mathbf{x}} + \frac{\partial \dot{\boldsymbol{\lambda}}}{\partial \boldsymbol{\lambda}} \right) = \operatorname{Tr} \left( \frac{\partial \mathbf{F}}{\partial \mathbf{x}} - \frac{\partial}{\partial \boldsymbol{\lambda}} \frac{\partial H}{\partial \mathbf{x}} \right) = 0$$

because

$$\frac{\partial}{\partial \boldsymbol{\lambda}} \frac{\partial H}{\partial \mathbf{x}} = \frac{\partial}{\partial \boldsymbol{\lambda}} \left( \frac{\partial L(\mathbf{x}, \mathbf{u})}{\partial \mathbf{x}} + \frac{\partial \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})}{\partial \mathbf{x}} \right) = \frac{\partial \mathbf{F}}{\partial \mathbf{x}}$$

## Conservation of volumes in the $\mathbf{x}, \boldsymbol{\lambda}$ space

Consider a compact domain  $D(t)$  in the  $\mathbf{x}(t), \boldsymbol{\lambda}(t)$  space, with boundary  $\partial D(t)$ . The volume of  $D(t)$  say  $\rho(D(t)) \in \mathbb{R}$  is given by:

$$\rho(D(t)) = \int_{D(t)} dD$$

and evolves with time as the integration of  $\mathbf{x}(t), \boldsymbol{\lambda}(t)$  proceeds. Its time-derivative reads as:

$$\frac{d}{dt} \rho(D(t)) = \int_{\partial D(t)} \left\langle \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix}, \mathbf{n}_{\partial D} \right\rangle d\partial D \stackrel{\text{Green's th.}}{\cong} \int_{D(t)} \operatorname{div} \left( \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix} \right) dD$$

where  $\mathbf{n}_{\partial D}$  the normal to the surface  $\partial D$ , and  $\operatorname{div}$  the divergence operator. Here it yields:

$$\operatorname{div} \left( \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix} \right) = \operatorname{Tr} \left( \frac{\partial \dot{\mathbf{x}}}{\partial \mathbf{x}} + \frac{\partial \dot{\boldsymbol{\lambda}}}{\partial \boldsymbol{\lambda}} \right) = \operatorname{Tr} \left( \frac{\partial \mathbf{F}}{\partial \mathbf{x}} - \frac{\partial}{\partial \boldsymbol{\lambda}} \frac{\partial H}{\partial \mathbf{x}} \right) = 0$$

because

$$\frac{\partial}{\partial \boldsymbol{\lambda}} \frac{\partial H}{\partial \mathbf{x}} = \frac{\partial}{\partial \boldsymbol{\lambda}} \left( \frac{\partial L(\mathbf{x}, \mathbf{u})}{\partial \mathbf{x}} + \frac{\partial \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})}{\partial \mathbf{x}} \right) = \frac{\partial \mathbf{F}}{\partial \mathbf{x}}$$

The volume of  $D(t)$  is constant\* throughout the integration of  $\mathbf{x}, \boldsymbol{\lambda}$  !!

## Conservation of volumes in the $\mathbf{x}, \boldsymbol{\lambda}$ space

Consider a compact domain  $D(t)$  in the  $\mathbf{x}(t), \boldsymbol{\lambda}(t)$  space, with boundary  $\partial D(t)$ . The volume of  $D(t)$  say  $\rho(D(t)) \in \mathbb{R}$  is given by:

$$\rho(D(t)) = \int_{D(t)} dD$$

and evolves with time as the integration of  $\mathbf{x}(t), \boldsymbol{\lambda}(t)$  proceeds. Its time-derivative reads as:

$$\frac{d}{dt} \rho(D(t)) = \int_{\partial D(t)} \left\langle \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix}, \mathbf{n}_{\partial D} \right\rangle d\partial D \stackrel{\text{Green's th.}}{\cong} \int_{D(t)} \operatorname{div} \left( \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix} \right) dD$$

where is  $\mathbf{n}_{\partial D}$  the normal to the surface  $\partial D$ , and  $\operatorname{div}$  the divergence operator. Here it yields:

$$\operatorname{div} \left( \begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\boldsymbol{\lambda}} \end{bmatrix} \right) = \operatorname{Tr} \left( \frac{\partial \dot{\mathbf{x}}}{\partial \mathbf{x}} + \frac{\partial \dot{\boldsymbol{\lambda}}}{\partial \boldsymbol{\lambda}} \right) = \operatorname{Tr} \left( \frac{\partial \mathbf{F}}{\partial \mathbf{x}} - \frac{\partial}{\partial \boldsymbol{\lambda}} \frac{\partial H}{\partial \mathbf{x}} \right) = 0$$

because

$$\frac{\partial}{\partial \boldsymbol{\lambda}} \frac{\partial H}{\partial \mathbf{x}} = \frac{\partial}{\partial \boldsymbol{\lambda}} \left( \frac{\partial L(\mathbf{x}, \mathbf{u})}{\partial \mathbf{x}} + \frac{\partial \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})}{\partial \mathbf{x}} \right) = \frac{\partial \mathbf{F}}{\partial \mathbf{x}}$$

The volume of  $D(t)$  is constant\* throughout the integration of  $\mathbf{x}, \boldsymbol{\lambda}$  !!

\* See Liouville's Theorem on phase-space distribution functions in Hamiltonian mechanics

## Conservation of volumes in the $x, \lambda$ space (cont')

Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

Hamiltonian is minimised by  $u = -\lambda$ .

The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda_f = \lambda(t_f) = 0$

## Conservation of volumes in the $x, \lambda$ space (cont')


Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

Hamiltonian is minimised by  $u = -\lambda$ .



The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$

## Conservation of volumes in the $x, \lambda$ space (cont')


Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

Hamiltonian is minimised by  $u = -\lambda$ .



The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$

## Conservation of volumes in the $x, \lambda$ space (cont')


Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

Hamiltonian is minimised by  $u = -\lambda$ .



The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$

## Conservation of volumes in the $x, \lambda$ space (cont')


Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

Hamiltonian is minimised by  $u = -\lambda$ .



The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$

## Conservation of volumes in the $x, \lambda$ space (cont')


Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

Hamiltonian is minimised by  $u = -\lambda$ .



The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$

## Conservation of volumes in the $x, \lambda$ space (cont')


Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

Hamiltonian is minimised by  $u = -\lambda$ .



The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$

## Conservation of volumes in the $x, \lambda$ space (cont')


Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

Hamiltonian is minimised by  $u = -\lambda$ .



The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$

## Conservation of volumes in the $x, \lambda$ space (cont')


Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

Hamiltonian is minimised by  $u = -\lambda$ .



The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$

## Conservation of volumes in the $x, \lambda$ space (cont')


Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$

Hamiltonian is minimised by  $u = -\lambda$ .



The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$

## Conservation of volumes in the $x, \lambda$ space (cont')

Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$


Hamiltonian is minimised by  $u = -\lambda$ .

The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$



**The punchline:**

## Conservation of volumes in the $x, \lambda$ space (cont')

Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$


Hamiltonian is minimised by  $u = -\lambda$ .

The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$



**The punchline:**

the states  $x$  and costates  $\lambda$  cannot be *both* stable, as it would imply  $\rho(D(t)) \rightarrow 0$  !!

## Conservation of volumes in the $x, \lambda$ space (cont')

Problem:

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

Yields

$$H(x, u, \lambda) = \frac{1}{2} (x^2 + u^2) + \lambda (u - \sin(x))$$


Hamiltonian is minimised by  $u = -\lambda$ .

The dynamics become

$$\dot{x} = -\lambda - \sin(x)$$

$$\dot{\lambda} = \lambda \cos(x) - x$$

Find  $\lambda_0$  to enforce  $\lambda(t_f) = 0$



### The punchline:

the states  $x$  and costates  $\lambda$  cannot be *both* stable, as it would imply  $\rho(D(t)) \rightarrow 0$  !!

Important consequence: the TPBVP is hard to solve numerically !!

## How to actually use PMP then ?

**PMP equations:**  $\mathbf{u}^* = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$  with:

$$\text{States : } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(\mathbf{t}_0) = \bar{\mathbf{x}}_0$$

$$\text{Costates : } \dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}), \quad \boldsymbol{\lambda}(\mathbf{t}_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$$

**Key idea:** apply the multiple-shooting principle to the PMP equations

## How to actually use PMP then ?

**PMP equations:**  $\mathbf{u}^* = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$  with:

$$\text{States : } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(\mathbf{t}_0) = \bar{\mathbf{x}}_0$$

$$\text{Costates : } \dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}), \quad \boldsymbol{\lambda}(\mathbf{t}_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$$

**Key idea:** apply the multiple-shooting principle to the PMP equations

State-costate

$$\mathbf{s} = \begin{bmatrix} \mathbf{x} \\ \boldsymbol{\lambda} \end{bmatrix}$$

Optimal control

$$\mathbf{u}(\mathbf{s}) = \arg \min_{\mathbf{u}} H(\mathbf{u}, \mathbf{s})$$

## How to actually use PMP then ?

**PMP equations:**  $\mathbf{u}^* = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$  with:

$$\text{States : } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(\mathbf{t}_0) = \bar{\mathbf{x}}_0$$

$$\text{Costates : } \dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}), \quad \boldsymbol{\lambda}(\mathbf{t}_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$$

**Key idea:** apply the multiple-shooting principle to the PMP equations

State-costate

$$\mathbf{s} = \begin{bmatrix} \mathbf{x} \\ \boldsymbol{\lambda} \end{bmatrix}$$

Optimal control

$$\mathbf{u}(\mathbf{s}) = \arg \min_{\mathbf{u}} H(\mathbf{u}, \mathbf{s})$$

Integrator  $\xi(s_k)$  over the time intervals  $t \in [t_k, t_{k+1}]$  with:

$$\dot{\mathbf{s}} = \begin{bmatrix} \mathbf{F}(\mathbf{x}, \mathbf{u}) \\ -\nabla_{\mathbf{x}} H(\mathbf{u}, \mathbf{s}) \end{bmatrix}, \quad \text{with } \mathbf{s}(t_k) = \mathbf{s}_k$$

## How to actually use PMP then ?

**PMP equations:**  $\mathbf{u}^* = \arg \min_{\mathbf{u}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$  with:

$$\text{States : } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \bar{\mathbf{x}}_0$$

$$\text{Costates : } \dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}), \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$$

**Key idea:** apply the multiple-shooting principle to the PMP equations

State-costate

$$\mathbf{s} = \begin{bmatrix} \mathbf{x} \\ \boldsymbol{\lambda} \end{bmatrix}$$

Optimal control

$$\mathbf{u}(\mathbf{s}) = \arg \min_{\mathbf{u}} H(\mathbf{u}, \mathbf{s})$$

Integrator  $\xi(s_k)$  over the time intervals  $t \in [t_k, t_{k+1}]$  with:

$$\dot{\mathbf{s}} = \begin{bmatrix} \mathbf{F}(\mathbf{x}, \mathbf{u}) \\ -\nabla_{\mathbf{x}} H(\mathbf{u}, \mathbf{s}) \end{bmatrix}, \quad \text{with} \quad \mathbf{s}(t_k) = \mathbf{s}_k$$

**Root-finding** problem over the variables  $s_0, \dots, s_N$ :

$$\mathbf{r}(\mathbf{s}) = \begin{bmatrix} \mathbf{x}_0 - \bar{\mathbf{x}}_0 \\ \xi(s_0) - s_1 \\ \vdots \\ \xi(s_{N-1}) - s_N \\ \boldsymbol{\lambda}_N - \nabla_{\mathbf{x}} \phi(\mathbf{x}_N) \end{bmatrix} = 0$$

## Indirect multiple-shooting

**Problem:**



$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$
$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 1 (single shooting)



## Indirect multiple-shooting

**Problem:**



$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$
$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 1 (single shooting)



## Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$



PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 1 (single shooting)



## Indirect multiple-shooting

**Problem:**



$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$
$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 1 (single shooting)



## Indirect multiple-shooting

**Problem:**



$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$
$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 1 (single shooting)



## Indirect multiple-shooting

**Problem:**



$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$
$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 1 (single shooting)



## Indirect multiple-shooting

**Problem:**



$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$
$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 3



## Indirect multiple-shooting

**Problem:**



$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$
$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 3



## Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$



PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 3



## Indirect multiple-shooting

**Problem:**



$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$
$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 3



# Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$



PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 3



## Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :



$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,

**Final time:**  $t_f = 5$ , **full Newton steps**

# of shooting interval = 3



## Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :



$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1, \lambda = 0$ ,

**Final time:**  $t_f = 5$ , **full Newton steps**

**# of shooting interval = 5**



## Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :



$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1, \lambda = 0$ ,

**Final time:**  $t_f = 5$ , **full Newton steps**

**# of shooting interval = 5**



## Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$



PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 5



# Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$



PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 5



## Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$



PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 5



## Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :



$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1, \lambda = 0$ ,

**Final time:**  $t_f = 5$ , **full Newton steps**

**# of shooting interval = 10**



## Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :



$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,

**Final time:**  $t_f = 5$ , **full Newton steps**

**# of shooting interval** = 10



# Indirect multiple-shooting

**Problem:**



$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :

$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$
$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,    **Final time:**  $t_f = 5$ ,    **full Newton steps**

# of shooting interval = 10



## Indirect multiple-shooting

**Problem:**

$$\min_{x,u} \frac{1}{2} \int_0^{t_f} (x^2 + u^2) dt$$
$$\dot{x} = u - \sin(x), \quad x(0) = 1$$

PMP equations with  $u = -\lambda$ :



$$\dot{x} = -\lambda - \sin(x), \quad x(0) = 1$$

$$\dot{\lambda} = \lambda \cos(x) - x, \quad \lambda(5) = 0$$

**Initial guess:**  $x = 1$ ,  $\lambda = 0$ ,

**Final time:**  $t_f = 5$ , **full Newton steps**

**# of shooting interval = 10**



# Outline

- 1 Introduction to the Pontryagin Maximum Principle (PMP)
- 2 Interpretation of  $H_u$
- 3 Input bounds in Indirect Optimal Control
- 4 Singular Optimal Control problems
- 5 General constraints in Indirect Optimal Control

## Interpretation of $H_u$

Consider the **functional**:

$$J[\mathbf{u}(\cdot)] = \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}, \mathbf{u}) dt$$

s.t.  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

that maps an input profile  $\mathbf{u}(\cdot)$  into the corresponding cost.

## Interpretation of $H_u$

Consider the **functional**:

$$J[\mathbf{u}(\cdot)] = \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}, \mathbf{u}) dt$$

s.t.  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

that maps an input profile  $\mathbf{u}(\cdot)$  into the corresponding cost.

**Gateaux derivative** (see Calculus of Variations)

$$dJ[\mathbf{u}(\cdot), \xi(\cdot)] = \lim_{\tau \rightarrow 0} \frac{J[\mathbf{u}(\cdot) + \tau \xi(\cdot)] - J[\mathbf{u}(\cdot)]}{\tau}$$

## Interpretation of $H_u$

Consider the **functional**:

$$J[\mathbf{u}(\cdot)] = \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}, \mathbf{u}) dt$$

s.t.  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

that maps an input profile  $\mathbf{u}(\cdot)$  into the corresponding cost.

**Gateaux derivative** (see Calculus of Variations)

$$dJ[\mathbf{u}(\cdot), \xi(\cdot)] = \lim_{\tau \rightarrow 0} \frac{J[\mathbf{u}(\cdot) + \tau \xi(\cdot)] - J[\mathbf{u}(\cdot)]}{\tau}$$

Note: Gateaux derivatives are directional derivatives for functionals ("direction"  $\xi$ )

## Interpretation of $H_u$

Consider the **functional**:

$$J[u(\cdot)] = \phi(x(t_f)) + \int_{t_0}^{t_f} L(x, u) dt$$




s.t.  $\dot{x} = F(x, u)$ ,  $x(t_0) = x_0$

that maps an input profile  $u(\cdot)$  into the corresponding cost.

**Gateaux derivative** (see Calculus of Variations)

$$dJ[u(\cdot), \xi(\cdot)] = \lim_{\tau \rightarrow 0} \frac{J[u(\cdot) + \tau \xi(\cdot)] - J[u(\cdot)]}{\tau}$$

Note: Gateaux derivatives are directional derivatives for functionals ("direction"  $\xi$ )



# Interpretation of $H_u$

Consider the **functional**:

$$J[\mathbf{u}(\cdot)] = \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}, \mathbf{u}) dt$$

s.t.  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

that maps an input profile  $\mathbf{u}(\cdot)$  into the corresponding cost.




**Gateaux derivative** (see Calculus of Variations)

$$dJ[\mathbf{u}(\cdot), \xi(\cdot)] = \lim_{\tau \rightarrow 0} \frac{J[\mathbf{u}(\cdot) + \tau \xi(\cdot)] - J[\mathbf{u}(\cdot)]}{\tau}$$

**Optimality:**

$$dJ[\mathbf{u}^*(\cdot), \xi(\cdot)] = 0, \quad \forall \xi(\cdot)$$

Note: Gateaux derivatives are directional derivatives for functionals ("direction"  $\xi$ )



## Interpretation of $H_u$

Consider the **functional**:

$$J[\mathbf{u}(\cdot)] = \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}, \mathbf{u}) dt$$

s.t.  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

that maps an input profile  $\mathbf{u}(\cdot)$  into the corresponding cost.

**Optimality:**

$$dJ[\mathbf{u}^*(\cdot), \xi(\cdot)] = 0, \quad \forall \xi(\cdot)$$

**Gateaux derivative** (see Calculus of Variations)

$$dJ[\mathbf{u}(\cdot), \xi(\cdot)] = \lim_{\tau \rightarrow 0} \frac{J[\mathbf{u}(\cdot) + \tau \xi(\cdot)] - J[\mathbf{u}(\cdot)]}{\tau}$$

Note: Gateaux derivatives are directional derivatives for functionals ("direction"  $\xi$ )

**Interpretation of  $H_u$**  (fundamental Lemma of Calculus of Variations):

$$dJ[\mathbf{u}(\cdot), \xi(\cdot)] = \int_0^{t_f} H_u(\mathbf{x}(t), \lambda(t), \mathbf{u}(t)) \cdot \xi(t) dt$$

## Interpretation of $H_u$

Consider the **functional**:

$$J[\mathbf{u}(\cdot)] = \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}, \mathbf{u}) dt$$

s.t.  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

that maps an input profile  $\mathbf{u}(\cdot)$  into the corresponding cost.

**Optimality:**

$$dJ[\mathbf{u}^*(\cdot), \xi(\cdot)] = 0, \quad \forall \xi(\cdot)$$

**Requires**

$$H_u = 0, \quad \forall t$$

at the solution.

**Gateaux derivative** (see Calculus of Variations)

$$dJ[\mathbf{u}(\cdot), \xi(\cdot)] = \lim_{\tau \rightarrow 0} \frac{J[\mathbf{u}(\cdot) + \tau \xi(\cdot)] - J[\mathbf{u}(\cdot)]}{\tau}$$

Note: Gateaux derivatives are directional derivatives for functionals ("direction"  $\xi$ )

**Interpretation of  $H_u$**  (fundamental Lemma of Calculus of Variations):

$$dJ[\mathbf{u}(\cdot), \xi(\cdot)] = \int_0^{t_f} H_u(\mathbf{x}(t), \lambda(t), \mathbf{u}(t)) \cdot \xi(t) dt$$

## Interpretation of $H_u$

Consider the **functional**:

$$J[\mathbf{u}(\cdot)] = \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}, \mathbf{u}) dt$$

s.t.  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

that maps an input profile  $\mathbf{u}(\cdot)$  into the corresponding cost.

**Gateaux derivative** (see Calculus of Variations)

$$dJ[\mathbf{u}(\cdot), \xi(\cdot)] = \lim_{\tau \rightarrow 0} \frac{J[\mathbf{u}(\cdot) + \tau \xi(\cdot)] - J[\mathbf{u}(\cdot)]}{\tau}$$

**Optimality:**

$$dJ[\mathbf{u}^*(\cdot), \xi(\cdot)] = 0, \quad \forall \xi(\cdot)$$

**Requires**

$$H_u = 0, \quad \forall t$$

at the solution.

Note: Gateaux derivatives are directional derivatives for functionals ("direction"  $\xi$ )

**Interpretation of  $H_u$  (fundamental Lemma of Calculus of Variations):**

$$dJ[\mathbf{u}(\cdot), \xi(\cdot)] = \int_0^{t_f} H_u(\mathbf{x}(t), \lambda(t), \mathbf{u}(t)) \cdot \xi(t) dt$$

**What if  $\mathbf{u}(\cdot)$  is restricted to some (Banach) space ?** E.g. piecewise-constant...  
... then  $\xi(\cdot)$  is restricted to the same space !

## Interpretation of $H_u$ (cont')

Consider a piecewise-constant parametrization of  $u(\cdot)$ ...

... akin to a restriction of  $u(\cdot)$  to that Banach space !

## Interpretation of $H_u$ (cont')

Consider a piecewise-constant parametrization of  $\mathbf{u}(.)$ ...

... akin to a restriction of  $\mathbf{u}(.)$  to that Banach space !

Piecewise-constant parametrization

$$\mathbf{u}(t) = \mathbf{u}_k \quad \forall t \in [t_k, t_{k+1}]$$

$$\xi(t) = \xi_k \quad \forall t \in [t_k, t_{k+1}]$$

## Interpretation of $H_u$ (cont')

Consider a piecewise-constant parametrization of  $\mathbf{u}(.)$ ...

... akin to a restriction of  $\mathbf{u}(.)$  to that Banach space !

Piecewise-constant parametrization

Functional  $\rightarrow$  function

$$\mathbf{u}(t) = \mathbf{u}_k \quad \forall t \in [t_k, t_{k+1}]$$

$$J[\mathbf{u}(.)] \equiv \bar{J}(\mathbf{u}_1, \dots, \mathbf{u}_{N-1})$$

$$\xi(t) = \xi_k \quad \forall t \in [t_k, t_{k+1}]$$

## Interpretation of $H_u$ (cont')

Consider a piecewise-constant parametrization of  $\mathbf{u}(\cdot)$ ...

... akin to a restriction of  $\mathbf{u}(\cdot)$  to that Banach space !

Piecewise-constant parametrization

Functional  $\rightarrow$  function

$$\mathbf{u}(t) = \mathbf{u}_k \quad \forall t \in [t_k, t_{k+1}]$$

$$J[\mathbf{u}(\cdot)] \equiv \bar{J}(\mathbf{u}_1, \dots, \mathbf{u}_{N-1})$$

$$\xi(t) = \xi_k \quad \forall t \in [t_k, t_{k+1}]$$

Then optimality requires:

$$\begin{aligned} dJ[\mathbf{u}^*(\cdot), \xi(\cdot)] &= \int_0^{t_f} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}^*(t)) \cdot \xi(t) dt = \\ &\sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k^*) \cdot \xi_k dt = 0, \quad \forall \xi_k \end{aligned}$$

## Interpretation of $H_u$ (cont')

Consider a piecewise-constant parametrization of  $\mathbf{u}(\cdot)$ ...

... akin to a restriction of  $\mathbf{u}(\cdot)$  to that Banach space !

Piecewise-constant parametrization

Functional  $\rightarrow$  function

$$\mathbf{u}(t) = \mathbf{u}_k \quad \forall t \in [t_k, t_{k+1}]$$

$$J[\mathbf{u}(\cdot)] \equiv \bar{J}(\mathbf{u}_1, \dots, \mathbf{u}_{N-1})$$

$$\xi(t) = \xi_k \quad \forall t \in [t_k, t_{k+1}]$$

Then optimality requires:

$$\begin{aligned} dJ[\mathbf{u}^*(\cdot), \xi(\cdot)] &= \int_0^{t_f} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}^*(t)) \cdot \xi(t) dt = \\ &\sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k^*) \cdot \xi_k dt = 0, \quad \forall \xi_k \end{aligned}$$

Hence **optimality condition** is  $\forall k$ :

$$\int_{t_k}^{t_{k+1}} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k^*) dt = 0$$

## Interpretation of $H_u$ (cont')

Consider a piecewise-constant parametrization of  $\mathbf{u}(\cdot)$ ...

... akin to a restriction of  $\mathbf{u}(\cdot)$  to that Banach space !

Piecewise-constant parametrization

Functional  $\rightarrow$  function

$$\mathbf{u}(t) = \mathbf{u}_k \quad \forall t \in [t_k, t_{k+1}]$$

$$J[\mathbf{u}(\cdot)] \equiv \bar{J}(\mathbf{u}_1, \dots, \mathbf{u}_{N-1})$$

$$\xi(t) = \xi_k \quad \forall t \in [t_k, t_{k+1}]$$

Then optimality requires:

$$\begin{aligned} dJ[\mathbf{u}^*(\cdot), \xi(\cdot)] &= \int_0^{t_f} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}^*(t)) \cdot \xi(t) dt = \\ &\sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k^*) \cdot \xi_k dt = 0, \quad \forall \xi_k \end{aligned}$$

Hence **optimality condition** is  $\forall k$ :

$$\int_{t_k}^{t_{k+1}} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k^*) dt = 0$$

Gateaux  $\rightarrow$  classic derivative

$$\frac{\partial \bar{J}}{\partial \mathbf{u}_k} = \int_{t_k}^{t_{k+1}} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k) dt$$

## Interpretation of $H_u$ (cont')

Consider a piecewise-constant parametrization of  $\mathbf{u}(\cdot)$ ...

... akin to a restriction of  $\mathbf{u}(\cdot)$  to that Banach space !

Piecewise-constant parametrization

Functional  $\rightarrow$  function

$$\mathbf{u}(t) = \mathbf{u}_k \quad \forall t \in [t_k, t_{k+1}]$$

$$J[\mathbf{u}(\cdot)] \equiv \bar{J}(\mathbf{u}_1, \dots, \mathbf{u}_{N-1})$$

$$\xi(t) = \xi_k \quad \forall t \in [t_k, t_{k+1}]$$

Then optimality requires:

$$\begin{aligned} dJ[\mathbf{u}^*(\cdot), \xi(\cdot)] &= \int_0^{t_f} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}^*(t)) \cdot \xi(t) dt = \\ &\sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k^*) \cdot \xi_k dt = 0, \quad \forall \xi_k \end{aligned}$$

Hence **optimality condition** is  $\forall k$ :

$$\int_{t_k}^{t_{k+1}} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k^*) dt = 0$$

Gateaux  $\rightarrow$  classic derivative

$$\frac{\partial \bar{J}}{\partial \mathbf{u}_k} = \int_{t_k}^{t_{k+1}} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k) dt$$

When solving an OCP using Direct Methods, one can see the NLP solver as trying to get  $\int_{t_k}^{t_{k+1}} H_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k) dt = 0$  !!

# Outline

- 1 Introduction to the Pontryagin Maximum Principle (PMP)
- 2 Interpretation of  $H_u$
- 3 Input bounds in Indirect Optimal Control
- 4 Singular Optimal Control problems
- 5 General constraints in Indirect Optimal Control

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \underbrace{\begin{bmatrix} \mathbf{u}_{\min} - \mathbf{u} \\ \mathbf{u} - \mathbf{u}_{\max} \end{bmatrix}}_{\leq 0}$$

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \underbrace{\begin{bmatrix} \mathbf{u}_{\min} - \mathbf{u} \\ \mathbf{u} - \mathbf{u}_{\max} \end{bmatrix}}_{\leq 0}$$

Get the input from:  $\mathbf{u} = \underset{\mathbf{u}}{\operatorname{argmin}} H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u})$ , with:

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \underbrace{\begin{bmatrix} \mathbf{u}_{\min} - \mathbf{u} \\ \mathbf{u} - \mathbf{u}_{\max} \end{bmatrix}}_{\leq 0}$$

Get the input from:  $\mathbf{u} = \underset{\mathbf{u}}{\operatorname{argmin}} H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u})$ , with:

States :  $\dot{\mathbf{x}} = \mathbf{F}, \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H, \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \underbrace{\begin{bmatrix} \mathbf{u}_{\min} - \mathbf{u} \\ \mathbf{u} - \mathbf{u}_{\max} \end{bmatrix}}_{\leq 0}$$

Get the input from:  $\mathbf{u} = \underset{\mathbf{u}}{\operatorname{argmin}} H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u})$ , with:

States :  $\dot{\mathbf{x}} = \mathbf{F}, \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H, \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

Feasibility :  $\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}, \quad \boldsymbol{\mu} \geq 0$

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \underbrace{\begin{bmatrix} \mathbf{u}_{\min} - \mathbf{u} \\ \mathbf{u} - \mathbf{u}_{\max} \end{bmatrix}}_{\leq 0}$$

Get the input from:  $\mathbf{u} = \underset{\mathbf{u}}{\operatorname{argmin}} H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u})$ , with:

States :  $\dot{\mathbf{x}} = \mathbf{F}, \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H, \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

Feasibility :  $\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}, \quad \boldsymbol{\mu} \geq 0$

Complementarity :  $\boldsymbol{\mu}^\top \begin{bmatrix} \mathbf{u}_{\min} - \mathbf{u} \\ \mathbf{u} - \mathbf{u}_{\max} \end{bmatrix} = 0$

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \underbrace{\begin{bmatrix} \mathbf{u}_{\min} - \mathbf{u} \\ \mathbf{u} - \mathbf{u}_{\max} \end{bmatrix}}_{\leq 0}$$

Get the input from:  $\mathbf{u} = \underset{\mathbf{u}}{\operatorname{argmin}} H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u})$ , with:

States :  $\dot{\mathbf{x}} = \mathbf{F}, \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H, \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

Feasibility :  $\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}, \quad \boldsymbol{\mu} \geq 0$

Complementarity :  $\boldsymbol{\mu}^\top \begin{bmatrix} \mathbf{u}_{\min} - \mathbf{u} \\ \mathbf{u} - \mathbf{u}_{\max} \end{bmatrix} = 0$

When  $\mathbf{u}$  hits the bounds,  
 $\boldsymbol{\mu}$  "creates a gradient" in  
 $H$  to enforce feasibility

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

An equivalent but simpler approach: define the **Lagrange** function

$$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

An equivalent but simpler approach: define the **Lagrange** function

$$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$

Get the input from:  $\mathbf{u} = \underset{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}}{\operatorname{argmin}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$ , with:

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

An equivalent but simpler approach: define the **Lagrange** function

$$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$

Get the input from:  $\mathbf{u} = \underset{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}}{\operatorname{argmin}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$ , with:

States :  $\dot{\mathbf{x}} = \mathbf{F}$ ,  $\mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} \mathcal{L}$ ,  $\boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

An equivalent but simpler approach: define the **Lagrange** function

$$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$

Get the input from:  $\mathbf{u} = \underset{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}}{\operatorname{argmin}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{u})$ , with:

States :  $\dot{\mathbf{x}} = \mathbf{F}$ ,  $\mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} \mathcal{L}$ ,  $\boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

Note: optimality now reads as...

$$\int_0^{t_f} \mathcal{L}_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}(t)) \cdot \boldsymbol{\xi}(t) dt \geq 0 \quad \text{for any feasible direction } \boldsymbol{\xi}(\cdot)$$

## Input bounds in indirect optimal control

OCP with input bounds:

$$\min_{\mathbf{x}, \mathbf{u}} \quad \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

Note:  $\mathbf{u}^*$  is now a **non-smooth** function of  $\mathbf{x}, \lambda$ . Must be handled carefully when solving TPBVP via Newton !!

An equivalent but simpler approach: define the **Lagrange** function

$$\mathcal{L}(\mathbf{x}, \lambda, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \lambda^\top \mathbf{F}(\mathbf{x}, \mathbf{u})$$

Get the input from:  $\mathbf{u} = \underset{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}}{\operatorname{argmin}} \mathcal{L}(\mathbf{x}, \lambda, \mathbf{u})$ , with:

States :  $\dot{\mathbf{x}} = \mathbf{F}$ ,  $\mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\lambda} = -\nabla_{\mathbf{x}} \mathcal{L}$ ,  $\lambda(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

Note: optimality now reads as...

$$\int_0^{t_f} \mathcal{L}_u(\mathbf{x}(t), \lambda(t), \mathbf{u}(t)) \cdot \xi(t) dt \geq 0 \quad \text{for any feasible direction } \xi(\cdot)$$

# Outline

- 1 Introduction to the Pontryagin Maximum Principle (PMP)
- 2 Interpretation of  $H_u$
- 3 Input bounds in Indirect Optimal Control
- 4 Singular Optimal Control problems
- 5 General constraints in Indirect Optimal Control

# Singular Optimal Control problems

Consider

$$\min_{\mathbf{u}, \mathbf{x}} \phi(\mathbf{x}(t_f))$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(0) = \bar{\mathbf{x}}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

# Singular Optimal Control problems

Consider

$$\min_{\mathbf{u}, \mathbf{x}} \phi(\mathbf{x}(t_f))$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(0) = \bar{\mathbf{x}}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

PMP equations with  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{F}$

$$\mathbf{u} = \operatorname{argmin}_{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}} \mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$$

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}^*)$$

$$\dot{\boldsymbol{\lambda}} = -\mathcal{L}_x(\mathbf{x}, \mathbf{u}^*, \boldsymbol{\lambda})$$

# Singular Optimal Control problems

Consider

$$\min_{\mathbf{u}, \mathbf{x}} \phi(\mathbf{x}(t_f))$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(0) = \bar{\mathbf{x}}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

PMP equations with  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{F}$

$$\mathbf{u} = \operatorname{argmin}_{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}} \mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$$

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}^*)$$

$$\dot{\boldsymbol{\lambda}} = -\mathcal{L}_x(\mathbf{x}, \mathbf{u}^*, \boldsymbol{\lambda})$$

What if  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$  is affine in  $\mathbf{u}$  ?

# Singular Optimal Control problems

Consider

$$\min_{\mathbf{u}, \mathbf{x}} \phi(\mathbf{x}(t_f))$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(0) = \bar{\mathbf{x}}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

PMP equations with  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{F}$

$$\mathbf{u} = \operatorname{argmin}_{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}} \mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$$

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}^*)$$

$$\dot{\boldsymbol{\lambda}} = -\mathcal{L}_x(\mathbf{x}, \mathbf{u}^*, \boldsymbol{\lambda})$$

What if  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$  is affine in  $\mathbf{u}$ ? E.g.

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}) + \mathbf{g}(\mathbf{x}) \mathbf{u}$$

# Singular Optimal Control problems

Consider

$$\min_{\mathbf{u}, \mathbf{x}} \phi(\mathbf{x}(t_f))$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(0) = \bar{\mathbf{x}}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

PMP equations with  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{F}$

$$\mathbf{u} = \underset{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}}{\operatorname{argmin}} \mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$$

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}^*)$$

$$\dot{\boldsymbol{\lambda}} = -\mathcal{L}_x(\mathbf{x}, \mathbf{u}^*, \boldsymbol{\lambda})$$

What if  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$  is affine in  $\mathbf{u}$ ? E.g.

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}) + \mathbf{g}(\mathbf{x}) \mathbf{u}$$

yields  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{f}(\mathbf{x}) + \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) \mathbf{u}$

# Singular Optimal Control problems

Consider

$$\min_{\mathbf{u}, \mathbf{x}} \phi(\mathbf{x}(t_f))$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(0) = \bar{\mathbf{x}}_0$$


$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

PMP equations with  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{F}$

$$\mathbf{u} = \underset{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}}{\operatorname{argmin}} \mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$$

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}^*)$$

$$\dot{\boldsymbol{\lambda}} = -\mathcal{L}_x(\mathbf{x}, \mathbf{u}^*, \boldsymbol{\lambda})$$



What if  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$  is affine in  $\mathbf{u}$ ? E.g.

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}) + \mathbf{g}(\mathbf{x}) \mathbf{u}$$

yields  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{f}(\mathbf{x}) + \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) \mathbf{u}$

$$\text{then } \mathbf{u} = \begin{cases} \mathbf{u}_{\max} & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) < 0 \end{cases}$$

# Singular Optimal Control problems

Consider

$$\min_{\mathbf{u}, \mathbf{x}} \phi(\mathbf{x}(t_f))$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(0) = \bar{\mathbf{x}}_0$$


$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

PMP equations with  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{F}$

$$\mathbf{u} = \underset{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}}{\operatorname{argmin}} \mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$$

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}^*)$$

$$\dot{\boldsymbol{\lambda}} = -\mathcal{L}_x(\mathbf{x}, \mathbf{u}^*, \boldsymbol{\lambda})$$



What if  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$  is affine in  $\mathbf{u}$ ? E.g.

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}) + \mathbf{g}(\mathbf{x}) \mathbf{u}$$

yields  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{f}(\mathbf{x}) + \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) \mathbf{u}$

$$\text{then } \mathbf{u} = \begin{cases} \mathbf{u}_{\max} & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) < 0 \\ \mathbf{u}_{\min} & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) > 0 \end{cases}$$

# Singular Optimal Control problems

Consider

$$\min_{\mathbf{u}, \mathbf{x}} \phi(\mathbf{x}(t_f))$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(0) = \bar{\mathbf{x}}_0$$


$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

PMP equations with  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{F}$

$$\mathbf{u} = \underset{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}}{\operatorname{argmin}} \mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$$

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}^*)$$

$$\dot{\boldsymbol{\lambda}} = -\mathcal{L}_x(\mathbf{x}, \mathbf{u}^*, \boldsymbol{\lambda})$$



What if  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$  is affine in  $\mathbf{u}$ ? E.g.

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}) + \mathbf{g}(\mathbf{x}) \mathbf{u}$$

yields  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{f}(\mathbf{x}) + \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) \mathbf{u}$

$$\text{then } \mathbf{u} = \begin{cases} \mathbf{u}_{\max} & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) < 0 \\ \mathbf{u}_{\min} & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) > 0 \\ ? & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) = 0 \end{cases}$$

# Singular Optimal Control problems

Consider

$$\begin{aligned} \min_{\mathbf{u}, \mathbf{x}} \quad & \phi(\mathbf{x}(t_f)) \\ \text{s.t.} \quad & \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(0) = \bar{\mathbf{x}}_0 \\ & \mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max} \end{aligned}$$

PMP equations with  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{F}$

$$\mathbf{u} = \underset{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}}{\operatorname{argmin}} \mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$$

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}^*)$$


$$\dot{\boldsymbol{\lambda}} = -\mathcal{L}_x(\mathbf{x}, \mathbf{u}^*, \boldsymbol{\lambda})$$

What if  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$  is affine in  $\mathbf{u}$ ? E.g.

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}) + \mathbf{g}(\mathbf{x}) \mathbf{u}$$

yields  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{f}(\mathbf{x}) + \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) \mathbf{u}$

$$\text{then } \mathbf{u} = \begin{cases} \mathbf{u}_{\max} & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) < 0 \\ \mathbf{u}_{\min} & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) > 0 \\ ? & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) = 0 \end{cases}$$



Find  $\mathbf{u}$  when  $\boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) = 0$  ?

# Singular Optimal Control problems

Consider

$$\min_{\mathbf{u}, \mathbf{x}} \phi(\mathbf{x}(t_f))$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(0) = \bar{\mathbf{x}}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

PMP equations with  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{F}$

$$\mathbf{u} = \underset{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}}{\operatorname{argmin}} \mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$$

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}^*)$$


$$\dot{\boldsymbol{\lambda}} = -\mathcal{L}_x(\mathbf{x}, \mathbf{u}^*, \boldsymbol{\lambda})$$

What if  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$  is affine in  $\mathbf{u}$ ? E.g.

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}) + \mathbf{g}(\mathbf{x}) \mathbf{u}$$

yields  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{f}(\mathbf{x}) + \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) \mathbf{u}$

then  $\mathbf{u} = \begin{cases} \mathbf{u}_{\max} & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) < 0 \\ \mathbf{u}_{\min} & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) > 0 \\ ? & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) = 0 \end{cases}$



Find  $\mathbf{u}$  when  $\boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) = 0$ ?

When  $\mathbf{u}_{\min} < \mathbf{u} < \mathbf{u}_{\max}$  we want

$$\mathcal{L}_u = \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) = 0, \quad \forall t$$

Hence we want  $\frac{d}{dt} \mathcal{L}_u = 0, \quad \forall t$

# Singular Optimal Control problems

Consider

$$\min_{\mathbf{u}, \mathbf{x}} \phi(\mathbf{x}(t_f))$$

$$\text{s.t. } \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(0) = \bar{\mathbf{x}}_0$$

$$\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}$$

PMP equations with  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{F}$

$$\mathbf{u} = \underset{\mathbf{u}_{\min} \leq \mathbf{u} \leq \mathbf{u}_{\max}}{\operatorname{argmin}} \mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$$

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}^*)$$


$$\dot{\boldsymbol{\lambda}} = -\mathcal{L}_x(\mathbf{x}, \mathbf{u}^*, \boldsymbol{\lambda})$$

What if  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda})$  is affine in  $\mathbf{u}$ ? E.g.

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}) + \mathbf{g}(\mathbf{x}) \mathbf{u}$$

yields  $\mathcal{L}(\mathbf{x}, \mathbf{u}, \boldsymbol{\lambda}) = \boldsymbol{\lambda}^\top \mathbf{f}(\mathbf{x}) + \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) \mathbf{u}$

then  $\mathbf{u} = \begin{cases} \mathbf{u}_{\max} & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) < 0 \\ \mathbf{u}_{\min} & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) > 0 \\ ? & \text{if } \boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) = 0 \end{cases}$



Find  $\mathbf{u}$  when  $\boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x}) = 0$ ?

For some  $i$ ,  $\mathbf{u}$  appears in  $\frac{d^i}{dt^i} \mathcal{L}_u$ ,  
then solve

$$\frac{d^i}{dt^i} \mathcal{L}_u = 0, \quad i > 0$$

for  $\mathbf{u}$  !!

## Singular Optimal Control - Example

$$\begin{aligned} & \min_{x(\cdot), u(\cdot)} \quad \frac{1}{2} \int_0^1 x_1^2 dt \\ \text{s.t.} \quad & \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \\ & -5 \leq u \leq 5 \end{aligned}$$

## Singular Optimal Control - Example

$$\min_{x(\cdot), u(\cdot)} \frac{1}{2} \int_0^1 x_1^2 dt$$

$$\text{s.t. } \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$
$$-5 \leq u \leq 5$$

$$\mathcal{L}_u = \lambda_2 \quad \frac{d}{dt} \mathcal{L}_u = -\lambda_1 \quad \frac{d^2}{dt^2} \mathcal{L}_u = x_1 \quad \frac{d^3}{dt^3} \mathcal{L}_u = x_2 \quad \frac{d^4}{dt^4} \mathcal{L}_u = u$$

## Singular Optimal Control - Example

$$\min_{x(\cdot), u(\cdot)} \frac{1}{2} \int_0^1 x_1^2 dt$$

$$\text{s.t. } \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$
$$-5 \leq u \leq 5$$

$$\mathcal{L}_u = \lambda_2 \quad \frac{d}{dt} \mathcal{L}_u = -\lambda_1 \quad \frac{d^2}{dt^2} \mathcal{L}_u = x_1 \quad \frac{d^3}{dt^3} \mathcal{L}_u = x_2 \quad \frac{d^4}{dt^4} \mathcal{L}_u = u$$

**Optimal input:**

$$u^* = \begin{cases} u_{\min} & \text{if } \lambda_2 > 0 \\ u_{\max} & \text{if } \lambda_2 < 0 \\ 0 & \text{if } \frac{d^i}{dt^i} \mathcal{L}_u = 0 \end{cases}$$

## Singular Optimal Control - Example

$$\min_{x(\cdot), u(\cdot)} \frac{1}{2} \int_0^1 x_1^2 dt$$

$$\text{s.t. } \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$
$$-5 \leq u \leq 5$$

$$\mathcal{L}_u = \lambda_2 \quad \frac{d}{dt} \mathcal{L}_u = -\lambda_1 \quad \frac{d^2}{dt^2} \mathcal{L}_u = x_1 \quad \frac{d^3}{dt^3} \mathcal{L}_u = x_2 \quad \frac{d^4}{dt^4} \mathcal{L}_u = u$$

**Optimal input:**

• Input is either in the bounds or zero !!

$$u^* = \begin{cases} u_{\min} & \text{if } \lambda_2 > 0 \\ u_{\max} & \text{if } \lambda_2 < 0 \\ 0 & \text{if } \frac{d^i}{dt^i} \mathcal{L}_u = 0 \end{cases}$$

## Singular Optimal Control - Example

$$\min_{x(\cdot), u(\cdot)} \frac{1}{2} \int_0^1 x_1^2 dt$$

$$\text{s.t. } \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$
$$-5 \leq u \leq 5$$

$$\mathcal{L}_u = \lambda_2 \quad \frac{d}{dt} \mathcal{L}_u = -\lambda_1 \quad \frac{d^2}{dt^2} \mathcal{L}_u = x_1 \quad \frac{d^3}{dt^3} \mathcal{L}_u = x_2 \quad \frac{d^4}{dt^4} \mathcal{L}_u = u$$

**Optimal input:**

$$u^* = \begin{cases} u_{\min} & \text{if } \lambda_2 > 0 \\ u_{\max} & \text{if } \lambda_2 < 0 \\ 0 & \text{if } \frac{d^i}{dt^i} \mathcal{L}_u = 0 \end{cases}$$

- Input is either in the bounds or zero !!
- Bang-bang input until  $x, \lambda = 0$

## Singular Optimal Control - Example

$$\min_{x(\cdot), u(\cdot)} \frac{1}{2} \int_0^1 x_1^2 dt$$

$$\text{s.t. } \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$
$$-5 \leq u \leq 5$$

$$\mathcal{L}_u = \lambda_2 \quad \frac{d}{dt} \mathcal{L}_u = -\lambda_1 \quad \frac{d^2}{dt^2} \mathcal{L}_u = x_1 \quad \frac{d^3}{dt^3} \mathcal{L}_u = x_2 \quad \frac{d^4}{dt^4} \mathcal{L}_u = u$$

**Optimal input:**

$$u^* = \begin{cases} u_{\min} & \text{if } \lambda_2 > 0 \\ u_{\max} & \text{if } \lambda_2 < 0 \\ 0 & \text{if } \frac{d^i}{dt^i} \mathcal{L}_u = 0 \end{cases}$$

- Input is either in the bounds or zero !!
- Bang-bang input until  $x, \lambda = 0$
- $\mathcal{L}_u$  is "controlled" via its 4<sup>th</sup>-order derivative

## Singular Optimal Control - Example

$$\min_{x(\cdot), u(\cdot)} \frac{1}{2} \int_0^1 x_1^2 dt$$

$$\text{s.t. } \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$
$$-5 \leq u \leq 5$$

$$\mathcal{L}_u = \lambda_2 \quad \frac{d}{dt} \mathcal{L}_u = -\lambda_1 \quad \frac{d^2}{dt^2} \mathcal{L}_u = x_1 \quad \frac{d^3}{dt^3} \mathcal{L}_u = x_2 \quad \frac{d^4}{dt^4} \mathcal{L}_u = u$$

**Optimal input:**

$$u^* = \begin{cases} u_{\min} & \text{if } \lambda_2 > 0 \\ u_{\max} & \text{if } \lambda_2 < 0 \\ 0 & \text{if } \frac{d^i}{dt^i} \mathcal{L}_u = 0 \end{cases}$$

- Input is either in the bounds or zero !!
- Bang-bang input until  $x, \lambda = 0$
- $\mathcal{L}_u$  is "controlled" via its 4<sup>th</sup>-order derivative
- Problem has a *degree of singularity* of  $\frac{4}{2} = 2$  (# of derivatives to get  $u$  is always even)

## Singular Optimal Control - Example

$$\min_{x(\cdot), u(\cdot)} \frac{1}{2} \int_0^1 x_1^2 dt$$

$$\text{s.t. } \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$
$$-5 \leq u \leq 5$$

$$\mathcal{L}_u = \lambda_2 \quad \frac{d}{dt} \mathcal{L}_u = -\lambda_1 \quad \frac{d^2}{dt^2} \mathcal{L}_u = x_1 \quad \frac{d^3}{dt^3} \mathcal{L}_u = x_2 \quad \frac{d^4}{dt^4} \mathcal{L}_u = u$$

**Optimal input:**

$$u^* = \begin{cases} u_{\min} & \text{if } \lambda_2 > 0 \\ u_{\max} & \text{if } \lambda_2 < 0 \\ 0 & \text{if } \frac{d^i}{dt^i} \mathcal{L}_u = 0 \end{cases}$$

- Input is either in the bounds or zero !!
- Bang-bang input until  $x, \lambda = 0$
- $\mathcal{L}_u$  is "controlled" via its 4<sup>th</sup>-order derivative
- Problem has a *degree of singularity* of  $\frac{4}{2} = 2$  (# of derivatives to get  $u$  is always even)

## Singular Optimal Control - Example

$$\min_{x(\cdot), u(\cdot)} \frac{1}{2} \int_0^1 x_1^2 dt$$

$$\text{s.t. } \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$
$$-5 \leq u \leq 5$$

$$\mathcal{L}_u = \lambda_2 \quad \frac{d}{dt} \mathcal{L}_u = -\lambda_1 \quad \frac{d^2}{dt^2} \mathcal{L}_u = x_1 \quad \frac{d^3}{dt^3} \mathcal{L}_u = x_2 \quad \frac{d^4}{dt^4} \mathcal{L}_u = u$$

**Optimal input:**

$$u^* = \begin{cases} u_{\min} & \text{if } \lambda_2 > 0 \\ u_{\max} & \text{if } \lambda_2 < 0 \\ 0 & \text{if } \frac{d^i}{dt^i} \mathcal{L}_u = 0 \end{cases}$$




- Input is either in the bounds or zero !!
- Bang-bang input until  $x, \lambda = 0$
- $\mathcal{L}_u$  is "controlled" via its 4<sup>th</sup>-order derivative
- Problem has a *degree of singularity* of  $\frac{4}{2} = 2$  (# of derivatives to get  $u$  is always even)

# Singular Optimal Control - Example

$$\min_{x(\cdot), u(\cdot)} \frac{1}{2} \int_0^1 x_1^2 dt$$

$$\text{s.t. } \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$
$$-5 \leq u \leq 5$$

## Optimal solution




## Singular Optimal Control - Example

$$\min_{x(\cdot), u(\cdot)} \frac{1}{2} \int_0^1 x_1^2 dt$$

$$\text{s.t. } \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$
$$-5 \leq u \leq 5$$


Optimal solution vs. solution from multiple-shooting ( $t_{k+1} - t_k = 0.01$ )



## Singular Optimal Control - Example


$$\begin{aligned} \min_{x(\cdot), u(\cdot)} \quad & \frac{1}{2} \int_0^1 x_1^2 dt \\ \text{s.t.} \quad & \dot{x} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u, \quad x(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \\ & -5 \leq u \leq 5 \end{aligned}$$

Optimal solution vs. solution from multiple-shooting ( $t_{k+1} - t_k = 0.01$ )



Very common behavior of numerical optimal control for singular problems.  
What is going on ?!?

## Singular optimal control with direct methods




Consider the problem:

$$\begin{aligned} J(\mathbf{u}) &= \phi(\mathbf{x}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t. } \dot{\mathbf{x}}(t) &= \mathbf{F}(\mathbf{x}(t), \mathbf{u}(t)) \\ \mathbf{x}(t_0) &= \mathbf{x}_0 \\ \mathbf{u}_{\min} &\leq \mathbf{u} \leq \mathbf{u}_{\max} \end{aligned}$$

with  $\mathbf{u}(t) = \mathbf{u}_k$  for  $t \in [t_k, t_{k+1}]$ .

## Singular optimal control with direct methods



Consider the problem:


$$\begin{aligned} J(\mathbf{u}) &= \phi(\mathbf{x}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t. } \dot{\mathbf{x}}(t) &= \mathbf{F}(\mathbf{x}(t), \mathbf{u}(t)) \\ \mathbf{x}(t_0) &= \mathbf{x}_0 \\ \mathbf{u}_{\min} &\leq \mathbf{u} \leq \mathbf{u}_{\max} \end{aligned}$$

with  $\mathbf{u}(t) = \mathbf{u}_k$  for  $t \in [t_k, t_{k+1}]$ . Then

$$\frac{\partial J}{\partial \mathbf{u}_k} = \int_{t_k}^{t_{k+1}} \mathcal{L}_{\mathbf{u}}(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k) dt$$

is zero when  $\mathbf{u}_k$  is off the bounds.

# Singular optimal control with direct methods




Consider the problem:

$$\begin{aligned} J(\mathbf{u}) &= \phi(\mathbf{x}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t. } \dot{\mathbf{x}}(t) &= \mathbf{F}(\mathbf{x}(t), \mathbf{u}(t)) \\ \mathbf{x}(t_0) &= \mathbf{x}_0 \\ \mathbf{u}_{\min} &\leq \mathbf{u} \leq \mathbf{u}_{\max} \end{aligned}$$


with  $\mathbf{u}(t) = \mathbf{u}_k$  for  $t \in [t_k, t_{k+1}]$ . Then

$$\frac{\partial J}{\partial \mathbf{u}_k} = \int_{t_k}^{t_{k+1}} \mathcal{L}_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k) dt$$

is zero when  $\mathbf{u}_k$  is off the bounds.



# Singular optimal control with direct methods




Consider the problem:

$$\begin{aligned} J(\mathbf{u}) &= \phi(\mathbf{x}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t. } \dot{\mathbf{x}}(t) &= \mathbf{F}(\mathbf{x}(t), \mathbf{u}(t)) \\ \mathbf{x}(t_0) &= \mathbf{x}_0 \\ \mathbf{u}_{\min} \leq \mathbf{u} &\leq \mathbf{u}_{\max} \end{aligned}$$


with  $\mathbf{u}(t) = \mathbf{u}_k$  for  $t \in [t_k, t_{k+1}]$ . Then

$$\frac{\partial J}{\partial \mathbf{u}_k} = \int_{t_k}^{t_{k+1}} \mathcal{L}_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k) dt$$

is zero when  $\mathbf{u}_k$  is off the bounds.



# Singular optimal control with direct methods




Consider the problem:

$$\begin{aligned} J(\mathbf{u}) &= \phi(\mathbf{x}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t. } \dot{\mathbf{x}}(t) &= \mathbf{F}(\mathbf{x}(t), \mathbf{u}(t)) \\ \mathbf{x}(t_0) &= \mathbf{x}_0 \\ \mathbf{u}_{\min} \leq \mathbf{u} &\leq \mathbf{u}_{\max} \end{aligned}$$

with  $\mathbf{u}(t) = \mathbf{u}_k$  for  $t \in [t_k, t_{k+1}]$ . Then

$$\frac{\partial J}{\partial \mathbf{u}_k} = \int_{t_k}^{t_{k+1}} \mathcal{L}_u(\mathbf{x}(t), \boldsymbol{\lambda}(t), \mathbf{u}_k) dt$$

is zero when  $\mathbf{u}_k$  is off the bounds.



# Outline

- 1 Introduction to the Pontryagin Maximum Principle (PMP)
- 2 Interpretation of  $H_u$
- 3 Input bounds in Indirect Optimal Control
- 4 Singular Optimal Control problems
- 5 General constraints in Indirect Optimal Control

## State constraints in indirect optimal control

OCP with state (mixed) constraints:

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{u}} \quad & \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt \\ \text{s.t.} \quad & \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0 \\ & \mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0 \end{aligned}$$

## State constraints in indirect optimal control

OCP with state (mixed) constraints:

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{u}} \quad & \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt \\ \text{s.t.} \quad & \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0 \\ & \mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0 \end{aligned}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \mathbf{h}(\mathbf{x}, \mathbf{u})$$

## State constraints in indirect optimal control

OCP with state (mixed) constraints:

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{u}} \quad & \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt \\ \text{s.t.} \quad & \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0 \\ & \mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0 \end{aligned}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \mathbf{h}(\mathbf{x}, \mathbf{u})$$

Get the optimal control solution from:  $\mathbf{u}^* = \underset{\mathbf{u}}{\operatorname{argmin}} H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u})$ , with:

## State constraints in indirect optimal control

OCP with state (mixed) constraints:

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{u}} \quad & \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt \\ \text{s.t.} \quad & \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0 \\ & \mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0 \end{aligned}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \mathbf{h}(\mathbf{x}, \mathbf{u})$$

Get the optimal control solution from:  $\mathbf{u}^* = \underset{\mathbf{u}}{\operatorname{argmin}} H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u})$ , with:

$$\text{States : } \dot{\mathbf{x}} = \mathbf{F}, \quad \mathbf{x}(t_0) = \mathbf{x}_0$$

$$\text{Costates : } \dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H, \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$$

## State constraints in indirect optimal control

OCP with state (mixed) constraints:

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{u}} \quad & \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt \\ \text{s.t.} \quad & \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0 \\ & \mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0 \end{aligned}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \mathbf{h}(\mathbf{x}, \mathbf{u})$$

Get the optimal control solution from:  $\mathbf{u}^* = \underset{\mathbf{u}}{\operatorname{argmin}} H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u})$ , with:

States :  $\dot{\mathbf{x}} = \mathbf{F}, \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H, \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

Feasibility :  $\mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0, \quad \boldsymbol{\mu} \geq 0$

## State constraints in indirect optimal control

OCP with state (mixed) constraints:

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{u}} \quad & \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt \\ \text{s.t.} \quad & \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0 \\ & \mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0 \end{aligned}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \mathbf{h}(\mathbf{x}, \mathbf{u})$$

Get the optimal control solution from:  $\mathbf{u}^* = \underset{\mathbf{u}}{\operatorname{argmin}} H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u})$ , with:

States :  $\dot{\mathbf{x}} = \mathbf{F}, \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H, \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

Feasibility :  $\mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0, \quad \boldsymbol{\mu} \geq 0$

Complementary slack. :  $\boldsymbol{\mu}^\top \mathbf{h}(\mathbf{x}, \mathbf{u}) = 0$

## State constraints in indirect optimal control

OCP with state (mixed) constraints:

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{u}} \quad & \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt \\ \text{s.t.} \quad & \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0 \\ & \mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0 \end{aligned}$$

Define the Hamiltonian function

$$H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = L(\mathbf{x}, \mathbf{u}) + \boldsymbol{\lambda}^\top \mathbf{F}(\mathbf{x}, \mathbf{u}) + \boldsymbol{\mu}^\top \mathbf{h}(\mathbf{x}, \mathbf{u})$$

Get the optimal control solution from:  $\mathbf{u}^* = \underset{\mathbf{u}}{\operatorname{argmin}} H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u})$ , with:

States :  $\dot{\mathbf{x}} = \mathbf{F}, \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H, \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

Feasibility :  $\mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0, \quad \boldsymbol{\mu} \geq 0$

Complementary slack. :  $\boldsymbol{\mu}^\top \mathbf{h}(\mathbf{x}, \mathbf{u}) = 0$

The PMP equations can be hard to solve in general. No good PMP-based general-purpose solver available.

## State constraints in indirect optimal control

OCP with state (mixed) constraints:

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{u}} \quad & \phi(\mathbf{x}(t_f)) + \int_{t_0}^{t_f} L(\mathbf{x}(t), \mathbf{u}(t)) dt \\ \text{s.t.} \quad & \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0 \\ & \mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0 \end{aligned}$$

Tentative solutions based on IP method

Stationarity :  $H_u(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}) = 0$

States :  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}), \quad \mathbf{x}(t_0) = \mathbf{x}_0$

Costates :  $\dot{\boldsymbol{\lambda}} = -\nabla_{\mathbf{x}} H(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\mu}, \mathbf{u}), \quad \boldsymbol{\lambda}(t_f) = \nabla_{\mathbf{x}} \phi(\mathbf{x}(t_f))$

Feasibility :  $\mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0, \quad \boldsymbol{\mu} \geq 0$

Complementary slack. :  $\boldsymbol{\mu}^\top \mathbf{h}(\mathbf{x}, \mathbf{u}) = \tau$

- Handle dynamics + constraints  $H_u = 0$  and  $\boldsymbol{\mu}^\top \mathbf{h} = \tau$  as a DAE
- Handle  $\mathbf{h}(\mathbf{x}, \mathbf{u}) \leq 0$  and  $\boldsymbol{\mu} \geq 0$  via step length (c.f. IP lecture)
- Also done using Primal IP approach (move  $\mathbf{h}$  in the cost using log barrier)