

# Numerical Optimal Control

## Lecture 12: Optimal Control with DAEs

Sébastien Gros

ITK, NTNU

NTNU PhD course

# Outline

1 Introduction

2 Differential Index

3 Index Reduction & Consistency

4 Multiple Shooting & Direct Collocation with DAEs

# Optimal Control with ODEs / DAEs

## ODE-constrained

$$\begin{aligned} \min_{\mathbf{x}(\cdot), \mathbf{u}(\cdot)} \quad & \phi(\mathbf{x}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t.} \quad & \dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}, t) \\ & \mathbf{h}(\mathbf{u}(t), \mathbf{x}(t), t) \leq 0 \\ & \mathbf{b}(\mathbf{x}(t_0), \mathbf{x}(t_f)) = 0 \end{aligned}$$

## DAE-constrained

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{z}, \mathbf{u}} \quad & \phi(\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t.} \quad & \mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0, \\ & \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u}) = 0 \\ & \mathbf{h}(\mathbf{u}(t), \mathbf{x}(t), \mathbf{z}(t)) \leq 0 \\ & \mathbf{b}(\mathbf{x}(t_0), \mathbf{x}(t_f)) = 0 \end{aligned}$$

- Intuitive definition: some **variables** in the dynamics are not time-differentiated
- Size of  $\mathbf{F}$  + size of  $\mathbf{G}$  = size of  $\mathbf{x}$  + size of  $\mathbf{z}$

Why do we use DAEs ? Where do they come from?

# Why do we use DAEs ? Where do they come from?


## Large systems made of many subsystems:

- ODEs describe each subsystems independently
- Algebraic relationships describe e.g. balance equations, flow, etc...
- DAE model is easier to develop, maintain, modify

# Why do we use DAEs ? Where do they come from?

## Large systems made of many subsystems:


- ODEs describe each subsystems independently
- Algebraic relationships describe e.g. balance equations, flow, etc...
- DAE model is easier to develop, maintain, modify



# Why do we use DAEs ? Where do they come from?

## Large systems made of many subsystems:

- ODEs describe each subsystems independently
- Algebraic relationships describe e.g. balance equations, flow, etc...
- DAE model is easier to develop, maintain, modify




## Mechanical/Robotic systems:

- DAE models can be much simpler than ODE models
- Modelling procedure is often easier using DAEs
- Often yields more "natural" representations of the system

# Why do we use DAEs ? Where do they come from?

## Large systems made of many subsystems:

- ODEs describe each subsystems independently
- Algebraic relationships describe e.g. balance equations, flow, etc...
- DAE model is easier to develop, maintain, modify



## Mechanical/Robotic systems:

- DAE models can be much simpler than ODE models
- Modelling procedure is often easier using DAEs
- Often yields more "natural" representations of the system

Delta robot





# A brief taxonomy of DAEs

## A brief taxonomy of DAEs

### Fully-implicit DAE:

$$\mathbf{F}(\mathbf{x}, \dot{\mathbf{x}}, \mathbf{u}) = 0$$

with  $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$  rank-deficient.

## A brief taxonomy of DAEs

### Fully-implicit DAE:

$$\mathbf{F}(\mathbf{x}, \dot{\mathbf{x}}, \mathbf{u}) = 0$$

with  $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$  rank-deficient.

Usually in the form:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

when some variables are clearly algebraic.

## A brief taxonomy of DAEs

### Fully-implicit DAE:

$$\mathbf{F}(\mathbf{x}, \dot{\mathbf{x}}, \mathbf{u}) = 0$$

with  $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$  rank-deficient.

Usually in the form:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

when some variables are clearly algebraic.

### Semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

## A brief taxonomy of DAEs

### Fully-implicit DAE:

$$\mathbf{F}(\mathbf{x}, \dot{\mathbf{x}}, \mathbf{u}) = 0$$

with  $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$  rank-deficient.

Usually in the form:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

when some variables are clearly algebraic.

### Semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

## A brief taxonomy of DAEs

### Fully-implicit DAE:

$$\mathbf{F}(\mathbf{x}, \dot{\mathbf{x}}, \mathbf{u}) = 0$$

with  $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$  rank-deficient.

Usually in the form:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

when some variables are clearly **algebraic**.

### Semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

**Any fully implicit DAE can be transformed into a semi-explicit one.**

## A brief taxonomy of DAEs

### Fully-implicit DAE:

$$\mathbf{F}(\mathbf{x}, \dot{\mathbf{x}}, \mathbf{u}) = 0$$

with  $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$  rank-deficient.

Usually in the form:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

when some variables are clearly **algebraic**.

### Semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$


$$0 = \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

**Any fully implicit DAE can be transformed into a semi-explicit one.** Though, it is not always wise to do so. The transformation can turn a simple set of equation into a very complex one !!

## An example of a DAE - 3D pendulum

**Position** given by  $\mathbf{p} \in \mathbb{R}^3$ , dynamics:

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3$$






## An example of a DAE - 3D pendulum

**Position** given by  $\mathbf{p} \in \mathbb{R}^3$ , dynamics:

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$$




**Force in the cable:** direction given by  $-\mathbf{p}$ , amplitude given by algebraic variable  $z \in \mathbb{R}_+$

## An example of a DAE - 3D pendulum

**Position** given by  $\mathbf{p} \in \mathbb{R}^3$ , dynamics:

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$$



**Force in the cable:** direction given by  $-\mathbf{p}$ , amplitude given by algebraic variable  $z \in \mathbb{R}_+$

Then  $z$  must be chosen such that:


$$\mathbf{G}(\mathbf{p}) = \mathbf{p}^\top \mathbf{p} - L^2 = 0$$

holds at all time.

## An example of a DAE - 3D pendulum

**Position** given by  $\mathbf{p} \in \mathbb{R}^3$ , dynamics:

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$$



**Force in the cable:** direction given by  $-\mathbf{p}$ , amplitude given by algebraic variable  $z \in \mathbb{R}_+$

Then  $z$  must be chosen such that:

$$\mathbf{G}(\mathbf{p}) = \mathbf{p}^\top \mathbf{p} - L^2 = 0$$

holds at all time.

Using  $\mathbf{v} = \dot{\mathbf{p}}$ , the **DAE** reads as:

$$\dot{\mathbf{p}} = \mathbf{v}$$


$$\dot{\mathbf{v}} = \frac{\mathbf{u}}{m} - g\mathbf{e}_3 - \frac{z}{m}\mathbf{p}$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2$$

## An example of a DAE - 3D pendulum

**Position** given by  $\mathbf{p} \in \mathbb{R}^3$ , dynamics:

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$$



**Force in the cable:** direction given by  $-\mathbf{p}$ , amplitude given by algebraic variable  $z \in \mathbb{R}_+$

Then  $z$  must be chosen such that:

$$\mathbf{G}(\mathbf{p}) = \mathbf{p}^\top \mathbf{p} - L^2 = 0$$

holds at all time.

Using  $\mathbf{v} = \dot{\mathbf{p}}$ , the **DAE** reads as:

$$\dot{\mathbf{p}} = \mathbf{v}$$

$$\dot{\mathbf{v}} = \frac{\mathbf{u}}{m} - g\mathbf{e}_3 - \frac{z}{m}\mathbf{p}$$


$$0 = \mathbf{p}^\top \mathbf{p} - L^2$$

What kind of DAE is that !?!

## An example of a DAE - 3D pendulum

**Position** given by  $\mathbf{p} \in \mathbb{R}^3$ , dynamics:

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$$



**Force in the cable:** direction given by  $-\mathbf{p}$ , amplitude given by algebraic variable  $z \in \mathbb{R}_+$

Then  $z$  must be chosen such that:

$$\mathbf{G}(\mathbf{p}) = \mathbf{p}^\top \mathbf{p} - L^2 = 0$$

holds at all time.

Using  $\mathbf{v} = \dot{\mathbf{p}}$ , the **DAE** reads as:

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= \frac{\mathbf{u}}{m} - g\mathbf{e}_3 - \frac{z}{m}\mathbf{p} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

What kind of DAE is that !?!

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{u}, z)$$

$$0 = \mathbf{G}(\mathbf{x})$$

**Semi-explicit** with  $\mathbf{G}$  independent of  $z$ ...

# Outline

- 1 Introduction
- 2 **Differential Index**
- 3 Index Reduction & Consistency
- 4 Multiple Shooting & Direct Collocation with DAEs

## Easy & Hard DAEs

**DAE:**

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

“At any time instant, for given  $\mathbf{x}, \mathbf{u}$ , the DAE equation provides  $\dot{\mathbf{x}}, \mathbf{z}$ , generating the trajectories.”

## Easy & Hard DAEs

### DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

“At any time instant, for given  $\mathbf{x}, \mathbf{u}$ , the DAE equation provides  $\dot{\mathbf{x}}, \mathbf{z}$ , generating the trajectories.”

What is a “well-behaved” DAE ?

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

... i.e. when can we compute  $\dot{\mathbf{x}}, \mathbf{z}$  “easily” ?



## Easy & Hard DAEs

### DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

“At any time instant, for given  $\mathbf{x}, \mathbf{u}$ , the DAE equation provides  $\dot{\mathbf{x}}, \mathbf{z}$ , generating the trajectories.”

What is a “well-behaved” DAE ?

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

... i.e. when can we compute  $\dot{\mathbf{x}}, \mathbf{z}$  “easily” ?

Consider  $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$  as a **root-finding problem** in  $\dot{\mathbf{x}}, \mathbf{z}$ . When can we find  $\dot{\mathbf{x}}, \mathbf{z}$ , (typ. using Newton) ?

## Easy & Hard DAEs

### DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

“At any time instant, for given  $\mathbf{x}, \mathbf{u}$ , the DAE equation provides  $\dot{\mathbf{x}}, \mathbf{z}$ , generating the trajectories.”

### Implicit Function Theorem:

$$\begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \end{bmatrix}$$

full-rank at  $\mathbf{x}, \mathbf{u}$  guarantees that the DAE is “solvable” at  $\dot{\mathbf{x}}, \mathbf{z}$ .

What is a “well-behaved” DAE ?

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

... i.e. when can we compute  $\dot{\mathbf{x}}, \mathbf{z}$  “easily” ?

Consider  $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$  as a **root-finding problem** in  $\dot{\mathbf{x}}, \mathbf{z}$ . When can we find  $\dot{\mathbf{x}}, \mathbf{z}$ , (typ. using Newton) ?

## Easy & Hard DAEs

### DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

“At any time instant, for given  $\mathbf{x}, \mathbf{u}$ , the DAE equation provides  $\dot{\mathbf{x}}, \mathbf{z}$ , generating the trajectories.”

### Implicit Function Theorem:

$$\begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \end{bmatrix}$$

full-rank at  $\mathbf{x}, \mathbf{u}$  guarantees that the DAE is “solvable” at  $\dot{\mathbf{x}}, \mathbf{z}$ .

For semi-explicit DAEs

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$\frac{\partial \mathbf{G}}{\partial \mathbf{z}}$  full rank makes the DAE “solvable”.

What is a “well-behaved” DAE ?

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

... i.e. when can we compute  $\dot{\mathbf{x}}, \mathbf{z}$  “easily” ?

Consider  $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$  as a **root-finding problem** in  $\dot{\mathbf{x}}, \mathbf{z}$ . When can we find  $\dot{\mathbf{x}}, \mathbf{z}$ , (typ. using Newton) ?

# Semi-explicit DAEs - Differential Index

## Definition

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE

# Semi-explicit DAEs - Differential Index

## Definition

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE

**Remark:** for an index-1 semi-explicit DAE:

$$\frac{d}{dt} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u}) = \frac{\partial \mathbf{G}}{\partial \mathbf{x}} \mathbf{F} + \frac{\partial \mathbf{G}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{G}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

yields a pure ODE.

# Semi-explicit DAEs - Differential Index

## Definition

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE

**Remark:** for an index-1 semi-explicit DAE:

$$\frac{d}{dt} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u}) = \frac{\partial \mathbf{G}}{\partial \mathbf{x}} \mathbf{F} + \frac{\partial \mathbf{G}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{G}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

yields a pure ODE. Then:

$$\dot{\mathbf{z}} = -\frac{\partial \mathbf{G}}{\partial \mathbf{z}}^{-1} \left( \frac{\partial \mathbf{G}}{\partial \mathbf{x}} \mathbf{F} + \frac{\partial \mathbf{G}}{\partial \mathbf{u}} \dot{\mathbf{u}} \right)$$

is well-defined, i.e.  $\frac{\partial \mathbf{G}}{\partial \mathbf{z}}$  is full rank !!

# Semi-explicit DAEs - Differential Index

## Definition

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE

**Remark:** for an index-1 semi-explicit DAE:

$$\frac{d}{dt} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u}) = \frac{\partial \mathbf{G}}{\partial \mathbf{x}} \mathbf{F} + \frac{\partial \mathbf{G}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{G}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

yields a pure ODE. Then:

$$\dot{\mathbf{z}} = -\frac{\partial \mathbf{G}}{\partial \mathbf{z}}^{-1} \left( \frac{\partial \mathbf{G}}{\partial \mathbf{x}} \mathbf{F} + \frac{\partial \mathbf{G}}{\partial \mathbf{u}} \dot{\mathbf{u}} \right)$$

is well-defined, i.e.  $\frac{\partial \mathbf{G}}{\partial \mathbf{z}}$  is full rank !!

**Index-1 DAEs**

$\equiv$

**“easy” DAEs**

## Differential Index - 3D pendulum

**Example:** 3D pendulum


$$\begin{aligned} m\ddot{\mathbf{p}} &= \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p} \\ 0 &= \underbrace{\frac{1}{2}(\mathbf{p}^\top \mathbf{p} - L^2)}_{G(\mathbf{x})} \end{aligned}$$

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE





## Differential Index - 3D pendulum

**Example:** 3D pendulum

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$$
$$0 = \underbrace{\frac{1}{2}(\mathbf{p}^\top \mathbf{p} - L^2)}_{G(\mathbf{x})}$$

Perform **two** time differentiations on  $G$  yields:


$$\ddot{G} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE



## Differential Index - 3D pendulum

**Example:** 3D pendulum

$$\begin{aligned} m\ddot{\mathbf{p}} &= \mathbf{u} - mge_3 - z\mathbf{p} \\ 0 &= \underbrace{\frac{1}{2}(\mathbf{p}^\top \mathbf{p} - L^2)}_{G(\mathbf{x})} \end{aligned}$$

Perform **two** time differentiations on  $G$  yields:

$$\ddot{G} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

Substitute  $\ddot{\mathbf{p}}$  from  $m\ddot{\mathbf{p}} = \mathbf{u} - mge_3 - z\mathbf{p}$  yields:


$$\mathbf{p}^\top \left( \frac{1}{m}\mathbf{u} - ge_3 - \frac{1}{m}z\mathbf{p} \right) + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE



## Differential Index - 3D pendulum

**Example:** 3D pendulum

$$\begin{aligned} m\ddot{\mathbf{p}} &= \mathbf{u} - mge_3 - z\mathbf{p} \\ 0 &= \underbrace{\frac{1}{2}(\mathbf{p}^\top \mathbf{p} - L^2)}_{G(\mathbf{x})} \end{aligned}$$

Perform **two** time differentiations on  $G$  yields:

$$\ddot{G} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

Substitute  $\ddot{\mathbf{p}}$  from  $m\ddot{\mathbf{p}} = \mathbf{u} - mge_3 - z\mathbf{p}$  yields:

$$\mathbf{p}^\top \left( \frac{1}{m}\mathbf{u} - ge_3 - \frac{1}{m}z\mathbf{p} \right) + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

which is solvable for  $z$ :


$$z = \frac{1}{\mathbf{p}^\top \mathbf{p}} \left( \mathbf{p}^\top \mathbf{u} - mg\mathbf{p}^\top e_3 + m\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \right)$$

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE



## Differential Index - 3D pendulum

**Example:** 3D pendulum

$$\begin{aligned} m\ddot{\mathbf{p}} &= \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p} \\ 0 &= \underbrace{\frac{1}{2}(\mathbf{p}^\top \mathbf{p} - L^2)}_{G(\mathbf{x})} \end{aligned}$$

Perform **two** time differentiations on  $G$  yields:

$$\ddot{G} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

Substitute  $\ddot{\mathbf{p}}$  from  $m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$  yields:

$$\mathbf{p}^\top \left( \frac{1}{m}\mathbf{u} - g\mathbf{e}_3 - \frac{1}{m}z\mathbf{p} \right) + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

which is solvable for  $z$ :

$$z = \frac{1}{\mathbf{p}^\top \mathbf{p}} \left( \mathbf{p}^\top \mathbf{u} - mg\mathbf{p}^\top \mathbf{e}_3 + m\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \right)$$


one extra time differentiation yields an ODE ( $\dot{z}$  appears)

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE



## Differential Index - 3D pendulum

**Example:** 3D pendulum

$$\begin{aligned} m\ddot{\mathbf{p}} &= \mathbf{u} - mge_3 - z\mathbf{p} \\ 0 &= \underbrace{\frac{1}{2}(\mathbf{p}^\top \mathbf{p} - L^2)}_{G(\mathbf{x})} \end{aligned}$$

Perform **two** time differentiations on  $G$  yields:

$$\ddot{G} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

Substitute  $\ddot{\mathbf{p}}$  from  $m\ddot{\mathbf{p}} = \mathbf{u} - mge_3 - z\mathbf{p}$  yields:

$$\mathbf{p}^\top \left( \frac{1}{m}\mathbf{u} - ge_3 - \frac{1}{m}z\mathbf{p} \right) + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

which is solvable for  $z$ :

$$z = \frac{1}{\mathbf{p}^\top \mathbf{p}} \left( \mathbf{p}^\top \mathbf{u} - mg\mathbf{p}^\top e_3 + m\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \right)$$


one extra time differentiation yields an ODE ( $\dot{z}$  appears)

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE



**The 3D pendulum is an index-3 DAE !!** Note: this is a general observation for mechanical systems modelled via DAEs

## Differential Index - 3D pendulum

**Example:** 3D pendulum

$$m\ddot{\mathbf{p}} = \mathbf{u} - mge_3 - z\mathbf{p}$$
$$0 = \underbrace{\frac{1}{2}(\mathbf{p}^\top \mathbf{p} - L^2)}_{G(\mathbf{x})}$$

Perform **two** time differentiations on  $G$  yields:

$$\ddot{\mathbf{G}} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

Assemble:


$$m\ddot{\mathbf{p}} + z\mathbf{p} = \mathbf{u} - mge_3$$
$$\mathbf{p}^\top \ddot{\mathbf{p}} = -\dot{\mathbf{p}}^\top \dot{\mathbf{p}}$$

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE



**The 3D pendulum is an index-3 DAE !!** Note: this is a general observation for mechanical systems modelled via DAEs

## Differential Index - 3D pendulum

**Example:** 3D pendulum

$$\begin{aligned} m\ddot{\mathbf{p}} &= \mathbf{u} - m\mathbf{g}\mathbf{e}_3 - z\mathbf{p} \\ 0 &= \underbrace{\frac{1}{2}(\mathbf{p}^\top \mathbf{p} - L^2)}_{G(\mathbf{x})} \end{aligned}$$

Perform **two** time differentiations on  $G$  yields:

$$\ddot{G} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

Assemble:

$$\begin{aligned} m\ddot{\mathbf{p}} + z\mathbf{p} &= \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ \mathbf{p}^\top \ddot{\mathbf{p}} &= -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{aligned}$$

in matrix form yields:


$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE



**The 3D pendulum is an index-3 DAE !!** Note: this is a general observation for mechanical systems modelled via DAEs

## Differential Index - 3D pendulum

**Example:** 3D pendulum

$$\begin{aligned} m\ddot{\mathbf{p}} &= \mathbf{u} - m\mathbf{g}\mathbf{e}_3 - z\mathbf{p} \\ 0 &= \underbrace{\frac{1}{2}(\mathbf{p}^\top \mathbf{p} - L^2)}_{G(\mathbf{x})} \end{aligned}$$

Perform **two** time differentiations on  $G$  yields:

$$\ddot{G} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

Assemble:

$$\begin{aligned} m\ddot{\mathbf{p}} + z\mathbf{p} &= \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ \mathbf{p}^\top \ddot{\mathbf{p}} &= -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{aligned}$$

in matrix form yields:

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$


**This is an index-1 (i.e. “easy”) DAE (implicit) !!**

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE



**The 3D pendulum is an index-3 DAE !!** Note: this is a general observation for mechanical systems modelled via DAEs



## Differential Index - 3D pendulum

**Example:** 3D pendulum

$$\begin{aligned} m\ddot{\mathbf{p}} &= \mathbf{u} - m\mathbf{g}\mathbf{e}_3 - z\mathbf{p} \\ 0 &= \underbrace{\frac{1}{2}(\mathbf{p}^\top \mathbf{p} - L^2)}_{G(\mathbf{x})} \end{aligned}$$

Perform **two** time differentiations on  $G$  yields:

$$\ddot{G} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$

Assemble:

$$\begin{aligned} m\ddot{\mathbf{p}} + z\mathbf{p} &= \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ \mathbf{p}^\top \ddot{\mathbf{p}} &= -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{aligned}$$

in matrix form yields:

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$


**This is an index-1 (i.e. “easy”) DAE (implicit) !!**

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE



**We have converted the index-3 DAE into an index-1 DAE !!**

## Differential Index - 3D pendulum

**Example:** 3D pendulum

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - \underbrace{\mathbf{z}\mathbf{p}}_{\mathbf{G}(\mathbf{x})}$$
$$0 = \frac{1}{2} \underbrace{(\mathbf{p}^\top \mathbf{p} - L^2)}_{\mathbf{G}(\mathbf{x})}$$

Perform **two** time differentiations on  $\mathbf{G}$  yields:

$$\ddot{\mathbf{G}} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}} = 0$$


Transforming a high-index DAE into an equivalent lower-index one is labelled **index reduction**. This is the most popular approach to tackle high-order DAEs numerically.

For a **semi-explicit DAE** the differential index is the minimum  $i$  such that:

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \frac{d^i}{dt^i} \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

is an ODE




**We have converted the index-3 DAE into an index-1 DAE !!**

# Outline

- 1 Introduction
- 2 Differential Index
- 3 Index Reduction & Consistency**
- 4 Multiple Shooting & Direct Collocation with DAEs

## DAE Consistency - 3D pendulum

Does the index reduction really yield equivalent models ?




### Index-3 DAE

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$$
$$0 = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2)$$


### Index-1 DAE

$$\begin{bmatrix} \frac{m}{l} & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - mg\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$



## DAE Consistency - 3D pendulum

Does the index reduction really yield equivalent models ?





### Index-3 DAE

$$m\ddot{\mathbf{p}} = \mathbf{u} - m\mathbf{g}\mathbf{e}_3 - \mathbf{z}\mathbf{p}$$
$$0 = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2)$$


### Index-1 DAE

$$\begin{bmatrix} mI & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ \mathbf{z} \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$



## DAE Consistency - 3D pendulum

Does the index reduction really yield equivalent models ?





### Index-3 DAE

$$\begin{aligned} m\ddot{\mathbf{p}} &= \mathbf{u} - m\mathbf{g}\mathbf{e}_3 - \mathbf{z}\mathbf{p} \\ 0 &= \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - l^2) \end{aligned}$$

### Index-1 DAE

$$\begin{bmatrix} mI & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$




What is going on ??

# DAE Consistency - 3D pendulum

## Index-3 DAE

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - \lambda\mathbf{p}$$

$$0 = \frac{1}{2} \left( \mathbf{p}^\top \mathbf{p} - L^2 \right)$$



# DAE Consistency - 3D pendulum

## Index-3 DAE

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$$


$$0 = \frac{1}{2} \left( \mathbf{p}^\top \mathbf{p} - L^2 \right)$$

## Index reduction

$$\mathbf{G} = \frac{1}{2} \left( \mathbf{p}^\top \mathbf{p} - L^2 \right)$$

$$\dot{\mathbf{G}} = \mathbf{p}^\top \dot{\mathbf{p}}$$

$$\ddot{\mathbf{G}} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}}$$





# DAE Consistency - 3D pendulum

## Index-3 DAE

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$$

$$0 = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2)$$

## Index reduction

$$\mathbf{G} = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2)$$


$$\dot{\mathbf{G}} = \mathbf{p}^\top \dot{\mathbf{p}}$$

$$\ddot{\mathbf{G}} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}}$$

## Index-1 DAE

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - mg\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

... is built to impose  $\ddot{\mathbf{G}} = 0$  at all time.



# DAE Consistency - 3D pendulum

## Index-3 DAE

$$m\ddot{\mathbf{p}} = \mathbf{u} - m\mathbf{g}\mathbf{e}_3 - z\mathbf{p}$$

$$0 = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2)$$

## Index reduction

$$\mathbf{G} = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2)$$

$$\dot{\mathbf{G}} = \mathbf{p}^\top \dot{\mathbf{p}}$$


$$\ddot{\mathbf{G}} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}}$$

## Index-1 DAE

$$\begin{bmatrix} m/l & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

... is built to impose  $\ddot{\mathbf{G}} = 0$  at all time. But it does not ensure

$$\dot{\mathbf{G}} = 0 \quad \text{and} \quad \mathbf{G} = 0 !!$$



# DAE Consistency - 3D pendulum

## Index-3 DAE

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$$

$$0 = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2)$$

## Index reduction

$$\mathbf{G} = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2)$$

$$\dot{\mathbf{G}} = \mathbf{p}^\top \dot{\mathbf{p}}$$




$$\ddot{\mathbf{G}} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}}$$

## Index-1 DAE

$$\begin{bmatrix} mI & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - mg\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

... is built to impose  $\ddot{\mathbf{G}} = 0$  at all time. But it does not ensure

$$\dot{\mathbf{G}} = 0 \quad \text{and} \quad \mathbf{G} = 0 !!$$



# DAE Consistency - 3D pendulum

## Index-3 DAE

$$m\ddot{\mathbf{p}} = \mathbf{u} - mg\mathbf{e}_3 - z\mathbf{p}$$

$$0 = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2)$$

## Index reduction

$$\mathbf{G} = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2)$$

$$\dot{\mathbf{G}} = \mathbf{p}^\top \dot{\mathbf{p}}$$

$$\ddot{\mathbf{G}} = \mathbf{p}^\top \ddot{\mathbf{p}} + \dot{\mathbf{p}}^\top \dot{\mathbf{p}}$$




## Index-1 DAE

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - mg\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

... is built to impose  $\ddot{\mathbf{G}} = 0$  at all time. But it does not ensure

$$\dot{\mathbf{G}} = 0 \quad \text{and} \quad \mathbf{G} = 0 !!$$

How can we address that ??






# DAE Consistency - 3D pendulum

## Index-1 DAE

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

... is built to impose  $\ddot{\mathbf{G}} = 0$  at all time.






# DAE Consistency - 3D pendulum

## Index-1 DAE

$$\begin{bmatrix} m/l & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

... is built to impose  $\ddot{\mathbf{G}} = 0$  at all time.

Then if  $\mathbf{G} = 0$  and  $\dot{\mathbf{G}} = 0$  are satisfied at **any** time on the trajectory, then they are satisfied at **all** time.



# DAE Consistency - 3D pendulum

## Index-1 DAE




$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - mg\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

... is built to impose  $\ddot{\mathbf{G}} = 0$  at all time.

Then if  $\mathbf{G} = 0$  and  $\dot{\mathbf{G}} = 0$  are satisfied at **any** time on the trajectory, then they are satisfied at **all** time.

An index-reduced DAE must come with **consistency conditions**. E.g. for the 3D pendulum, the index-1 DAE should be given as:

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - mg\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$



# DAE Consistency - 3D pendulum

## Index-1 DAE

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

... is built to impose  $\ddot{\mathbf{G}} = 0$  at all time.

Then if  $\mathbf{G} = 0$  and  $\dot{\mathbf{G}} = 0$  are satisfied at **any** time on the trajectory, then they are satisfied at **all** time.




An index-reduced DAE must come with **consistency conditions**. E.g. for the 3D pendulum, the index-1 DAE should be given as:

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

with the consistency conditions:

$$\mathbf{G} = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2) = 0, \quad \dot{\mathbf{G}} = \mathbf{p}^\top \dot{\mathbf{p}} = 0$$

... to be satisfied e.g. at  $t_0$ .





# DAE Consistency - 3D pendulum

## Index-1 DAE

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

... is built to impose  $\ddot{\mathbf{G}} = 0$  at all time.

Then if  $\mathbf{G} = 0$  and  $\dot{\mathbf{G}} = 0$  are satisfied at **any** time on the trajectory, then they are satisfied at **all** time.




An index-reduced DAE must come with **consistency conditions**. E.g. for the 3D pendulum, the index-1 DAE should be given as:

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$

with the consistency conditions:

$$\mathbf{G} = \frac{1}{2} (\mathbf{p}^\top \mathbf{p} - L^2) = 0, \quad \dot{\mathbf{G}} = \mathbf{p}^\top \dot{\mathbf{p}} = 0$$

... to be satisfied e.g. at  $t_0$ .



# Outline

- 1 Introduction
- 2 Differential Index
- 3 Index Reduction & Consistency
- 4 Multiple Shooting & Direct Collocation with DAEs**

# Formulating an OCP

## Semi-explicit DAE-constrained OCP

$$\begin{aligned} \min_{\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)} \quad & \phi(\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t.} \quad & \dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \\ & 0 = \mathbf{G}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \\ & \mathbf{h}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \leq 0 \\ & \mathbf{x}(0) - \bar{\mathbf{x}}_0 = 0 \end{aligned}$$

# Formulating an OCP

## Semi-explicit DAE-constrained OCP

$$\begin{aligned} \min_{\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)} \quad & \phi(\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t.} \quad & \dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \\ & 0 = \mathbf{G}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \\ & \mathbf{h}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \leq 0 \\ & \mathbf{x}(0) - \bar{\mathbf{x}}_0 = 0 \end{aligned}$$

## Fully implicit DAE-constrained OCP

$$\begin{aligned} \min_{\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)} \quad & \phi(\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t.} \quad & \mathbf{F}(\dot{\mathbf{x}}(t), \mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) = 0 \\ & \mathbf{h}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \leq 0 \\ & \mathbf{x}(0) - \bar{\mathbf{x}}_0 = 0 \end{aligned}$$

# Formulating an OCP

## Semi-explicit DAE-constrained OCP

$$\begin{aligned} \min_{\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)} \quad & \phi(\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t.} \quad & \dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \\ & 0 = \mathbf{G}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \\ & \mathbf{h}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \leq 0 \\ & \mathbf{x}(0) - \bar{\mathbf{x}}_0 = 0 \end{aligned}$$

## Fully implicit DAE-constrained OCP

$$\begin{aligned} \min_{\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)} \quad & \phi(\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t.} \quad & \mathbf{F}(\dot{\mathbf{x}}(t), \mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) = 0 \\ & \mathbf{h}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \leq 0 \\ & \mathbf{x}(0) - \bar{\mathbf{x}}_0 = 0 \end{aligned}$$

- OCPs based on **index-1 DAEs** are the most common

# Formulating an OCP

## Semi-explicit DAE-constrained OCP

$$\begin{aligned} \min_{\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)} \quad & \phi(\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t.} \quad & \dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \\ & 0 = \mathbf{G}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \\ & \mathbf{h}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \leq 0 \\ & \mathbf{x}(0) - \bar{\mathbf{x}}_0 = 0 \end{aligned}$$

## Fully implicit DAE-constrained OCP

$$\begin{aligned} \min_{\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)} \quad & \phi(\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t.} \quad & \mathbf{F}(\dot{\mathbf{x}}(t), \mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) = 0 \\ & \mathbf{h}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \leq 0 \\ & \mathbf{x}(0) - \bar{\mathbf{x}}_0 = 0 \end{aligned}$$

- OCPs based on **index-1 DAEs** are the most common
- The selected initial condition  $\bar{\mathbf{x}}_0$  has to be **consistent**

# Formulating an OCP


## Semi-explicit DAE-constrained OCP

$$\begin{aligned} \min_{\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)} \quad & \phi(\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t.} \quad & \dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \\ & 0 = \mathbf{G}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \\ & \mathbf{h}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \leq 0 \\ & \mathbf{x}(0) - \bar{\mathbf{x}}_0 = 0 \end{aligned}$$

## Fully implicit DAE-constrained OCP

$$\begin{aligned} \min_{\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)} \quad & \phi(\mathbf{x}(\cdot), \mathbf{z}(\cdot), \mathbf{u}(\cdot)) \\ \text{s.t.} \quad & \mathbf{F}(\dot{\mathbf{x}}(t), \mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) = 0 \\ & \mathbf{h}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) \leq 0 \\ & \mathbf{x}(0) - \bar{\mathbf{x}}_0 = 0 \end{aligned}$$

- OCPs based on **index-1 DAEs** are the most common
- The selected initial condition  $\bar{\mathbf{x}}_0$  has to be **consistent**
- If “exotic” boundary conditions are needed (e.g. point-to-point or periodic motion), then they ought to be imposed very carefully<sup>†</sup> !!



<sup>†</sup> *Numerical Periodic Optimal Control in the Presence of Invariants*, Trans. on Automatic Control, S. Gros, M. Zanon



# Multiple-Shooting for DAE-constrained OCPs



Integrator for index-1 DAE:

$$F(\dot{x}(t), z(t), x(t), u(t)) = 0$$

Provides the function:

$$f(x_k, u_k)$$

delivering the integration of the DAE over a time interval  $[t_k, t_{k+1}]$ .



Integrations on the time intervals  $[t_k, t_{k+1}]$

# Multiple-Shooting for DAE-constrained OCPs

**Integrator for index-1 DAE:**

$$\mathbf{F}(\dot{\mathbf{x}}(t), \mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) = 0$$

**Provides** the function:



$$\mathbf{f}(\mathbf{x}_k, \mathbf{u}_k)$$

delivering the integration of the DAE over a time interval  $[t_k, t_{k+1}]$ .

**E.g. semi-explicit DAE:**

$$\dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t))$$

$$0 = \mathbf{G}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t))$$



**Integrations on the time intervals  $[t_k, t_{k+1}]$**

# Multiple-Shooting for DAE-constrained OCPs


**Integrator for index-1 DAE:**

$$\mathbf{F}(\dot{\mathbf{x}}(t), \mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) = 0$$

**Provides** the function:

$$\mathbf{f}(\mathbf{x}_k, \mathbf{u}_k)$$

delivering the integration of the DAE over a time interval  $[t_k, t_{k+1}]$ .



**E.g. semi-explicit DAE:**

$$\dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t))$$


$$0 = \mathbf{G}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t))$$

**with one-step implicit Euler:**

- Solve for  $\mathbf{x}_{k+1}, \mathbf{z}_{k+1}$ :

$$\mathbf{x}_{k+1} = \mathbf{x}_k + h\mathbf{F}(\mathbf{x}_{k+1}, \mathbf{z}_{k+1}, \mathbf{u}_k)$$

$$0 = \mathbf{G}(\mathbf{x}_{k+1}, \mathbf{z}_{k+1}, \mathbf{u}_k)$$



**Integrations on the time intervals  $[t_k, t_{k+1}]$**

# Multiple-Shooting for DAE-constrained OCPs


**Integrator for index-1 DAE:**

$$\mathbf{F}(\dot{\mathbf{x}}(t), \mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) = 0$$

**Provides** the function:

$$\mathbf{f}(\mathbf{x}_k, \mathbf{u}_k)$$

delivering the integration of the DAE over a time interval  $[t_k, t_{k+1}]$ .



**E.g. semi-explicit DAE:**

$$\dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t))$$

$$0 = \mathbf{G}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t))$$


**with one-step implicit Euler:**

- Solve for  $\mathbf{x}_{k+1}, \mathbf{z}_{k+1}$ :

$$\mathbf{x}_{k+1} = \mathbf{x}_k + h\mathbf{F}(\mathbf{x}_{k+1}, \mathbf{z}_{k+1}, \mathbf{u}_k)$$

$$0 = \mathbf{G}(\mathbf{x}_{k+1}, \mathbf{z}_{k+1}, \mathbf{u}_k)$$

- Return  $\mathbf{f}(\mathbf{x}_k, \mathbf{u}_k) \equiv \mathbf{x}_+$



**Integrations on the time intervals  $[t_k, t_{k+1}]$**

# Multiple-Shooting for DAE-constrained OCPs

**Integrator for index-1 DAE:**


$$\mathbf{F}(\dot{\mathbf{x}}(t), \mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) = 0$$

**Provides** the function:

$$\mathbf{f}(\mathbf{x}_k, \mathbf{u}_k)$$

delivering the integration of the DAE over a time interval  $[t_k, t_{k+1}]$ .

Note that the integrator "eliminates" the algebraic variables  $\mathbf{z}(\cdot)$  by treating them "internally". We have "hidden" the complexity !



**E.g. semi-explicit DAE:**

$$\dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t))$$

$$0 = \mathbf{G}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t))$$


**with one-step implicit Euler:**

- Solve for  $\mathbf{x}_{k+1}, \mathbf{z}_{k+1}$ :

$$\mathbf{x}_{k+1} = \mathbf{x}_k + h\mathbf{F}(\mathbf{x}_{k+1}, \mathbf{z}_{k+1}, \mathbf{u}_k)$$

$$0 = \mathbf{G}(\mathbf{x}_{k+1}, \mathbf{z}_{k+1}, \mathbf{u}_k)$$

- Return  $\mathbf{f}(\mathbf{x}_k, \mathbf{u}_k) \equiv \mathbf{x}_{k+1}$



**Integrations on the time intervals  $[t_k, t_{k+1}]$**

# Multiple-Shooting for DAE-constrained OCPs

Integrator for index-1 DAE:


$$\mathbf{F}(\dot{\mathbf{x}}(t), \mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t)) = 0$$

Provides the function:

$$\mathbf{f}(\mathbf{x}_k, \mathbf{u}_k)$$

delivering the integration of the DAE over a time interval  $[t_k, t_{k+1}]$ .

Note that the integrator "eliminates" the algebraic variables  $\mathbf{z}(\cdot)$  by treating them "internally". We have "hidden" the complexity !  
**Is that all ??**



E.g. semi-explicit DAE:

$$\dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t))$$

$$0 = \mathbf{G}(\mathbf{z}(t), \mathbf{x}(t), \mathbf{u}(t))$$


with one-step implicit Euler:

- Solve for  $\mathbf{x}_{k+1}, \mathbf{z}_{k+1}$ :

$$\mathbf{x}_{k+1} = \mathbf{x}_k + h\mathbf{F}(\mathbf{x}_{k+1}, \mathbf{z}_{k+1}, \mathbf{u}_k)$$

$$0 = \mathbf{G}(\mathbf{x}_{k+1}, \mathbf{z}_{k+1}, \mathbf{u}_k)$$





- Return  $\mathbf{f}(\mathbf{x}_k, \mathbf{u}_k) \equiv \mathbf{x}_{k+1}$



Integrations on the time intervals  $[t_k, t_{k+1}]$





# Algebraic variables & discrete inputs

## 3D pendulum with discretized inputs: (force on the mass)



# Algebraic variables & discrete inputs

## 3D pendulum with discretized inputs: (force on the **mass**)







Index-1 DAE:

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$




# Algebraic variables & discrete inputs

## 3D pendulum with discretized inputs: (force on the mass)







### Index-1 DAE:

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$



# Algebraic variables & discrete inputs


## 3D pendulum with discretized inputs: (force on the mass)



### Index-1 DAE:

$$\begin{bmatrix} ml & \mathbf{p} \\ \mathbf{p}^\top & 0 \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{p}} \\ z \end{bmatrix} = \begin{bmatrix} \mathbf{u} - m\mathbf{g}\mathbf{e}_3 \\ -\dot{\mathbf{p}}^\top \dot{\mathbf{p}} \end{bmatrix}$$


When using a discontinuous input parametrization, the algebraic variables **can** also be discontinuous !!



# Direct Collocation for DAE-constrained problems

On each interval  $[t_k, t_{k+1}]$  with:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$



# Direct Collocation for DAE-constrained problems


On each interval  $[t_k, t_{k+1}]$  with:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

Integration is approximated using:

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \underbrace{\boldsymbol{\theta}_{k,i}}_{\text{parameters}} \cdot \underbrace{P_{k,i}(t)}_{\text{polynomials}}$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \underbrace{\mathbf{z}_{k,i}}_{\text{parameters}} \cdot \underbrace{P_{k,i}(t)}_{\text{polynomials}}$$



# Direct Collocation for DAE-constrained problems


On each interval  $[t_k, t_{k+1}]$  with:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

Integration is approximated using:

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \underbrace{\boldsymbol{\theta}_{k,i}}_{\text{parameters}} \cdot \underbrace{P_{k,i}(t)}_{\text{polynomials}}$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \underbrace{\mathbf{z}_{k,i}}_{\text{parameters}} \cdot \underbrace{P_{k,i}(t)}_{\text{polynomials}}$$



Note:

- $\mathbf{x}(\boldsymbol{\theta}_k, t_{k,i}) = \boldsymbol{\theta}_{k,i}$
- $\mathbf{z}(\mathbf{z}_k, t_{k,i}) = \mathbf{z}_{k,i}$
- $K + 1$  d.o.f. per differential state
- $K$  d.o.f. per algebraic state

# Direct Collocation for DAE-constrained problems

On each interval  $[t_k, t_{k+1}]$  with:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$



Integration is approximated using:

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \underbrace{\boldsymbol{\theta}_{k,i}}_{\text{parameters}} \cdot \underbrace{P_{k,i}(t)}_{\text{polynomials}}$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \underbrace{\mathbf{z}_{k,i}}_{\text{parameters}} \cdot \underbrace{P_{k,i}(t)}_{\text{polynomials}}$$

Note:

- $\mathbf{x}(\boldsymbol{\theta}_k, t_{k,i}) = \boldsymbol{\theta}_{k,i}$
- $\mathbf{z}(\mathbf{z}_k, t_{k,i}) = \mathbf{z}_{k,i}$
- $K + 1$  d.o.f. per differential state
- $K$  d.o.f. per algebraic state



# Direct Collocation for DAE-constrained problems

On each interval  $[t_k, t_{k+1}]$  with:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$



Integration is approximated using:

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \underbrace{\boldsymbol{\theta}_{k,i}}_{\text{parameters}} \cdot \underbrace{P_{k,i}(t)}_{\text{polynomials}}$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \underbrace{\mathbf{z}_{k,i}}_{\text{parameters}} \cdot \underbrace{P_{k,i}(t)}_{\text{polynomials}}$$

Note:

- $\mathbf{x}(\boldsymbol{\theta}_k, t_{k,i}) = \boldsymbol{\theta}_{k,i}$
- $\mathbf{z}(\mathbf{z}_k, t_{k,i}) = \mathbf{z}_{k,i}$
- $K + 1$  d.o.f. per differential state
- $K$  d.o.f. per algebraic state





**Why different d.o.f ?** The differential states need an extra degree of freedom (hence  $K + 1$ ) for continuity (i.e. to close the shooting gaps). Algebraic states can be discontinuous and therefore need only  $K$  degrees of freedom !

# Direct Collocation for DAE-constrained problems

## Fully implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$





# Direct Collocation for DAE-constrained problems



## Fully implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

## Interpolation:

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \boldsymbol{\theta}_{k,i} P_{k,i}(t)$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$



# Direct Collocation for DAE-constrained problems

Fully implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

Interpolation:

$$\mathbf{x}(\theta_k, t) = \sum_{i=0}^K \theta_{k,i} P_{k,i}(t)$$



$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$

Collocation uses the constraints:

$$\mathbf{x}(\theta_k, t_{k+1}) - \mathbf{x}(\theta_{k+1}, t_k) = 0 \quad \text{continuity}$$

$$\mathbf{F}\left(\frac{\partial}{\partial t} \mathbf{x}(\theta_k, t_{k,i}), \mathbf{x}(\theta_k, t_{k,i}), \mathbf{z}_{k,i}, \mathbf{u}_k\right) = 0 \quad \text{dynamics}$$

with  $k = 0, \dots, N-1$ , and  $i = 1, \dots, K$ .



# Direct Collocation for DAE-constrained problems

Fully implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

Interpolation:

$$\mathbf{x}(\theta_k, t) = \sum_{i=0}^K \theta_{k,i} P_{k,i}(t)$$



$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$

Collocation uses the constraints:

$$\theta_{k,K} - \theta_{k+1,0} = 0 \quad \text{continuity}$$

$$\mathbf{F}\left(\sum_{j=0}^K \theta_{k,j} \dot{P}_{k,j}(t_{k,i}), \theta_{k,i}, \mathbf{z}_{k,i}, \mathbf{u}_k\right) = 0 \quad \text{dynamics}$$

with  $k = 0, \dots, N-1$ , and  $i = 1, \dots, K$ .



# Direct Collocation for DAE-constrained problems

Fully implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

Interpolation:

$$\mathbf{x}(\theta_k, t) = \sum_{i=0}^K \theta_{k,i} P_{k,i}(t)$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$



Collocation uses the constraints:

$$\theta_{k,K} - \theta_{k+1,0} = 0 \quad \text{continuity}$$

$$\mathbf{F}\left(\sum_{j=0}^K \theta_{k,j} \dot{P}_{k,j}(t_{k,i}), \theta_{k,i}, \mathbf{z}_{k,i}, \mathbf{u}_k\right) = 0 \quad \text{dynamics}$$

with  $k = 0, \dots, N-1$ , and  $i = 1, \dots, K$ .

**Note:** algebraic states appear only in the **dynamics** ( $i = 1, \dots, K$  hence  $K$  equations !!), hence only  $K$  are needed.





# Direct Collocation for DAE-constrained problems

## Semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u}_k)$$

$$0 = \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u}_k)$$



# Direct Collocation for DAE-constrained problems

## Semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u}_k)$$

$$0 = \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u}_k)$$



## Collocation uses the constraints:

$$0 = \mathbf{x}(\boldsymbol{\theta}_k, t_{k+1}) - \mathbf{x}(\boldsymbol{\theta}_{k+1}, t_k) \quad \text{continuity}$$

$$\frac{\partial}{\partial t} \mathbf{x}(\boldsymbol{\theta}_k, t_{k,i}) = \mathbf{F}(\mathbf{x}(\boldsymbol{\theta}_k, t_{k,i}), \mathbf{z}_{k,i}, \mathbf{u}_k) \quad \text{dynamics}$$

$$0 = \mathbf{G}(\mathbf{x}(\boldsymbol{\theta}_k, t_{k,i}), \mathbf{z}_{k,i}, \mathbf{u}_k) \quad \text{algebraic}$$

with  $k = 0, \dots, N-1$ , and  $i = 1, \dots, K$ .



# Direct Collocation for DAE-constrained problems

## Semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x}, \mathbf{z}, \mathbf{u}_k)$$

$$0 = \mathbf{G}(\mathbf{x}, \mathbf{z}, \mathbf{u}_k)$$



Collocation uses the constraints:

$$0 = \boldsymbol{\theta}_{k,K} - \boldsymbol{\theta}_{k+1,0} \quad \text{continuity}$$

$$\frac{\partial}{\partial t} \mathbf{x}(\boldsymbol{\theta}_k, t_{k,i}) = \mathbf{F}(\boldsymbol{\theta}_{k,i}, \mathbf{z}_{k,i}, \mathbf{u}_k) \quad \text{dynamics}$$

$$0 = \mathbf{G}(\boldsymbol{\theta}_{k,i}, \mathbf{z}_{k,i}, \mathbf{u}_k) \quad \text{algebraic}$$

with  $k = 0, \dots, N-1$ , and  $i = 1, \dots, K$ .



## Selection of the collocation points for DAEs

**What collocation scheme to use for DAEs ?!?**



## Selection of the collocation points for DAEs

What collocation scheme to use for DAEs ?!?


K	Legendre	Radau
1	0.5	1.0
2	0.211325 0.788675	0.333333 1.000000
3	0.112702 0.500000 0.887298	0.155051 0.644949 1.000000
4	0.069432 0.330009 0.669991 0.930568	0.088588 0.409467 0.787659 1.000000
5	0.046910 0.230765 0.500000 0.769235 0.953090	0.057104 0.276843 0.583590 0.860240 1.000000

c.f. Lecture "Direct Collocation"


# Selection of the collocation points for DAEs

What collocation scheme to use for DAEs ?!

E.g. Legendre,  $K = 3$




E.g. Radau,  $K = 3$



# Selection of the collocation points for DAEs


What collocation scheme to use for DAEs ?!

E.g. Legendre,  $K = 3$



- has a collocation point at  $t_k$  all others inside  $[t_k, t_{k+1}]$

E.g. Radau,  $K = 3$




- has collocation points at  $t_k$  and  $t_{k+1}$

# Selection of the collocation points for DAEs


What collocation scheme to use for DAEs ?!

E.g. Legendre,  $K = 3$



- has a collocation point at  $t_k$  all others inside  $[t_k, t_{k+1}]$
- integration order  $2K = 6$

E.g. Radau,  $K = 3$




- has collocation points at  $t_k$  and  $t_{k+1}$
- integration order  $2K - 1 = 5$

# Selection of the collocation points for DAEs


What collocation scheme to use for DAEs !?

E.g. Legendre,  $K = 3$



- has a collocation point at  $t_k$  all others inside  $[t_k, t_{k+1}]$
- integration order  $2K = 6$
- has **A-stability** (stable for eigenvalues  $\rightarrow -\infty$ )

E.g. Radau,  $K = 3$




- has collocation points at  $t_k$  and  $t_{k+1}$
- integration order  $2K - 1 = 5$
- has **L-stability** (stable for eigenvalues at  $-\infty$ )

# Selection of the collocation points for DAEs


What collocation scheme to use for DAEs !?

E.g. Legendre,  $K = 3$



- has a collocation point at  $t_k$  all others inside  $[t_k, t_{k+1}]$
- integration order  $2K = 6$
- has **A-stability** (stable for eigenvalues  $\rightarrow -\infty$ )
- best suited for stiff ODEs

E.g. Radau,  $K = 3$




- has collocation points at  $t_k$  and  $t_{k+1}$
- integration order  $2K - 1 = 5$
- has **L-stability** (stable for eigenvalues at  $-\infty$ )
- best suited for DAEs

# Selection of the collocation points for DAEs


What collocation scheme to use for DAEs !?

E.g. Legendre,  $K = 3$



- has a collocation point at  $t_k$  all others inside  $[t_k, t_{k+1}]$
- integration order  $2K = 6$
- has **A-stability** (stable for eigenvalues  $\rightarrow -\infty$ )
- best suited for stiff ODEs

E.g. Radau,  $K = 3$



- has collocation points at  $t_k$  and  $t_{k+1}$
- integration order  $2K - 1 = 5$
- has **L-stability** (stable for eigenvalues at  $-\infty$ )
- best suited for DAEs

**Careful:** using a very high-order collocation setup *can* deteriorate the conditioning of your KKT matrices and hinder the linear algebra underlying the NLP solver !!

# NLP from Direct Collocation for DAE-constrained OCPs

**Fully implicit DAE:**

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$



# NLP from Direct Collocation for DAE-constrained OCPs



Fully implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

Interpolation:

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \boldsymbol{\theta}_{k,i} P_{k,i}(t)$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$



# NLP from Direct Collocation for DAE-constrained OCPs

**Fully implicit DAE:**

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

**NLP with direct collocation**



$$\min_{\mathbf{w}} \Phi(\mathbf{w})$$

**Interpolation:**

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \boldsymbol{\theta}_{k,i} P_{k,i}(t)$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$

$$\text{s.t. } \mathbf{g}(\mathbf{w}) =$$



# NLP from Direct Collocation for DAE-constrained OCPs

**Fully implicit DAE:**

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

**NLP** with direct collocation

$$\min_{\mathbf{w}} \Phi(\mathbf{w})$$



**Interpolation:**

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \boldsymbol{\theta}_{k,i} P_{k,i}(t)$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$

$$\text{s.t. } \mathbf{g}(\mathbf{w}) = \begin{bmatrix} \boldsymbol{\theta}_{0,0} - \bar{\mathbf{x}}_0 \end{bmatrix}$$

Initial conditions  $\bar{\mathbf{x}}_0$



# NLP from Direct Collocation for DAE-constrained OCPs

**Fully implicit DAE:**

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

**NLP** with direct collocation

$$\min_{\mathbf{w}} \Phi(\mathbf{w})$$



**Interpolation:**

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \boldsymbol{\theta}_{k,i} P_{k,i}(t)$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$

$$\text{s.t. } \mathbf{g}(\mathbf{w}) = \begin{bmatrix} \boldsymbol{\theta}_{0,0} - \bar{\mathbf{x}}_0 \\ \boldsymbol{\theta}_{0,K} - \boldsymbol{\theta}_{1,0} \end{bmatrix}$$

Continuity constraints ( $\equiv$  shooting gaps)



# NLP from Direct Collocation for DAE-constrained OCPs

**Fully implicit DAE:**

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

**NLP** with direct collocation

$$\min_{\mathbf{w}} \Phi(\mathbf{w})$$



**Interpolation:**

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \boldsymbol{\theta}_{k,i} P_{k,i}(t)$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$

$$\text{s.t. } \mathbf{g}(\mathbf{w}) = \begin{bmatrix} \boldsymbol{\theta}_{0,0} - \bar{\mathbf{x}}_0 \\ \boldsymbol{\theta}_{0,K} - \boldsymbol{\theta}_{1,0} \\ \mathbf{F}\left(\frac{\partial}{\partial t} \mathbf{x}(\boldsymbol{\theta}_k, t_{k,0}), \boldsymbol{\theta}_{k,0}, \mathbf{z}_{k,0}, \mathbf{u}_k\right) \end{bmatrix}$$

Integration constraints for  $k = 0$



# NLP from Direct Collocation for DAE-constrained OCPs

Fully implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

NLP with direct collocation

$$\min_{\mathbf{w}} \Phi(\mathbf{w})$$



Interpolation:

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \boldsymbol{\theta}_{k,i} P_{k,i}(t)$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$

$$\text{s.t. } \mathbf{g}(\mathbf{w}) = \begin{bmatrix} \boldsymbol{\theta}_{0,0} - \bar{\mathbf{x}}_0 \\ \boldsymbol{\theta}_{0,K} - \boldsymbol{\theta}_{1,0} \\ \mathbf{F}\left(\frac{\partial}{\partial t} \mathbf{x}(\boldsymbol{\theta}_k, t_{k,0}), \boldsymbol{\theta}_{k,0}, \mathbf{z}_{k,0}, \mathbf{u}_k\right) \\ \dots \\ \boldsymbol{\theta}_{k,K} - \boldsymbol{\theta}_{k+1,0} \\ \mathbf{F}\left(\frac{\partial}{\partial t} \mathbf{x}(\boldsymbol{\theta}_k, t_{k,K}), \boldsymbol{\theta}_{k,i}, \mathbf{z}_{k,K}, \mathbf{u}_k\right) \\ \dots \end{bmatrix}$$

Remaining integration constraints  $k = 1, \dots, N-1$



# NLP from Direct Collocation for DAE-constrained OCPs

Fully implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

NLP with direct collocation

$$\min_{\mathbf{w}} \Phi(\mathbf{w})$$

Interpolation:



$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \boldsymbol{\theta}_{k,i} P_{k,i}(t)$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$

$$\text{s.t. } \mathbf{g}(\mathbf{w}) = \begin{bmatrix} \boldsymbol{\theta}_{0,0} - \bar{\mathbf{x}}_0 \\ \boldsymbol{\theta}_{0,K} - \boldsymbol{\theta}_{1,0} \\ \mathbf{F}\left(\frac{\partial}{\partial t} \mathbf{x}(\boldsymbol{\theta}_k, t_{k,0}), \boldsymbol{\theta}_{k,0}, \mathbf{z}_{k,0}, \mathbf{u}_k\right) \\ \dots \\ \boldsymbol{\theta}_{k,K} - \boldsymbol{\theta}_{k+1,0} \\ \mathbf{F}\left(\frac{\partial}{\partial t} \mathbf{x}(\boldsymbol{\theta}_k, t_{k,K}), \boldsymbol{\theta}_{k,i}, \mathbf{z}_{k,K}, \mathbf{u}_k\right) \\ \dots \end{bmatrix}$$

Decision variables ( $k = 0, \dots, N-1$ )

$$\mathbf{w} = \{\dots, \boldsymbol{\theta}_{k,0}, \boldsymbol{\theta}_{k,1}, \mathbf{z}_{k,1}, \dots, \boldsymbol{\theta}_{k,K}, \mathbf{z}_{k,K}, \mathbf{u}_k, \dots\}$$



# NLP from Direct Collocation for DAE-constrained OCPs

**Fully implicit DAE:**

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}_k) = 0$$

**Interpolation:**

$$\mathbf{x}(\boldsymbol{\theta}_k, t) = \sum_{i=0}^K \boldsymbol{\theta}_{k,i} P_{k,i}(t)$$

$$\mathbf{z}(\mathbf{z}_k, t) = \sum_{i=1}^K \mathbf{z}_{k,i} P_{k,i}(t)$$

**Note: for  $\mathbf{z}$ , the interpolation plays no role in the collocation equations !**



**NLP** with direct collocation

$$\min_{\mathbf{w}} \Phi(\mathbf{w})$$

$$\text{s.t. } \mathbf{g}(\mathbf{w}) = \begin{bmatrix} \boldsymbol{\theta}_{0,0} - \bar{\mathbf{x}}_0 \\ \boldsymbol{\theta}_{0,K} - \boldsymbol{\theta}_{1,0} \\ \mathbf{F}\left(\frac{\partial}{\partial t} \mathbf{x}(\boldsymbol{\theta}_k, t_{k,0}), \boldsymbol{\theta}_{k,0}, \mathbf{z}_{k,0}, \mathbf{u}_k\right) \\ \dots \\ \boldsymbol{\theta}_{k,K} - \boldsymbol{\theta}_{k+1,0} \\ \mathbf{F}\left(\frac{\partial}{\partial t} \mathbf{x}(\boldsymbol{\theta}_k, t_{k,K}), \boldsymbol{\theta}_{k,i}, \mathbf{z}_{k,K}, \mathbf{u}_k\right) \\ \dots \end{bmatrix}$$

Decision variables ( $k = 0, \dots, N-1$ )

$$\mathbf{w} = \{\dots, \boldsymbol{\theta}_{k,0}, \boldsymbol{\theta}_{k,1}, \mathbf{z}_{k,1}, \dots, \boldsymbol{\theta}_{k,K}, \mathbf{z}_{k,K}, \mathbf{u}_k, \dots\}$$






# Direct Methods for DAE-based OCPs - Wrap up



## Multiple-shooting

- Hides the algebraic variables  $z$  in the integrator
- If they are needed in the constraints/cost, the integrator needs to report them back to the NLP solver, with sensitivities.



## Direct Collocation:

- Collocation equations are *almost* the same as for ODEs
- A discrete instance of the algebraic variables exists at every collocation time but the first one (associated to the continuity conditions)
- Use the Radau collocation times
- Careful about very high orders in the collocation polynomial !





## What about an “Optimal control discussion group” at NTNU?

- Regular meetings (e.g. monthly) where people can share questions / results related to optimal control in their research
- Act as a support group to
  - ▶ Unlock difficulties in research
  - ▶ Promote “good practices” in optimal control
  - ▶ Make sure that NTNU use state-of-the-art approaches in its research
- Act as a platform to communicate ideas