

# Specialization project

Jonas Bueie

2021

## Contents

<b>1</b>	<b>Theory</b>	<b>2</b>
1.1	Theoretical description of hard-sphere fluids . . . . .	2
1.2	Simulation hard-sphere fluids . . . . .	2
	<b>References</b>	<b>4</b>

# 1 Theory

## 1.1 Theoretical description of hard-sphere fluids

Theoretical models of hard-sphere fluids are important descriptions of fluids. Of particular interest for this project is the Enskog equation for the viscosity of fluids, and its generalization to hard-sphere fluid mixtures.

## 1.2 Simulation hard-sphere fluids

Several methods exist for simulating fluids consisting of hard spheres. While Monte Carlo methods are relatively simple to use with hard sphere potentials – that is, spherical particles (molecules) via non-continuous interaction potential of the form

$$u = \begin{cases} \infty, & r < \sigma \\ 0, & r > \sigma, \end{cases} \quad (1)$$

molecular dynamics methods allow calculating dynamical and out-of-equilibrium properties of a system (Allen et al. 1989).

Furthermore, Jover et. al (Jover et al. 2012) has shown that a pseudo hard Mie (or generalized Lennard-Jones) potential

$$u_{\text{Mie}}(r) = \frac{\lambda_r}{\lambda_r - \lambda_a} \left( \frac{\lambda_r}{\lambda_a} \right)^{\frac{\lambda_a}{\lambda_r - \lambda_a}} \epsilon \left[ \left( \frac{\sigma}{r} \right)^{\lambda_r} - \left( \frac{\sigma}{r} \right)^{\lambda_a} \right], \quad (2)$$

can approximate a hard-sphere interaction potential well, when the repulsive part of the Mie potential is isolated, by shifting it upwards by its minimal value (the well depth  $\epsilon$ ), and cutting it off here, setting the potential to zero once it has reached its minimum. This gives a steep non-negative potential of the form

$$u_{(\lambda_a, \lambda_b)}(r) = \begin{cases} \frac{\lambda_r}{\lambda_r - \lambda_a} \left( \frac{\lambda_r}{\lambda_a} \right)^{\frac{\lambda_a}{\lambda_r - \lambda_a}} \epsilon \left[ \left( \frac{\sigma}{r} \right)^{\lambda_r} - \left( \frac{\sigma}{r} \right)^{\lambda_a} \right] + \epsilon, & r < \sigma \left( \frac{\lambda_r}{\lambda_a} \right)^{\frac{1}{\lambda_r - \lambda_a}} \\ 0, & r > \sigma \left( \frac{\lambda_r}{\lambda_a} \right)^{\frac{1}{\lambda_r - \lambda_a}}, \end{cases} \quad (3)$$

closely resembling that of an infinitely steep hard wall potential. Jover et. al chose the exponents  $(\lambda_r, \lambda_a) = (50, 49)$  as a compromise between the faithfulness of the pseudo hard representation and computational speed: Higher exponents will give steeper potentials, but at the expense of making simulations more time-consuming, because shorter time steps are needed. For clarity, the Mie (50, 49) potential has the form

$$u_{(50,49)}(r) = \begin{cases} 50 \left( \frac{50}{49} \right)^{49} \epsilon \left[ \left( \frac{\sigma}{r} \right)^{50} - \left( \frac{\sigma}{r} \right)^{49} \right] + \epsilon, & r < \frac{50}{49} \sigma \\ 0, & r > \frac{50}{49} \sigma. \end{cases} \quad (4)$$

Pousaneh and de Wijn (Pousaneh et al. 2020) have shown that such a pseudo-hard wall potential can be used to model viscosity for a one-component, and that the obtained viscosity is in agreement with Enskog theory.

## References

- Allen, M.P., D. Frenkel, and J. Talbot (1989). “Molecular dynamics simulation using hard particles”. In: *Computer Physics Reports* 9.6, pp. 301–353. ISSN: 0167-7977. DOI: [https://doi.org/10.1016/0167-7977\(89\)90009-9](https://doi.org/10.1016/0167-7977(89)90009-9). URL: <https://www.sciencedirect.com/science/article/pii/0167797789900099>.
- Jover, J. et al. (2012). “Pseudo hard-sphere potential for use in continuous molecular-dynamics simulation of spherical and chain molecules”. In: *The Journal of Chemical Physics* 137.14, p. 144505. DOI: 10.1063/1.4754275. eprint: <https://doi.org/10.1063/1.4754275jkk>. URL: <https://doi.org/10.1063/1.4754275>.
- Pousaneh, Faezeh and Astrid S. de Wijn (2020). “Shear viscosity of pseudo hard-spheres”. In: *Molecular Physics* 118.4, p. 1622050. DOI: 10.1080/00268976.2019.1622050. eprint: <https://doi.org/10.1080/00268976.2019.1622050>. URL: <https://doi.org/10.1080/00268976.2019.1622050>.