

COMMENTS CONCERNING THE USE OF
URANIUM-BEARING MINERALS
AND CHLORINE-36 TO RECONSTRUCT WATER
MOVEMENT AT YUCCA MOUNTAIN

by

Stanley N. Davis
Professor of Hydrology Emeritus
Department of Hydrology and Water Resources
University of Arizona
Tucson, Arizona 85721

Presentation to the United States
Nuclear Waste Technical Review Board

Denver, Colorado
July, 1996

①

OUTLINE

Introduction

Material to be covered

Water is not "dated"

History of nuclides

Use of Uranium

Disequilibrium

Distance of movement not determined

Use of Chlorine-36

Origin and distribution

Importance of bromide

Possible sources of anomalies

In situ production

Dissolution of surface rocks




Effects of microhydrology

Conclusions

③

PRODUCTION OF CHLORINE-36

(half-life = 301,000 yrs)



Atmospheric origin: reaction 1

Land surface: reactions 2, 3, & 4

Deep subsurface: reaction 2

④





Figure 3. Estimated $(\text{Cl-36}/\text{Cl}) \times 10\text{E-15}$ ratios in prebomb ground water based on available measurements of soil and ground-water samples.


Key:
 ○ Well water samples
 ● Soil samples

②

ACTIVITY RATIOS



NORTHEASTERN ARIZONA Chloride vs chlorine-36 USGS Data



⑦ SOME $^{38}\text{Cl}/\text{total Cl}$ RATIOS

Bomb fallout

Yucca Mountain, NV	$3,821 \times 10^{-15}$
West Texas	$6,560 \times 10^{-15}$
New Mexico	$7,040 \times 10^{-15}$
New Mexico	$9,700 \times 10^{-15}$
New Mexico	$6,000 \times 10^{-15}$

Atmospheric constituents

Stratospheric HCl	$3,260 \times 10^{-15}$
-------------------	-------------------------


Uranium ore

Koongarra, Australia	$128,500 \times 10^{-15}$
Cigar Lake, Canada	$47,800 \times 10^{-15}$


Surface rocks

Meteor Crater	$1,400 \times 10^{-15}$
Boulder Bishop Creek	$8,500 \times 10^{-15}$

⑥ Median values of Cl⁻/Br⁻ ratios related to the Cl⁻ concentration in ground water.



⑧ Number



Pluralites non est ponenda sine necessitate.

"Multiplicity ought not to be posited without necessity."

— William Ockham (1280-1349)

10.

POSSIBLE SOURCES OF CHLORINE-36 ANOMALIES

1. Testing nuclear explosives
2. Fluctuations of cosmic-ray production
3. In situ natural production
4. Dissolution of surface rocks
5. Variations in total chloride deposition
6. Variations of ^{36}Cl in troposphere related to annual recharge
7. Atomic reactor sources
8. Contamination of sample (Lucas Heights)
9. Analytical problems
10. Gas-phase transport
11. Prehistoric supernovas

11.

CHLORINE-36 FROM DISSOLUTION OF SURFACE ROCKS

Assume

Rock with 100 mg/kg chloride

Chloride with $^{36}\text{Cl}/\text{Cl}$ ratio of 10^{-11}

Available water per $1.0 \text{ cm}^2 = 200 \text{ ml}$

Inwash cancels erosion

TDS from rock dissolution = 100 mg/L

Precipitation has 0.5 mg/L Cl^- and $^{36}\text{Cl}/\text{Cl}$ ratio of 5×10^{-13}

Result

3.39×10^5 atoms $^{36}\text{Cl}/\text{cm}^2 \text{ yr}$ from rock

8.48×10^5 atoms $^{36}\text{Cl}/\text{cm}^2 \text{ yr}$ from rain

Expected average from atmosphere, about 10^5 atoms

$^{36}\text{Cl}/\text{cm}^2 \text{ yr}$

12.

