

**Exercise 0.1** (8.6). Let  $A \in \text{End}(V)$  be nilpotent, and  $U \subset V$  invariant. Show that the quotient map  $\bar{A} \in \text{End}(V/U)$  is nilpotent.

*Proof.* Suppose  $A^k = 0$  for some  $k > 0$ . We claim that  $\bar{A}^k = 0$  for the same  $k$ . We recall by lemma 2.16 that  $\bar{A} \in \text{End}(V/U)$  is the unique endomorphism making  $\bar{A} \circ \pi = \pi \circ A$  commute where  $\pi: V \rightarrow V/U$  is the quotient map. It thus immediately follows that  $\bar{A}^k = 0$  since this satisfies the commutative criterion. Now, we claim that suppose that for  $N$  we have shown  $\bar{A}^N \circ \pi = \pi \circ A^N$ . Then we get

$$\pi \circ A^{N+1} = (\pi \circ A) \circ A^N = \bar{A} \circ \pi \circ A^N = \bar{A}^{N+1} \circ \pi$$

so since the case for  $N = 1$  was shown, we get by induction that  $\bar{A}^k \circ \pi = \pi \circ A^k = 0$ . Now,  $\pi$  is surjective by lemma 2.9, so given some  $\bar{x} \in V/U$ , let  $x \in V$  be such that  $\pi(x) = \bar{x}$ . Then  $\bar{A}^k \bar{x} = \bar{A}^k (\pi(x)) = \pi \circ A^k(x) = \pi(0) = \bar{0}$ . So indeed  $\bar{A}^k$  is equal to the zero endomorphism in  $\text{End}(V/U)$ . Thus  $\bar{A}$  is nilpotent. □