

MATH 5301 Elementary Analysis - Homework 3

Jonas Wagner

2021, September 17

Problem 1

Let X denote the universal set. Two subsets A and B are said to have the same cardinality if there is a bijection $f : A \rightarrow B$. Notation: $|A| = |B|$.

1a) Prove that $|A| = |B|$ is an equivalence relation on the power set of X

The relation $\mathcal{R} = “|A| = |B|”$ ¹ is defined as:

$$\mathcal{R} = “|A| = |B|” = \{(A, B) : \exists(f : A \xrightarrow{B} B)\}$$

Equivalence can be demonstrated by proven by demonstrating: (i) Reflexivity, (ii) Symmetry, and (iii) Transitivity

i) Reflexive:

$$ARA = \{(A, A) : \exists(f : A \xrightarrow{B} A)\}$$

Since $f : A \xrightarrow{B} A = a \in A$ is true $\forall A \in 2^X$, \mathcal{R} is Reflexive.

ii) Symmetric:

$$ARB \implies BRA$$
$$\{(A, B) : \exists(f : A \xrightarrow{B} B)\} \implies \{(B, A) : \exists(f : B \xrightarrow{B} A)\}$$

Since $f : A \xrightarrow{B} B \implies g : B \xrightarrow{B} A = f^{-1}$ is true $\forall A, B \in 2^X$, \mathcal{R} is Symmetric.
(Essentially if f is bijective one way, f^{-1} is bijective for the other way)

iii) Transitive:

$$(ARB) \wedge (BRC) \implies ARC$$
$$(\{(A, B) : \exists(f : A \xrightarrow{B} B)\}) \wedge \{(B, C) : \exists(g : B \xrightarrow{B} C)\} \implies \{(A, C) : \exists(h : A \xrightarrow{B} C)\}$$

Since $(f : A \xrightarrow{B} B) \wedge (g : B \xrightarrow{B} C) \implies (h : A \xrightarrow{B} C = A \xrightarrow{f} B \xrightarrow{g} C)$ is true $\forall A, B, C \in 2^X$, \mathcal{R} is Transitive.

Therefore $\mathcal{R} = “|A| = |B|”$ is an equivalence relation over 2^X .

¹using \mathcal{R} for simplicity/reusability

1b) Is it true that if $|A_1| = |B_1|$ and $|A_2| = |B_2|$ then $|A_1 \cup A_2| = |B_1 \cup B_2|$?

$$(|A_1| = |B_1|) \wedge (|A_2| = |B_2|) \implies |A_1 \cup A_2| = |B_1 \cup B_2|$$

$$(A_1 \mathcal{R} B_1) \wedge (A_2 \mathcal{R} B_2) \implies (A_1 \cup A_2) \mathcal{R} (B_1 \cup B_2)$$

$$\{(A_1, B_1) : \exists(f_1 : A_1 \xrightarrow{\mathcal{B}} B_1)\} \wedge \{(A_2, B_2) : \exists(f_2 : A_2 \xrightarrow{\mathcal{B}} B_2)\} \implies \{((A_1 \cup A_2), (B_1 \cup B_2)) : \exists(f : (A_1 \cup A_2) \xrightarrow{\mathcal{B}} (B_1 \cup B_2))\}$$

This itself is false, as in the case when

$$(A_1 \cap A_2 \neq \emptyset) \wedge (B_1 \cap B_2 = \emptyset) \implies (f : (A_1 \cup A_2) \xrightarrow{\mathcal{I}} (B_1 \cup B_2)) \wedge (f^{-1} : (B_1 \cup B_2) \xrightarrow{\mathcal{S}} (A_1 \cup A_2))$$

But, f is not surjective and f^{-1} is not injective, so f cannot be bijective.

Problem 2

Finish the proof of the Cantor-Bernstein theorem: For the sets A and B , such that $|A| \leq |B|$ and $|B| \leq |A|$ define A_∞ as the set of all elements of A having infinite order, A_0 as the set of all elements of A having even order, and A_1 the set of all elements of A having odd order. Similarly for B .

Let

$$A, B : (|A| \leq |B|) \wedge (|B| \leq |A|)$$

Define

$$A_\infty = \{a \in A : \mathcal{O}(a) = \infty\}$$

$$A_0 = \{a \in A : \mathcal{O}(a) \dot{=} 2\}$$

$$A_1 = \{a \in A : (\mathcal{O}(a) + 1) \dot{=} 2\}$$

$$B_\infty = \{b \in B : \mathcal{O}(b) = \infty\}$$

$$B_0 = \{b \in B : \mathcal{O}(b) \dot{=} 2\}$$

$$B_1 = \{b \in B : (\mathcal{O}(b) + 1) \dot{=} 2\}$$

2a) Show that $|A_\infty| = |B_\infty|$.

Since,

$$|A| \leq |B| \implies \exists (f_{AB} : A \xrightarrow{1} B) \therefore f_{AB}(A_\infty) = B_\infty$$

Similarly,

$$|B| \leq |A| \implies \exists (f_{BA} : B \xrightarrow{1} A) \therefore f_{BA}(B_\infty) = A_\infty$$

Since,

$$(\exists f_{AB} : A \xrightarrow{1} B) \wedge (\exists f_{BA} : B \xrightarrow{1} A) \implies (\exists f : A \xrightarrow{B} B)$$

Therefore, since a bijective mapping exists,

$$|A_\infty| = |B_\infty|$$

2b) Construct an injective mapping $A_1 \rightarrow B_0$.

From the definition of A_1 , it is known that A_1 contains all elements whose ancestors (for f and g) are of odd order.

Similarly, from the definition of B_0 it is known that B_0 contains all elements whose ancestors (for f and g) are of even order.

Since all elements of B_0 are of even order, then that implies that according to the mappings f and g ,

$$a \in \{a \in A : \exists b \in B_0 : f(a) = b\} \implies a \in A_1$$

therefore, f is a direct mapping from A_1 to B_0 (which is technically bijective so it is also injective)

2c) Show that this mapping is also surjective.

Since, from the previous argument, f is bijective so it is also surjective.

Problem 3

Set A is called countable if $|A| \leq |\mathbb{N}|$. Prove that the following sets are countable.

3a) Set \mathbb{Z}_+ of all non-negative integer numbers

Define $f_0 : \mathbb{N} \rightarrow \mathbb{Z}_+$,

$$f_0(x) = x - 1$$

Claim $f_0 : \mathbb{N} \xrightarrow{s} \mathbb{Z}_+$ (surjective). This is true because $\forall x \in \mathbb{N} \exists f_0(x) \in \mathbb{Z}_+$.

When (i) $x \in \mathbb{N} = 1$, $f_0(x) = x - 1 = 0$ which is in \mathbb{Z}_+ .

Similarly, when (ii) $x \in \mathbb{N} > 1$, $f_0(x) = x - 1$ which is in \mathbb{Z}_+ .

Since f_0 is in fact surjective, \mathbb{Z}_+ is countable. (i.e)

$$f_0 : \mathbb{N} \xrightarrow{s} \mathbb{Z}_+ \implies |\mathbb{Z}_+| \leq |\mathbb{N}|$$

3b) Set $2\mathbb{N}$ of all even numbers

Define $f : \mathbb{N} \rightarrow 2\mathbb{N}$,

$$f(x) = 2x$$

Claim $f : \mathbb{N} \xrightarrow{s} 2\mathbb{N}$ (surjective). This is true because $\forall x \in \mathbb{N} \exists f(x) \in 2\mathbb{N}$.

When $x \in \mathbb{N}$, $f(x) = 2x$ which is in $2\mathbb{N}$.

Since f is surjective, $2\mathbb{N}$ is countable. (i.e)

$$f : \mathbb{N} \xrightarrow{s} 2\mathbb{N} \implies |2\mathbb{N}| \leq |\mathbb{N}|$$

3c) Set \mathbb{Z}^2 of all ordered pairs of integer numbers

Define $g : \mathbb{N} \rightarrow \mathbb{Z}^2$,

$$g(x) = \begin{cases} (0, 0) & x = 1 \\ (1, 0) & x = 2 \\ (1, 1) & x = 3 \\ (0, 1) & x = 4 \\ (-1, 1) & x = 5 \\ (-1, 0) & x = 6 \\ (-1, -1) & x = 7 \\ (0, -1) & x = 8 \\ (1, -1) & x = 9 \\ (2, -1) & x = 10 \\ (2, 0) & x = 11 \\ \vdots & \vdots \end{cases}$$

Claim $g : \mathbb{N} \xrightarrow{s} \mathbb{Z}^2$ (surjective). This is true because $\forall x \in \mathbb{N} \exists g(x) \in \mathbb{Z}^2$ as is clearly evident in the spiraling mapping described by g mapping to all \mathbb{Z}^2 .

Since g is surjective, \mathbb{Z}^2 is countable. (i.e)

$$g : \mathbb{N} \xrightarrow{s} \mathbb{Z}^2 \implies |\mathbb{Z}^2| \leq |\mathbb{N}|$$

3d) Set \mathbb{Q} of all rational numbers

Define $h : \mathbb{N} \rightarrow \mathbb{Q}$,

$$h(x) = \begin{cases} 0 & x = 1 \\ 1 & x = 2 \\ -1 & x = 3 \\ 2 & x = 4 \\ \frac{3}{2} & x = 5 \\ \frac{1}{2} & x = 6 \\ -\frac{1}{2} & x = 7 \\ -\frac{3}{2} & x = 8 \\ -2 & x = 9 \\ 3 & x = 10 \\ \frac{8}{3} & x = 11 \\ \frac{5}{2} & x = 12 \\ \frac{7}{3} & x = 13 \\ \frac{5}{3} & x = 14 \\ \frac{4}{3} & x = 15 \\ \frac{2}{3} & x = 16 \\ \frac{1}{3} & x = 17 \\ -\frac{1}{3} & x = 18 \\ \vdots & \vdots \end{cases}$$

Claim $h : \mathbb{N} \xrightarrow{s} \mathbb{Q}$ (surjective). This is true because $\forall x \in \mathbb{N} \exists h(x) \in \mathbb{Q}$ as is clearly evident in the spiraling mapping described by h mapping to all \mathbb{Q} .

Since h is surjective, \mathbb{Q} is countable. (i.e)

$$g : \mathbb{N} \xrightarrow{s} \mathbb{Q} \implies |\mathbb{Q}| \leq |\mathbb{N}|$$

3e) Set \mathbb{Q}^2 of all ordered pairs of rational numbers

Define $m : \mathbb{N} \rightarrow \mathbb{Q}^2$,

$$m(x) = \begin{cases} (0, 0) & x = 1 \\ (0, 1) & x = 2 \\ (1, 0) & x = 3 \\ (1, 1) & x = 4 \\ (0, -1) & x = 5 \\ (1, -1) & x = 6 \\ (-1, 0) & x = 7 \\ (-1, 1) & x = 8 \\ (-1, -1) & x = 9 \\ (0, 2) & x = 10 \\ \vdots & \vdots \\ (2, 2) & \vdots \\ (0, \frac{3}{2}) & \vdots \\ \vdots & \vdots \end{cases}$$

Note: $m(x)$ continues to spiral following the pattern in a way that ultimately combines the mappings of f_0 , g , and h .

Claim $m : \mathbb{N} \xrightarrow{s} \mathbb{Q}^2$ (surjective). This is true because $\forall x \in \mathbb{N} \exists m(x) \in \mathbb{Q}^2$ as is clearly evident in the spiraling mapping described by m mapping to all \mathbb{Q}^2 .

Since m is surjective, \mathbb{Q}^2 is countable. (i.e)

$$g : \mathbb{N} \xrightarrow{s} \mathbb{Q}^2 \implies |\mathbb{Q}^2| \leq |\mathbb{N}|$$

Problem 4

Prove that the following sets are countable.

4a) Set $P_5(\mathbb{Z})$ of all polynomials of degree 4 with integer coefficients

4b) Any collection of non-intersecting discs on a plane

4c) Any collection of non-intersecting T-shapes on a plane

Note: T-shape consists of two perpendicular line segments such that one of the segments is attached by one of its endpoints to the center of the other segment. The lengths of these segments can be arbitrary. The orientation of the T-shape can be arbitrary.

4d) Set \mathbb{P} of all prime numbers

4e) Set \mathbb{A} of all algebraic numbers


Note: Algebraic numbers are numbers which are roots of some polynomials with integer coefficients.

Problem 5

Prove that for any infinite set A there exists $B \subset A$, so that $|B| = |\mathbb{N}|$.

Problem 6

Prove that the following sets have the same cardinality.



These sets can all be represented as a set of real number ordered pairs.

These are constructed with arbitrary constants: $x_a, x_b, x_c, x_d, x_e, x_f, y_a, y_b, y_c, y_d, y_e, y_f, x_s, y_s, r_s, m_l, b_l$

$$\begin{aligned}
 AB &= \{(x, y) \in \mathbb{R}^2, t \in \mathbb{R} : (x = (1-t)x_a + tx_b) \wedge (y = (1-t)y_a + ty_b) \wedge (0 \leq t \leq 1)\} \\
 CD &= \{(x, y) \in \mathbb{R}^2, t \in \mathbb{R} : (x = (1-t)x_c + tx_d) \wedge (y = (1-t)y_c + ty_d) \wedge (0 \leq t < 1)\} \\
 EF &= \{(x, y) \in \mathbb{R}^2, t \in \mathbb{R} : (x = (1-t)x_e + tx_f) \wedge (y = (1-t)y_e + ty_f) \wedge (0 < t < 1)\} \\
 \mathbb{S} &= \{(x, y) \in \mathbb{R}^2, t \in \mathbb{R} : (x = r_s \cos(2\pi t) + x_s) \wedge (y = \sin(2\pi t) + y_s) \wedge (0 \leq t < 1)\} \\
 l &= \{(x, y) \in \mathbb{R}^2, t \in \mathbb{R} : (x = t) \wedge (y = m_l t + b_l)\}
 \end{aligned}$$

It is clear that each set is defined parametrically with bijective equations mapping the parameter t into a 2-D coordinates (x, y) , thus proving that the cardinality of each set parameter is sufficient to showing the each set has the same cardinality.

Let $T_i = \{t \in A_i\}$ for each of the sets A_i , then (from the reasoning above) the following can be said:

$$\begin{aligned}
 T_{AB} &= \{t \in \mathbb{R} : 0 \leq t \leq 1\}, & |T_{AB}| &= |AB| \\
 T_{CD} &= \{t \in \mathbb{R} : 0 \leq t < 1\}, & |T_{CD}| &= |CD| \\
 T_{EF} &= \{t \in \mathbb{R} : 0 < t < 1\}, & |T_{EF}| &= |EF| \\
 T_{\mathbb{S}} &= \{t \in \mathbb{R} : 0 \leq t < 1\}, & |T_{\mathbb{S}}| &= |\mathbb{S}| \\
 T_l &= \{t \in \mathbb{R}\}, & |T_l| &= |l|
 \end{aligned}$$

Clearly, the equivalent definition of T_{CD} and $T_{\mathbb{S}}$ indicates

$$|CD| = |T_{CD}| = |T_{\mathbb{S}}| = |\mathbb{S}|$$

The equivalence of the other sets is more difficult then via definition.

First, the baseline cardinality can be shown to be \aleph_1 as (by definition of T_l)

$$|T_l| = |\mathbb{R}| = \aleph_1$$

Next, the equivalence of T_{EF} and T_l can be shown with the biforjective mapping

$$f_1 : \mathbb{R} \rightarrow T_{EF} = \frac{2\pi \tan^{-1}(x) + 1}{2}$$

Therefore,

$$|EF| = |T_{EF}| = |T_l| = |\mathbb{R}| = \aleph_1$$

Next, due to the nature of infinite sets, the addition of $t = 0$ from T_{EF} to T_{CD} does not affect the overall cardinality of T_{CD} , thus

$$|CD| = |T_{CD}| = |T_{EF}| = |\mathbb{R}| = \aleph_1$$

Similarly, the addition of $t = 1$ from T_{CD} to T_{AB} will stil result in

$$|AB| = |T_{AB}| = |T_{CD}| = |\mathbb{R}| = \aleph_1$$

Ultimently this means that

$$|AB| = |CD| = |EF| = |\mathbb{S}| = |l| = |\mathbb{R}| = \aleph_1$$