



Geometry › Projective Geometry › General Projective Geometry ›

Affine Transformation

An affine transformation is any transformation that preserves collinearity (i.e., all points lying on a line initially still lie on a line after transformation) and ratios of distances (e.g., the midpoint of a line segment remains the midpoint after transformation). In this sense, affine indicates a special class of projective transformations that do not move any objects from the affine space \mathbb{R}^3 to the plane at infinity or conversely. An affine transformation is also called an affinity.

Geometric contraction, expansion, dilation, reflection, rotation, shear, similarity transformations, spiral similarities, and translation are all affine transformations, as are their combinations. In general, an affine transformation is a composition of rotations, translations, dilations, and shears.

While an affine transformation preserves *proportions* on lines, it does not necessarily preserve angles or lengths. Any triangle can be transformed into any other by an affine transformation, so all triangles are affine and, in this sense, affine is a generalization of congruent and similar.

A particular example combining rotation and expansion is the rotation-enlargement transformation

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = s \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} x - x_0 \\ y - y_0 \end{bmatrix} \quad (1)$$

$$= s \begin{bmatrix} \cos \alpha (x - x_0) + \sin \alpha (y - y_0) \\ -\sin \alpha (x - x_0) + \cos \alpha (y - y_0) \end{bmatrix}. \quad (2)$$

Separating the equations,

$$x' = (s \cos \alpha) x + (s \sin \alpha) y - s (x_0 \cos \alpha + y_0 \sin \alpha) \quad (3)$$

$$y' = (-s \sin \alpha) x + (s \cos \alpha) y + s (x_0 \sin \alpha - y_0 \cos \alpha). \quad (4)$$

This can be also written as

$$x' = a x - b y + c \quad (5)$$

$$y' = b x + a y + d, \quad (6)$$

where



About MathWorld | MathWorld Classroom | Contribute | MathWorld Book

The scale factor s is then defined by

13,154 Entries | Last Updated: Thu May 30 2024 | ©1999–2024 Wolfram Research, Inc. | Terms of Use

$$s \equiv \sqrt{a^2 + b^2}, \quad (9)$$

wolfram.com | Wolfram Language | Mathematica | Wolfram Demonstrations | Wolfram for Education

and the rotation angle by

$$\theta = \arctan\left(\frac{b}{a}\right)$$

Created, developed and nurtured by Eric Weisstein at Wolfram Research

An affine transformation of \mathbb{R}^n is a map $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ of the form

$$F(\mathbf{p}) = \mathbf{A}\mathbf{p} + \mathbf{q} \quad (11)$$

for all $\mathbf{p} \in \mathbb{R}^n$, where \mathbf{A} is a linear transformation of \mathbb{R}^n . If $\det(\mathbf{A}) > 0$, the transformation is orientation-preserving; if $\det(\mathbf{A}) < 0$, it is orientation-reversing.

SEE ALSO

Affine, Affine Complex Plane, Affine Equation, Affine Geometry, Affine Group, Affine Hull, Affine Plane, Affine Space, Equiaffinity, Euclidean Motion, Special Affine Transformation

EXPLORE WITH WOLFRAM|ALPHA



3,2 TM rule 2139050



More things to try:

= 3,2 TM rule 2139050

= Catalan number

= minimal polynomial sqrt(2)+sqrt(3)

REFERENCES

Croft, H. T.; Falconer, K. J.; and Guy, R. K. *Unsolved Problems in Geometry*. New York: Springer-Verlag, p. 3, 1991.

Gray, A. *Modern Differential Geometry of Curves and Surfaces with Mathematica, 2nd ed.* Boca Raton, FL: CRC Press, p. 130, 1997.

Zwillinger, D. (Ed.). "Affine Transformations." §4.3.2 in *CRC Standard Mathematical Tables and Formulae*. Boca Raton, FL: CRC Press, pp. 265-266, 1995.

REFERENCED ON WOLFRAM|ALPHA

Affine Transformation

CITE THIS AS:

Weisstein, Eric W. "Affine Transformation." From *MathWorld*--A Wolfram Web Resource. <https://mathworld.wolfram.com/AffineTransformation.html>

SUBJECT CLASSIFICATIONS

