

Here's what I put together for the R code with the data given by Christian.

-Jason

```
#STAT 201
```

```
#Group Final Project
```

```
#Christian Price, Joseph Lyon, Jason Biesinger, Jesse Van Horn, Jon Wilson, Rex Henretta
```

```
#Fall 2015
```

```
#Data
```

```
knocking=read.table(header=TRUE,text="
```

```
Run AirFuelRatio OctaneLevel AmountOfKnock
```

```
1 Lean Booster1 30
```

```
2 Rich Booster2 58
```

```
3 Stoichiometric Regular 82
```

```
4 Lean Regular 75
```

```
5 Stoichiometric Booster1 70
```

```
6 Lean Booster1 30
```

```
7 Lean Booster2 55
```

```
8 Stoichiometric Booster2 75
```

```
9 Rich Booster1 46
```

```
10 Lean Regular 72
```

```
11 Rich Regular 61
```

```
12 Rich Regular 64
```

```
13 Stoichiometric Booster2 76
```

```
14 Lean Booster2 58
```

```
15 Rich Booster1 43
```

```
16 Stoichiometric Regular 85
```

```
17 Stoichiometric Booster1 70
```

```
18 Rich Booster2 57
```

```
")
```

```
#fit the model
```

```
out.knocking=aov(AmountOfKnock~AirFuelRatio+OctaneLevel+AirFuelRatio:OctaneLevel,data=knocking)
```

```
#Inference
```

```
anova(out.knocking)
```

```
#95% Conf Interval
```

```
TukeyHSD(out.knocking)
```

```
#Mean by factor
```

```
by(knocking$AmountOfKnock,knocking$AirFuelRatio,mean)
```

```
by(knocking$AmountOfKnock,knocking$OctaneLevel,mean)
```

```
#Standard Deviation by factor
```

```
by(knocking$AmountOfKnock,knocking$AirFuelRatio,sd)
```

```
by(knocking$AmountOfKnock,knocking$OctaneLevel,sd)
```

```
#Best plot for Air/Fuel Ratio
```

```
main.effect<-c(53.33333,54.83333,76.33333) # from
```

```
by(knocking$AmountOfKnock,knocking$AirFuelRatio,mean)
```

```
se<-sqrt( 993.50 / 2) # Mean Sq Residuals / number of replicates
```

```
mp<-barplot(main.effect,names=c("Lean","Rich","Stoichiometric"),
```

```
col=c("blue","green","black"),
```

```
ylab="Amount of Knock",xlab="Air / Fuel Ratio",
```

```
ylim=c(0,100))
```

```
arrows(mp,main.effect-se,mp,main.effect+se,
```

```
code=3,angle=90,col="gray")
```

```
#Best plot for Octane Level
```

```
main.effect<-c(73.16667,48.16667,63.16667) #from
```

```
by(knocking$AmountOfKnock,knocking$OctaneLevel,mean)
```

```
se<-sqrt( 950.00 / 2) # Mean Sq Residuals / number of replicates
```

```
mp<-barplot(main.effect,names=c("Regular","Booster #1","Booster #2"),
```

```
col=c("red","brown","yellow"),
```

```
ylab="Amount of Knocking",xlab="Octane Level",
```

```
ylim=c(0,100))
```

```
arrows(mp,main.effect-se,mp,main.effect+se,
```

```
code=3,angle=90,col="gray")
```

```
#QQ plot
```

```
qqnorm(resid(out.knocking))
```

```
resid(out.knocking)
```

```
#Interaction
```

```
interaction.plot(knocking$AirFuelRatio,knocking$OctaneLevel,knocking$AmountOfKnock,type='b')
```

```

+ 11 Rich Regular 61
+ 12 Rich Regular 64
+ 13 Stoichiometric Booster2 76
+ 14 Lean Booster2 58
+ 15 Rich Booster1 43
+ 16 Stoichiometric Regular 85
+ 17 Stoichiometric Booster1 70
+ 18 Rich Booster2 57
+ ")
> out.knocking=aov(AmountOfKnock~AirFuelRatio+OctaneLevel+AirFuelRatio:OctaneLe$
> anova(out.knocking)
Analysis of Variance Table

Response: AmountOfKnock
          Df Sum Sq Mean Sq F value    Pr(>F)
AirFuelRatio      2 1987.0   993.50 380.489 2.018e-09 ***
OctaneLevel       2 1900.0   950.00 363.830 2.463e-09 ***
AirFuelRatio:OctaneLevel  4  552.0   138.00  52.851 2.987e-06 ***
Residuals        9    23.5     2.61
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

the p-values are all less than 0.05

```
R File Edit Packages Windows Help
#STAT 201
#Group Final Project
#Christian Price, Joseph Lyon, Jason Biesinger, Jesse Van Horn, Jon Wilson, Rex Henretta
#Fall 2015

#Data
knocking=read.table(header=TRUE,text="
Run AirFuelRatio OctaneLevel AmountOfKnock
1 Lean Booster1 30
2 Rich Booster2 58
3 Stoichiometric Regular 82
4 Lean Regular 75
5 Stoichiometric Booster1 70
6 Lean Booster1 30
7 Lean Booster2 55
8 Stoichiometric Booster2 75
9 Rich Booster1 46
10 Lean Regular 72
11 Rich Regular 61
12 Rich Regular 64
13 Stoichiometric Booster2 76
14 Lean Booster2 58
15 Rich Booster1 43
16 Stoichiometric Regular 85
17 Stoichiometric Booster1 70
18 Rich Booster2 57
")

#fit the model
out.knocking=aov(AmountOfKnock~AirFuelRatio+OctaneLevel+AirFuelRatio:OctaneLevel,data=knocking)

#Inference
anova(out.knocking)

#95% Conf Interval
TukeyHSD(out.knocking)

#Mean by factor
by(knocking$AmountOfKnock, knocking$AirFuelRatio, mean)
by(knocking$AmountOfKnock, knocking$OctaneLevel, mean)
```



```

#95% Conf Interval
TukeyHSD(out.knocking)

#Mean by factor
by(knocking$AmountOfKnock, knocking$AirFuelRatio, mean)
by(knocking$AmountOfKnock, knocking$OctaneLevel, mean)

#Standard Deviation by factor
by(knocking$AmountOfKnock, knocking$AirFuelRatio, sd)
by(knocking$AmountOfKnock, knocking$OctaneLevel, sd)

#Best plot for Air/Fuel Ratio
main.effect<-c(53.33333, 54.83333, 76.33333) # from by(knocking$AmountOfKnock, knocking$AirFuelRatio, mean)
se<-sqrt( 993.50 / 2) # Mean Sq Residuals / number of replicates
mp<-barplot(main.effect, names=c("Lean", "Rich", "Stoichiometric"),
            col=c("blue", "green", "black"),
            ylab="Amount of Knock", xlab="Air / Fuel Ratio",
            ylim=c(0, 100))
arrows(mp, main.effect-se, mp, main.effect+se,
       code=3, angle=90, col="gray")

#Best plot for Octane Level
main.effect<-c(73.16667, 48.16667, 63.16667) # from by(knocking$AmountOfKnock, knocking$OctaneLevel, mean)
se<-sqrt( 950.00 / 2) # Mean Sq Residuals / number of replicates
mp<-barplot(main.effect, names=c("Regular", "Booster #1", "Booster #2"),
            col=c("red", "brown", "yellow"),
            ylab="Amount of Knocking", xlab="Octane Level",
            ylim=c(0, 100))
arrows(mp, main.effect-se, mp, main.effect+se,
       code=3, angle=90, col="gray")

#QQ plot
qqnorm(resid(out.knocking))
resid(out.knocking)

#Interaction
interaction.plot(knocking$AirFuelRatio, knocking$OctaneLevel, knocking$AmountOfKnock, type='b')

```