

# Update: Trilinear reweighting

Jonathon Langford

Imperial College London

IC Hgg  
23 Mar. 2018

# Status

- Successfully generated LO and  $\mathcal{O}(\lambda_3)$  events via madgraph. Can calculate  $C_1$  in bins of kinematic distribution.
- Building up analysis: ZHLeptonic
  - ▶ Use ZHLeptonicTag on LHC MC samples
  - ▶ Dumper configured to output info of reconstructed + gen-level diphotons and leptons
  - ▶ End goal: fit for  $\kappa_\lambda$  using  $C_1$  values determined from madgraph samples. Sensitivity at  $3000 \text{ fb}^{-1}$ ?






diagram 1

HIW=1, QCD=0, QED=4




# Discrepancy at low $p_T(\gamma\gamma)$

- $p_T(\gamma\gamma)$  spectrum of ZHLeptonicTag events (from LHC sample) has deficit at low  $p_T(\gamma\gamma)$  compared to madgraph generated events.




- Gen-level Higgs:  $p_T(H)$  for full VHTOGG LHC sample has same shape as madgraph events. Suggests there is **no significant difference** in intrinsic kinematics.
- Selection removing low  $p_T(H)$  events?
  - ▶  $p_T(\gamma)$
  - ▶ Diphoton MVA: tried fully relaxing constraint and no significant change in event numbers (+2). Preselection?

# Discrepancy at low $p_T(\gamma\gamma)$



# Discrepancy at low $p_T(\gamma\gamma)$



# Why lack of statistics in MC ZHLeptonicTag?

