

# Neural circuits underlying the generation of theta oscillations

Michele Pignatelli\*, Anna Beyeler, Xavier Leinekugel\*

Institut des Maladies Neurodégénératives (IMN), UMR 5293, CNRS and Université Bordeaux 1 & 2, Avenue des Facultés, Bat. B2, Talence, France

## ARTICLE INFO

### Article history:

Available online 22 September 2011

### Keywords:

Theta  
Oscillations  
Hippocampus  
Cholinergic system  
Brainstem  
Limbic system  
Neuronal networks

## ABSTRACT

Theta oscillations represent the neural network configuration underlying active awake behavior and paradoxical sleep. This major EEG pattern has been extensively studied, from physiological to anatomical levels, for more than half a century. Nevertheless the cellular and network mechanisms accountable for the theta generation are still not fully understood. This review synthesizes the current knowledge on the circuitry involved in the generation of theta oscillations, from the hippocampus to extra hippocampal structures such as septal complex, entorhinal cortex and pedunculopontine tegmentum, a main trigger of theta state through direct and indirect projections to the septal complex. We conclude with a short overview of the perspectives offered by technical advances for deciphering more precisely the different neural components underlying the emergence of theta oscillations.

© 2011 Elsevier Ltd. All rights reserved.

## 1. Introduction

The development of multi-electrode recordings allowed to study the temporal relationships between neuronal discharges and EEG patterns. This led to the current view that EEG oscillations serve as a temporal reference for information coding by spike times. Nevertheless, the mechanisms and circuits underlying the generation of EEG oscillations are still a matter of active investigation. In vertebrates, active brain state is associated with high cholinergic activity, cortical desynchronization and hippocampal theta rhythm (Buzsáki et al., 2003; Moruzzi and Magoun, 1949; Vanderwolf, 1969; Zhang et al., 2010). Hippocampal theta rhythm (Buzsáki et al., 2003; Green and Arduini, 1954; Saper et al., 2010; Vanderwolf, 1969) is a well known EEG pattern characterized by coherent oscillations in the 4–10 Hz frequency band observable during active awake behavior and paradoxical sleep (Bland, 1986; Buzsáki, 2002). In addition to the consistent amount of observations reported in rodents, theta oscillations can be detected as well in humans and primates through intracranial EEG (Kahana

et al., 2001), being reported in both cortical network and hippocampus under condition of active navigation (Kahana et al., 1999) or paradoxical sleep (Cantero et al., 2003), the latter characterized by short bursts of 4–7 Hz oscillations. In the animal, hippocampal theta oscillations are detected under urethane anesthesia as well, either spontaneously or after noxious stimulation and accompanied by muscle atonia and whisker twitching as during paradoxical sleep (Clement et al., 2008). Following extensive multisite recordings in behaving and anesthetized rodents, the expression of theta oscillations found its cardinal generator in the hippocampus (Green and Arduini, 1954) from which theta waves propagate through volume conduction to a large part of the brain (Sirota et al., 2008). It is even possible to induce oscillations in the theta frequency band in the isolated rat hippocampal circuit *in vitro* (Bland et al., 1988; Goutagny et al., 2009; Konopacki et al., 1987a,b). The hippocampus seems to be constituted by coupled theta oscillators disposed in series along the septo-temporal axis and giving rise *in vivo* to traveling septo-temporal theta waves (Lubenov and Siapas, 2009). However, these hippocampal oscillators are largely driven by extra hippocampal inputs. In this review, we will describe the interplay between the different structures involved in the generation of theta oscillations in rodents, from the intra-hippocampal currents and main hippocampal inputs to the deep structures that trigger and sustain the theta-associated, active brain state.

## 2. Theta oscillations in the hippocampus

Laminar multisite recording of hippocampal theta oscillations in the rat allowed the investigation of the distribution of the synaptic currents involved in the genesis of the hippocampal theta


**Abbreviations:** Amy, amygdala; DR, dorsal raphe; EC, entorhinal cortex; HDB, horizontal limb of the diagonal band of Broca; HT, hypothalamus; LDT, laterodorsal tegmentum; LEC, lateral entorhinal cortex; MB, mammillary bodies; MEC, medial entorhinal cortex; mPFC, medial prefrontal cortex; MS, medial septum; NI, nucleus incertus; PC, preceruleus region; PH, posterior hypothalamic nucleus; PPT, pedunculopontine tegmentum; RE, reuniens nucleus of the thalamus; RPO, reticularis pontis oralis nucleus; SC, septal complex; SLD, sublateralodorsal nucleus; Sub, subiculum; SUM, supramammillary nucleus; VDB, vertical limb of the diagonal band of Broca; VTn, ventral tegmental nucleus of Gudden; Ach, acetylcholine; 5HT, serotonin;  $I_h$ , hyperpolarization activated cationic current;  $I_{pNa+}$ , persistent sodium current.

\* Corresponding authors.

E-mail addresses: [pignatelli.michele@gmail.com](mailto:pignatelli.michele@gmail.com) (M. Pignatelli), [xlkg@loadinc.net](mailto:xlkg@loadinc.net) (X. Leinekugel).

waves (see Fig. 1). Specifically, oscillatory activation of the perforant path arising from layer II–III cells of the entorhinal cortex (Amaral and Witter, 1989), generates prominent theta oscillations in the molecular layer of the dentate gyrus, accompanied by phased locked firing of granule cells (Muñoz et al., 1990) and interneurons (Sik et al., 1997). Current source density analysis of the laminar profile of CA1 field potentials recorded during theta oscillations revealed the presence of current sources in the pyramidal cell layer coupled to current sinks in the stratum lacunosum moleculare, suggesting synchronous somatic inhibition and dendritic excitation (Kamondi et al., 1998). Because lesion of the entorhinal cortex suppressed the sink in the stratum lacunosum moleculare (Kamondi et al., 1998) and around the hippocampal fissure (Ylinen et al., 1995), it has been suggested that while parvalbumin positive interneurons, in particular axoaxonic basket cells (Klausberger et al., 2003), send inhibitory inputs to the pyramidal

cell layer, the entorhinal cortex provides periodic excitatory inputs to the apical dendrites of the pyramidal cells (Kamondi et al., 1998). In addition, because oriens-lacunosum molecular interneurons and pyramidal cells share similar preferred theta phase of firing (Klausberger et al., 2003) CA1 pyramidal cells receive rhythmic somatic hyperpolarization coupled to rhythmic dendritic depolarization, resulting in a net decrease of the input resistance during theta oscillations (Buzsáki, 2002; Kamondi et al., 1998). However, while both deep and superficial CA1 pyramidal cells fire preferentially at the trough of theta oscillations during active exploration, only deep pyramidal cells shift their preferred phase of firing to the peak of theta during paradoxical sleep (Mizuseki et al., 2011). Interestingly, CA1 and CA3 pyramidal cells are active on opposite theta phases suggesting that interleaving CA3 neuronal sequences may bind CA1 assemblies (Dragoi and Buzsáki, 2006). In addition to the laminar characteristics of theta



**Fig. 1.** Hippocampal circuitry underlying theta oscillations. (a) Excitatory (black) and inhibitory (blue) transmission involved in electrogeneration of theta oscillations. The medial septum (MS) provides rhythmic inhibitory inputs to CA1 basket cells (soma and dendrites in black plus axon in blue), which in turn provide rhythmic perisomatic inhibition to CA1 pyramidal cells. The entorhinal cortex (EC) provides rhythmic excitatory inputs at theta frequency through the perforant path to the granule cells of the dentate gyrus (DG) and to the apical dendrites of CA1 and CA3 pyramidal cells. The rhythmic activity from DG granule cells is conveyed through the mossy fibers to the CA3 pyramidal cells and then, through the Schaffer collaterals, to CA1 pyramidal cells. (b) Current source density analysis of theta oscillations in the CA1 region during REM sleep, obtained with a silicone probe (16 sites, 50  $\mu\text{m}$  spaced). Note the presence of rhythmic somatic hyperpolarization (source in stratum pyramidale) coupled to rhythmic dendritic depolarization (sink in stratum lacunosum moleculare). Current sources in red and sinks in blue. Abbreviations: or stratum oriens, pyr stratum pyramidale, rad stratum radiatum, slm stratum lacunosum moleculare.

oscillations, *in vivo* multisite electrophysiological recordings have shown that the coherence of theta oscillations (Royer et al., 2010; Sabolek et al., 2009) and the relationship between theta power and locomotion speed (Hinman et al., 2011) decreases along the septo-temporal axis.

### 3. Hippocampus-associated regions implicated in theta oscillations

#### 3.1. Medial Septum (MS)

In the behaving rat, hippocampal theta oscillations are accompanied by high cholinergic tone (Jasper and Tessier, 1971; Zhang et al., 2010). In order to tune the network state into theta regime, the hippocampal activity diverges from a resting state characterized by sharp-wave-ripple complexes (Csicsvari et al., 2000) also known as large irregular activity. This transition is triggered and sustained by the pacemaker activity of the medial septum (Petsche et al., 1962), a structure well known for its direct cholinergic projections to the hippocampus. The MS together with the vertical and horizontal limb of the diagonal band of Broca (VDB and HDB) is a midline structure part of a vast region known as basal forebrain (Zaborszky et al., 2005). The MS projects to specific hippocampal subregions (Crutcher et al., 1981) such as: (i) dentate gyrus (the supragranular zone of the septal pole and the hilus), (ii) CA1 (stratum oriens and stratum lacunosum moleculare), (iii) CA2–CA3 (stratum oriens and stratum radiatum). The subiculum, presubiculum and parasubiculum are targeted as well (Crutcher et al., 1981). The MS contains an heterogeneous population of projecting cells characterized by cholinergic (Mesulam et al., 1983), GABAergic (Freund, 1989; Kiss et al., 1990; Leranthy and Frotscher, 1989) or glutamatergic (Colom et al., 2005; Manns et al., 2001) neurotransmission. Interestingly, only GABAergic and glutamatergic cells display pacemaker properties *in vitro* (Huh et al., 2010; Markram and Segal, 1990; Varga et al., 2008). *In vivo*, MS cells are known to fire at theta frequency, in phase with the negative peak of the theta signal recorded in the CA1 pyramidal layer (Dragoi et al., 1999). In addition, the firing of the MS cells is suppressed during sharp-waves ripples (Dragoi et al., 1999). Inhibition of the MS cell activity, through targeted lidocaine (Koenig et al., 2011; Mizumori et al., 1990; Winson, 1978) or muscimol (Bland et al., 1996; Brandon et al., 2011) injection, leads to complete suppression of the theta oscillations. However, while i.p. injection of the muscarinic antagonist, atropine (Bradley and Nicholson, 1962), suppresses the theta oscillations associated with paradoxical sleep, it leaves almost intact the one related to locomotion, usually absent under urethane anesthesia (Kramis et al., 1975). Furthermore, direct injection of atropine into the hippocampus leads to a decrease of the theta power (Li et al., 2007) and selective lesion of MS cholinergic neurons with IgG-saporin (Lee et al., 1994), or MS GABAergic neurons with kainic acid (Yoder and Pang, 2005), significantly reduces theta power, suggesting that both populations of neurons are involved in the generation of theta oscillations. Septo-hippocampal cholinergic afferents are known to establish synaptic contacts in the dentate gyrus (Clarke, 1985; Frotscher, 1991), preferentially with neuropeptide-Y interneurons (Dougherty and Milner, 1999) and cholinergic activity can trigger intrinsic theta oscillations in the hippocampus as demonstrated by *in vitro* intracellular recordings under bath perfusion with the cholinergic agonist, carbachol (Bland et al., 1988). The MS GABAergic cells expressing parvalbumin (Freund, 1989) and targeting the hippocampal basket cells (Freund and Antal, 1988; Tóth et al., 1997), have been reported to drive hippocampal activity (Hangya et al., 2009) and the hippocampal feedback to the medial (Tóth et al., 1993) and lateral septum (Leranthy and Frotscher, 1989) may

support the synchronization between the two structures. Interestingly, MS glutamatergic cells expressing the vesicular glutamate transporter 2 (VGLUT-2) (Hajszan et al., 2004) have been reported to project to the hippocampus and to produce rhythmic discharges in the theta range (Huh et al., 2010) but their role in the genesis of the hippocampal theta oscillations remains to be established.

#### 3.2. Entorhinal Cortex (EC)

The EC provides much of the cortical information to the hippocampal formation. This important region is identified by its projections targeting the dentate gyrus of the hippocampus: the perforant path. In the hippocampus, the septo-temporal organization of the EC fibers is associated to specific subregions of the EC (Dolorfo and Amaral, 1998): (i) the septal part of the hippocampus receives afferents from a portion of the lateral EC (LEC) adjacent to the rhinal fissure and from the medial EC (MEC), (ii) the central part is innervated by the medial portion of the LEC and by rostral portion of the MEC, whereas (iii) the temporal part is innervated by the most medial portion of the LEC and by most rostral portion of the MEC. The EC sends direct glutamatergic inputs to the stratum lacunosum moleculare of CA1, CA3 and to the molecular layer of the dentate gyrus (Amaral and Witter, 1989; Van Groen et al., 2003) but only the MEC seems strongly involved in the generation of the hippocampal theta oscillations (Deshmukh et al., 2010). MEC lesion reduces hippocampal theta oscillations in drug free animals and in atropine treated animals (Montoya and Sainsbury, 1985), suggesting that atropine resistant theta oscillations involve the MEC. However, septal lesion or septal injections of lidocaine or muscimol abolish theta oscillations in the MEC of freely behaving animals (Brandon et al., 2011; Koenig et al., 2011; Mitchell et al., 1982). Combined septal injection of kainic acid (preferentially eliminating GABAergic cells) and IgG-saporin (preferentially eliminating cholinergic cells) suppresses locomotion-related theta oscillations (Yoder and Pang, 2005), suggesting that septal GABAergic cells may drive the MEC theta inputs to the hippocampus. Nevertheless, the MEC circuitry generates local theta oscillations that reverse in layer II and MEC units are phase locked to the local theta signal (Dickson et al., 1995; Mitchell and Ranck, 1980; Mizuseki et al., 2009; Stewart et al., 1992). Moreover, layer II stellate cells and deep layer cells (Schmitz et al., 1998) expressing  $I_{pNa+}$  and  $I_h$  currents (Alonso and Llinás, 1989; Dickson et al., 2000), display subthreshold oscillatory activity within the theta range and the cholinergic agonist, carbachol, can induce theta oscillations in MEC slices (Gołbiewski et al., 1994; Konopacki and Gołbiewski, 1992). Interestingly, layer V pyramidal cells, the link between the hippocampus and many cortical regions, respond to consecutive stimuli with graded changes in firing frequency linked to cholinergic muscarinic receptor activation and characterized by sustained levels of firing frequency that can be either increased or decreased in an input-specific manner (Egorov et al., 2002). As the hippocampus, the MEC circuit usually displays a resting state and its transition towards theta state is triggered and supported by the activation of MS, VDB and HDB (Jeffery et al., 1995; Mizumori et al., 1992). Therefore, the theta activity of both EC and hippocampus seems to rely on MS, VDB and HDB although with a certain degree of specialization. In fact, while the EC inputs are mainly attributed to VDB and HDB, the hippocampal inputs seem restricted to MS and VDB (Gaykema et al., 1990).

#### 3.3. Subicular Complex (Sub)


The output of CA1 is characterized by columnar projections towards the subiculum, a structure known to project directly and indirectly, through the presubiculum and the parasubiculum, to the EC layer II–III cells (Amaral and Witter, 1989). Subicular



pyramidal cells (Wang et al., 2006) and layer II parasubicular cells (Glasgow and Chapman, 2007, 2008) display intrinsic subthreshold oscillations in the theta range, which depend on  $I_{pNa+}$  and  $I_h$  currents. In addition, the whole subicular complex is under the control of the septal complex (Crutcher et al., 1981). As a potential consequence, locally generated atropine-sensitive theta oscillations can be observed in the parasubiculum, in phase with the theta oscillations recorded in the stratum lacunosum moleculare, and display phase reversal between the superficial and deep layers (Glasgow and Chapman, 2007). Cellular activity is usually phase locked to the theta signal (Anderson and O'Mara, 2003; Boccara et al., 2010; Cacucci et al., 2004). Interestingly, a more detailed investigation has revealed that only the spiking activity of the superficial layers is phase locked to theta oscillations whereas deep layers fire tonically during theta state (Chrobak and Buzsáki, 1994).

#### 4. Control of the theta pacemaker

The septal complex (MS, VDB and HDB) is under the control of different deep structures (Bland and Oddie, 1998; Vertes and Kocsis, 1997). This control is specifically operated by the cholinergic nuclei of the brainstem, the pedunculo pontine tegmentum and the laterodorsal tegmentum (Mesulam et al., 1983), in two different ways (see Fig. 2). The first, direct pathway, projects directly to the septal complex (Hallanger and Wainer, 1988; Woolf and Butcher, 1986). The second, indirect pathway, is mainly mediated by the hypothalamus (Woodnorth et al., 2003) through its supramammillary nuclei (Kirk and McNaughton, 1991) and posterior hypothalamic nuclei (Oddie et al., 1994). Importantly, the two brainstem nuclei, which are known to mediate cortical desynchronization (Datta and Siwek, 1997), support in their caudal part,



**Fig. 2.** Neural circuitry involved in the generation of theta oscillations. (a) Layer specific afferents supporting hippocampal theta oscillations (or: stratum oriens, pyr: stratum pyramidale, rad: stratum radiatum, slm: stratum lacunosum moleculare, m: stratum moleculare, gr: stratum granulare). Neurotransmitters: acetylcholine (ACh), gamma amino butyric acid (GABA), glutamate (glu). Abbreviations: EC, entorhinal cortex, SUM, supramammillary nucleus. (b) Direct (dotted line) and indirect pathways mediating the activation of the septal complex (SC). Both of them pass through the medial forebrain bundle (MFB) and the fornix. HT: hypothalamus, BS: brainstem.

composed by the sublaterodorsal nucleus and precoeruleus region, the generation of paradoxical sleep (Lu et al., 2006). Instantaneous increase in power and frequency of the theta oscillations during paradoxical sleep, known as phasic theta (Montgomery et al., 2008), is correlated with phasic pontine activity (Karashima et al., 2005).

#### 4.1. Brainstem control of theta oscillations

##### 4.1.1. Pedunculopontine Tegmentum (PPT) and Laterodorsal Tegmentum (LDT)

The PPT and the LDT contain a heterogeneous cellular population characterized by cholinergic, GABAergic or glutamatergic neurotransmission (Lee et al., 1988). The cells fire tonically during the active state (Steriade et al., 1990). Direct stimulation of PPT-LDT elicits cortical desynchronization, hippocampal theta oscillations accompanied by wakefulness or paradoxical sleep (Datta and Siwek, 1997) and can induce locomotion, as other components of the mesencephalic locomotor region (Milner and Mogenson, 1988; Skinner and Garcia-Rill, 1984). Suppression of the PPT activity, through local injection of procaine, abolishes hippocampal theta rhythm in urethane anaesthetized rats (Nowacka et al., 2002). The anatomical substrates of these effects are represented by two groups of projections: (i) a descending pathway innervating the brainstem and the spinal cord (Rye et al., 1988) and (ii) an ascending pathway projecting to the diencephalon and to the basal forebrain (Hallanger et al., 1987). The descending projections are characterized by cholinergic transmission up to the beginning of the spinal cord and then by non-cholinergic terminals (Rye et al., 1988). The ascending component is mainly characterized by cholinergic transmission, specifically targeting the thalamus, lateral septum, MS, VLB and HLB, amygdala, globus pallidus (internal segment and entopeduncular nucleus), substantia nigra, subthalamic nucleus, zona incerta, lateral hypothalamus and superior colliculus (Hallanger et al., 1987; Hallanger and Wainer, 1988; Inglis and Winn, 1995; Rye et al., 1987; Woolf and Butcher, 1986). The nucleus reticularis pontis oralis is targeted by the PPT as well (Mitani et al., 1988; Semba et al., 1990; Shiromani et al., 1988). In addition, the PPT is directly modulated by the activity of the following structures: nucleus accumbens, ventral pallidum, caudate putamen (ventrolateral), globus pallidus (medial), subthalamic nucleus, substantia nigra, locus coeruleus, raphe nuclei and finally the parabrachial nucleus and the contralateral PPT (Inglis and Winn, 1995; Semba and Fibiger, 1992; Steininger et al., 1992). Different neuromodulators are involved in the control of the PPT-LDT activity and in the modulation of theta oscillations: (i) noradrenergic inputs from the locus coeruleus leads to theta activity (Berridge and Foote, 1991) and, (ii) serotonergic inputs from the median raphe (MR) suppress theta oscillations desynchronizing hippocampal activity (Jackson et al., 2008), whereas inactivation of the MR through muscimol injection can elicit theta oscillations (Kinney et al., 1995).

##### 4.1.2. Sublaterodorsal nucleus (SLD) and Precoeruleus region (PC)

The SLD and PC are two structures located caudally to the LDT and their inactivation is responsible for a significant decrease in paradoxical sleep occurrence together with a reduction of the associated theta oscillations (Lu et al., 2006; Saper et al., 2010). These nuclei contain glutamatergic and GABAergic cells, as well as cholinergic cells in the SLD (Allen-Brain-Atlas). Injection of retrograde tracers into the ventral horn of the spinal cord has revealed direct projections from SLD cells which, importantly, express VGLUT-2 (Fuller et al., 2007). Animals with lesion of the SLD do not display muscle atonia. Therefore SLD, by exciting glycinergic/GABAergic cells of the spinal cord, could elicit the muscle atonia observed during paradoxical sleep. In addition, injection of a retrograde tracer in

the MS has revealed labeled cells in the PC region and selective lesion of the PC suppresses sleep-related hippocampal theta oscillations without affecting muscle atonia and cortical desynchronization (Fuller et al., 2007; Lu et al., 2006).

##### 4.1.3. Nucleus Reticularis Pontis Oralis (RPO)

The RPO is a pontine structure directly innervated by the PPT (Semba et al., 1990) and known to project mainly to the supramammillary nucleus (Vertes et al., 1986). Neurons in the RPO display tonic spiking activity during theta state (Vertes, 1977, 1979). Under urethane anesthesia, stimulation of the RPO leads to theta activity while its inhibition by local procaine injection suppresses the spontaneous and tail-pinch evoked theta oscillations (Bland and Oddie, 1998).

##### 4.1.4. Nucleus Incertus (NI)

The NI is a small nucleus embedded in the periaqueductal gray adjacent to the dorsal tegmental nucleus. It is composed of longitudinal columns of cells coexpressing GABA, relaxin 3 and neuromedin B (Ryan et al., 2011). The NI integrates synaptic inputs from prefrontal and premotor cortex, hippocampus, septal complex, lateral habenula, interpeduncular nucleus, posterior hypothalamus, periaqueductal gray, dorsal raphe and retro-LDT, targeting the output on the prefrontal cortex, premotor cortex, hippocampus, medio-dorsal and ventromedial thalamic nucleus, septal complex (in particular to medial part), posterior hypothalamus, supramammillary nucleus, periaqueductal gray, dorsal raphe and retro-LDT (Goto et al., 2001). Electrical stimulation of the NI elicits theta oscillations in urethane-anaesthetized rats while injection of muscimol prevents theta oscillations evoked by RPO stimulation (Nuñez et al., 2006).

##### 4.1.5. Ventral tegmental nucleus of Gudden (VTn)

The VTn is located in the medial part of the pontine tegmental area. It is related to Papez's circuit through its GABAergic connections with the medial mammillary nucleus (Allen and Hopkins, 1990) and, in addition, it receives direct afferents from the LDT (Irle et al., 1984). The VTn cells are reported to rhythmically discharge during theta activity, in awake behaving and urethane-anaesthetized rats (Bassant and Poindessous-Jazat, 2001; Kocsis et al., 2001).

##### 4.1.6. Dorsal Raphe (DR)

The DR is the major serotonergic nucleus of the brain (Steinbusch and Nieuwenhuys, 1981). The DR projects to the piriform, insular, frontal, occipital, entorhinal, perirhinal, frontal orbital, anterior cingulate, and infralimbic cortices, hippocampal formation, as well as the intralaminar nuclei of the thalamus, septal complex (in particular the lateral septum, VDB and HDB), SUM, ventral tegmental area and the PPT (Vertes, 1991). The main afferents arise from orbital, cingulate, infralimbic, dorsal peduncular, and insular cortices, ventral pallidum, lateral habenula, claustrum, zona incerta, central nucleus of the amygdala, lateral, dorsal and posterior hypothalamic areas (Peyron et al., 1998). Stimulation of the DR can elicit scopolamine sensitive theta oscillations, suggesting that the effect is mediated by the cholinergic system (Peck and Vanderwolf, 1991). Around 55% of the cellular population of the DR is reported to rhythmically fire in phase with theta oscillations in the behaving rat (Kocsis and Vertes, 1992).

#### 4.2. Hypothalamic control of theta oscillations

##### 4.2.1. Supramammillary nucleus (SUM)

SUM cells receive direct innervations from the nucleus reticularis pontis oralis (Vertes et al., 1986), the septal complex and the LDT (Gonzalo-Ruiz et al., 1999), and project to the septal

complex (Borhegyi et al., 1998) and to the hippocampus (Vertes, 1992), innervating the granule cell layer of the dentate gyrus and the pyramidal cell layer of the CA2-CA3 region (Maglóczy et al., 1994). In the rat, its cellular population is composed of glutamatergic cells (Kiss et al., 2002) producing synchronous discharges phase-locked to the theta signal (Kirk and McNaughton, 1991). Electrical stimulation of the nucleus reticularis pontis oralis, following procaine injection in the SUM, leads to a reduced theta power and frequency in urethane-anesthetized rats, whereas injection of procaine in the MS reduces exclusively theta power (Kirk and McNaughton, 1993), suggesting that the SUM could exert a direct control on the frequency of theta oscillations. However, selective lesions of the SUM have failed to suppress hippocampal theta activity in the behaving animal (Thinschmidt et al., 1995), suggesting the restricted role of a theta modulator. Even so, based on the existence of reciprocal connections between SUM and septo-hippocampal systems (Borhegyi and Freund, 1998; Borhegyi et al., 1998), it is proposed a state-dependent bidirectional coupling characterized by alternated leadership between the two oscillators (Kocsis, 2006).

#### 4.2.2. Mammillary bodies (MB)

The SUM can also be considered as a component of the mammillary bodies (MB), a central relay of the limbic system (Vann and Aggleton, 2004), which receives direct inputs from the subiculum (Meibach and Siegel, 1975) and which cellular activity is phase-locked to the theta signal (Kocsis and Vertes, 1994). However, these structures are remarkably different both anatomically and physiologically. The anatomical difference lies in the thalamic target of the projections: (i) the SUM projects preferentially to the nucleus reuniens, a site which provides direct innervation to the CA1 stratum lacunosum moleculare (Vertes, 1992; Vertes et al., 2006) and characterized by tonic discharges during theta state (Morales et al., 2007), whereas (ii) the MB projects to the anterior thalamic nuclei (Allen and Hopkins, 1990), in which rhythmic discharges phase locked to the theta rhythm have been reported (Vertes et al., 2001). The physiological difference between SUM and MB appears after septal inactivation, leading to the persistence of theta activity in the SUM but its suppression in the MB (Kirk et al., 1996). Overall, these experimental evidences indicate that the MS drives the MB whereas the SUM is independent.

#### 4.2.3. Posterior nucleus of the hypothalamus (PH)

The PH is a nucleus adjacent to the MB. Reciprocally connected to the PPT (Hallanger and Wainer, 1988; Steininger et al., 1992), it also innervates the SUM, the septal complex and the reuniens nucleus of the thalamus (Vertes et al., 1995). The nucleus mainly contains glutamatergic cells expressing VGLUT-2 (Ziegler et al., 2002) and displaying tonic firing during theta state (Bland et al., 1995). In addition, selective inactivation of the PH by muscimol (Bocian and Konopacki, 2007) or procaine injection (Oddie et al., 1994), in rats under urethane anesthesia, impairs theta oscillations.

### 5. Additional regions involved in theta oscillations

#### 5.1. Medial Prefrontal Cortex (mPFC)

The mPFC is one of the principal targets of the CA1 pyramidal cells from the ventral part of the hippocampus (Swanson, 1981). The CA1 pyramidal cell terminals, distributed in layer I and layer V, provide excitation to both mPFC pyramidal cells and interneurons (Gabbott et al., 2002). The mPFC is typically identified by the thalamic afferents originating in the mediodorsal nucleus (Condé et al., 1990) and receives, in addition, afferent projections from the agranular insular cortex, the entorhinal and piriform

cortices, the subiculum, the endopiriform nucleus, the amygdalo-piriform transition, the amygdalohippocampal area, the lateral tegmentum, the parabrachial nucleus (Condé et al., 1995) and importantly from the septal complex (Gaykema et al., 1990), the ventral tegmental area (Sesack et al., 1998) and the LDT (Sato and Fibiger, 1986). Therefore, since the mPFC is under the control of the hippocampus, EC, septal complex and LDT, the detection of synaptically driven theta oscillations in this structure is not surprising. In fact, the presence of pyramidal cells and parvalbumin positive interneurons phase locked to theta oscillations has been reported in the rat mPFC (Hartwich et al., 2009; Hyman et al., 2005; Jones and Wilson, 2005a; Sirota et al., 2008; Tsujimoto et al., 2006), although delayed relative to CA1 pyramidal cells (Siapas et al., 2005). The efferent projections of the mPFC are targeted on cingulate, entorhinal and perirhinal cortical structures, the dorsomedial and ventral striatum, basal forebrain nuclei, basolateral amygdala, lateral hypothalamus, mediodorsal, midline and intralaminar thalamic nuclei, periaqueductal gray region, ventral midbrain tegmentum, LDT, PPT, and raphe nuclei (Apergis-Schoute et al., 2006; Gabbott et al., 2005; Gaykema et al., 1991; Sesack et al., 1989; Vertes, 2002). Interestingly, the ventral striatum, one of the principal targets of the mPFC, contains theta modulated cells (van der Meer and Redish, 2011). In summary, the mPFC is reciprocally connected to most of the regions involved in theta generation and can be considered as a global modulator of theta activity. Simultaneous multisite electrophysiological recordings, targeting both hippocampus and mPFC, have shown: (i) a correlation between the two structures paralleled by the increased coupling in the 7 Hz range during behaviors involving spatial working memory (Jensen, 2005; Jones and Wilson, 2005b; Paz et al., 2008), and (ii) an increase in the coherence between the two theta signals, which reaches a maximum at the choice point of a Y-maze, in particular following acquisition of the task (Benchenane et al., 2010).

#### 5.2. Amygdala

The amygdala is a complex anatomical structure critical for producing appropriate emotional and behavioral responses to biologically relevant sensory stimuli, representing a cardinal link between the sensory systems, the limbic system and all the sub-cortical structures involved in emotional responses such as the hypothalamus, the brainstem and the striatum (McDonald, 1998). In the cat, the projection cells and the interneurons of the basolateral amygdala are significantly phase locked to theta oscillations (Paré and Gaudreau, 1996). To support this observation, the amygdala receives direct innervation from different regions implicated in theta oscillations such as: ventral hippocampus (Kishi et al., 2006), prefrontal cortex (Cassell and Wright, 1986; McDonald et al., 1996), lateral entorhinal cortex (McDonald and Mascagni, 1997), ventral subiculum (Canteras and Swanson, 1992), HDB (Meibach and Siegel, 1977), parabrachial area (Bernard et al., 1993) and a small contribution from the PPT and LDT (Hallanger and Wainer, 1988). The efferent projections are as well directed to most of the regions implicated in theta oscillations: (i) from the basomedial nucleus to the HDB, the mPFC and the lateral and ventromedial nuclei of the hypothalamus (Jolkkonen et al., 2002; Petrovich et al., 1996), (ii) from the basal nucleus to the stratum oriens and stratum radiatum of CA3 and CA1 and from lateral nucleus to layer III of the entorhinal cortex (Pikkarainen et al., 1999), (iii) from the posterior nucleus to the stratum lacunosum-moleculare of the temporal CA1 subfield, the adjacent molecular layer of the proximal temporal subiculum, the parahippocampal region, and the entorhinal cortex (Canteras et al., 1992; Kemppainen et al., 2002), (iv) from the medial and central nuclei to the midline thalamic nuclei (McKenna and Vertes, 2004; Otake et al., 1994) and (v) from the central nucleus to the PPT (Semba



and Fibiger, 1992). Synchronization of theta activities in the amygdalohippocampal network represents a neuronal correlate of conditioned fear, potentially facilitating neuronal communication during memory retrieval (Seidenbecher et al., 2003). While theta coupling increases between all areas during retrieval of conditioned fear, it decreases during extinction learning (Lesting et al., 2011). Furthermore, overnight bidirectional changes in fear memory are selectively correlated with modifications in theta coherence between the amygdala, medial prefrontal cortex, and hippocampus during paradoxical sleep (Popa et al., 2010). Finally, septal lesions were found to potentiate the freezing response elicited by contextual stimuli (Sparks and LeDoux, 1995).

### 5.3. Reuniens nucleus of the thalamus (RE)

The RE is a midline structure known to mediate arousal and awareness, receiving direct cholinergic afferents from the PPT (Hallanger et al., 1987; Van der Werf et al., 2002; Vertes et al., 1995). Furthermore, it is considered as an interface between the prefrontal cortex and the hippocampus since pyramidal cells of the prefrontal cortex innervate RE cells projecting to the stratum lacunosum moleculare of CA1 (Vertes, 2002; Vertes et al., 2007). In the hippocampus, RE terminals establish asymmetric synapses with both GABAergic and non-GABAergic dendrites (Dolleman-Van der Weel and Witter, 2000). The RE sends region specific glutamatergic fibers to the septal complex (Bokor et al., 2002), EC, subiculum and the amygdala (Dolleman-Van der Weel and Witter, 1996; Su and Bentivoglio, 1990; Vertes et al., 2006). The RPO, PH, SUM and PPT all innervate the RE (Hallanger et al., 1987; Newman and Ginsberg, 1994). RE cells coexpress calretinin (Cicchetti et al., 1998; Wouterlood et al., 2008) and VGLUT-2 (Wouterlood et al., 2008) and display tonic firing during theta state in rats under urethane anesthesia (Morales et al., 2007).

## 6. Future directions

The dynamic of the extracellular currents along the hippocampal septo-temporal axis and the relative contribution of each

independent hippocampal oscillator to the theta field still represents an open question that could be addressed by selective inactivation of hippocampal or entorhinal subregions associated with multisite recordings along the longitudinal axis and network modeling (Buzsáki, 2002).

The power of the theta oscillations exhibits fluctuations correlated to the behavioral state of the animal (Wyble et al., 2004) and phasic theta episodes are reported during paradoxical sleep (Montgomery et al., 2008). The dynamic interplay between all the extra hippocampal oscillators and the resulting integration at the hippocampal level provides an intricate substrate to modulate theta power. Alternatively, the significant correlation between theta power and pontine waves occurrence (Karashima et al., 2004) may suggest a pivotal modulation of the PPT on the extra hippocampal theta generators. Moreover, because sensory-related theta has a slower frequency than locomotion-related theta (4–9 Hz against 6–12 Hz) (Kramis et al., 1975), their expected coexpression during behavior is a potential source of oscillatory interference. From the signal theory, the interference between two summed theta waves of slightly different frequencies would result in a rhythmically modulated theta wave. Therefore, one might expect theta power to be rhythmically modulated during spontaneous behavior. Simultaneous recordings of hippocampal theta together with different anatomical elements of its generating circuit during controlled behavioral protocols should help understand the role played by these structures in the generation and modulation of hippocampal theta, allowing to explore new dimensions of theta oscillations and potentially helping us understand the temporal organization of sensory-motor processes.

Interestingly, the dorsocaudal region of the rat MEC contains a topographically organized neural map of the spatial environment constituted by the *grid cell*, which is activated whenever the animal position coincides with any vertex of a regular grid of equilateral triangles spanning the surface of the environment (Hafting et al., 2005). The map is anchored to external landmarks, but persists in their absence. Grid cells exhibit firing at regular spatial locations and temporal modulation with theta oscillation. It has been recently shown in rats, that MS inactivation, through muscimol

**Table 1**  
Synaptic and network properties of the regions implicated in theta oscillations.

	Neurotransmitters				Network properties				Brain states		
	ACh	5HT	Glutamate	GABA	Pacemaker	Oscillator	Driver	Modulator	Awake	REM	Urethane
CA1 pyr			+			+			+	+	+
CA3 pyr			+			+			+	+	+
DG			+			+			+	+	+
SC	+		+	+	+	+			+	+	+
EC			+			+			+	+	+
Sub			+			+			+	+	+
SUM			+			+			ND	ND	+
MB			+			+			ND	ND	+
PH			+					+	ND	ND	+
PPT											
LDT	+						+		+	+	+
SLD											
PO			+				+			+	
RPO			+					+	ND	ND	+
NI				+				+	ND	ND	+
VTn				+		+			+	+	+
DR		+				+	+		+	+	ND
PFC			+			+			+	+	+
Amy			+			+			+	+	ND
RE			+					+	ND	ND	+

+, reported in the literature.

ND, expression of theta not demonstrated in this structure in the corresponding brain state.

Pacemaker, displays intrinsic theta activity.

Oscillator, displays inducible theta activity upon adequate stimulation.

Driver, required for theta oscillations in other structures.

Modulator, able to modulate the power or frequency of theta oscillations in other structures.

or lidocaine, leads to suppression in the spatial periodicity of the grid cells, unaffected the spatial representation of the hippocampal place cells (Brandon et al., 2011; Koenig et al., 2011). In addition, grid cells persist after hippocampal inactivation (Hafting et al., 2008). It would be interesting to understand how the pacing activity of the MS affects the interplay occurring between the cortical and the subcortical inputs in the generation of spatial representation of the MEC and to address, through selective lesion of the septo-entorhinal pathway, what is the impact of a disorganized entorhinal input on theta activity of the hippocampus.

Other critical issues are to decipher the exact sequential activation of all the structures involved in the theta generation (see Table 1) as well as the mechanisms involved in the synchronization of the theta oscillators during specific behavioral tasks (Benchenane et al., 2010; Jensen, 2005; Seidenbecher et al., 2003). These issues could be examined by combining simultaneously, in different configurations, multiple intracellular (Poulet and Petersen, 2008) and massive extracellular recordings (Csicsvari et al., 2003). The precise timing between circuit elements may highlight potential targets useful to address specific behavioral paradigms. Finally, despite the considerable effort reported by all the past anatomical studies, a methodic dissection aimed to quantify the structural and ultrastructural contribution of all the deep structures to the theta generation can still provide useful information about the theta circuit. Detailed information relative to the projections and synaptic targets of the cholinergic and glutamatergic cells of the PPT and LDT, the glutamatergic cells of the MS, the SUM and the PH, the noradrenergic cells of the locus coeruleus and the serotonergic cells of the raphe nuclei can still provide important insights.

## 7. Conclusions

The progress of our understanding of theta generators closely correlates with technological advances. The identification of theta generators was primarily based on electrophysiological recordings such as unit activity and field potential followed by electrolytic lesions to allow histological verification. The spiking activity was classified in relation to theta signal as (i) theta-phase modulated (units phase locked to theta signal), (ii) theta-ON (tonic activity during theta) and (iii) theta-OFF (no activity during theta) (Colom and Bland, 1987). Electrical stimulation and surgical or chemical lesions allowed a systematic inspection of the regions involved in theta generation (Robinson and Vanderwolf, 1978). In addition, anatomical studies based on combined applications of retrograde and anterograde tracers coupled to ultrastructural observation provided the necessary morphological support to the physiological results. Nevertheless, a real progress appears with the use of pharmacological tools first applied systemically (atropine, scopolamine, physostigmine, urethane, ...) and then locally by targeted injections. Glutamate was employed to provide excitation of local populations of neurons, and lidocaine, procaine or muscimol to provide their inhibition. Furthermore, the advent of the immunocytochemistry has lead rapidly towards a targeted cellular dissection of single theta generators. However, the precision of both anatomical and pharmacological techniques was often limited by the relative control of the volume of tracers or drugs injected. With such techniques, selective stimulation of a specific cellular population was out of reach. In fact, electrical stimulation and lidocaine injection can alter not only the somas but also passing axons, resulting in less straightforward interpretation.

Today, the advent of molecular biology offers new perspectives to the field. Selective tagging coupled to the expression of optogenetic tools (Gradinaru et al., 2010) allow the stimulation or inhibition of selective cellular populations, increasing the precision from

the regional to the single cell scale. The convergence of physiological, anatomical and molecular tools finally allows the perspective of a new generation of experimental investigations.

## Acknowledgments

We are grateful to Thibault Maviel for comments and discussion. This work was supported by the following funding sources: INSERM (X.L.), FRM (X.L.), CNRS (A.B., M.P., X.L.), Région Aquitaine (X.L.), ENI-Net (X.L.), ANR (A.B., X.L.), ERA-Net (M.P., X.L.).

## References

- Allen-Brain-Atlas.
- Allen, G.V., Hopkins, D.A., 1990. Topography and synaptology of mamillary body projections to the mesencephalon and pons in the rat. *J. Comp. Neurol.* 301 (2), 214–231.
- Alonso, A., Llinás, R.R., 1989. Subthreshold  $\text{Na}^+$ -dependent theta-like rhythmicity in stellate cells of entorhinal cortex layer II. *Nature* 342 (6246), 175–177.
- Amaral, D.G., Witter, M.P., 1989. The three-dimensional organization of the hippocampal formation: a review of anatomical data. *Neuroscience* 31 (3), 571–591.
- Anderson, M.I., O'Mara, S.M., 2003. Analysis of recordings of single-unit firing and population activity in the dorsal subiculum of unrestrained, freely moving rats. *J. Neurophysiol.* 90 (2), 655–665.
- Apergis-Schoute, J., Pinto, A., Paré, D., 2006. Ultrastructural organization of medial prefrontal inputs to the rhinal cortices. *Eur. J. Neurosci.* 24 (1), 135–144.
- Bassant, M.H., Poindessous-Jazat, F., 2001. Ventral tegmental nucleus of Gudden: a pontine hippocampal theta generator? *Hippocampus* 11 (6), 809–813.
- Benchenane, K., Peyrache, A., Khamassi, M., Tierney, P.L., Gioanni, Y., Battaglia, F.P., Wiener, S.I., 2010. Coherent theta oscillations and reorganization of spike timing in the hippocampal–prefrontal network upon learning. *Neuron* 66 (6), 921–936.
- Bernard, J.F., Alden, M., Besson, J.M., 1993. The organization of the efferent projections from the pontine parabrachial area to the amygdaloid complex: a Phaseolus vulgaris leucoagglutinin (PHA-L) study in the rat. *J. Comp. Neurol.* 329 (2), 201–229.
- Berridge, C.W., Foote, S.L., 1991. Effects of locus coeruleus activation on electroencephalographic activity in neocortex and hippocampus. *J. Neurosci.* 11 (10), 3135–3145.
- Bland, B.H., 1986. The physiology and pharmacology of hippocampal formation theta rhythms. *Prog. Neurobiol.* 26 (1), 1–54.
- Bland, B.H., Colom, L.V., Konopacki, J., Roth, S.H., 1988. Intracellular records of carbachol-induced theta rhythm in hippocampal slices. *Brain Res.* 447 (2), 364–368.
- Bland, B.H., Konopacki, J., Kirk, I.J., Oddie, S.D., Dickson, C.T., 1995. Discharge patterns of hippocampal theta-related cells in the caudal diencephalon of the urethane-anesthetized rat. *J. Neurophysiol.* 74 (1), 322–333.
- Bland, B.H., Oddie, S.D., 1998. Anatomical, electrophysiological and pharmacological studies of ascending brainstem hippocampal synchronizing pathways. *Neurosci. Biobehav. Rev.* 22 (2), 259–273.
- Bland, B.H., Trepel, C., Oddie, S.D., Kirk, I.J., 1996. Intraseptal microinfusion of muscimol: effects on hippocampal formation theta field activity and phasic theta-ON cell discharges. *Exp. Neurol.* 138 (2), 286–297.
- Boccard, C.N., Sargolini, F., Thoresen, V.H., Solstad, T., Witter, M.P., Moser, E.I., Moser, M.-B., 2010. Grid cells in pre- and parasubiculum. *Nat. Neurosci.* 13 (8), 987–994.
- Bocian, R., Konopacki, J., 2007. Posterior hypothalamic GABAergic mediation of hippocampal theta in the cat. *Brain Res. Bull.* 73 (4–6), 289–300.
- Bokor, H., Csáki, A., Kocsis, K., Kiss, J., 2002. Cellular architecture of the nucleus reuniens thalami and its putative aspartatergic/glutamatergic projection to the hippocampus and medial septum in the rat. *Eur. J. Neurosci.* 16 (7), 1227–1239.
- Borhegyi, Z., Freund, T.F., 1998. Dual projection from the medial septum to the supramammillary nucleus in the rat. *Brain Res. Bull.* 46 (5), 453–459.
- Borhegyi, Z., Maglóczy, Z., Acsády, L., Freund, T.F., 1998. The supramammillary nucleus innervates cholinergic and GABAergic neurons in the medial septum-diagonal band of Broca complex. *Neuroscience* 82 (4), 1053–1065.
- Bradley, P.B., Nicholson, A.N., 1962. The effect of some drugs on hippocampal arousal. *Electroencephalogr. Clin. Neurophysiol.* 14, 824–834.
- Brandon, M.P., Bogaard, A.R., Libby, C.P., Connerney, M.A., Gupta, K., Hasselmo, M.E., 2011. Reduction of theta rhythm dissociates grid cell spatial periodicity from directional tuning. *Science* 332 (6029), 595–599.
- Buzsáki, G., 2002. Theta oscillations in the hippocampus. *Neuron*.
- Buzsáki, G., Buhl, D.L., Harris, K.D., Csicsvari, J., Czéh, B., Morozov, A., 2003. Hippocampal network patterns of activity in the mouse. *Neuroscience* 116 (1), 201–211.
- Cacucci, F., Lever, C., Wills, T.J., Burgess, N., O'Keefe, J., 2004. Theta-modulated place-by-direction cells in the hippocampal formation in the rat. *J. Neurosci.* 24 (38), 8265–8277.
- Canteras, N.S., Simerly, R.B., Swanson, L.W., 1992. Connections of the posterior nucleus of the amygdala. *J. Comp. Neurol.* 324 (2), 143–179.



- Canteras, N.S., Swanson, L.W., 1992. Projections of the ventral subiculum to the amygdala, septum, and hypothalamus: a PHAL anterograde tract-tracing study in the rat. *J. Comp. Neurol.* 324 (2), 180–194.
- Cantero, J.L., Atienza, M., Stickgold, R., Kahana, M.J., Madsen, J.R., Kocsis, B., 2003. Sleep-dependent theta oscillations in the human hippocampus and neocortex. *J. Neurosci.* 23 (34), 10897–10903.
- Cassell, M.D., Wright, D.J., 1986. Topography of projections from the medial prefrontal cortex to the amygdala in the rat. *Brain Res. Bull.* 17 (3), 321–333.
- Chrobak, J.J., Buzsáki, G., 1994. Selective activation of deep layer (V–VI) retrohippocampal cortical neurons during hippocampal sharp waves in the behaving rat. *J. Neurosci.* 14 (10), 6160–6170.
- Cicchetti, F., Lacroix, S., Beach, T.G., Parent, A., 1998. Calretinin gene expression in the human thalamus. *Brain Res. Mol. Brain Res.* 54 (1), 1–12.
- Clarke, D.J., 1985. Cholinergic innervation of the rat dentate gyrus: an immunocytochemical and electron microscopical study. *Brain Res.* 360 (1–2), 349–354.
- Clement, E.A., Richard, A., Thwaites, M., Ailon, J., Peters, S., Dickson, C.T., 2008. Cyclic and sleep-like spontaneous alternations of brain state under urethane anaesthesia. *PLoS ONE* 3 (4), e2004.
- Colom, L.V., Bland, B.H., 1987. State-dependent spike train dynamics of hippocampal formation neurons: evidence for theta-on and theta-off cells. *Brain Res.* 422 (2), 277–286.
- Colom, L.V., Castaneda, M.T., Reyna, T., Hernandez, S., Garrido-Sanabria, E., 2005. Characterization of medial septal glutamatergic neurons and their projection to the hippocampus. *Synapse* 58 (3), 151–164.
- Condé, F., Audinat, E., Maire-Lepoivre, E., Crépel, F., 1990. Afferent connections of the medial frontal cortex of the rat. A study using retrograde transport of fluorescent dyes. I. Thalamic afferents. *Brain Res. Bull.* 24 (3), 341–354.
- Condé, F., Maire-Lepoivre, E., Audinat, E., Crépel, F., 1995. Afferent connections of the medial frontal cortex of the rat. II. Cortical and subcortical afferents. *J. Comp. Neurol.* 352 (4), 567–593.
- Crutcher, K.A., Madison, R., Davis, J.N., 1981. A study of the rat septohippocampal pathway using anterograde transport of horseradish peroxidase. *Neuroscience* 6 (10), 1961–1973.
- Csicsvari, J., Henze, D.A., Jamieson, B., Harris, K.D., Sirota, A., Barthó, P., Wise, K.D., Buzsáki, G., 2003. Massively parallel recording of unit and local field potentials with silicon-based electrodes. *J. Neurophysiol.* 90 (2), 1314–1323.
- Csicsvari, J., Hirase, H., Mamiya, A., Buzsáki, G., 2000. Ensemble patterns of hippocampal CA3–CA1 neurons during sharp wave-associated population events. *Neuron* 28 (2), 585–594.
- Datta, S., Siwek, D.F., 1997. Excitation of the brain stem pedunculopontine tegmentum cholinergic cells induces wakefulness and ... *J. Neurophysiol.*
- Deshmukh, S.S., Yoganarasimha, D., Voicu, H., Knierim, J.J., 2010. Theta modulation in the medial and the lateral entorhinal cortices. *J. Neurophysiol.* 104 (2), 994–1006.
- Dickson, C.T., Kirk, I.J., Oddie, S.D., Bland, B.H., 1995. Classification of theta-related cells in the entorhinal cortex: cell discharges are controlled by the ascending brainstem synchronizing pathway in parallel with hippocampal theta-related cells. *Hippocampus* 5 (4), 306–319.
- Dickson, C.T., Magistretti, J., Shalinsky, M.H., Fransén, E., Hasselmo, M.E., Alonso, A., 2000. Properties and role of I(h) in the pacing of subthreshold oscillations in entorhinal cortex layer II neurons. *J. Neurophysiol.* 83 (5), 2562–2579.
- Dolleman-Van der Weel, M.J., Witter, M.P., 1996. Projections from the nucleus reuniens thalami to the entorhinal cortex, hippocampal field CA1, and the subiculum in the rat arise from different populations of neurons. *J. Comp. Neurol.* 364 (4), 637–650.
- Dolleman-Van der Weel, M.J., Witter, M.P., 2000. Nucleus reuniens thalami innervates gamma aminobutyric acid positive cells in hippocampal field CA1 of the rat. *Neurosci. Lett.* 278 (3), 145–148.
- Dolorfo, C.L., Amaral, D.G., 1998. Entorhinal cortex of the rat: topographic organization of the cells of origin of the perforant path projection to the dentate gyrus. *J. Comp. Neurol.* 398 (1), 25–48.
- Dougherty, K.D., Milner, T.A., 1999. Cholinergic septal afferent terminals preferentially contact neuropeptide Y-containing interneurons compared to parvalbumin-containing interneurons in the rat dentate gyrus. *J. Neurosci.* 19 (22), 10140–10152.
- Dragoi, G., Buzsáki, G., 2006. Temporal encoding of place sequences by hippocampal cell assemblies. *Neuron* 50 (1), 145–157.
- Dragoi, G., Carpi, D., Recce, M., Csicsvari, J., Buzsáki, G., 1999. Interactions between hippocampus and medial septum during sharp waves and theta oscillation in the behaving rat. *J. Neurosci.* 19 (14), 6191–6199.
- Egorov, A.V., Hamam, B.N., Fransén, E., Hasselmo, M.E., Alonso, A.A., 2002. Graded persistent activity in entorhinal cortex neurons. *Nature* 420 (6912), 173–178.
- Freund, T.F., 1989. GABAergic septohippocampal neurons contain parvalbumin. *Brain Res.* 478 (2), 375–381.
- Freund, T.F., Antal, M., 1988. GABA-containing neurons in the septum control inhibitory interneurons in the hippocampus. *Nature* 336 (6195), 170–173.
- Frotscher, M., 1991. Target cell specificity of synaptic connections in the hippocampus. *Hippocampus* 1 (2), 123–130.
- Fuller, P.M., Saper, C.B., Lu, J., 2007. The pontine REM switch: past and present. *J. Physiol. (Lond.)* 584 (Pt. 3), 735–741.
- Gabbott, P., Headlam, A., Busby, S., 2002. Morphological evidence that CA1 hippocampal afferents monosynaptically innervate PV-containing neurons and NADPH-diaphorase reactive cells in the medial prefrontal cortex (Areas 25/32) of the rat. *Brain Res.* 946 (2), 314–322.
- Gabbott, P.L.A., Warner, T.A., Jays, P.R.L., Salway, P., Busby, S.J., 2005. Prefrontal cortex in the rat: projections to subcortical autonomic, motor, and limbic centers. *J. Comp. Neurol.* 492 (2), 145–177.
- Gaykema, R.P., Luiten, P.G., Nyakas, C., Traber, J., 1990. Cortical projection patterns of the medial septum-diagonal band complex. *J. Comp. Neurol.* 293 (1), 103–124.
- Gaykema, R.P., van Weeghel, R., Hersh, L.B., Luiten, P.G., 1991. Prefrontal cortical projections to the cholinergic neurons in the basal forebrain. *J. Comp. Neurol.* 303 (4), 563–583.
- Glasgow, S.D., Chapman, C.A., 2007. Local generation of theta-frequency EEG activity in the parasubiculum. *J. Neurophysiol.* 97 (6), 3868–3879.
- Glasgow, S.D., Chapman, C.A., 2008. Conductances mediating intrinsic theta-frequency membrane potential oscillations in layer II parasubicular neurons. *J. Neurophysiol.* 100 (5), 2746–2756.
- Golebiewski, H., Eckersdorf, B., Błaszczyk, M., Grabowski, R., Konopacki, J., 1994. Muscarinic (M1) mediation of carbachol-induced theta in the rat entorhinal cortex in vitro. *Neuroreport* 5 (15), 1989–1992.
- Gonzalo-Ruiz, A., Morte, L., Flecha, J.M., Sanz, J.M., 1999. Neurotransmitter characteristics of neurons projecting to the supramammillary nucleus of the rat. *Anat. Embryol.* 200 (4), 377–392.
- Goto, M., Swanson, L.W., Canteras, N.S., 2001. Connections of the nucleus incertus. *J. Comp. Neurol.* 438 (1), 86–122.
- Goutagny, R., Jackson, J., Williams, S., 2009. Self-generated theta oscillations in the hippocampus. *Nat. Neurosci.* 12 (12), 1491–1493.
- Gradinaru, V., Zhang, F., Ramakrishnan, C., Mattis, J., Prakash, R., Diester, I., Goshen, I., Thompson, K.R., Deisseroth, K., 2010. Molecular and cellular approaches for diversifying and extending optogenetics. *Cell* 141 (1), 154–165.
- Green, J.D., Arduini, A.A., 1954. Hippocampal electrical activity in arousal. *J. Neurophysiol.* 17 (6), 533–557.
- Hafting, T., Fyhn, M., Bonnevie, T., Moser, M.-B., Moser, E.I., 2008. Hippocampus-independent phase precession in entorhinal grid cells. *Nature* 453 (7199), 1248–1252.
- Hafting, T., Fyhn, M., Molden, S., Moser, M.-B., Moser, E.I., 2005. Microstructure of a spatial map in the entorhinal cortex. *Nature* 436 (7052), 801–806.
- Hajszan, T., Alreja, M., Leranth, C., 2004. Intrinsic vesicular glutamate transporter 2-immunoreactive input to septohippocampal parvalbumin-containing neurons: novel glutamatergic local circuit cells. *Hippocampus* 14 (4), 499–509.
- Hallanger, A.E., Levey, A.I., Lee, H.J., Rye, D.B., Wainer, B.H., 1987. The origins of cholinergic and other subcortical afferents to the thalamus in the rat. *J. Comp. Neurol.* 262 (1), 105–124.
- Hallanger, A.E., Wainer, B.H., 1988. Ascending projections from the pedunculopontine tegmental nucleus and the adjacent mesopontine tegmentum in the rat. *J. Comp. Neurol.* 274 (4), 483–515.
- Hangya, B., Borhegyi, Z., Szilágyi, N., Freund, T.F., Varga, V., 2009. GABAergic neurons of the medial septum lead the hippocampal network during theta activity. *J. Neurosci.* 29 (25), 8094–8102.
- Hartwich, K., Pollak, T., Klausberger, T., 2009. Distinct firing patterns of identified basket and dendrite-targeting interneurons in the prefrontal cortex during hippocampal theta and local spindle oscillations. *J. Neurosci.* 29 (30), 9563–9574.
- Hinman, J.R., Penley, S.C., Long, L.L., Escabi, M.A., Chrobak, J.J., 2011. Septotemporal variation in dynamics of theta: speed and habituation. *J. Neurophysiol.* 105 (6), 2675–2686.
- Huh, C.Y.L., Goutagny, R., Williams, S., 2010. Glutamatergic neurons of the mouse medial septum and diagonal band of Broca synaptically drive hippocampal pyramidal cells: relevance for hippocampal theta rhythm. *J. Neurosci.* 30 (47), 15951–15961.
- Hyman, J.M., Zilli, E.A., Paley, A.M., Hasselmo, M.E., 2005. Medial prefrontal cortex cells show dynamic modulation with the hippocampal theta rhythm dependent on behavior. *Hippocampus* 15 (6), 739–749.
- Inglis, W.L., Winn, P., 1995. The pedunculopontine tegmental nucleus: where the striatum meets the reticular formation. *Prog. Neurobiol.* 47 (1), 1–29.
- Irlé, E., Sarter, M., Guldin, W.O., Markowitsch, H.J., 1984. Afferents to the ventral tegmental nucleus of Gudden in the mouse, rat, and cat. *J. Comp. Neurol.* 228 (4), 509–541.
- Jackson, J., Dickson, C.T., Bland, B.H., 2008. Median raphe stimulation disrupts hippocampal theta via rapid inhibition and state-dependent phase reset of theta-related neural circuitry. *J. Neurophysiol.* 99 (6), 3009–3026.
- Jasper, H.H., Tessier, J., 1971. Acetylcholine liberation from cerebral cortex during paradoxical (REM) sleep. *Science* 172 (983), 601–602.
- Jeffery, K.J., Donnet, J.G., O'Keefe, J., 1995. Medial septal control of theta-correlated unit firing in the entorhinal cortex of awake rats. *Neuroreport* 6 (16), 2166–2170.
- Jensen, O., 2005. Reading the hippocampal code by theta phase-locking. *Trends Cogn. Sci. (Regul. Ed.)* 9 (12), 551–553.
- Jolkkonen, E., Miettinen, R., Pikkarainen, M., Pitkänen, A., 2002. Projections from the amygdaloid complex to the magnocellular cholinergic basal forebrain in rat. *Neuroscience* 111 (1), 133–149.
- Jones, M.W., Wilson, M.A., 2005a. Phase precession of medial prefrontal cortical activity relative to the hippocampal theta rhythm. *Hippocampus* 15 (7), 867–873.
- Jones, M.W., Wilson, M.A., 2005b. Theta rhythms coordinate hippocampal–prefrontal interactions in a spatial memory task. *PLoS Biol* 3 (12), e402.
- Kahana, M.J., Seelig, D., Madsen, J.R., 2001. Theta returns. *Curr. Opin. Neurobiol.* 11 (6), 739–744.

- Kahana, M.J., Sekuler, R., Caplan, J.B., Kirschen, M., Madsen, J.R., 1999. Human theta oscillations exhibit task dependence during virtual maze navigation. *Nature* 399 (6738), 781–784.
- Kamondi, A., Acsády, L., Wang, X.J., Buzsáki, G., 1998. Theta oscillations in somata and dendrites of hippocampal pyramidal cells in vivo: activity-dependent phase-precession of action potentials. *Hippocampus* 8 (3), 244–261.
- Karashima, A., Nakao, M., Honda, K., Iwasaki, N., Katayama, N., Yamamoto, M., 2004. Theta wave amplitude and frequency are differentially correlated with pontine waves and rapid eye movements during REM sleep in rats. *Neurosci. Res.* 50 (3), 283–289.
- Karashima, A., Nakao, M., Katayama, N., Honda, K., 2005. Instantaneous acceleration and amplification of hippocampal theta wave coincident with phasic pontine activities during REM sleep. *Brain Res.* 1051 (1–2), 50–56.
- Kempainen, S., Jolkkonen, E., Pitkänen, A., 2002. Projections from the posterior cortical nucleus of the amygdala to the hippocampal formation and parahippocampal region in rat. *Hippocampus* 12 (6), 735–755.
- Kinney, G.G., Kocsis, B., Vertes, R.P., 1995. Injections of muscimol into the median raphe nucleus produce hippocampal theta rhythm in the urethane anesthetized rat. *Psychopharmacology (Berl.)* 120 (3), 244–248.
- Kirk, I.J., McNaughton, N., 1991. Supramammillary cell firing and hippocampal rhythmic slow activity. *Neuroreport* 2 (11), 723–725.
- Kirk, I.J., McNaughton, N., 1993. Mapping the differential effects of procaine on frequency and amplitude of rhythmically elicited hippocampal rhythmic slow activity. *Hippocampus* 3 (4), 517–525.
- Kirk, I.J., Oddie, S.D., Konopacki, J., Bland, B.H., 1996. Evidence for differential control of posterior hypothalamic, supramammillary, and medial mammillary theta-related cellular discharge by ascending and descending pathways. *J. Neurosci.* 16 (17), 5547–5554.
- Kishi, T., Tsumori, T., Yokota, S., Yasui, Y., 2006. Topographical projection from the hippocampal formation to the amygdala: a combined anterograde and retrograde tracing study in the rat. *J. Comp. Neurol.* 496 (3), 349–368.
- Kiss, J., Csáki, A., Bokor, H., Kocsis, K., Kocsis, B., 2002. Possible glutamatergic/aspartatergic projections to the supramammillary nucleus and their origins in the rat studied by selective [(3)H]D-aspartate labelling and immunocytochemistry. *Neuroscience* 111 (3), 671–691.
- Kiss, J., Patel, A.J., Baimbridge, K.G., Freund, T.F., 1990. Topographical localization of neurons containing parvalbumin and choline acetyltransferase in the medial septum-diagonal band region of the rat. *Neuroscience* 36 (1), 61–72.
- Klausberger, T., Magill, P.J., Márton, L.F., Roberts, J.D.B., Cobden, P.M., Buzsáki, G., Somogyi, P., 2003. Brain-state- and cell-type-specific firing of hippocampal interneurons in vivo. *Nature* 421 (6925), 844–848.
- Kocsis, B., 2006. The effect of descending theta rhythmic input from the septohippocampal system on firing in the supramammillary nucleus. *Brain Res.* 1086 (1), 92–97.
- Kocsis, B., Di Prisco, G.V., Vertes, R.P., 2001. Theta synchronization in the limbic system: the role of Gudden's tegmental nuclei. *Eur. J. Neurosci.* 13 (2), 381–388.
- Kocsis, B., Vertes, R.P., 1992. Dorsal raphe neurons: synchronous discharge with the theta rhythm of the hippocampus in the freely behaving rat. *J. Neurophysiol.* 68 (4), 1463–1467.
- Kocsis, B., Vertes, R.P., 1994. Characterization of neurons of the supramammillary nucleus and mammillary body that discharge rhythmically with the hippocampal theta rhythm in the rat. *J. Neurosci.* 14 (11 Pt. 2), 7040–7052.
- Koenig, J., Linder, A.N., Leutgeb, J.K., Leutgeb, S., 2011. The spatial periodicity of grid cells is not sustained during reduced theta oscillations. *Science* 332 (6029), 592–595.
- Konopacki, J., Bland, B.H., MacIver, M.B., Roth, S.H., 1987a. Cholinergic theta rhythm in transected hippocampal slices: independent CA1 and dentate generators. *Brain Res.* 436 (2), 217–222.
- Konopacki, J., Gołebiewski, H., 1992. Theta rhythms in the rat medial entorhinal cortex in vitro: evidence for involvement of muscarinic receptors. *Neurosci. Lett.* 141 (1), 93–96.
- Konopacki, J., MacIver, M.B., Bland, B.H., Roth, S.H., 1987b. Carbachol-induced EEG 'theta' activity in hippocampal brain slices. *Brain Res.* 405 (1), 196–198.
- Kramis, R., Vanderwolf, C.H., Bland, B.H., 1975. Two types of hippocampal rhythmic slow activity in both the rabbit and the rat: relations to behavior and effects of atropine, diethyl ether, urethane, and pentobarbital. *Exp. Neurol.* 49 (1 Pt. 1), 58–85.
- Lee, H.J., Rye, D.B., Hallanger, A.E., Levey, A.I., Wainer, B.H., 1988. Cholinergic vs. noncholinergic efferents from the mesopontine tegmentum to the extrapyramidal motor system nuclei. *J. Comp. Neurol.* 275 (4), 469–492.
- Lee, M.G., Chrobak, J.J., Sik, A., Wiley, R.G., Buzsáki, G., 1994. Hippocampal theta activity following selective lesion of the septal cholinergic system. *Neuroscience* 62 (4), 1033–1047.
- Leranth, C., Frotscher, M., 1989. Organization of the septal region in the rat brain: cholinergic-GABAergic interconnections and the termination of hippocampo-septal fibers. *J. Comp. Neurol.* 289 (2), 304–314.
- Lesting, J., Narayanan, R.T., Kluge, C., Sangha, S., Seidenbecher, T., Pape, H.-C., 2011. Patterns of coupled theta activity in amygdala–hippocampal–prefrontal cortical circuits during fear extinction. *PLoS ONE* 6 (6), e21714.
- Li, S., Topchy, I., Kocsis, B., 2007. The effect of atropine administered in the medial septum or hippocampus on high- and low-frequency theta rhythms in the hippocampus of urethane anesthetized rats. *Synapse* 61 (6), 412–419.
- Lu, J., Sherman, D., Devor, M., Saper, C.B., 2006. A putative flip-flop switch for control of REM sleep. *Nature* 441 (7093), 589–594.
- Lubenov, E.V., Siapas, A.G., 2009. Hippocampal theta oscillations are travelling waves. *Nature* 459 (7246), 534–539.
- Maglóczy, Z., Acsády, L., Freund, T.F., 1994. Principal cells are the postsynaptic targets of supramammillary afferents in the hippocampus of the rat. *Hippocampus* 4 (3), 322–334.
- Manns, I.D., Mainville, L., Jones, B.E., 2001. Evidence for glutamate, in addition to acetylcholine and GABA, neurotransmitter synthesis in basal forebrain neurons projecting to the entorhinal cortex. *Neuroscience* 107 (2), 249–263.
- Markram, H., Segal, M., 1990. Electrophysiological characteristics of cholinergic and non-cholinergic neurons in the rat medial septum-diagonal band complex. *Brain Res.* 513 (1), 171–174.
- McDonald, A.J., 1998. Cortical pathways to the mammalian amygdala. *Prog. Neurobiol.* 55 (3), 257–332.
- McDonald, A.J., Mascagni, F., 1997. Projections of the lateral entorhinal cortex to the amygdala: a Phaseolus vulgaris leucoagglutinin study in the rat. *Neuroscience* 77 (2), 445–459.
- McDonald, A.J., Mascagni, F., Guo, L., 1996. Projections of the medial and lateral prefrontal cortices to the amygdala: a Phaseolus vulgaris leucoagglutinin study in the rat. *Neuroscience* 71 (1), 55–75.
- McKenna, J.T., Vertes, R.P., 2004. Afferent projections to nucleus reuniens of the thalamus. *J. Comp. Neurol.* 480 (2), 115–142.
- Meibach, R.C., Siegel, A., 1975. The origin of fornix fibers which project to the mammillary bodies in the rat: a horseradish peroxidase study. *Brain Res.* 88 (3), 508–512.
- Meibach, R.C., Siegel, A., 1977. Efferent connections of the septal area in the rat: an analysis utilizing retrograde and anterograde transport methods. *Brain Res.* 119 (1), 1–20.
- Mesulam, M.M., Mufson, E.J., Wainer, B.H., Levey, A.I., 1983. Central cholinergic pathways in the rat: an overview based on an alternative nomenclature (Ch1–Ch6). *Neuroscience* 10 (4), 1185–1201.
- Milner, K.L., Mogenson, G.J., 1988. Electrical and chemical activation of the mesencephalic and subthalamic locomotor regions in freely moving rats. *Brain Res.* 452 (1–2), 273–285.
- Mitani, A., Ito, K., Hallanger, A.E., Wainer, B.H., Kataoka, K., McCarley, R.W., 1988. Cholinergic projections from the laterodorsal and pedunculopontine tegmental nuclei to the pontine gigantocellular tegmental field in the cat. *Brain Res.* 451 (1–2), 397–402.
- Mitchell, S.J., Ranck, J.B., 1980. Generation of theta rhythm in medial entorhinal cortex of freely moving rats. *Brain Res.* 189 (1), 49–66.
- Mitchell, S.J., Rawlins, J.N., Steward, O., Olton, D.S., 1982. Medial septal area lesions disrupt theta rhythm and cholinergic staining in medial entorhinal cortex and produce impaired radial arm maze behavior in rats. *J. Neurosci.* 2 (3), 292–302.
- Mizumori, S.J., Perez, G.M., Alvarado, M.C., Barnes, C.A., McNaughton, B.L., 1990. Reversible inactivation of the medial septum differentially affects two forms of learning in rats. *Brain Res.* 528 (1), 12–20.
- Mizumori, S.J., Ward, K.E., Lavoie, A.M., 1992. Medial septal modulation of entorhinal single unit activity in anesthetized and freely moving rats. *Brain Res.* 570 (1–2), 188–197.
- Mizuseki, K., Diba, K., Pastalkova, E., Buzsáki, G., 2011. Hippocampal CA1 pyramidal cells form functionally distinct sublayers. *Nat. Neurosci.* 14 (9), 1174–1181.
- Mizuseki, K., Sirota, A., Pastalkova, E., Buzsáki, G., 2009. Theta oscillations provide temporal windows for local circuit computation in the entorhinal–hippocampal loop. *Neuron* 64 (2), 267–280.
- Montgomery, S.M., Sirota, A., Buzsáki, G., 2008. Theta and gamma coordination of hippocampal networks during waking and rapid eye movement sleep. *J. Neurosci.* 28 (26), 6731–6741.
- Montoya, C.P., Sainsbury, R.S., 1985. The effects of entorhinal cortex lesions on type 1 and type 2 theta. *Physiol. Behav.* 35 (1), 121–126.
- Morales, G.J., Ramcharan, E.J., Sundaraman, N., Morgera, S.D., Vertes, R.P., 2007. Analysis of the actions of nucleus reuniens and the entorhinal cortex on EEG and evoked population behavior of the hippocampus. In: Conference Proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society IEEE Engineering in Medicine and Biology Society Conference 2007, pp. 2480–2484.
- Moruzzi, G., Magoun, H.W., 1949. Brain stem reticular formation and activation of the EEG. *Electroencephalogr. Clin. Neurophysiol.* 1 (4), 455–473.
- Muñoz, M.D., Núñez, A., García-Austt, E., 1990. In vivo intracellular analysis of rat dentate granule cells. *Brain Res.* 509 (1), 91–98.
- Newman, D.B., Ginsberg, C.Y., 1994. Brainstem reticular nuclei that project to the thalamus in rats: a retrograde tracer study. *Brain Behav. Evol.* 44 (1), 1–39.
- Nowacka, A., Jurkowlaniec, E., Trojnar, W., 2002. Microinjection of procaine into the pedunculopontine tegmental nucleus suppresses hippocampal theta rhythm in urethane-anesthetized rats. *Brain Res. Bull.* 58 (4), 377–384.
- Núñez, A., Cervera-Ferri, A., Olucha-Bordonau, F., Ruiz-Torner, A., Teruel, V., 2006. Nucleus incertus contribution to hippocampal theta rhythm generation. *Eur. J. Neurosci.* 23 (10), 2731–2738.
- Oddie, S.D., Bland, B.H., Colom, L.V., Vertes, R.P., 1994. The midline posterior hypothalamic region comprises a critical part of the ascending brainstem hippocampal synchronizing pathway. *Hippocampus* 4 (4), 454–473.
- Otake, K., Reis, D.J., Ruggiero, D.A., 1994. Afferents to the midline thalamus issue collaterals to the nucleus tractus solitarius: an anatomical basis for thalamic and visceral reflex integration. *J. Neurosci.* 14 (9), 5694–5707.
- Paré, D., Gaudreau, H., 1996. Projection cells and interneurons of the lateral and basolateral amygdala: distinct firing patterns and differential relation to theta and delta rhythms in conscious cats. *J. Neurosci.* 16 (10), 3334–3350.
- Paz, R., Bauer, E.P., Paré, D., 2008. Theta synchronizes the activity of medial prefrontal neurons during learning. *Learn. Mem.* 15 (7), 524–531.

- Peck, B.K., Vanderwolf, C.H., 1991. Effects of raphe stimulation on hippocampal and neocortical activity and behaviour. *Brain Res.* 568 (1–2), 244–252.
- Petrovich, G.D., Risold, P.Y., Swanson, L.W., 1996. Organization of projections from the basomedial nucleus of the amygdala: a PHAL study in the rat. *J. Comp. Neurol.* 374 (3), 387–420.
- Petsche, H., Stumpf, C., Gogolak, G., 1962. The significance of the rabbit's septum as a relay station between the midbrain and the hippocampus. I. The control of hippocampus arousal activity by the septum cells. *Electroencephalogr. Clin. Neurophysiol.* 14, 202–211.
- Peyron, C., Petit, J.M., Rampon, C., Jouvet, M., Luppi, P.H., 1998. Forebrain afferents to the rat dorsal raphe nucleus demonstrated by retrograde and anterograde tracing methods. *Neuroscience* 82 (2), 443–468.
- Pikkarainen, M., Rönkkö, S., Savander, V., Insausti, R., Pitkänen, A., 1999. Projections from the lateral, basal, and accessory basal nuclei of the amygdala to the hippocampal formation in rat. *J. Comp. Neurol.* 403 (2), 229–260.
- Popa, D., Duvarci, S., Popescu, A.T., Léna, C., Paré, D., 2010. Coherent amygdalocortical theta promotes fear memory consolidation during paradoxical sleep. *Proc. Natl. Acad. Sci. USA* 107 (14), 6516–6519.
- Poulet, J., Petersen, C., 2008. Internal brain state regulates membrane potential synchrony in barrel cortex of behaving mice. *Nature*.
- Robinson, T.E., Vanderwolf, C.H., 1978. Electrical stimulation of the brain stem in freely moving rats: II. Effects on hippocampal and neocortical electrical activity, and relations to behavior. *Exp. Neurol.* 61 (3), 485–515.
- Royer, S., Sirota, A., Patel, J., Buzsáki, G., 2010. Distinct representations and theta dynamics in dorsal and ventral hippocampus. *J. Neurosci.* 30 (5), 1777–1787.
- Ryan, P.J., Ma, S., Olucha-Bordonau, F.E., Gundlach, A.L., 2011. Nucleus incertus – an emerging modulatory role in arousal, stress and memory. *Neurosci. Biobehav. Rev.* 35 (6), 1326–1341.
- Rye, D.B., Lee, H.J., Saper, C.B., Wainer, B.H., 1988. Medullary and spinal efferents of the pedunculopontine tegmental nucleus and adjacent mesopontine tegmentum in the rat. *J. Comp. Neurol.* 269 (3), 315–341.
- Rye, D.B., Saper, C.B., Lee, H.J., Wainer, B.H., 1987. Pedunculopontine tegmental nucleus of the rat: cytoarchitecture, cytochemistry, and some extrapyramidal connections of the mesopontine tegmentum. *J. Comp. Neurol.* 259 (4), 483–528.
- Sabolek, H.R., Penley, S.C., Hinman, J.R., Bunce, J.G., Markus, E.J., Escabi, M., Chrobak, J.J., 2009. Theta and gamma coherence along the septotemporal axis of the hippocampus. *J. Neurophysiol.* 101 (3), 1192–1200.
- Saper, C.B., Fuller, P.M., Pedersen, N.P., Lu, J., Scammell, T.E., 2010. Sleep state switching. *Neuron* 68 (6), 1023–1042.
- Satoh, K., Fibiger, H.C., 1986. Cholinergic neurons of the laterodorsal tegmental nucleus: efferent and afferent connections. *J. Comp. Neurol.* 253 (3), 277–302.
- Schmitz, D., Gloveli, T., Behr, J., Dugladze, T., Heinemann, U., 1998. Subthreshold membrane potential oscillations in neurons of deep layers of the entorhinal cortex. *Neuroscience* 85 (4), 999–1004.
- Seidenbecher, T., Laxmi, T.R., Stork, O., Pape, H.-C., 2003. Amygdalar and hippocampal theta rhythm synchronization during fear memory retrieval. *Science* 301 (5634), 846–850.
- Semba, K., Fibiger, H.C., 1992. Afferent connections of the laterodorsal and the pedunculopontine tegmental nuclei in the rat: a retro- and antero-grade transport and immunohistochemical study. *J. Comp. Neurol.* 323 (3), 387–410.
- Semba, K., Reiner, P.B., Fibiger, H.C., 1990. Single cholinergic mesopontine tegmental neurons project to both the pontine reticular formation and the thalamus in the rat. *Neuroscience* 38 (3), 643–654.
- Sesack, S.R., Deutch, A.Y., Roth, R.H., Bunney, B.S., 1989. Topographical organization of the efferent projections of the medial prefrontal cortex in the rat: an anterograde tract-tracing study with Phaseolus vulgaris leucoagglutinin. *J. Comp. Neurol.* 290 (2), 213–242.
- Sesack, S.R., Hawrylyk, V.A., Matus, C., Guido, M.A., Levey, A.I., 1998. Dopamine axon varicosities in the prelimbic division of the rat prefrontal cortex exhibit sparse immunoreactivity for the dopamine transporter. *J. Neurosci.* 18 (7), 2697–2708.
- Shiromani, P.J., Armstrong, D.M., Gillin, J.C., 1988. Cholinergic neurons from the dorsolateral pons project to the medial pons: a WGA-HRP and choline acetyltransferase immunohistochemical study. *Neurosci. Lett.* 95 (1–3), 19–23.
- Siapas, A.G., Lubenov, E.V., Wilson, M.A., 2005. Prefrontal phase locking to hippocampal theta oscillations. *Neuron* 46 (1), 141–151.
- Sik, A., Penttonen, M., Buzsáki, G., 1997. Interneurons in the hippocampal dentate gyrus: an in vivo intracellular study. *Eur. J. Neurosci.* 9 (3), 573–588.
- Sirota, A., Montgomery, S., Fujisawa, S., Isomura, Y., Zugaro, M., Buzsáki, G., 2008. Entrainment of neocortical neurons and gamma oscillations by the hippocampal theta rhythm. *Neuron* 60 (4), 683–697.
- Skinner, R.D., Garcia-Rill, E., 1984. The mesencephalic locomotor region (MLR) in the rat. *Brain Res.* 323 (2), 385–389.
- Sparks, P.D., LeDoux, J.E., 1995. Septal lesions potentiate freezing behavior to contextual but not to phasic conditioned stimuli in rats. *Behav. Neurosci.* 109 (1), 184–188.
- Steinbusch, H.W., Nieuwenhuys, R., 1981. Localization of serotonin-like immunoreactivity in the central nervous system and pituitary of the rat, with special references to the innervation of the hypothalamus. *Adv. Exp. Med. Biol.* 133, 7–35.
- Steininger, T.L., Rye, D.B., Wainer, B.H., 1992. Afferent projections to the cholinergic pedunculopontine tegmental nucleus and adjacent midbrain extrapyramidal area in the albino rat. I. Retrograde tracing studies. *J. Comp. Neurol.* 321 (4), 515–543.
- Steriade, M., Datta, S., Paré, D., Oakson, G., Curró Dossi, R.C., 1990. Neuronal activities in brain-stem cholinergic nuclei related to tonic activation processes in thalamocortical systems. *J. Neurosci.* 10 (8), 2541–2559.
- Stewart, M., Quirk, G.J., Barry, M., Fox, S.E., 1992. Firing relations of medial entorhinal neurons to the hippocampal theta rhythm in urethane anesthetized and walking rats. *Exp. Brain Res. Experimentelle Hirnforschung Expérimentation cérébrale* 90 (1), 21–28.
- Su, H.S., Bentivoglio, M., 1990. Thalamic midline cell populations projecting to the nucleus accumbens, amygdala, and hippocampus in the rat. *J. Comp. Neurol.* 297 (4), 582–593.
- Swanson, L.W., 1981. A direct projection from Ammon's horn to prefrontal cortex in the rat. *Brain Res.* 217 (1), 150–154.
- Thinschmidt, J.S., Kinney, G.G., Kocsis, B., 1995. The supramammillary nucleus: is it necessary for the mediation of hippocampal theta rhythm? *Neuroscience* 67 (2), 301–312.
- Tóth, K., Borhegyi, Z., Freund, T.F., 1993. Postsynaptic targets of GABAergic hippocampal neurons in the medial septum-diagonal band of Broca complex. *J. Neurosci.* 13 (9), 3712–3724.
- Tóth, K., Freund, T.F., Miles, R., 1997. Disinhibition of rat hippocampal pyramidal cells by GABAergic afferents from the septum. *J. Physiol. (Lond.)* 500 (Pt. 2), 463–474.
- Tsujimoto, T., Shimazu, H., Isomura, Y., 2006. Direct recording of theta oscillations in primate prefrontal and anterior cingulate cortices. *J. Neurophysiol.* 95 (5), 2987–3000.
- van der Meer, M.A.A., Redish, A.D., 2011. Theta phase precession in rat ventral striatum links place and reward information. *J. Neurosci.* 31 (8), 2843–2854.
- Van der Werf, Y.D., Witter, M.P., Groenewegen, H.J., 2002. The intralaminar and midline nuclei of the thalamus. Anatomical and functional evidence for participation in processes of arousal and awareness. *Brain Res. Brain Res. Rev.* 39 (2–3), 107–140.
- Van Groen, T., Miettinen, P., Kadish, I., 2003. The entorhinal cortex of the mouse: organization of the projection to the hippocampal formation. *Hippocampus* 13 (1), 133–149.
- Vanderwolf, C.H., 1969. Hippocampal electrical activity and voluntary movement in the rat. *Electroencephalogr. Clin. Neurophysiol.* 26 (4), 407–418.
- Vann, S.D., Aggleton, J.P., 2004. The mammillary bodies: two memory systems in one? *Nat. Rev. Neurosci.* 5 (1), 35–44.
- Varga, V., Hangya, B., Kránitz, K., Ludányi, A., Zemankovics, R., Katona, I., Shigemoto, R., Freund, T.F., Borhegyi, Z., 2008. The presence of pacemaker HCN channels identifies theta rhythmic GABAergic neurons in the medial septum. *J. Physiol. (Lond.)* 586 (16), 3893–3915.
- Vertes, R.P., 1977. Selective firing of rat pontine gigantocellular neurons during movement and REM sleep. *Brain Res.* 128 (1), 146–152.
- Vertes, R.P., 1979. Brain stem gigantocellular neurons: patterns of activity during behavior and sleep in the freely moving rat. *J. Neurophysiol.* 42 (1 Pt. 1), 214–228.
- Vertes, R.P., 1991. A PHA-L analysis of ascending projections of the dorsal raphe nucleus in the rat. *J. Comp. Neurol.* 313 (4), 643–668.
- Vertes, R.P., 1992. PHA-L analysis of projections from the supramammillary nucleus in the rat. *J. Comp. Neurol.* 326 (4), 595–622.
- Vertes, R.P., 2002. Analysis of projections from the medial prefrontal cortex to the thalamus in the rat, with emphasis on nucleus reuniens. *J. Comp. Neurol.* 442 (2), 163–187.
- Vertes, R.P., Albo, Z., Viana Di Prisco, G., 2001. Theta-rhythmically firing neurons in the anterior thalamus: implications for mnemonic functions of Papez's circuit. *Neuroscience* 104 (3), 619–625.
- Vertes, R.P., Crane, A.M., Colom, L.V., Bland, B.H., 1995. Ascending projections of the posterior nucleus of the hypothalamus: PHA-L analysis in the rat. *J. Comp. Neurol.* 359 (1), 90–116.
- Vertes, R.P., Hoover, W.B., Do Valle, A.C., Sherman, A., Rodriguez, J.J., 2006. Efferent projections of reuniens and rhomboid nuclei of the thalamus in the rat. *J. Comp. Neurol.* 499 (5), 768–796.
- Vertes, R.P., Hoover, W.B., Szigeti-Buck, K., Leranthy, C., 2007. Nucleus reuniens of the midline thalamus: link between the medial prefrontal cortex and the hippocampus. *Brain Res. Bull.* 71 (6), 601–609.
- Vertes, R.P., Kocsis, B., 1997. Brainstem-diencephalo-septohippocampal systems controlling the theta rhythm of the hippocampus. *Neuroscience* 81 (4), 893–926.
- Vertes, R.P., Martin, G.F., Waltzer, R., 1986. An autoradiographic analysis of ascending projections from the medullary reticular formation in the rat. *Neuroscience* 19 (3), 873–898.
- Wang, W.-T., Wan, Y.-H., Zhu, J.-L., Lei, G.-S., Wang, Y.-Y., Zhang, P., Hu, S.-J., 2006. Theta-frequency membrane resonance and its ionic mechanisms in rat subicular pyramidal neurons. *Neuroscience* 140 (1), 45–55.
- Winson, J., 1978. Loss of hippocampal theta rhythm results in spatial memory deficit in the rat. *Science* 201 (4351), 160–163.
- Woodnorth, M.-A., Kyd, R.J., Logan, B.J., Long, M.A., McNaughton, N., 2003. Multiple hypothalamic sites control the frequency of hippocampal theta rhythm. *Hippocampus* 13 (3), 361–374.
- Woolf, N.J., Butcher, L.L., 1986. Cholinergic systems in the rat brain: III. Projections from the pontomesencephalic tegmentum to the thalamus, tectum, basal ganglia, and basal forebrain. *Brain Res. Bull.* 16 (5), 603–637.
- Wouterlood, F.G., Aliane, V., Boekel, A.J., Hur, E.E., Zaborszky, L., Barroso-Chinea, P., Härtig, W., Lanciego, J.L., Witter, M.P., 2008. Origin of calretinin-containing, vesicular glutamate transporter 2-coexpressing fiber terminals in the entorhinal cortex of the rat. *J. Comp. Neurol.* 506 (2), 359–370.
- Wyble, B.P., Hyman, J.M., Rossi, C.A., Hasselmo, M.E., 2004. Analysis of theta power in hippocampal EEG during bar pressing and running behavior in rats during distinct behavioral contexts. *Hippocampus* 14 (5), 662–674.



- Ylinen, A., Soltész, I., Bragin, A., Penttonen, M., Sik, A., Buzsáki, G., 1995. Intracellular correlates of hippocampal theta rhythm in identified pyramidal cells, granule cells, and basket cells. *Hippocampus* 5 (1), 78–90.
- Yoder, R.M., Pang, K.C.H., 2005. Involvement of GABAergic and cholinergic medial septal neurons in hippocampal theta rhythm. *Hippocampus* 15 (3), 381–392.
- Zaborszky, L., Buhl, D.L., Pabalashingham, S., Bjaalie, J.G., Nadasdy, Z., 2005. Three-dimensional chemoarchitecture of the basal forebrain: spatially specific association of cholinergic and calcium binding protein-containing neurons. *Neuroscience* 136 (3), 697–713.
- Zhang, H., Lin, S.-C., Nicolelis, M.A.L., 2010. Spatiotemporal coupling between hippocampal acetylcholine release and theta oscillations in vivo. *J. Neurosci.* 30 (40), 13431–13440.
- Ziegler, D.R., Cullinan, W.E., Herman, J.P., 2002. Distribution of vesicular glutamate transporter mRNA in rat hypothalamus. *J. Comp. Neurol.* 448 (3), 217–229.