
November: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (135 mornings; 15 days missing from the archive)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in November


% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**86%**
2	**Punta Hermosa** Peru	**75%**
3	**Piha** New Zealand	**74%**
4	**Elands Bay** South Africa	**68%**
5	**Short Sands** Oregon	**68%**
6	**Mavericks** California	**68%**
7	**Lakey Peak** Sumbawa, Indonesia	**67%**
8	**Imsouane** Morocco	**63%**
9	**Chicama** Peru	**61%**
10	**Todos Santos** Baja California	**58%**
11	**Uluwatu** Bali, Indonesia	**57%**
12	**Punta de Lobos** Chile	**50%**
13	**Puerto Escondido** Oaxaca, Mexico	**49%**
14	**Hossegor** France	**48%**
15	**Shipstern Bluff** Tasmania, Australia	**48%**
16	**Bundoran** Ireland	**46%**
17	**Santa Catalina** Panama	**46%**
18	**Safi** Morocco	**46%**
19	**Rincon** California	**44%**
20	**Skeleton Bay** Namibia	**44%**
21	**Jeffreys Bay** South Africa	**44%**
22	**Ericeira** Portugal	**44%**
23	**Punta Roca** El Salvador	**43%**
24	**Montañita** Ecuador	**39%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Open-Meteo has no swell peak period on the Great Lakes; it is taken as 1.29× the mean swell period, the median ratio in the other seas.

December: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (155 mornings)



Best named breaks in December


% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**81%**
2	**Imsouane** Morocco	**74%**
3	**Elands Bay** South Africa	**71%**
4	**Punta Hermosa** Peru	**69%**
5	**Piha** New Zealand	**68%**
6	**Hossegor** France	**64%**
7	**Lakey Peak** Sumbawa, Indonesia	**63%**
8	**Short Sands** Oregon	**63%**
9	**Desert Point** Lombok, Indonesia	**62%**
10	**Mavericks** California	**60%**
11	**Todos Santos** Baja California	**60%**
12	**G-Land** Java, Indonesia	**59%**
13	**Chicama** Peru	**58%**
14	**Safi** Morocco	**54%**
15	**Ericeira** Portugal	**52%**
16	**Nias** Sumatra, Indonesia	**52%**
17	**Rincon** California	**48%**
18	**Jeffreys Bay** South Africa	**43%**
19	**Bundoran** Ireland	**42%**
20	**La Santa** Lanzarote, Canaries	**39%**
21	**Shipstern Bluff** Tasmania, Australia	**39%**
22	**Punta de Lobos** Chile	**38%**
23	**Skeleton Bay** Namibia	**37%**
24	**Margaret River** Western Australia	**35%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Open-Meteo has no swell peak period on the Great Lakes; it is taken as 1.29× the mean swell period, the median ratio in the other seas.

January: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Best named breaks in January


% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**86%**
2	**Short Sands** Oregon	**79%**
3	**Punta Hermosa** Peru	**77%**
4	**Todos Santos** Baja California	**77%**
5	**Anchor Point** Taghazout, Morocco	**71%**
6	**Mavericks** California	**70%**
7	**Elands Bay** South Africa	**68%**
8	**Rincon** California	**66%**
9	**Skeleton Bay** Namibia	**63%**
10	**Piha** New Zealand	**63%**
11	**Chicama** Peru	**59%**
12	**Hossegor** France	**58%**
13	**Bundoran** Ireland	**55%**
14	**Safi** Morocco	**52%**
15	**Lakey Peak** Sumbawa, Indonesia	**52%**
16	**Nias** Sumatra, Indonesia	**51%**
17	**Honolua Bay** Maui, Hawaii	**50%**
18	**Hanalei Bay** Kauai, Hawaii	**48%**
19	**Shipstern Bluff** Tasmania, Australia	**48%**
20	**Pipeline** Oahu, Hawaii	**47%**
21	**Sayulita** Nayarit, Mexico	**47%**
22	**Desert Point** Lombok, Indonesia	**47%**
23	**Montañita** Ecuador	**45%**
24	**Punta de Lobos** Chile	**44%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Open-Meteo has no swell peak period on the Great Lakes; it is taken as 1.29× the mean swell period, the median ratio in the other seas.

February: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (141 mornings; -1 days missing from the archive)




Best named breaks in February


% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**88%**
2	**Hossegor** France	**81%**
3	**Punta Hermosa** Peru	**80%**
4	**Short Sands** Oregon	**79%**
5	**Piha** New Zealand	**78%**
6	**Imsouane** Morocco	**73%**
7	**Elands Bay** South Africa	**71%**
8	**Todos Santos** Baja California	**70%**
9	**Chicama** Peru	**65%**
10	**Skeleton Bay** Namibia	**65%**
11	**Mavericks** California	**63%**
12	**Ericeira** Portugal	**57%**
13	**Rincon** California	**52%**
14	**Safi** Morocco	**51%**
15	**Bundoran** Ireland	**49%**
16	**Nias** Sumatra, Indonesia	**48%**
17	**Shipstern Bluff** Tasmania, Australia	**46%**
18	**Lakey Peak** Sumbawa, Indonesia	**43%**
19	**Tofino** British Columbia	**43%**
20	**Uluwatu** Bali, Indonesia	**43%**
21	**Jeffreys Bay** South Africa	**42%**
22	**Punta de Lobos** Chile	**40%**
23	**Sayulita** Nayarit, Mexico	**37%**
24	**Hanalei Bay** Kauai, Hawaii	**35%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Open-Meteo has no swell peak period on the Great Lakes; it is taken as 1.29× the mean swell period, the median ratio in the other seas.

March: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)


→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in March


% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**95%**
2	**Punta Hermosa** Peru	**84%**
3	**Chicama** Peru	**80%**
4	**Piha** New Zealand	**78%**
5	**Imsouane** Morocco	**75%**
6	**Hossegor** France	**69%**
7	**Short Sands** Oregon	**65%**
8	**Santa Catalina** Panama	**59%**
9	**Ericeira** Portugal	**58%**
10	**Lakey Peak** Sumbawa, Indonesia	**57%**
11	**Elands Bay** South Africa	**56%**
12	**Punta Roca** El Salvador	**54%**
13	**Punta de Lobos** Chile	**54%**
14	**Desert Point** Lombok, Indonesia	**53%**
15	**G-Land** Java, Indonesia	**51%**
16	**Bundoran** Ireland	**50%**
17	**Mavericks** California	**49%**
18	**Pavones** Costa Rica	**49%**
19	**Shipstern Bluff** Tasmania, Australia	**49%**
20	**Todos Santos** Baja California	**49%**
21	**Safi** Morocco	**48%**
22	**Nias** Sumatra, Indonesia	**48%**
23	**Montañita** Ecuador	**48%**
24	**Tofino** British Columbia	**46%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Enclosed-sea data for this month is still downloading (Open-Meteo free-tier quota), so those seas show as no data until the PDF is rebuilt.

April: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)


→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in April


% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**92%**
2	**Punta Hermosa** Peru	**85%**
3	**Nias** Sumatra, Indonesia	**76%**
4	**Santa Catalina** Panama	**75%**
5	**Punta Roca** El Salvador	**74%**
6	**Piha** New Zealand	**72%**
7	**Chicama** Peru	**71%**
8	**Elands Bay** South Africa	**67%**
9	**Pascuales** Colima, Mexico	**67%**
10	**Playa Hermosa** Costa Rica	**65%**
11	**Pavones** Costa Rica	**63%**
12	**Lakey Peak** Sumbawa, Indonesia	**61%**
13	**Montañita** Ecuador	**58%**
14	**Sayulita** Nayarit, Mexico	**57%**
15	**Punta de Lobos** Chile	**55%**
16	**Short Sands** Oregon	**54%**
17	**Puerto Escondido** Oaxaca, Mexico	**53%**
18	**Imsouane** Morocco	**53%**
19	**Skeleton Bay** Namibia	**51%**
20	**Busua** Ghana	**50%**
21	**Ericeira** Portugal	**49%**
22	**Hossegor** France	**48%**
23	**G-Land** Java, Indonesia	**47%**
24	**Shipstern Bluff** Tasmania, Australia	**47%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Enclosed-sea data for this month is still downloading (Open-Meteo free-tier quota), so those seas show as no data until the PDF is rebuilt.

May: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)


Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in May


% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**90%**
2	**Punta Hermosa** Peru	**82%**
3	**Punta Roca** El Salvador	**79%**
4	**Elands Bay** South Africa	**78%**
5	**Santa Catalina** Panama	**75%**
6	**Piha** New Zealand	**73%**
7	**Pascuales** Colima, Mexico	**71%**
8	**Punta de Lobos** Chile	**68%**
9	**Montañita** Ecuador	**68%**
10	**Playa Hermosa** Costa Rica	**68%**
11	**Skeleton Bay** Namibia	**66%**
12	**Nias** Sumatra, Indonesia	**66%**
13	**Sayulita** Nayarit, Mexico	**65%**
14	**Chicama** Peru	**64%**
15	**Pavones** Costa Rica	**61%**
16	**Puerto Escondido** Oaxaca, Mexico	**58%**
17	**Short Sands** Oregon	**52%**
18	**Busua** Ghana	**51%**
19	**Jeffreys Bay** South Africa	**49%**
20	**Witch's Rock** Costa Rica	**47%**
21	**Robertsport** Liberia	**47%**
22	**Mentawais** Sumatra, Indonesia	**44%**
23	**Shipstern Bluff** Tasmania, Australia	**41%**
24	**Margaret River** Western Australia	**40%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Enclosed-sea data for this month is still downloading (Open-Meteo free-tier quota), so those seas show as no data until the PDF is rebuilt.

June: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in June

% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**83%**
2	**Nias** Sumatra, Indonesia	**75%**
3	**Skeleton Bay** Namibia	**64%**
4	**Punta Hermosa** Peru	**63%**
5	**Elands Bay** South Africa	**62%**
6	**Punta de Lobos** Chile	**61%**
7	**Mentawais** Sumatra, Indonesia	**61%**
8	**Santa Catalina** Panama	**58%**
9	**Pascuales** Colima, Mexico	**57%**
10	**Piha** New Zealand	**57%**
11	**Punta Roca** El Salvador	**56%**
12	**Sayulita** Nayarit, Mexico	**53%**
13	**Montañita** Ecuador	**51%**
14	**Puerto Escondido** Oaxaca, Mexico	**50%**
15	**Jeffreys Bay** South Africa	**46%**
16	**Playa Hermosa** Costa Rica	**45%**
17	**Máncora** Peru	**42%**
18	**Chicama** Peru	**41%**
19	**Pavones** Costa Rica	**40%**
20	**Busua** Ghana	**38%**
21	**Margaret River** Western Australia	**38%**
22	**Shipstern Bluff** Tasmania, Australia	**36%**
23	**Witch's Rock** Costa Rica	**36%**
24	**Lakey Peak** Sumbawa, Indonesia	**35%**

Great Lakes

Open-Meteo data for this month is still downloading

Mediterranean & Black Sea

Open-Meteo data for this month is still downloading

Caspian Sea

Open-Meteo data for this month is still downloading


Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Enclosed-sea data for this month is still downloading (Open-Meteo free-tier quota), so those seas show as no data until the PDF is rebuilt.

July: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (154 mornings; 1 days missing from the archive)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)


Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in July


% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**88%**
2	**Nias** Sumatra, Indonesia	**76%**
3	**Mentawais** Sumatra, Indonesia	**70%**
4	**Punta Hermosa** Peru	**69%**
5	**Elands Bay** South Africa	**67%**
6	**Puerto Escondido** Oaxaca, Mexico	**65%**
7	**Santa Catalina** Panama	**65%**
8	**Punta Roca** El Salvador	**62%**
9	**Skeleton Bay** Namibia	**61%**
10	**Punta de Lobos** Chile	**59%**
11	**Pascuales** Colima, Mexico	**54%**
12	**Pavones** Costa Rica	**51%**
13	**Playa Hermosa** Costa Rica	**50%**
14	**Sayulita** Nayarit, Mexico	**49%**
15	**Piha** New Zealand	**48%**
16	**Chicama** Peru	**48%**
17	**Montañita** Ecuador	**47%**
18	**Jeffreys Bay** South Africa	**45%**
19	**Teahupo'o** Tahiti	**40%**
20	**Shipstern Bluff** Tasmania, Australia	**36%**
21	**Lakey Peak** Sumbawa, Indonesia	**32%**
22	**Rapa Nui** Easter Island	**32%**
23	**Busua** Ghana	**32%**
24	**North Malé Atoll** Maldives	**30%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Enclosed-sea data for this month is still downloading (Open-Meteo free-tier quota), so those seas show as no data until the PDF is rebuilt.

August: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)


→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in August


% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**81%**
2	**Nias** Sumatra, Indonesia	**73%**
3	**Elands Bay** South Africa	**71%**
4	**Punta Roca** El Salvador	**70%**
5	**Puerto Escondido** Oaxaca, Mexico	**68%**
6	**Skeleton Bay** Namibia	**67%**
7	**Punta Hermosa** Peru	**66%**
8	**Mentawais** Sumatra, Indonesia	**65%**
9	**Piha** New Zealand	**62%**
10	**Pascuales** Colima, Mexico	**61%**
11	**Lakey Peak** Sumbawa, Indonesia	**58%**
12	**Punta de Lobos** Chile	**55%**
13	**Sayulita** Nayarit, Mexico	**54%**
14	**Montañita** Ecuador	**53%**
15	**Santa Catalina** Panama	**52%**
16	**Playa Hermosa** Costa Rica	**46%**
17	**Chicama** Peru	**46%**
18	**Jeffreys Bay** South Africa	**45%**
19	**Máncora** Peru	**43%**
20	**Busua** Ghana	**41%**
21	**Teahupo'o** Tahiti	**39%**
22	**Rapa Nui** Easter Island	**38%**
23	**Pavones** Costa Rica	**37%**
24	**Shipstern Bluff** Tasmania, Australia	**36%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Enclosed-sea data for this month is still downloading (Open-Meteo free-tier quota), so those seas show as no data until the PDF is rebuilt.

September: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)


→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in September


% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**84%**
2	**Lakey Peak** Sumbawa, Indonesia	**77%**
3	**Santa Catalina** Panama	**65%**
4	**Elands Bay** South Africa	**64%**
5	**Punta Hermosa** Peru	**59%**
6	**Punta de Lobos** Chile	**59%**
7	**Punta Roca** El Salvador	**58%**
8	**Piha** New Zealand	**57%**
9	**Nias** Sumatra, Indonesia	**55%**
10	**Skeleton Bay** Namibia	**53%**
11	**Puerto Escondido** Oaxaca, Mexico	**53%**
12	**Mentawais** Sumatra, Indonesia	**52%**
13	**Montañita** Ecuador	**51%**
14	**Playa Hermosa** Costa Rica	**49%**
15	**Pavones** Costa Rica	**47%**
16	**Busua** Ghana	**46%**
17	**Pascuales** Colima, Mexico	**46%**
18	**Chicama** Peru	**43%**
19	**G-Land** Java, Indonesia	**42%**
20	**Short Sands** Oregon	**42%**
21	**Sayulita** Nayarit, Mexico	**42%**
22	**Jeffreys Bay** South Africa	**40%**
23	**Hossegor** France	**39%**
24	**Nazaré** Portugal	**39%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Enclosed-sea data for this month is still downloading (Open-Meteo free-tier quota), so those seas show as no data until the PDF is rebuilt.

October: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (148 mornings; 7 days missing from the archive)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)


→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in October


% of mornings that are good+ with swell inside the break's exposure window; best break per 100 miles of coast

1	**Arica** Chile	**93%**
2	**Lakey Peak** Sumbawa, Indonesia	**91%**
3	**Punta Hermosa** Peru	**82%**
4	**G-Land** Java, Indonesia	**70%**
5	**Desert Point** Lombok, Indonesia	**66%**
6	**Puerto Escondido** Oaxaca, Mexico	**65%**
7	**Chicama** Peru	**65%**
8	**Elands Bay** South Africa	**64%**
9	**Punta Roca** El Salvador	**64%**
10	**Piha** New Zealand	**63%**
11	**Montañita** Ecuador	**61%**
12	**Nias** Sumatra, Indonesia	**61%**
13	**Santa Catalina** Panama	**60%**
14	**Pascuales** Colima, Mexico	**59%**
15	**Imsouane** Morocco	**59%**
16	**Short Sands** Oregon	**59%**
17	**Margaret River** Western Australia	**57%**
18	**Mavericks** California	**57%**
19	**Punta de Lobos** Chile	**56%**
20	**Hossegor** France	**54%**
21	**Ericeira** Portugal	**54%**
22	**Skeleton Bay** Namibia	**50%**
23	**Sayulita** Nayarit, Mexico	**49%**
24	**Mentawais** Sumatra, Indonesia	**47%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea



Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Enclosed-sea data for this month is still downloading (Open-Meteo free-tier quota), so those seas show as no data until the PDF is rebuilt.