
Year-round: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 1,803 mornings, Oct 2021 to Sep 2026



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks year-round

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt S	**87%**
2	**Iquique** Chile · 7 kt S	**82%**
3	**Punta Hermosa** Peru · 7 kt SSE	**74%**
4	**Krui** Lampung, Sumatra · 10 kt SE	**66%**
5	**Elands Bay** South Africa · 9 kt SSE	**66%**
6	**Piha** New Zealand · 14 kt SW	**65%**
7	**Mentawais** Sumatra, Indonesia · 6 kt SE	**64%**
8	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**62%**
9	**Nias** Sumatra, Indonesia · 6 kt SE	**58%**
10	**Chicama** Peru · 9 kt SSE	**58%**
11	**Skeleton Bay** Namibia · 10 kt S	**55%**
12	**Santa Catalina** Panama · 7 kt NE	**54%**
13	**Punta de Lobos** Chile · 12 kt S	**53%**
14	**Punta Roca** El Salvador · 7 kt ENE	**52%**
15	**Lakey Peak** Sumbawa, Indonesia · 8 kt E	**51%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 9 kt N	**80%**
◆2	**Camaná coast** Peru · 10 kt SE	**79%**
◆3	**Antofagasta coast** Chile · 8 kt S	**79%**
◆4	**La Ligua coast** Chile · 12 kt S	**73%**
◆5	**Toltén coast** Chile · 13 kt S	**70%**
◆7	**Chañaral coast** Chile · 6 kt SSW	**69%**
◆6	**Isabela coast** Galápagos, Ecuador · 10 kt SSE	**69%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea



Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.29× the mean swell period.

Popular breaks, month by month

The 40 well-known breaks with the strongest three-month season (best one per 100 miles of coast), listed by year-round consistency. Breaks in the enclosed seas are on the map insets.

#	Break	Region	J	F	M	A	M	J	J	A	S	O	N	D	Year	Worst	Best 3 months	Typical good morning	Morning wind
1	**Arica**	Chile	86	88	92	92	90	81	88	81	83	93	85	81	**87%**	81%	Mar–May 91%	5 ft @ 13 s SSW, surf ~8 ft	4 kt S
2	**Iquique**	Chile	86	84	86	89	87	72	82	73	72	89	81	80	**82%**	72%	Mar–May 87%	6 ft @ 13 s SSW, surf ~9 ft	7 kt S
3	**Punta Hermosa**	Peru	77	80	84	85	82	63	69	66	59	82	75	69	**74%**	59%	Mar–May 84%	5 ft @ 14 s SSW, surf ~8 ft	7 kt SSE
4	**Krui**	Lampung, Sumatra	50	48	53	75	80	82	80	77	79	78	38	52	**66%**	38%	May–Jul 81%	4 ft @ 14 s SSW, surf ~7 ft	10 kt SE
5	**Elands Bay**	South Africa	68	71	55	66	78	59	65	70	61	64	67	71	**66%**	55%	Dec–Feb 70%	5 ft @ 13 s SW, surf ~8 ft	9 kt SSE
6	**Piha**	New Zealand	63	77	76	71	72	54	47	61	55	63	73	68	**65%**	47%	Feb–Apr 75%	7 ft @ 13 s SW, surf ~10 ft	14 kt SW
7	**Mentawais**	Sumatra, Indonesia	50	55	54	74	69	81	80	79	64	74	28	50	**64%**	28%	Jun–Aug 80%	5 ft @ 14 s SSW, surf ~7 ft	6 kt SE
8	**Cimaja**	Pelabuhan Ratu, West Java	58	56	66	73	72	73	63	59	55	62	43	57	**62%**	43%	Apr–Jun 73%	4 ft @ 14 s SSW, surf ~7 ft	4 kt E
9	**Nias**	Sumatra, Indonesia	51	48	46	76	65	75	76	72	54	61	16	52	**58%**	16%	Jun–Aug 74%	4 ft @ 14 s SSW, surf ~7 ft	6 kt SE
10	**Chicama**	Peru	59	65	79	71	63	39	47	45	41	65	60	58	**58%**	39%	Feb–Apr 71%	4 ft @ 14 s SSW, surf ~7 ft	9 kt SSE
11	**Skeleton Bay**	Namibia	63	65	42	51	65	63	59	66	52	50	43	37	**55%**	37%	Jun–Aug 63%	6 ft @ 12 s SSW, surf ~9 ft	10 kt S
12	**Santa Catalina**	Panama	30	30	58	75	75	58	65	52	65	60	46	32	**54%**	30%	Mar–May 70%	4 ft @ 14 s SSW, surf ~6 ft	7 kt NE
13	**Punta de Lobos**	Chile	44	40	53	55	68	59	58	55	58	55	49	38	**53%**	38%	May–Jul 62%	8 ft @ 13 s SW, surf ~11 ft	12 kt S
14	**Punta Roca**	El Salvador	17	30	54	74	78	55	62	70	58	64	43	21	**52%**	17%	Apr–Jun 69%	3 ft @ 15 s SSW, surf ~6 ft	7 kt ENE
15	**Lakey Peak**	Sumbawa, Indonesia	58	65	57	50	28	23	27	43	51	82	55	73	**51%**	23%	Oct–Dec 70%	4 ft @ 14 s SSW, surf ~7 ft	8 kt E
16	**Short Sands**	Oregon	79	79	65	54	52	27	11	12	42	59	68	63	**51%**	11%	Jan–Mar 74%	7 ft @ 12 s W, surf ~10 ft	14 kt N
17	**Cactus Beach**	South Australia	32	45	43	69	62	46	50	57	54	52	37	40	**49%**	32%	Apr–Jun 59%	6 ft @ 14 s SW, surf ~10 ft	12 kt SE
18	**Montañita**	Ecuador	45	35	48	58	68	51	47	53	51	61	39	27	**49%**	27%	Apr–Jun 59%	3 ft @ 15 s SW, surf ~6 ft	8 kt SSW
19	**Desert Point**	Lombok, Indonesia	54	48	62	57	33	22	16	32	45	74	63	64	**47%**	16%	Oct–Dec 67%	5 ft @ 14 s SSW, surf ~8 ft	10 kt ESE
20	**Puerto Escondido**	Oaxaca, Mexico	17	32	35	53	56	48	64	67	52	65	49	25	**47%**	17%	Aug–Oct 61%	4 ft @ 15 s SSW, surf ~6 ft	6 kt WNW
21	**G-Land**	Java, Indonesia	48	43	58	57	32	19	17	33	47	81	63	63	**47%**	17%	Oct–Dec 69%	5 ft @ 14 s SSW, surf ~9 ft	10 kt ESE
22	**Imsouane**	Morocco	71	73	75	53	25	14	7	13	32	59	63	74	**46%**	7%	Jan–Mar 73%	6 ft @ 13 s NW, surf ~9 ft	11 kt N
23	**Hossegor**	France	57	80	69	47	31	29	18	28	38	53	46	61	**46%**	18%	Jan–Mar 69%	5 ft @ 13 s WNW, surf ~8 ft	9 kt W
24	**Sayulita**	Nayarit, Mexico	47	37	41	57	65	53	49	54	42	49	31	26	**46%**	26%	Apr–Jun 59%	3 ft @ 15 s SW, surf ~6 ft	7 kt N
25	**Pascuales**	Colima, Mexico	8	32	39	67	71	57	54	61	46	59	36	15	**45%**	8%	Apr–Jun 65%	4 ft @ 15 s SSW, surf ~6 ft	6 kt E
26	**Montara**	California	70	62	48	35	32	21	8	8	31	57	67	60	**41%**	8%	Nov–Jan 66%	7 ft @ 13 s WNW, surf ~10 ft	12 kt NW
27	**Jeffreys Bay**	South Africa	42	42	38	35	48	43	43	43	38	34	43	43	**41%**	34%	May–Jul 45%	8 ft @ 12 s SW, surf ~11 ft	13 kt W
28	**Pavones**	Costa Rica	17	28	49	63	61	40	51	37	47	40	28	20	**40%**	17%	Mar–May 58%	4 ft @ 14 s SSW, surf ~6 ft	5 kt E
29	**Playa Hermosa**	Costa Rica	14	21	40	65	68	45	50	46	49	38	25	17	**40%**	14%	Apr–Jun 59%	3 ft @ 14 s SSW, surf ~6 ft	3 kt SE
30	**Todos Santos**	Baja California	77	70	48	35	34	21	10	9	15	35	58	60	**39%**	9%	Dec–Feb 69%	4 ft @ 13 s WNW, surf ~7 ft	8 kt NW
31	**Shipstern Bluff**	Tasmania, Australia	48	45	46	45	39	34	34	35	31	25	46	37	**39%**	25%	Jan–Mar 46%	8 ft @ 13 s SW, surf ~12 ft	16 kt SW
32	**Lhoknga**	Aceh, Sumatra	28	32	33	45	37	53	26	49	42	34	15	38	**36%**	15%	Apr–Jun 45%	4 ft @ 15 s SSW, surf ~7 ft	11 kt S
33	**Ericeira**	Portugal	37	52	52	47	23	18	6	13	36	53	42	51	**36%**	6%	Feb–Apr 50%	7 ft @ 12 s NW, surf ~10 ft	13 kt N
34	**Busua**	Ghana	25	26	16	50	51	37	31	41	44	43	31	19	**35%**	16%	Apr–Jun 46%	4 ft @ 13 s SSW, surf ~6 ft	8 kt SSW
35	**Margaret River**	Western Australia	32	25	26	40	37	33	24	29	35	55	33	35	**34%**	24%	Sep–Nov 41%	9 ft @ 13 s SW, surf ~13 ft	13 kt ESE
36	**Bundoran**	Ireland	51	44	46	35	31	20	10	14	31	41	39	38	**33%**	10%	Jan–Mar 47%	7 ft @ 12 s WNW, surf ~10 ft	13 kt S
37	**Safi**	Morocco	51	51	47	33	22	9	4	10	30	44	46	53	**33%**	4%	Dec–Feb 52%	7 ft @ 13 s NW, surf ~10 ft	12 kt NNE
38	**Rincon**	California	66	52	35	25	28	19	8	8	15	24	44	48	**31%**	8%	Dec–Feb 56%	4 ft @ 13 s W, surf ~7 ft	8 kt NW
39	**Tofino**	British Columbia	36	43	45	36	38	15	6	5	33	39	32	34	**30%**	5%	Jan–Mar 41%	7 ft @ 12 s W, surf ~10 ft	11 kt ESE
40	**Robertsport**	Liberia	23	16	14	43	47	29	22	26	31	37	19	12	**27%**	12%	Apr–Jun 40%	4 ft @ 13 s S, surf ~6 ft	7 kt SE

Month cells use the map colours (% of mornings good+). Typical good morning = medians over the good+ mornings: swell height @ peak period and the direction it comes from, and estimated breaking surf (Komar–Gaughan).
Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward.

Each break is read from its nearest seaward model cell (0.5° in East and Southeast Asia, 1° elsewhere) with swell inside its exposure window. The rule has no wind; local reefs, points and sheltering are not modelled.

Under-the-radar stretches, month by month

Coast where mornings are often good+ with swell aimed at the shore (≥40% year-round, or ≥65% over its best three months),
at least 300 km from the well-known breaks and within 150 km of a town of 5,000+. Listed by year-round consistency; numbers match the ◆ markers on the maps.


#	Stretch	Region	J	F	M	A	M	J	J	A	S	O	N	D	Year	Worst	Best 3 months	Typical good morning	Morning wind
◆1	**Ancud coast**	Chile	89	82	77	79	78	77	67	77	72	89	82	90	**80%**	67%	Dec–Feb 87%	7 ft @ 13 s WSW, surf ~10 ft	9 kt N
◆2	**Camaná coast**	Peru	83	81	87	87	81	72	84	72	71	81	79	72	**79%**	71%	Mar–May 85%	6 ft @ 13 s SSW, surf ~9 ft	10 kt SE
◆3	**Antofagasta coast**	Chile	83	82	88	83	81	76	75	68	69	85	80	75	**79%**	68%	Jan–Mar 84%	6 ft @ 13 s SW, surf ~9 ft	8 kt S
◆4	**La Ligua coast**	Chile	72	75	75	77	77	79	76	70	69	75	71	65	**73%**	65%	Apr–Jun 78%	7 ft @ 13 s SW, surf ~10 ft	12 kt S
◆5	**Toltén coast**	Chile	81	72	70	69	72	65	60	73	67	72	73	71	**70%**	60%	Nov–Jan 75%	8 ft @ 13 s SW, surf ~11 ft	13 kt S
◆7	**Chañaral coast**	Chile	76	74	72	63	74	66	74	64	62	75	61	69	**69%**	61%	Jan–Mar 74%	5 ft @ 13 s SW, surf ~8 ft	6 kt SSW
◆6	**Isabela coast**	Galápagos, Ecuador	75	74	79	82	79	59	59	59	49	77	71	61	**69%**	49%	Mar–May 80%	4 ft @ 14 s SW, surf ~7 ft	10 kt SSE
◆8	**Soyo coast**	Angola	61	62	50	65	77	70	77	84	65	56	52	49	**64%**	49%	Jun–Aug 77%	4 ft @ 13 s SSW, surf ~7 ft	7 kt S
◆9	**Sumbe coast**	Angola	54	62	46	61	77	68	76	87	65	64	59	48	**64%**	46%	Jun–Aug 77%	4 ft @ 13 s SW, surf ~7 ft	5 kt S
◆10	**Pointe-Noire coast**	Congo (Brazzaville)	54	58	48	65	81	70	76	89	65	54	51	45	**63%**	45%	Jun–Aug 78%	4 ft @ 13 s SSW, surf ~7 ft	6 kt S
◆11	**Greymouth coast**	New Zealand	54	65	61	69	67	48	58	67	63	58	62	53	**60%**	48%	Mar–May 66%	5 ft @ 13 s WSW, surf ~9 ft	9 kt E
◆12	**La Serena coast**	Chile	64	60	68	65	65	61	55	46	49	65	56	60	**59%**	46%	Mar–May 66%	7 ft @ 13 s SW, surf ~10 ft	7 kt SSW
◆13	**Omboué coast**	Gabon	46	50	39	63	69	65	69	82	67	49	45	35	**57%**	35%	Jul–Sep 73%	4 ft @ 13 s SSW, surf ~7 ft	7 kt SSW
◆14	**Escuintla coast**	Mexico	15	34	45	69	77	56	68	72	55	59	45	23	**52%**	15%	Apr–Jun 67%	4 ft @ 15 s SSW, surf ~7 ft	5 kt ESE
◆16	**Acapulco coast**	Mexico	17	42	50	70	70	50	61	68	45	66	53	24	**51%**	17%	Apr–Jun 64%	4 ft @ 15 s SSW, surf ~6 ft	6 kt NW
◆15	**Marcona coast**	Peru	50	66	68	73	57	37	47	36	34	51	53	44	**51%**	34%	Feb–Apr 69%	6 ft @ 13 s SSW, surf ~9 ft	13 kt SE
◆17	**Port Lincoln coast**	Australia	49	47	51	54	54	45	48	55	55	53	36	56	**50%**	36%	Aug–Oct 54%	7 ft @ 14 s SW, surf ~10 ft	12 kt SE
◆18	**Luanda coast**	Angola	30	36	28	49	68	69	69	80	54	43	41	28	**50%**	28%	Jun–Aug 73%	4 ft @ 13 s SW, surf ~6 ft	6 kt S
◆19	**Meningie coast**	Australia	37	60	54	63	50	39	42	46	47	50	44	52	**49%**	37%	Feb–Apr 59%	7 ft @ 14 s SW, surf ~11 ft	10 kt S
◆20	**Oranjemund coast**	Namibia	54	47	38	36	62	60	51	61	45	43	41	42	**48%**	36%	May–Jul 58%	7 ft @ 13 s SW, surf ~10 ft	9 kt SE
◆21	**Blitar coast**	Indonesia	48	42	55	66	37	26	16	36	58	77	56	63	**48%**	16%	Oct–Dec 65%	5 ft @ 14 s SSW, surf ~9 ft	12 kt ESE
◆22	**Maintirano coast**	Madagascar	31	38	31	51	59	55	52	53	51	39	40	48	**46%**	31%	May–Jul 55%	4 ft @ 13 s SW, surf ~7 ft	7 kt SE
◆23	**Morombe coast**	Madagascar	30	43	37	55	54	56	54	50	49	40	39	39	**45%**	30%	Apr–Jun 55%	5 ft @ 13 s SSW, surf ~8 ft	10 kt SSE
◆24	**São Tomé coast**	Sao Tome and Principe	37	45	33	62	60	50	51	58	47	36	38	26	**45%**	26%	Apr–Jun 58%	4 ft @ 13 s SSW, surf ~7 ft	8 kt SSW
◆25	**Huarmey coast**	Peru	59	55	60	54	46	27	34	26	32	45	49	41	**44%**	26%	Jan–Mar 58%	5 ft @ 14 s SSW, surf ~8 ft	10 kt SSE
◆26	**Quellón coast**	Chile	61	52	37	41	39	32	29	39	30	51	50	56	**43%**	29%	Dec–Feb 56%	8 ft @ 13 s WSW, surf ~12 ft	13 kt W
◆27	**Rabat coast**	Morocco	62	61	71	53	26	16	5	11	31	56	55	68	**43%**	5%	Jan–Mar 65%	6 ft @ 13 s WNW, surf ~9 ft	8 kt NNE
◆28	**Juradó coast**	Colombia	15	25	37	55	66	50	53	52	48	50	33	22	**42%**	15%	Apr–Jun 57%	3 ft @ 15 s SW, surf ~6 ft	5 kt S
◆29	**Port Denison coast**	Australia	30	44	35	47	45	47	37	37	45	59	46	35	**42%**	30%	Sep–Nov 50%	9 ft @ 14 s SW, surf ~12 ft	12 kt SSE
◆30	**Kaitaia coast**	New Zealand	39	55	54	35	33	29	31	40	42	42	53	53	**42%**	29%	Oct–Dec 49%	7 ft @ 14 s SW, surf ~11 ft	13 kt S
◆31	**Nelson coast**	New Zealand	50	55	52	41	43	31	28	41	31	34	51	44	**42%**	28%	Jan–Mar 52%	7 ft @ 13 s WSW, surf ~11 ft	8 kt SSW
◆32	**San Cristóbal coast**	Galápagos, Ecuador	37	50	70	72	55	35	25	22	26	31	36	29	**41%**	22%	Mar–May 66%	4 ft @ 14 s SSW, surf ~7 ft	10 kt SSE
◆33	**Kangaroo Island coast**	Australia	44	44	48	53	43	35	29	34	27	41	39	46	**40%**	27%	Feb–Apr 48%	9 ft @ 13 s SW, surf ~12 ft	12 kt SE

Month cells use the map colours (% of mornings good+). Typical good morning = medians over the good+ mornings: swell height @ peak period and the direction it comes from, and estimated breaking surf (Komar–Gaughan).
Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward.

Each stretch is read from one WW3 cell 50–120 km offshore. The rule has no wind, which flatters cold, steady coasts (Chile, Peru, Namibia) and storm coasts (southern Chile, New Zealand's West Coast); check local wind and access before booking a trip.

October–March: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 889 mornings, Oct 2021 to Mar 2026, these months only



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks, October–March

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)

1	**Arica** Chile · 5 kt S	**88%**
2	**Iquique** Chile · 7 kt S	**84%**
3	**Punta Hermosa** Peru · 7 kt SSE	**78%**
4	**Piha** New Zealand · 13 kt SSW	**70%**
5	**Imsouane** Morocco · 8 kt NNE	**69%**
6	**Short Sands** Oregon · 14 kt S	**69%**
7	**Elands Bay** South Africa · 9 kt S	**66%**
8	**Lakey Peak** Sumbawa, Indonesia · 7 kt WNW	**65%**
9	**Chicama** Peru · 9 kt SSE	**64%**
10	**Hossegor** France · 10 kt SSW	**61%**
11	**Montara** California · 11 kt NNW	**61%**
12	**Desert Point** Lombok, Indonesia · 8 kt W	**61%**
13	**G-Land** Java, Indonesia · 9 kt W	**59%**
14	**Todos Santos** Baja California · 6 kt NW	**58%**
15	**Cimaja** Pelabuhan Ratu, West Java · 3 kt E	**57%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 8 kt SE	**85%**
◆3	**Antofagasta coast** Chile · 7 kt SSW	**82%**
◆2	**Camaná coast** Peru · 9 kt SE	**80%**
◆5	**Toltén coast** Chile · 13 kt S	**73%**
◆6	**Isabela coast** Galápagos, Ecuador · 10 kt SSE	**73%**
◆4	**La Ligua coast** Chile · 11 kt S	**72%**
◆7	**Chañaral coast** Chile · 5 kt SSW	**71%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.29× the mean swell period.

April–September: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 914 mornings, Apr 2022 to Sep 2026, these months only



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks, April–September

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt SSE	**86%**
2	**Iquique** Chile · 7 kt S	**79%**
3	**Krui** Lampung, Sumatra · 9 kt SE	**79%**
4	**Mentawais** Sumatra, Indonesia · 5 kt SE	**75%**
5	**Punta Hermosa** Peru · 8 kt S	**71%**
6	**Nias** Sumatra, Indonesia · 5 kt SE	**70%**
7	**Elands Bay** South Africa · 8 kt SSE	**67%**
8	**Punta Roca** El Salvador · 7 kt ESE	**66%**
9	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**66%**
10	**Santa Catalina** Panama · 7 kt SSE	**65%**
11	**Piha** New Zealand · 15 kt WSW	**60%**
12	**Skeleton Bay** Namibia · 10 kt S	**59%**
13	**Pascuales** Colima, Mexico · 6 kt E	**59%**
14	**Punta de Lobos** Chile · 12 kt S	**59%**
15	**Puerto Escondido** Oaxaca, Mexico · 8 kt WNW	**57%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Camaná coast** Peru · 11 kt SE	**78%**
◆3	**Antofagasta coast** Chile · 10 kt S	**75%**
◆1	**Ancud coast** Chile · 11 kt N	**75%**
◆4	**La Ligua coast** Chile · 13 kt S	**75%**
◆10	**Pointe-Noire coast** Congo (Brazzaville) · 6 kt S	**74%**
◆8	**Soyo coast** Angola · 7 kt S	**73%**
◆9	**Sumbe coast** Angola · 5 kt SSW	**72%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.28× the mean swell period.

November: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (135 mornings; 15 days missing from the archive)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in November

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)

1	**Arica** Chile · 5 kt S	**85%**
2	**Iquique** Chile · 7 kt S	**81%**
3	**Punta Hermosa** Peru · 6 kt SSE	**75%**
4	**Piha** New Zealand · 11 kt SW	**73%**
5	**Short Sands** Oregon · 11 kt SSE	**68%**
6	**Montara** California · 11 kt NNW	**67%**
7	**Elands Bay** South Africa · 9 kt S	**67%**
8	**G-Land** Java, Indonesia · 6 kt E	**63%**
9	**Imsouane** Morocco · 5 kt NNE	**63%**
10	**Desert Point** Lombok, Indonesia · 7 kt SE	**63%**
11	**Chicama** Peru · 9 kt SSE	**60%**
12	**Todos Santos** Baja California · 5 kt NW	**58%**
13	**Lakey Peak** Sumbawa, Indonesia · 6 kt E	**55%**
14	**Puerto Escondido** Oaxaca, Mexico · 5 kt NNE	**49%**
15	**Punta de Lobos** Chile · 11 kt SSW	**49%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 10 kt NNW	**82%**
◆3	**Antofagasta coast** Chile · 9 kt SSW	**80%**
◆2	**Camaná coast** Peru · 10 kt SE	**79%**
◆5	**Toltén coast** Chile · 11 kt S	**73%**
◆4	**La Ligua coast** Chile · 12 kt S	**71%**
◆6	**Isabela coast** Galápagos, Ecuador · 12 kt SSE	**71%**
◆11	**Greymouth coast** New Zealand · 5 kt ESE	**62%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.29× the mean swell period.

December: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (155 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in December

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)

1	**Arica** Chile · 5 kt S	**81%**
2	**Iquique** Chile · 7 kt S	**80%**
3	**Imsouane** Morocco · 7 kt NNE	**74%**
4	**Lakey Peak** Sumbawa, Indonesia · 7 kt WSW	**73%**
5	**Elands Bay** South Africa · 9 kt S	**71%**
6	**Punta Hermosa** Peru · 7 kt SSE	**69%**
7	**Piha** New Zealand · 11 kt N	**68%**
8	**Keramas** Bali, Indonesia · 7 kt W	**65%**
9	**Short Sands** Oregon · 14 kt SSE	**63%**
10	**Hossegor** France · 9 kt SE	**61%**
11	**Montara** California · 11 kt NNW	**60%**
12	**Todos Santos** Baja California · 5 kt NNW	**60%**
13	**Chicama** Peru · 9 kt SSE	**58%**
14	**Cimaja** Pelabuhan Ratu, West Java · 3 kt E	**57%**
15	**Safi** Morocco · 12 kt ENE	**53%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 8 kt SSW	**90%**
◆3	**Antofagasta coast** Chile · 7 kt S	**75%**
◆2	**Camaná coast** Peru · 9 kt SE	**72%**
◆5	**Toltén coast** Chile · 14 kt S	**71%**
◆7	**Chañaral coast** Chile · 5 kt SSW	**69%**
◆27	**Rabat coast** Morocco · 8 kt ENE	**68%**
◆4	**La Ligua coast** Chile · 12 kt S	**65%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.29× the mean swell period.

January: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Best named breaks in January

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)

1	**Arica** Chile · 5 kt S	**86%**
2	**Iquique** Chile · 6 kt S	**86%**
3	**Short Sands** Oregon · 14 kt S	**79%**
4	**Punta Hermosa** Peru · 7 kt SSE	**77%**
5	**Todos Santos** Baja California · 5 kt SE	**77%**
6	**Anchor Point** Taghazout, Morocco · 8 kt SE	**71%**
7	**Montara** California · 13 kt SSE	**70%**
8	**Elands Bay** South Africa · 8 kt SSE	**68%**
9	**Rincon** California · 7 kt NW	**66%**
10	**Skeleton Bay** Namibia · 7 kt S	**63%**
11	**Piha** New Zealand · 13 kt ENE	**63%**
12	**Chicama** Peru · 8 kt SSE	**59%**
13	**Lakey Peak** Sumbawa, Indonesia · 8 kt WSW	**58%**
14	**Cimaja** Pelabuhan Ratu, West Java · 3 kt ENE	**58%**
15	**Hossegor** France · 16 kt SSW	**57%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 8 kt SE	**89%**
◆3	**Antofagasta coast** Chile · 5 kt SSW	**83%**
◆2	**Camaná coast** Peru · 9 kt ESE	**83%**
◆5	**Toltén coast** Chile · 14 kt S	**81%**
◆7	**Chañaral coast** Chile · 5 kt S	**76%**
◆6	**Isabela coast** Galápagos, Ecuador · 9 kt SSE	**75%**
◆4	**La Ligua coast** Chile · 10 kt S	**72%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.29× the mean swell period.

February: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (141 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in February

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)

1	**Arica** Chile · 5 kt S	**88%**
2	**Iquique** Chile · 7 kt S	**84%**
3	**Punta Hermosa** Peru · 7 kt SSE	**80%**
4	**Hossegor** France · 8 kt SSW	**80%**
5	**Short Sands** Oregon · 13 kt SSE	**79%**
6	**Piha** New Zealand · 11 kt SSW	**77%**
7	**Imsouane** Morocco · 6 kt NNE	**73%**
8	**Elands Bay** South Africa · 9 kt S	**71%**
9	**Todos Santos** Baja California · 6 kt NW	**70%**
10	**Lakey Peak** Sumbawa, Indonesia · 7 kt WNW	**65%**
11	**Chicama** Peru · 8 kt SE	**65%**
12	**Skeleton Bay** Namibia · 9 kt S	**65%**
13	**Montara** California · 11 kt NNW	**62%**
14	**Cimaja** Pelabuhan Ratu, West Java · 3 kt NE	**56%**
15	**Mentawais** Sumatra, Indonesia · 7 kt NNW	**55%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 7 kt SE	**82%**
◆3	**Antofagasta coast** Chile · 6 kt SSW	**82%**
◆2	**Camaná coast** Peru · 8 kt ESE	**81%**
◆4	**La Ligua coast** Chile · 12 kt S	**75%**
◆6	**Isabela coast** Galápagos, Ecuador · 5 kt SSW	**74%**
◆7	**Chañaral coast** Chile · 5 kt SSW	**74%**
◆5	**Toltén coast** Chile · 14 kt S	**72%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.29× the mean swell period.

March: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in March

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)

1	**Arica** Chile · 5 kt S	**92%**
2	**Iquique** Chile · 7 kt S	**86%**
3	**Punta Hermosa** Peru · 7 kt SSE	**84%**
4	**Chicama** Peru · 10 kt SSE	**79%**
5	**Piha** New Zealand · 15 kt SSW	**76%**
6	**Imsouane** Morocco · 11 kt N	**75%**
7	**Hossegor** France · 13 kt W	**69%**
8	**Cimaja** Pelabuhan Ratu, West Java · 4 kt ENE	**66%**
9	**Short Sands** Oregon · 16 kt S	**65%**
10	**Desert Point** Lombok, Indonesia · 10 kt NE	**62%**
11	**Santa Catalina** Panama · 8 kt NE	**58%**
12	**G-Land** Java, Indonesia · 10 kt WNW	**58%**
13	**Lakey Peak** Sumbawa, Indonesia · 7 kt NE	**57%**
14	**Elands Bay** South Africa · 9 kt S	**55%**
15	**Punta Roca** El Salvador · 6 kt SE	**54%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆3	**Antofagasta coast** Chile · 7 kt S	**88%**
◆2	**Camaná coast** Peru · 9 kt ESE	**87%**
◆6	**Isabela coast** Galápagos, Ecuador · 4 kt SSE	**79%**
◆1	**Ancud coast** Chile · 9 kt ESE	**77%**
◆4	**La Ligua coast** Chile · 11 kt S	**75%**
◆7	**Chañaral coast** Chile · 5 kt S	**72%**
◆27	**Rabat coast** Morocco · 10 kt WSW	**71%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.28× the mean swell period.

April: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in April

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)

1	**Arica** Chile · 4 kt SSE	**92%**
2	**Iquique** Chile · 8 kt S	**89%**
3	**Punta Hermosa** Peru · 8 kt SSE	**85%**
4	**Nias** Sumatra, Indonesia · 5 kt SSE	**76%**
5	**Santa Catalina** Panama · 8 kt NE	**75%**
6	**Krui** Lampung, Sumatra · 6 kt ESE	**75%**
7	**Punta Roca** El Salvador · 6 kt ESE	**74%**
8	**Mentawais** Sumatra, Indonesia · 4 kt SSE	**74%**
9	**Cimaja** Pelabuhan Ratu, West Java · 4 kt ENE	**73%**
10	**Piha** New Zealand · 16 kt ENE	**71%**
11	**Chicama** Peru · 10 kt SE	**71%**
12	**Cactus Beach** South Australia · 11 kt ENE	**69%**
13	**Pascuales** Colima, Mexico · 5 kt ESE	**67%**
14	**Elands Bay** South Africa · 8 kt S	**66%**
15	**Playa Hermosa** Costa Rica · 3 kt SE	**65%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Camaná coast** Peru · 11 kt ESE	**87%**
◆3	**Antofagasta coast** Chile · 10 kt S	**83%**
◆6	**Isabela coast** Galápagos, Ecuador · 6 kt SSE	**82%**
◆1	**Ancud coast** Chile · 9 kt N	**79%**
◆4	**La Ligua coast** Chile · 12 kt S	**77%**
◆15	**Marcona coast** Peru · 14 kt SE	**73%**
◆32	**San Cristóbal coast** Galápagos, Ecuador · 7 kt SE	**72%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.28× the mean swell period.

May: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Best named breaks in May

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 3 kt SSE	**90%**
2	**Iquique** Chile · 7 kt S	**87%**
3	**Punta Hermosa** Peru · 7 kt S	**82%**
4	**Krui** Lampung, Sumatra · 9 kt ESE	**80%**
5	**Punta Roca** El Salvador · 5 kt ESE	**78%**
6	**Elands Bay** South Africa · 8 kt SSE	**78%**
7	**Santa Catalina** Panama · 7 kt SE	**75%**
8	**Piha** New Zealand · 15 kt S	**72%**
9	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**72%**
10	**Pascuales** Colima, Mexico · 5 kt NW	**71%**
11	**Mentawais** Sumatra, Indonesia · 5 kt WSW	**69%**
12	**Montañita** Ecuador · 8 kt SSW	**68%**
13	**Punta de Lobos** Chile · 11 kt S	**68%**
14	**Playa Hermosa** Costa Rica · 4 kt SE	**68%**
15	**Sayulita** Nayarit, Mexico · 6 kt NNE	**65%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Camaná coast** Peru · 11 kt SE	**81%**
◆3	**Antofagasta coast** Chile · 10 kt S	**81%**
◆10	**Pointe-Noire coast** Congo (Brazzaville) · 6 kt S	**81%**
◆6	**Isabela coast** Galápagos, Ecuador · 9 kt SSE	**79%**
◆1	**Ancud coast** Chile · 8 kt N	**78%**
◆4	**La Ligua coast** Chile · 12 kt S	**77%**
◆14	**Escuintla coast** Mexico · 5 kt NW	**77%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.27× the mean swell period.

June: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in June

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Krui** Lampung, Sumatra · 9 kt ESE	**82%**
2	**Arica** Chile · 3 kt SSE	**81%**
3	**Mentawais** Sumatra, Indonesia · 4 kt SE	**81%**
4	**Nias** Sumatra, Indonesia · 4 kt SE	**75%**
5	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**73%**
6	**Iquique** Chile · 7 kt S	**72%**
7	**Punta Hermosa** Peru · 8 kt S	**63%**
8	**Skeleton Bay** Namibia · 10 kt SSE	**63%**
9	**Punta de Lobos** Chile · 14 kt S	**59%**
10	**Elands Bay** South Africa · 8 kt SE	**59%**
11	**Santa Catalina** Panama · 6 kt WNW	**58%**
12	**Pascuales** Colima, Mexico · 6 kt NW	**57%**
13	**Punta Roca** El Salvador · 8 kt ESE	**55%**
14	**Piha** New Zealand · 15 kt NE	**54%**
15	**Sayulita** Nayarit, Mexico · 5 kt SSW	**53%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆4	**La Ligua coast** Chile · 13 kt S	**79%**
◆1	**Ancud coast** Chile · 12 kt N	**77%**
◆3	**Antofagasta coast** Chile · 10 kt S	**76%**
◆2	**Camaná coast** Peru · 11 kt SE	**72%**
◆8	**Soyo coast** Angola · 8 kt S	**70%**
◆10	**Pointe-Noire coast** Congo (Brazzaville) · 6 kt SSE	**70%**
◆18	**Luanda coast** Angola · 7 kt SSE	**69%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.27× the mean swell period.

July: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (154 mornings; 1 days missing from the archive)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in July

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt SSE	**88%**
2	**Iquique** Chile · 7 kt S	**82%**
3	**Mentawais** Sumatra, Indonesia · 4 kt SE	**80%**
4	**Krui** Lampung, Sumatra · 11 kt SE	**80%**
5	**Nias** Sumatra, Indonesia · 5 kt SE	**76%**
6	**Punta Hermosa** Peru · 8 kt SSE	**69%**
7	**Elands Bay** South Africa · 10 kt SSE	**65%**
8	**Santa Catalina** Panama · 6 kt W	**65%**
9	**Puerto Escondido** Oaxaca, Mexico · 6 kt NW	**64%**
10	**Cimaja** Pelabuhan Ratu, West Java · 5 kt E	**63%**
11	**Punta Roca** El Salvador · 7 kt ENE	**62%**
12	**Skeleton Bay** Namibia · 11 kt S	**59%**
13	**Punta de Lobos** Chile · 12 kt S	**58%**
14	**Pascuales** Colima, Mexico · 6 kt NE	**54%**
15	**Pavones** Costa Rica · 6 kt W	**51%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Camaná coast** Peru · 11 kt SE	**84%**
◆8	**Soyo coast** Angola · 7 kt S	**77%**
◆4	**La Ligua coast** Chile · 13 kt S	**76%**
◆9	**Sumbe coast** Angola · 4 kt S	**76%**
◆10	**Pointe-Noire coast** Congo (Brazzaville) · 6 kt S	**76%**
◆3	**Antofagasta coast** Chile · 9 kt S	**75%**
◆7	**Chañaral coast** Chile · 7 kt S	**74%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.28× the mean swell period.

August: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in August

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)

1	**Arica** Chile · 4 kt S	**81%**
2	**Mentawais** Sumatra, Indonesia · 9 kt SE	**79%**
3	**Krui** Lampung, Sumatra · 14 kt SE	**77%**
4	**Iquique** Chile · 8 kt S	**73%**
5	**Nias** Sumatra, Indonesia · 7 kt SE	**72%**
6	**Punta Roca** El Salvador · 6 kt ENE	**70%**
7	**Elands Bay** South Africa · 9 kt N	**70%**
8	**Puerto Escondido** Oaxaca, Mexico · 7 kt NW	**67%**
9	**Skeleton Bay** Namibia · 10 kt S	**66%**
10	**Punta Hermosa** Peru · 8 kt SSE	**66%**
11	**Pascuales** Colima, Mexico · 6 kt E	**61%**
12	**Piha** New Zealand · 14 kt ESE	**61%**
13	**Cimaja** Pelabuhan Ratu, West Java · 5 kt E	**59%**
14	**Cactus Beach** South Australia · 12 kt E	**57%**
15	**Punta de Lobos** Chile · 11 kt S	**55%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆10	**Pointe-Noire coast** Congo (Brazzaville) · 6 kt S	**89%**
◆9	**Sumbe coast** Angola · 4 kt S	**87%**
◆8	**Soyo coast** Angola · 7 kt S	**84%**
◆13	**Omboué coast** Gabon · 7 kt SSW	**82%**
◆18	**Luanda coast** Angola · 6 kt S	**80%**
◆1	**Ancud coast** Chile · 12 kt N	**77%**
◆5	**Toltén coast** Chile · 10 kt SSE	**73%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.29× the mean swell period.

September: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in September

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)

1	**Arica** Chile · 4 kt S	**83%**
2	**Krui** Lampung, Sumatra · 12 kt SE	**79%**
3	**Iquique** Chile · 7 kt S	**72%**
4	**Santa Catalina** Panama · 6 kt SSE	**65%**
5	**Mentawais** Sumatra, Indonesia · 9 kt SE	**64%**
6	**Elands Bay** South Africa · 8 kt SE	**61%**
7	**Punta Hermosa** Peru · 8 kt S	**59%**
8	**Punta Roca** El Salvador · 6 kt ESE	**58%**
9	**Punta de Lobos** Chile · 13 kt S	**58%**
10	**Piha** New Zealand · 15 kt WSW	**55%**
11	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**55%**
12	**Nias** Sumatra, Indonesia · 8 kt SSE	**54%**
13	**Cactus Beach** South Australia · 11 kt ENE	**54%**
14	**Puerto Escondido** Oaxaca, Mexico · 7 kt E	**52%**
15	**Skeleton Bay** Namibia · 10 kt S	**52%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 11 kt WNW	**72%**
◆2	**Camaná coast** Peru · 11 kt SE	**71%**
◆3	**Antofagasta coast** Chile · 10 kt S	**69%**
◆4	**La Ligua coast** Chile · 14 kt S	**69%**
◆13	**Omboué coast** Gabon · 8 kt SSW	**67%**
◆5	**Toltén coast** Chile · 11 kt N	**67%**
◆10	**Pointe-Noire coast** Congo (Brazzaville) · 6 kt SW	**65%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.29× the mean swell period.

October: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (148 mornings; 7 days missing from the archive)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in October

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)

1	**Arica** Chile · 4 kt S	**93%**
2	**Iquique** Chile · 7 kt S	**89%**
3	**Lakey Peak** Sumbawa, Indonesia · 6 kt E	**82%**
4	**Punta Hermosa** Peru · 7 kt SSE	**82%**
5	**G-Land** Java, Indonesia · 7 kt ESE	**81%**
6	**Krui** Lampung, Sumatra · 9 kt SE	**78%**
7	**Desert Point** Lombok, Indonesia · 8 kt ESE	**74%**
8	**Mentawais** Sumatra, Indonesia · 4 kt SE	**74%**
9	**Puerto Escondido** Oaxaca, Mexico · 6 kt E	**65%**
10	**Chicama** Peru · 9 kt SSE	**65%**
11	**Punta Roca** El Salvador · 7 kt ESE	**64%**
12	**Elands Bay** South Africa · 9 kt SSE	**64%**
13	**Piha** New Zealand · 14 kt NW	**63%**
14	**Cimaja** Pelabuhan Ratu, West Java · 3 kt E	**62%**
15	**Montañita** Ecuador · 8 kt SW	**61%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 7 kt SSW	**89%**
◆3	**Antofagasta coast** Chile · 7 kt S	**85%**
◆2	**Camaná coast** Peru · 11 kt SE	**81%**
◆6	**Isabela coast** Galápagos, Ecuador · 12 kt SSE	**77%**
◆21	**Blitar coast** Indonesia · 11 kt E	**77%**
◆4	**La Ligua coast** Chile · 14 kt S	**75%**
◆7	**Chañaral coast** Chile · 7 kt SSW	**75%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 5–16 ft (1.5–5 m, Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Météo-France MFWAM via Open-Meteo and Copernicus Marine, 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.


Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Where swell peak period is missing (Great Lakes, Copernicus years) it is 1.29× the mean swell period.

Lake Michigan: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with surfable windswell



#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Holland**	Michigan	23	17	13	7	2	0	1	0	5	9	22	19	**10%**	Nov–Jan 22%	0%	6 ft @ 5 s WNW	13 kt S
2	**Grand Haven**	Michigan	23	14	11	7	1	0	0	1	3	8	22	21	**9%**	Nov–Jan 22%	0%	6 ft @ 5 s W	13 kt S
3	**Ludington**	Michigan	15	15	12	7	1	1	1	1	3	7	21	21	**8%**	Nov–Jan 19%	0%	6 ft @ 5 s WSW	14 kt S
4	**New Buffalo**	Michigan	15	16	11	7	1	2	1	1	5	13	16	14	**8%**	Dec–Feb 15%	0%	5 ft @ 5 s NNW	13 kt SSW
5	**St. Joseph**	Michigan	17	15	13	6	1	1	2	1	5	11	19	12	**8%**	Nov–Jan 16%	0%	5 ft @ 5 s NW	13 kt SSW
6	**Frankfort**	Michigan	15	13	12	5	2	0	1	1	3	6	19	20	**8%**	Nov–Jan 18%	0%	6 ft @ 5 s WSW	14 kt S
7	**Portage**	Indiana	11	9	9	5	1	1	1	1	6	12	10	8	**6%**	Oct–Dec 10%	0%	4 ft @ 5 s N	13 kt SSW
8	**Gary**	Indiana	11	8	9	5	1	1	1	1	6	12	8	5	**6%**	Jan–Mar 9%	0%	4 ft @ 6 s N	12 kt NW
9	**Milwaukee**	Wisconsin	6	6	9	5	1	1	2	2	5	8	4	4	**4%**	Jan–Mar 7%	0%	5 ft @ 5 s NE	13 kt NW
10	**Whiting**	Indiana	8	4	8	4	1	1	1	1	5	9	7	3	**4%**	Sep–Nov 7%	0%	4 ft @ 6 s N	12 kt NW
11	**Sheboygan**	Wisconsin	5	9	8	7	0	1	1	2	5	7	4	5	**4%**	Feb–Apr 8%	0%	5 ft @ 5 s ENE	14 kt S
12	**Racine**	Wisconsin	6	5	7	2	1	1	1	1	4	5	1	2	**3%**	Jan–Mar 6%	0%	4 ft @ 6 s NE	13 kt WNW
13	**Chicago**	Illinois	5	4	6	3	1	1	1	1	3	6	1	1	**3%**	Jan–Mar 5%	0%	4 ft @ 6 s NNE	13 kt SSW


Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.
Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Mediterranean: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with surfable windswell



Mediterranean: coast by coast

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.


#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Ajaccio**	Corsica, France	54	45	35	31	23	18	23	19	24	21	53	37	**32%**	Nov–Jan 48%	3%	5 ft @ 7 s W	11 kt WNW
2	**Susa**	Cyrenaica, Libya	50	45	37	32	19	13	29	15	24	38	30	38	**31%**	Dec–Feb 44%	1%	4 ft @ 7 s NNW	8 kt W
3	**Capo Mannu**	Sardinia, Italy	47	40	35	29	26	13	22	22	20	25	53	37	**31%**	Nov–Jan 46%	2%	5 ft @ 7 s WNW	11 kt ESE
4	**Alexandria**	Egypt	35	45	39	30	26	19	33	21	21	27	21	33	**29%**	Jan–Mar 40%	1%	4 ft @ 7 s NW	9 kt NW
5	**Tabarka**	Tunisia	51	49	30	33	23	9	19	14	17	23	49	37	**29%**	Dec–Feb 46%	3%	5 ft @ 7 s NW	12 kt W
6	**Paphos**	Cyprus	37	43	36	32	28	15	19	25	27	19	21	34	**28%**	Jan–Mar 39%	1%	4 ft @ 6 s W	9 kt W
7	**Falasarna**	Crete, Greece	45	37	32	31	14	9	19	11	14	14	37	43	**25%**	Nov–Jan 41%	0%	5 ft @ 6 s W	14 kt NE
8	**Benghazi**	Libya	43	32	28	30	19	11	16	8	12	17	33	48	**25%**	Nov–Jan 41%	1%	5 ft @ 7 s NW	10 kt NW
9	**Algiers**	Algeria	34	40	41	29	16	7	11	7	12	16	36	32	**23%**	Jan–Mar 38%	1%	4 ft @ 7 s N	10 kt WSW
10	**Gaza**	Palestine	35	39	33	21	23	7	13	10	10	18	19	35	**22%**	Dec–Feb 36%	1%	4 ft @ 7 s WNW	7 kt WSW
11	**Bizerte**	Tunisia	39	35	23	27	15	5	17	11	11	20	30	31	**22%**	Dec–Feb 35%	1%	5 ft @ 7 s NNW	13 kt WNW
12	**Tripoli**	Libya	43	32	26	30	14	8	10	5	13	18	21	37	**21%**	Dec–Feb 37%	1%	4 ft @ 6 s N	10 kt ESE
13	**Beirut**	Lebanon	37	36	33	21	25	5	7	13	11	13	15	32	**21%**	Jan–Mar 35%	1%	4 ft @ 7 s W	10 kt SW
14	**Livorno**	Tuscany, Italy	33	26	21	19	14	9	13	12	23	15	37	27	**21%**	Nov–Jan 32%	1%	5 ft @ 7 s WSW	10 kt ENE
15	**Kalo Nero**	Messinia, Greece	40	35	21	20	8	3	5	3	9	17	39	39	**20%**	Nov–Jan 39%	1%	4 ft @ 7 s WSW	9 kt NNW
16	**Santa Marinella**	Lazio, Italy	26	23	25	20	12	7	8	10	15	17	27	27	**18%**	Nov–Jan 27%	0%	4 ft @ 6 s SW	11 kt NNE
17	**Marseille**	France	25	26	17	16	9	5	5	5	10	13	26	15	**14%**	Jan–Mar 23%	0%	5 ft @ 6 s WSW	16 kt NNW
18	**Barcelona**	Spain	23	19	37	17	2	3	0	5	8	14	17	12	**13%**	Jan–Mar 26%	1%	4 ft @ 6 s ESE	10 kt W
19	**Varazze**	Liguria, Italy	22	20	21	12	7	5	3	5	7	15	21	14	**12%**	Jan–Mar 21%	1%	4 ft @ 7 s SSW	10 kt N
20	**Kelibia**	Cap Bon, Tunisia	11	9	29	18	14	2	1	1	6	16	9	14	**11%**	Mar–May 20%	0%	4 ft @ 6 s ESE	13 kt NW
21	**Mondello**	Sicily, Italy	14	11	9	11	2	1	3	1	9	10	10	15	**8%**	Dec–Feb 13%	0%	5 ft @ 6 s NNW	9 kt WSW
22	**Hammamet**	Tunisia	6	7	25	15	8	2	1	1	1	9	1	7	**7%**	Mar–May 16%	1%	4 ft @ 6 s ESE	11 kt NW
23	**Kastro**	Kyllini, Greece	14	11	3	5	3	1	0	0	1	3	16	14	**6%**	Nov–Jan 15%	0%	4 ft @ 6 s SW	8 kt NNW
24	**Antalya**	Turkey	14	7	3	2	1	0	0	0	0	0	4	12	**4%**	Dec–Feb 11%	0%	4 ft @ 6 s SSE	4 kt NNW

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.
Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Greece: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.






#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Zakynthos west**	Ionian Islands	48	38	28	33	17	8	17	12	14	24	46	45	**27%**	Nov–Jan 46%	0%	5 ft @ 6 s W	11 kt NNW
2	**Falasarna**	West Crete	45	37	32	32	14	9	18	14	15	15	37	43	**26%**	Nov–Jan 41%	0%	5 ft @ 6 s W	14 kt NE
3	**Komos**	South Crete	36	33	28	27	15	5	8	7	7	11	28	40	**21%**	Dec–Feb 36%	1%	4 ft @ 6 s W	9 kt N
4	**Kalo Nero**	Messinia, W Peloponnese	40	35	25	22	10	5	7	3	9	16	39	39	**20%**	Nov–Jan 39%	1%	4 ft @ 7 s WSW	9 kt NNW
5	**Kokkini Hani**	North Crete	31	30	28	15	14	7	13	10	19	26	23	23	**20%**	Jan–Mar 30%	0%	4 ft @ 6 s N	11 kt W
6	**Chania**	North Crete	22	27	26	16	17	11	14	15	20	30	23	16	**20%**	Jan–Mar 25%	0%	5 ft @ 6 s N	12 kt N
7	**Agios Gordios**	Corfu, west coast	34	28	21	23	12	3	8	4	12	16	33	35	**19%**	Nov–Jan 34%	0%	4 ft @ 6 s WSW	12 kt NNW
8	**Kymi**	East Evia	22	24	23	11	8	1	3	9	15	18	16	17	**14%**	Jan–Mar 23%	0%	5 ft @ 6 s NE	13 kt NNW
9	**Tinos north**	Cyclades	19	23	20	10	9	5	6	8	17	22	16	11	**14%**	Jan–Mar 21%	0%	5 ft @ 5 s N	15 kt NNW
10	**Kastro**	Kyllini, NW Peloponnese	14	13	3	3	1	0	0	0	1	3	16	14	**6%**	Nov–Jan 15%	0%	4 ft @ 6 s SSW	8 kt NNW
11	**Ialysos**	Rhodes, northwest	6	5	4	1	2	0	0	0	0	0	4	2	**2%**	Jan–Mar 5%	0%	4 ft @ 6 s W	9 kt NNW


Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.
Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Caspian Sea: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with surfable windswell




#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Nabran**	Azerbaijan	37	26	19	15	9	5	6	8	17	26	29	36	**19%**	Nov–Jan 34%	0%	5 ft @ 6 s NE	12 kt NNW
2	**Ramsar**	Iran	17	18	15	8	8	7	11	5	15	20	13	14	**13%**	Jan–Mar 17%	0%	4 ft @ 6 s N	5 kt NNW
3	**Novkhani**	Absheron, Azerbaijan	17	19	14	9	6	5	8	8	10	16	18	14	**12%**	Dec–Feb 17%	0%	5 ft @ 6 s N	12 kt NNW
4	**Babolsar**	Iran	15	17	12	8	6	8	14	6	13	12	16	14	**12%**	Dec–Feb 16%	0%	4 ft @ 7 s NNW	7 kt E
5	**Derbent**	Dagestan, Russia	22	9	8	9	3	1	2	6	11	17	15	27	**11%**	Nov–Jan 21%	0%	4 ft @ 6 s E	12 kt NNW
6	**Awaza**	Turkmenistan	12	14	12	8	8	5	8	5	7	9	13	12	**9%**	Jan–Mar 13%	0%	4 ft @ 6 s NW	10 kt NNW
7	**Makhachkala**	Dagestan, Russia	22	8	7	5	1	0	1	1	7	12	15	27	**9%**	Nov–Jan 21%	0%	4 ft @ 6 s ESE	12 kt SE
8	**Aktau**	Kazakhstan	15	10	8	3	6	0	1	0	1	9	20	14	**7%**	Nov–Jan 16%	0%	5 ft @ 6 s W	13 kt ESE
9	**Lankaran**	Azerbaijan	8	8	4	1	1	0	1	1	2	6	4	3	**3%**	Jan–Mar 6%	0%	4 ft @ 6 s ENE	9 kt S
10	**Shikhov**	Baku, Azerbaijan	1	0	0	0	0	0	0	0	0	0	0	1	**0%**	Nov–Jan 1%	0%	4 ft @ 5 s SE	12 kt N


Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.
Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Black Sea: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with surfable windswell



#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Kilyos**	Istanbul, Turkey	33	36	26	18	18	9	12	34	38	34	24	35	**26%**	Aug–Oct 35%	0%	5 ft @ 6 s NNE	11 kt NNE
2	**Şile**	Istanbul, Turkey	32	35	23	15	14	7	10	27	33	33	19	33	**23%**	Dec–Feb 33%	0%	5 ft @ 6 s NNE	10 kt NE
3	**Vama Veche**	Romania	22	32	21	9	13	4	3	15	27	21	16	27	**17%**	Dec–Feb 27%	0%	5 ft @ 6 s ENE	13 kt NNW
4	**Sozopol**	Bulgaria	23	30	18	9	11	3	2	19	27	22	16	24	**17%**	Dec–Feb 26%	0%	4 ft @ 6 s ENE	12 kt N
5	**Batumi**	Georgia	25	27	15	15	5	7	6	3	13	14	19	19	**14%**	Dec–Feb 23%	0%	4 ft @ 7 s WNW	8 kt E
6	**Varna**	Bulgaria	19	25	16	8	10	1	1	14	19	16	9	21	**13%**	Dec–Feb 21%	0%	4 ft @ 6 s ENE	13 kt N
7	**Sochi**	Russia	31	28	9	13	4	4	3	1	6	9	24	25	**13%**	Dec–Feb 28%	0%	5 ft @ 6 s WSW	8 kt SE
8	**Anapa**	Russia	23	17	9	15	3	3	1	0	4	9	16	23	**10%**	Dec–Feb 21%	0%	5 ft @ 6 s WSW	13 kt ENE
9	**Odesa**	Ukraine	12	6	4	3	2	1	1	0	1	1	11	10	**4%**	Nov–Jan 11%	0%	4 ft @ 6 s SSE	12 kt NNE

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.

Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.