
Year-round: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 1,803 mornings, Nov 2021 to Oct 2026



Best named breaks year-round

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Iquique** Chile · 7 kt S	**77%**
2	**Arica** Chile · 4 kt S	**77%**
3	**Piha** New Zealand · 14 kt SW	**62%**
4	**Skeleton Bay** Namibia · 10 kt S	**59%**
5	**Punta Hermosa** Peru · 7 kt SSE	**57%**
6	**Elands Bay** South Africa · 9 kt SSE	**56%**
7	**Punta de Lobos** Chile · 12 kt S	**56%**
8	**Short Sands** Oregon · 14 kt N	**56%**
9	**Montara** California · 12 kt NW	**47%**
10	**Cactus Beach** South Australia · 12 kt SE	**46%**
11	**Krui** Lampung, Sumatra · 10 kt SE	**45%**
12	**Imsouane** Morocco · 11 kt N	**44%**
13	**Jeffreys Bay** South Africa · 13 kt W	**42%**
14	**Chicama** Peru · 9 kt SSE	**41%**
15	**Mentawais** Sumatra, Indonesia · 6 kt SE	**41%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 9 kt N	**77%**
◆2	**Antofagasta coast** Chile · 8 kt S	**77%**
◆3	**La Ligua coast** Chile · 12 kt S	**76%**
◆4	**Camaná coast** Peru · 10 kt SE	**74%**
◆5	**Toltén coast** Chile · 13 kt S	**72%**
◆6	**La Serena coast** Chile · 7 kt SSW	**63%**
◆7	**Chañaral coast** Chile · 6 kt SSW	**59%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s



Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

Popular breaks, month by month

The 40 well-known breaks with the strongest three-month season (best one per 100 miles of coast), listed by year-round consistency. Breaks in the enclosed seas are on the map insets.

#	Break	Region	J	F	M	A	M	J	J	A	S	O	N	D	Year	Worst	Best 3 months	Typical good morning	Morning wind
1	**Iquique**	Chile	73	81	79	83	87	71	77	74	74	82	73	68	**77%**	68%	Mar–May 83%	6 ft @ 13 s SSW, surf ~9 ft	7 kt S
2	**Arica**	Chile	69	72	79	83	85	74	82	76	78	85	72	65	**77%**	65%	Mar–May 83%	6 ft @ 13 s SSW, surf ~9 ft	4 kt S
3	**Piha**	New Zealand	55	68	73	69	68	53	53	58	53	60	73	65	**62%**	53%	Mar–May 70%	7 ft @ 13 s SW, surf ~10 ft	14 kt SW
4	**Skeleton Bay**	Namibia	63	70	47	55	68	61	61	65	57	54	59	53	**59%**	47%	May–Jul 63%	7 ft @ 12 s SSW, surf ~9 ft	10 kt S
5	**Punta Hermosa**	Peru	44	38	64	71	78	59	58	61	57	59	54	37	**57%**	37%	Mar–May 71%	5 ft @ 14 s SSW, surf ~8 ft	7 kt SSE
6	**Elands Bay**	South Africa	54	55	43	47	71	55	63	66	58	51	55	55	**56%**	43%	May–Jul 63%	6 ft @ 13 s SW, surf ~8 ft	9 kt SSE
7	**Punta de Lobos**	Chile	45	44	61	54	68	59	64	57	55	56	53	51	**56%**	44%	May–Jul 64%	8 ft @ 13 s SW, surf ~11 ft	12 kt S
8	**Short Sands**	Oregon	77	79	63	70	61	40	16	19	53	60	69	65	**56%**	16%	Dec–Feb 74%	7 ft @ 12 s WNW, surf ~10 ft	14 kt N
9	**Montara**	California	67	63	50	45	46	37	19	15	39	62	67	56	**47%**	15%	Nov–Jan 63%	7 ft @ 12 s WNW, surf ~10 ft	12 kt NW
10	**Cactus Beach**	South Australia	22	40	41	66	55	46	50	57	53	51	36	37	**46%**	22%	Apr–Jun 56%	7 ft @ 14 s SW, surf ~10 ft	12 kt SE
11	**Krui**	Lampung, Sumatra	23	23	23	42	45	69	77	75	65	64	17	21	**45%**	17%	Jun–Aug 74%	5 ft @ 14 s SSW, surf ~8 ft	10 kt SE
12	**Imsouane**	Morocco	68	60	66	51	25	15	15	16	38	47	57	72	**44%**	15%	Dec–Feb 66%	6 ft @ 12 s NW, surf ~9 ft	11 kt N
13	**Jeffreys Bay**	South Africa	43	43	38	35	49	44	42	45	40	38	44	46	**42%**	35%	May–Jul 45%	8 ft @ 12 s SW, surf ~11 ft	13 kt W
14	**Chicama**	Peru	19	21	47	54	57	41	44	46	51	48	44	25	**41%**	19%	Mar–May 53%	5 ft @ 14 s SSW, surf ~8 ft	9 kt SSE
15	**Mentawais**	Sumatra, Indonesia	24	26	23	42	50	68	74	74	53	9	10	24	**41%**	9%	Jun–Aug 72%	5 ft @ 14 s SSW, surf ~8 ft	6 kt SE
16	**Nias**	Sumatra, Indonesia	22	19	16	41	46	67	70	70	49	49	7	25	**40%**	7%	Jun–Aug 69%	5 ft @ 14 s SSW, surf ~8 ft	6 kt SE
17	**Hossegor**	France	43	73	60	40	21	23	19	26	40	43	43	54	**40%**	19%	Jan–Mar 59%	6 ft @ 13 s NW, surf ~9 ft	9 kt W
18	**G-Land**	Java, Indonesia	41	36	44	53	28	18	16	33	45	62	51	51	**39%**	16%	Oct–Dec 55%	6 ft @ 14 s SSW, surf ~9 ft	10 kt ESE
19	**Shipstern Bluff**	Tasmania, Australia	48	46	46	45	39	34	33	35	31	28	47	36	**39%**	28%	Jan–Mar 47%	8 ft @ 13 s SW, surf ~12 ft	16 kt SW
20	**Ericeira**	Portugal	39	50	53	45	29	25	14	21	37	53	44	52	**38%**	14%	Oct–Dec 50%	7 ft @ 12 s NW, surf ~10 ft	13 kt N
21	**Cimaja**	Pelabuhan Ratu, West Java	24	21	30	45	43	57	54	56	44	30	19	23	**38%**	19%	Jun–Aug 56%	5 ft @ 14 s SSW, surf ~8 ft	4 kt E
22	**Tofino**	British Columbia	37	48	48	47	41	29	9	13	44	42	35	35	**35%**	9%	Feb–Apr 47%	7 ft @ 11 s W, surf ~10 ft	11 kt ESE
23	**Desert Point**	Lombok, Indonesia	35	33	40	43	23	19	14	32	40	58	44	46	**35%**	14%	Oct–Dec 49%	5 ft @ 14 s SSW, surf ~8 ft	10 kt ESE
24	**Bundoran**	Ireland	55	43	40	36	32	28	18	19	36	38	41	38	**35%**	18%	Jan–Mar 46%	7 ft @ 12 s WNW, surf ~10 ft	13 kt S
25	**Safi**	Morocco	48	45	48	33	22	19	13	17	32	44	46	53	**35%**	13%	Nov–Jan 49%	7 ft @ 12 s NW, surf ~10 ft	12 kt NNE
26	**Margaret River**	Western Australia	31	26	26	39	37	33	24	29	36	58	33	35	**33%**	24%	Sep–Nov 42%	9 ft @ 13 s SW, surf ~13 ft	13 kt ESE
27	**Lakey Peak**	Sumbawa, Indonesia	39	33	28	26	19	15	24	41	43	79	24	33	**33%**	15%	Aug–Oct 54%	5 ft @ 14 s SSW, surf ~8 ft	8 kt E
28	**Rapa Nui**	Easter Island	26	28	23	33	37	28	34	37	34	43	35	20	**31%**	20%	Aug–Oct 38%	7 ft @ 14 s SW, surf ~10 ft	11 kt E
29	**Todos Santos**	Baja California	55	55	46	33	24	11	1	0	4	14	33	46	**27%**	0%	Jan–Mar 52%	5 ft @ 13 s WNW, surf ~8 ft	8 kt NW
30	**Teahupo'o**	Tahiti	8	13	22	22	34	32	38	36	21	21	19	13	**24%**	8%	Jun–Aug 35%	6 ft @ 13 s SSW, surf ~9 ft	13 kt E
31	**Saquarema**	Brazil	19	11	22	37	36	31	26	26	24	14	17	14	**23%**	11%	Apr–Jun 35%	6 ft @ 11 s S, surf ~9 ft	8 kt ENE
32	**La Santa**	Lanzarote, Canaries	38	32	34	22	9	7	4	4	15	26	21	39	**21%**	4%	Dec–Feb 36%	6 ft @ 12 s NW, surf ~9 ft	14 kt NNE
33	**Cloud 9**	Siargao, Philippines	30	34	32	39	10	1	1	3	3	6	24	48	**19%**	1%	Dec–Feb 37%	6 ft @ 10 s NE, surf ~8 ft	10 kt ENE
34	**Busua**	Ghana	8	2	1	22	37	32	27	34	33	21	7	3	**19%**	1%	May–Jul 32%	5 ft @ 12 s S, surf ~7 ft	8 kt SSW
35	**Rincon**	California	41	34	29	19	17	11	1	3	5	9	21	32	**19%**	1%	Dec–Feb 36%	5 ft @ 13 s WNW, surf ~8 ft	8 kt NW
36	**Lhoknga**	Aceh, Sumatra	8	3	6	15	26	39	22	39	23	22	1	15	**19%**	1%	Jun–Aug 33%	5 ft @ 15 s SW, surf ~8 ft	11 kt S
37	**Salsa Brava**	Caribbean Costa Rica	30	40	30	13	14	22	33	12	1	0	4	14	**18%**	0%	Jan–Mar 33%	6 ft @ 9 s NE, surf ~8 ft	6 kt WNW
38	**Hanalei Bay**	Kauai, Hawaii	47	38	21	12	7	1	0	1	5	28	20	33	**17%**	0%	Dec–Feb 39%	7 ft @ 13 s NNW, surf ~11 ft	15 kt E
39	**Pipeline**	Oahu, Hawaii	46	30	18	8	6	1	0	0	2	27	18	32	**15%**	0%	Dec–Feb 36%	7 ft @ 13 s NW, surf ~10 ft	15 kt E
40	**Honolua Bay**	Maui, Hawaii	48	28	16	8	6	1	0	1	3	26	21	27	**15%**	0%	Dec–Feb 34%	7 ft @ 13 s NNW, surf ~10 ft	15 kt E

Month cells use the map colours (% of mornings good+). Typical good morning = medians over the good+ mornings: swell height @ peak period and the direction it comes from, and estimated breaking surf (Komar–Gaughan).
Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward.

Each break is read from its nearest seaward model cell (0.5° in East and Southeast Asia, 1° elsewhere) with swell inside its exposure window. The rule has no wind; local reefs, points and sheltering are not modelled.

Under-the-radar stretches, month by month

Coast where mornings are often good+ with swell aimed at the shore (≥40% year-round, or ≥65% over its best three months),
at least 300 km from the well-known breaks and within 150 km of a town of 5,000+. Listed by year-round consistency; numbers match the ◆ markers on the maps.


#	Stretch	Region	J	F	M	A	M	J	J	A	S	O	N	D	Year	Worst	Best 3 months	Typical good morning	Morning wind
◆1	**Ancud coast**	Chile	85	77	75	77	75	73	65	72	72	85	83	90	**77%**	65%	Oct–Dec 86%	7 ft @ 13 s WSW, surf ~10 ft	9 kt N
◆2	**Antofagasta coast**	Chile	77	78	80	84	84	74	77	71	73	83	75	68	**77%**	68%	Mar–May 83%	6 ft @ 13 s SW, surf ~9 ft	8 kt S
◆3	**La Ligua coast**	Chile	72	79	82	79	79	74	77	69	72	83	76	77	**76%**	69%	Feb–Apr 80%	7 ft @ 13 s SW, surf ~10 ft	12 kt S
◆4	**Camaná coast**	Peru	66	74	79	82	81	69	81	72	72	77	71	58	**74%**	58%	Mar–May 80%	6 ft @ 13 s SSW, surf ~9 ft	10 kt SE
◆5	**Toltén coast**	Chile	79	75	72	72	74	67	63	69	73	76	75	74	**72%**	63%	Dec–Feb 76%	8 ft @ 13 s SW, surf ~11 ft	13 kt S
◆6	**La Serena coast**	Chile	69	68	73	72	66	60	55	48	51	72	61	68	**63%**	48%	Feb–Apr 71%	7 ft @ 13 s SW, surf ~10 ft	7 kt SSW
◆7	**Chañaral coast**	Chile	55	52	59	59	68	61	66	60	62	72	51	46	**59%**	46%	May–Jul 65%	5 ft @ 13 s SW, surf ~8 ft	6 kt SSW
◆8	**Greymouth coast**	New Zealand	34	46	50	55	60	49	56	62	60	58	51	47	**52%**	34%	Aug–Oct 60%	6 ft @ 13 s W, surf ~9 ft	9 kt E
◆9	**Marcona coast**	Peru	46	56	64	71	59	39	45	41	38	50	53	41	**50%**	38%	Mar–May 65%	6 ft @ 13 s SSW, surf ~9 ft	13 kt SE
◆10	**Port Lincoln coast**	Australia	40	41	47	51	50	45	47	54	55	54	34	52	**47%**	34%	Aug–Oct 54%	7 ft @ 14 s SW, surf ~10 ft	12 kt SE
◆11	**Meningie coast**	Australia	35	57	51	62	46	38	42	49	46	50	44	48	**47%**	35%	Feb–Apr 57%	7 ft @ 14 s SW, surf ~11 ft	10 kt S
◆12	**Oranjemund coast**	Namibia	52	41	37	33	61	54	50	56	41	38	40	40	**46%**	33%	May–Jul 55%	7 ft @ 13 s SW, surf ~10 ft	9 kt SE
◆14	**Quellón coast**	Chile	63	55	38	42	42	35	33	39	32	55	51	58	**45%**	32%	Dec–Feb 58%	8 ft @ 13 s WSW, surf ~12 ft	13 kt W
◆13	**Nzeto coast**	Angola	26	32	27	50	60	57	67	76	57	38	22	21	**45%**	21%	Jul–Sep 67%	5 ft @ 13 s SSW, surf ~8 ft	7 kt SSE
◆15	**Port Denison coast**	Australia	32	45	37	48	45	49	37	37	45	60	46	35	**42%**	32%	Sep–Nov 50%	9 ft @ 14 s SW, surf ~12 ft	12 kt SSE
◆17	**Nelson coast**	New Zealand	44	55	52	41	42	30	29	44	31	31	53	43	**41%**	29%	Jan–Mar 50%	8 ft @ 13 s WSW, surf ~11 ft	8 kt SSW
◆19	**Bredasdorp coast**	South Africa	46	44	39	44	48	35	37	32	38	32	47	46	**41%**	32%	Nov–Jan 46%	8 ft @ 12 s SW, surf ~12 ft	10 kt W
◆18	**Kaitaia coast**	New Zealand	36	49	52	36	32	27	33	39	41	41	53	50	**41%**	27%	Oct–Dec 48%	7 ft @ 14 s SW, surf ~11 ft	13 kt S
◆16	**Blitar coast**	Indonesia	43	34	42	53	31	25	17	35	56	70	44	47	**41%**	17%	Sep–Nov 57%	6 ft @ 14 s SSW, surf ~9 ft	12 kt ESE
◆20	**Pointe-Noire coast**	Congo (Brazzaville)	21	23	20	41	59	55	59	72	52	31	27	19	**40%**	19%	Jun–Aug 62%	5 ft @ 13 s SSW, surf ~8 ft	6 kt S
◆21	**Huarmey coast**	Peru	43	31	53	53	50	31	34	32	41	37	38	30	**40%**	30%	Mar–May 52%	6 ft @ 14 s SSW, surf ~9 ft	10 kt SSE
◆22	**Gwadar coast**	Pakistan	0	0	0	0	6	67	95	86	33	3	0	0	**25%**	0%	Jun–Aug 82%	8 ft @ 11 s SSW, surf ~10 ft	8 kt WSW
◆23	**Salalah coast**	Oman	0	0	0	0	7	55	95	66	21	0	0	0	**21%**	0%	Jun–Aug 72%	7 ft @ 10 s S, surf ~10 ft	6 kt SW
◆24	**Karachi coast**	Pakistan	0	0	0	0	2	43	86	69	28	1	0	0	**20%**	0%	Jun–Aug 66%	8 ft @ 11 s SW, surf ~11 ft	10 kt W
◆25	**Chabahar coast**	Iran	0	0	0	0	2	46	92	70	15	3	0	0	**20%**	0%	Jun–Aug 69%	7 ft @ 11 s S, surf ~9 ft	8 kt ESE

Month cells use the map colours (% of mornings good+). Typical good morning = medians over the good+ mornings: swell height @ peak period and the direction it comes from, and estimated breaking surf (Komar–Gaughan).
Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward.

Each stretch is read from one WW3 cell 50–120 km offshore. The rule has no wind, which flatters cold, steady coasts (Chile, Peru, Namibia) and storm coasts (southern Chile, New Zealand's West Coast); check local wind and access before booking a trip.

October–March: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 889 mornings, Nov 2021 to Oct 2026, these months only



Best named breaks, October–March

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Iquique** Chile · 7 kt S	**76%**
2	**Arica** Chile · 5 kt S	**73%**
3	**Short Sands** Oregon · 14 kt S	**69%**
4	**Piha** New Zealand · 13 kt SSW	**66%**
5	**Imsouane** Morocco · 8 kt NNE	**62%**
6	**Montara** California · 11 kt NNW	**61%**
7	**Skeleton Bay** Namibia · 8 kt S	**58%**
8	**Hossegor** France · 10 kt SSW	**53%**
9	**Elands Bay** South Africa · 9 kt S	**52%**
10	**Punta de Lobos** Chile · 12 kt SSW	**52%**
11	**Punta Hermosa** Peru · 7 kt SSE	**49%**
12	**Ericeira** Portugal · 14 kt NNW	**48%**
13	**Safi** Morocco · 11 kt ENE	**48%**
14	**G-Land** Java, Indonesia · 9 kt W	**47%**
15	**Todos Santos** Baja California · 6 kt NW	**43%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 8 kt SE	**82%**
◆3	**La Ligua coast** Chile · 11 kt S	**78%**
◆2	**Antofagasta coast** Chile · 7 kt SSW	**77%**
◆5	**Toltén coast** Chile · 13 kt S	**75%**
◆4	**Camaná coast** Peru · 9 kt SE	**71%**
◆6	**La Serena coast** Chile · 5 kt SSW	**69%**
◆7	**Chañaral coast** Chile · 5 kt SSW	**55%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

April–September: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 914 mornings, Apr 2022 to Sep 2026, these months only



Best named breaks, April–September

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt SSE	**80%**
2	**Iquique** Chile · 7 kt S	**78%**
3	**Punta Hermosa** Peru · 8 kt S	**64%**
4	**Krui** Lampung, Sumatra · 9 kt SE	**62%**
5	**Skeleton Bay** Namibia · 10 kt S	**61%**
6	**Mentawais** Sumatra, Indonesia · 5 kt SE	**60%**
7	**Elands Bay** South Africa · 8 kt SSE	**60%**
8	**Punta de Lobos** Chile · 12 kt S	**60%**
9	**Piha** New Zealand · 15 kt WSW	**59%**
10	**Nias** Sumatra, Indonesia · 5 kt SE	**57%**
11	**Cactus Beach** South Australia · 12 kt ENE	**55%**
12	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**50%**
13	**Chicama** Peru · 10 kt SSE	**49%**
14	**Short Sands** Oregon · 13 kt N	**43%**
15	**Jeffreys Bay** South Africa · 12 kt W	**43%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Antofagasta coast** Chile · 10 kt S	**77%**
◆4	**Camaná coast** Peru · 11 kt SE	**76%**
◆3	**La Ligua coast** Chile · 13 kt S	**75%**
◆1	**Ancud coast** Chile · 11 kt N	**72%**
◆5	**Toltén coast** Chile · 12 kt SSE	**70%**
◆7	**Chañaral coast** Chile · 8 kt SSW	**63%**
◆13	**Nzeto coast** Angola · 7 kt SSE	**61%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

November: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (135 mornings; 15 days missing from the archive)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in November

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Iquique** Chile · 7 kt S	**73%**
2	**Piha** New Zealand · 11 kt SW	**73%**
3	**Arica** Chile · 5 kt S	**72%**
4	**Short Sands** Oregon · 11 kt SSE	**69%**
5	**Montara** California · 11 kt NNW	**67%**
6	**Skeleton Bay** Namibia · 7 kt S	**59%**
7	**Imsouane** Morocco · 5 kt NNE	**57%**
8	**Elands Bay** South Africa · 9 kt S	**55%**
9	**Punta Hermosa** Peru · 6 kt SSE	**54%**
10	**Punta de Lobos** Chile · 11 kt SSW	**53%**
11	**G-Land** Java, Indonesia · 6 kt E	**51%**
12	**Shipstern Bluff** Tasmania, Australia · 15 kt WNW	**47%**
13	**Safi** Morocco · 11 kt NE	**46%**
14	**Nazaré** Portugal · 13 kt N	**44%**
15	**Jeffreys Bay** South Africa · 13 kt WSW	**44%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 10 kt NNW	**83%**
◆3	**La Ligua coast** Chile · 12 kt S	**76%**
◆2	**Antofagasta coast** Chile · 9 kt SSW	**75%**
◆5	**Toltén coast** Chile · 11 kt S	**75%**
◆4	**Camaná coast** Peru · 10 kt SE	**71%**
◆6	**La Serena coast** Chile · 8 kt SSW	**61%**
◆9	**Marcona coast** Peru · 12 kt SE	**53%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

December: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (155 mornings)



Best named breaks in December

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Imsouane** Morocco · 7 kt NNE	**72%**
2	**Iquique** Chile · 7 kt S	**68%**
3	**Arica** Chile · 5 kt S	**65%**
4	**Piha** New Zealand · 11 kt N	**65%**
5	**Short Sands** Oregon · 14 kt SSE	**65%**
6	**Montara** California · 11 kt NNW	**56%**
7	**Elands Bay** South Africa · 9 kt S	**55%**
8	**Hossegor** France · 9 kt SE	**54%**
9	**Safi** Morocco · 12 kt ENE	**53%**
10	**Skeleton Bay** Namibia · 9 kt S	**53%**
11	**Uluwatu** Bali, Indonesia · 9 kt W	**52%**
12	**Ericeira** Portugal · 14 kt ENE	**52%**
13	**Punta de Lobos** Chile · 12 kt SSW	**51%**
14	**Cloud 9** Siargao, Philippines · 10 kt ENE	**48%**
15	**Todos Santos** Baja California · 5 kt NNW	**46%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 8 kt SSW	**90%**
◆3	**La Ligua coast** Chile · 12 kt S	**77%**
◆5	**Toltén coast** Chile · 14 kt S	**74%**
◆2	**Antofagasta coast** Chile · 7 kt S	**68%**
◆6	**La Serena coast** Chile · 6 kt SSW	**68%**
◆4	**Camaná coast** Peru · 9 kt SE	**58%**
◆14	**Quellón coast** Chile · 13 kt SW	**58%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

January: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Best named breaks in January

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Short Sands** Oregon · 14 kt S	**77%**
2	**Iquique** Chile · 6 kt S	**73%**
3	**Arica** Chile · 5 kt S	**69%**
4	**Imsouane** Morocco · 9 kt NNE	**68%**
5	**Montara** California · 13 kt SSE	**67%**
6	**Skeleton Bay** Namibia · 7 kt S	**63%**
7	**Todos Santos** Baja California · 5 kt SE	**55%**
8	**Bundoran** Ireland · 13 kt SSE	**55%**
9	**Piha** New Zealand · 13 kt ENE	**55%**
10	**Elands Bay** South Africa · 8 kt SSE	**54%**
11	**Shipstern Bluff** Tasmania, Australia · 11 kt SW	**48%**
12	**Honolua Bay** Maui, Hawaii · 14 kt E	**48%**
13	**Safi** Morocco · 11 kt ENE	**48%**
14	**Hanalei Bay** Kauai, Hawaii · 12 kt E	**47%**
15	**Pipeline** Oahu, Hawaii · 14 kt ENE	**46%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 8 kt SE	**85%**
◆5	**Toltén coast** Chile · 14 kt S	**79%**
◆2	**Antofagasta coast** Chile · 5 kt SSW	**77%**
◆3	**La Ligua coast** Chile · 10 kt S	**72%**
◆6	**La Serena coast** Chile · 4 kt SSW	**69%**
◆4	**Camaná coast** Peru · 9 kt ESE	**66%**
◆14	**Quellón coast** Chile · 11 kt SSW	**63%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

February: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (141 mornings)



Best named breaks in February

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Iquique** Chile · 7 kt S	**81%**
2	**Short Sands** Oregon · 13 kt SSE	**79%**
3	**Hossegor** France · 8 kt SSW	**73%**
4	**Arica** Chile · 5 kt S	**72%**
5	**Skeleton Bay** Namibia · 9 kt S	**70%**
6	**Piha** New Zealand · 11 kt SSW	**68%**
7	**Montara** California · 11 kt NNW	**63%**
8	**Imsouane** Morocco · 6 kt NNE	**60%**
9	**Todos Santos** Baja California · 6 kt NW	**55%**
10	**Elands Bay** South Africa · 9 kt S	**55%**
11	**Nazaré** Portugal · 12 kt WNW	**52%**
12	**Tofino** British Columbia · 10 kt ESE	**48%**
13	**Shipstern Bluff** Tasmania, Australia · 16 kt SW	**46%**
14	**Safi** Morocco · 11 kt ENE	**45%**
15	**Punta de Lobos** Chile · 14 kt SSW	**44%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆3	**La Ligua coast** Chile · 12 kt S	**79%**
◆2	**Antofagasta coast** Chile · 6 kt SSW	**78%**
◆1	**Ancud coast** Chile · 7 kt SE	**77%**
◆5	**Toltén coast** Chile · 14 kt S	**75%**
◆4	**Camaná coast** Peru · 8 kt ESE	**74%**
◆6	**La Serena coast** Chile · 5 kt SSW	**68%**
◆11	**Meningie coast** Australia · 10 kt SE	**57%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

March: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Best named breaks in March

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Iquique** Chile · 7 kt S	**79%**
2	**Arica** Chile · 5 kt S	**79%**
3	**Piha** New Zealand · 15 kt SSW	**73%**
4	**Imsouane** Morocco · 11 kt N	**66%**
5	**Punta Hermosa** Peru · 7 kt SSE	**64%**
6	**Short Sands** Oregon · 16 kt S	**63%**
7	**Punta de Lobos** Chile · 14 kt SSW	**61%**
8	**Hossegor** France · 13 kt W	**60%**
9	**Ericeira** Portugal · 12 kt SSW	**53%**
10	**Montara** California · 13 kt NNW	**50%**
11	**Safi** Morocco · 13 kt NE	**48%**
12	**Tofino** British Columbia · 12 kt ESE	**48%**
13	**Skeleton Bay** Namibia · 11 kt S	**47%**
14	**Chicama** Peru · 10 kt SSE	**47%**
15	**Todos Santos** Baja California · 10 kt NNW	**46%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆3	**La Ligua coast** Chile · 11 kt S	**82%**
◆2	**Antofagasta coast** Chile · 7 kt S	**80%**
◆4	**Camaná coast** Peru · 9 kt ESE	**79%**
◆1	**Ancud coast** Chile · 9 kt ESE	**75%**
◆6	**La Serena coast** Chile · 6 kt S	**73%**
◆5	**Toltén coast** Chile · 15 kt S	**72%**
◆9	**Marcona coast** Peru · 12 kt SE	**64%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

April: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Best named breaks in April

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt SSE	**83%**
2	**Iquique** Chile · 8 kt S	**83%**
3	**Punta Hermosa** Peru · 8 kt SSE	**71%**
4	**Short Sands** Oregon · 15 kt N	**70%**
5	**Piha** New Zealand · 16 kt ENE	**69%**
6	**Cactus Beach** South Australia · 11 kt ENE	**66%**
7	**Skeleton Bay** Namibia · 10 kt S	**55%**
8	**Chicama** Peru · 10 kt SE	**54%**
9	**Punta de Lobos** Chile · 12 kt S	**54%**
10	**G-Land** Java, Indonesia · 9 kt ESE	**53%**
11	**Imsouane** Morocco · 10 kt N	**51%**
12	**Nazaré** Portugal · 14 kt N	**47%**
13	**Elands Bay** South Africa · 8 kt S	**47%**
14	**Tofino** British Columbia · 14 kt NW	**47%**
15	**Shipstern Bluff** Tasmania, Australia · 16 kt WSW	**45%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Antofagasta coast** Chile · 10 kt S	**84%**
◆4	**Camaná coast** Peru · 11 kt ESE	**82%**
◆3	**La Ligua coast** Chile · 12 kt S	**79%**
◆1	**Ancud coast** Chile · 9 kt N	**77%**
◆5	**Toltén coast** Chile · 13 kt SSE	**72%**
◆6	**La Serena coast** Chile · 7 kt S	**72%**
◆9	**Marcona coast** Peru · 14 kt SE	**71%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

May: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Best named breaks in May

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Iquique** Chile · 7 kt S	**87%**
2	**Arica** Chile · 3 kt SSE	**85%**
3	**Punta Hermosa** Peru · 7 kt S	**78%**
4	**Elands Bay** South Africa · 8 kt SSE	**71%**
5	**Piha** New Zealand · 15 kt S	**68%**
6	**Punta de Lobos** Chile · 11 kt S	**68%**
7	**Skeleton Bay** Namibia · 10 kt S	**68%**
8	**Short Sands** Oregon · 14 kt NNW	**61%**
9	**Chicama** Peru · 10 kt SE	**57%**
10	**Cactus Beach** South Australia · 11 kt E	**55%**
11	**Mentawais** Sumatra, Indonesia · 5 kt WSW	**50%**
12	**Jeffreys Bay** South Africa · 9 kt W	**49%**
13	**Nias** Sumatra, Indonesia · 5 kt W	**46%**
14	**Montara** California · 16 kt NW	**46%**
15	**Krui** Lampung, Sumatra · 9 kt ESE	**45%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Antofagasta coast** Chile · 10 kt S	**84%**
◆4	**Camaná coast** Peru · 11 kt SE	**81%**
◆3	**La Ligua coast** Chile · 12 kt S	**79%**
◆1	**Ancud coast** Chile · 8 kt N	**75%**
◆5	**Toltén coast** Chile · 14 kt SSE	**74%**
◆7	**Chañaral coast** Chile · 8 kt SSW	**68%**
◆6	**La Serena coast** Chile · 9 kt SSW	**66%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.27× the mean period.

June: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Best named breaks in June

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 3 kt SSE	**74%**
2	**Iquique** Chile · 7 kt S	**71%**
3	**Krui** Lampung, Sumatra · 9 kt ESE	**69%**
4	**Mentawais** Sumatra, Indonesia · 4 kt SE	**68%**
5	**Nias** Sumatra, Indonesia · 4 kt SE	**67%**
6	**Skeleton Bay** Namibia · 10 kt SSE	**61%**
7	**Punta de Lobos** Chile · 14 kt S	**59%**
8	**Punta Hermosa** Peru · 8 kt S	**59%**
9	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**57%**
10	**Elands Bay** South Africa · 8 kt SE	**55%**
11	**Piha** New Zealand · 15 kt NE	**53%**
12	**Cactus Beach** South Australia · 14 kt N	**46%**
13	**Jeffreys Bay** South Africa · 12 kt W	**44%**
14	**Chicama** Peru · 10 kt SSE	**41%**
15	**Short Sands** Oregon · 14 kt N	**40%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Antofagasta coast** Chile · 10 kt S	**74%**
◆3	**La Ligua coast** Chile · 13 kt S	**74%**
◆1	**Ancud coast** Chile · 12 kt N	**73%**
◆4	**Camaná coast** Peru · 11 kt SE	**69%**
◆5	**Toltén coast** Chile · 13 kt N	**67%**
◆22	**Gwadar coast** Pakistan · 8 kt WSW	**67%**
◆7	**Chañaral coast** Chile · 10 kt SSW	**61%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.27× the mean period.

July: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (154 mornings; 1 days missing from the archive)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in July

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt SSE	**82%**
2	**Iquique** Chile · 7 kt S	**77%**
3	**Krui** Lampung, Sumatra · 11 kt SE	**77%**
4	**Mentawais** Sumatra, Indonesia · 4 kt SE	**74%**
5	**Nias** Sumatra, Indonesia · 5 kt SE	**70%**
6	**Punta de Lobos** Chile · 12 kt S	**64%**
7	**Elands Bay** South Africa · 10 kt SSE	**63%**
8	**Skeleton Bay** Namibia · 11 kt S	**61%**
9	**Punta Hermosa** Peru · 8 kt SSE	**58%**
10	**Cimaja** Pelabuhan Ratu, West Java · 5 kt E	**54%**
11	**Piha** New Zealand · 15 kt SW	**53%**
12	**Cactus Beach** South Australia · 13 kt WSW	**50%**
13	**Chicama** Peru · 11 kt SSE	**44%**
14	**Jeffreys Bay** South Africa · 13 kt W	**42%**
15	**Teahupo'o** Tahiti · 14 kt ENE	**38%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆22	**Gwadar coast** Pakistan · 7 kt WSW	**95%**
◆23	**Salalah coast** Oman · 7 kt SW	**95%**
◆25	**Chabahar coast** Iran · 8 kt ESE	**92%**
◆24	**Karachi coast** Pakistan · 11 kt WSW	**86%**
◆4	**Camaná coast** Peru · 11 kt SE	**81%**
◆2	**Antofagasta coast** Chile · 9 kt S	**77%**
◆3	**La Ligua coast** Chile · 13 kt S	**77%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

August: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in August

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt S	**76%**
2	**Krui** Lampung, Sumatra · 14 kt SE	**75%**
3	**Mentawais** Sumatra, Indonesia · 9 kt SE	**74%**
4	**Iquique** Chile · 8 kt S	**74%**
5	**Nias** Sumatra, Indonesia · 7 kt SE	**70%**
6	**Elands Bay** South Africa · 9 kt N	**66%**
7	**Skeleton Bay** Namibia · 10 kt S	**65%**
8	**Punta Hermosa** Peru · 8 kt SSE	**61%**
9	**Piha** New Zealand · 14 kt ESE	**58%**
10	**Cactus Beach** South Australia · 12 kt E	**57%**
11	**Punta de Lobos** Chile · 11 kt S	**57%**
12	**Cimaja** Pelabuhan Ratu, West Java · 5 kt E	**56%**
13	**Chicama** Peru · 10 kt SSE	**46%**
14	**Jeffreys Bay** South Africa · 13 kt W	**45%**
15	**Lakey Peak** Sumbawa, Indonesia · 11 kt ESE	**41%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆22	**Gwadar coast** Pakistan · 8 kt WSW	**86%**
◆13	**Nzeto coast** Angola · 7 kt SSE	**76%**
◆4	**Camaná coast** Peru · 11 kt SE	**72%**
◆20	**Pointe-Noire coast** Congo (Brazzaville) · 6 kt S	**72%**
◆1	**Ancud coast** Chile · 12 kt N	**72%**
◆2	**Antofagasta coast** Chile · 11 kt S	**71%**
◆25	**Chabahar coast** Iran · 9 kt ESE	**70%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

September: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in September

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt S	**78%**
2	**Iquique** Chile · 7 kt S	**74%**
3	**Krui** Lampung, Sumatra · 12 kt SE	**65%**
4	**Elands Bay** South Africa · 8 kt SE	**58%**
5	**Skeleton Bay** Namibia · 10 kt S	**57%**
6	**Punta Hermosa** Peru · 8 kt S	**57%**
7	**Punta de Lobos** Chile · 13 kt S	**55%**
8	**Piha** New Zealand · 15 kt WSW	**53%**
9	**Cactus Beach** South Australia · 11 kt ENE	**53%**
10	**Short Sands** Oregon · 13 kt N	**53%**
11	**Mentawais** Sumatra, Indonesia · 9 kt SE	**53%**
12	**Chicama** Peru · 9 kt SSE	**51%**
13	**Nias** Sumatra, Indonesia · 8 kt SSE	**49%**
14	**G-Land** Java, Indonesia · 11 kt ESE	**45%**
15	**Tofino** British Columbia · 8 kt ESE	**44%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Antofagasta coast** Chile · 10 kt S	**73%**
◆5	**Toltén coast** Chile · 11 kt N	**73%**
◆1	**Ancud coast** Chile · 11 kt WNW	**72%**
◆3	**La Ligua coast** Chile · 14 kt S	**72%**
◆4	**Camaná coast** Peru · 11 kt SE	**72%**
◆7	**Chañaral coast** Chile · 7 kt SSW	**62%**
◆8	**Greymouth coast** New Zealand · 14 kt E	**60%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

October: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (148 mornings; 7 days missing from the archive)



Best named breaks in October

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt S	**85%**
2	**Iquique** Chile · 7 kt S	**82%**
3	**Lakey Peak** Sumbawa, Indonesia · 6 kt E	**79%**
4	**Krui** Lampung, Sumatra · 9 kt SE	**64%**
5	**Montara** California · 11 kt NNW	**62%**
6	**G-Land** Java, Indonesia · 7 kt ESE	**62%**
7	**Piha** New Zealand · 14 kt NW	**60%**
8	**Short Sands** Oregon · 16 kt S	**60%**
9	**Punta Hermosa** Peru · 7 kt SSE	**59%**
10	**Desert Point** Lombok, Indonesia · 8 kt ESE	**58%**
11	**Margaret River** Western Australia · 9 kt SSW	**58%**
12	**Punta de Lobos** Chile · 14 kt SSW	**56%**
13	**Skeleton Bay** Namibia · 9 kt S	**54%**
14	**Ericeira** Portugal · 13 kt NNE	**53%**
15	**Elands Bay** South Africa · 9 kt SSE	**51%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 7 kt SSW	**85%**
◆2	**Antofagasta coast** Chile · 7 kt S	**83%**
◆3	**La Ligua coast** Chile · 14 kt S	**83%**
◆4	**Camaná coast** Peru · 11 kt SE	**77%**
◆5	**Toltén coast** Chile · 12 kt S	**76%**
◆6	**La Serena coast** Chile · 7 kt SSW	**72%**
◆7	**Chañaral coast** Chile · 7 kt SSW	**72%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s



Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.


Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km.

Lake Michigan: when and where it's good

Good lake surf = total wave height 4–10 ft (1.22–3 m) with dominant period ≥ 6 s (taken as 1.29× the mean period), arriving within 75° of the shore's facing, on the ~08:00 sample. Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with good lake surf




#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical good morning	Morning wind
1	**Holland**	Michigan	30	22	17	11	3	2	1	1	5	–	28	26	pending			5 ft @ 7 s W	13 kt S
2	**Grand Haven**	Michigan	28	22	15	10	2	1	0	1	4	–	27	26	pending			5 ft @ 7 s W	13 kt S
3	**Ludington**	Michigan	23	21	12	9	3	1	1	1	3	–	28	27	pending			5 ft @ 7 s W	14 kt S
4	**Frankfort**	Michigan	23	20	14	9	3	1	1	1	4	–	26	25	pending			5 ft @ 7 s W	14 kt S
5	**St. Joseph**	Michigan	24	16	14	6	2	2	2	1	5	–	24	14	pending			5 ft @ 7 s NW	13 kt SSW
6	**New Buffalo**	Michigan	19	14	12	5	2	2	1	1	5	–	16	10	pending			5 ft @ 7 s NNW	13 kt SSW
7	**Portage**	Indiana	10	7	7	3	1	1	1	1	4	–	6	5	pending			5 ft @ 7 s N	13 kt SSW
8	**Milwaukee**	Wisconsin	6	7	8	6	1	1	1	2	5	–	5	5	pending			5 ft @ 7 s ENE	13 kt NW
9	**Sheboygan**	Wisconsin	6	7	9	7	1	1	0	1	4	–	4	5	pending			5 ft @ 7 s E	14 kt S
10	**Gary**	Indiana	8	5	6	3	1	1	1	1	3	–	5	4	pending			5 ft @ 7 s N	12 kt NW
11	**Racine**	Wisconsin	6	3	3	3	1	1	1	1	3	–	2	2	pending			5 ft @ 7 s ENE	13 kt WNW
12	**Whiting**	Indiana	5	3	6	1	0	1	0	1	3	–	1	2	pending			5 ft @ 7 s N	12 kt NW
13	**Chicago**	Illinois	3	2	4	2	0	1	0	1	2	–	0	0	pending			4 ft @ 7 s NNE	13 kt SSW

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical good morning = medians over good mornings: total wave height @ dominant period and the direction waves come from. Months marked – are still downloading.


Coasts are ranked by year-round good %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Lake Superior: when and where it's good

Good lake surf = total wave height 4–10 ft (1.22–3 m) with dominant period ≥ 6 s (taken as 1.29× the mean period), arriving within 75° of the shore's facing, on the ~08:00 sample. Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with good lake surf



#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical good morning	Morning wind
1	**Grand Marais MI**	Michigan	28	18	13	8	1	3	0	3	6	–	27	23	pending			5 ft @ 7 s NW	14 kt W
2	**Whitefish Point**	Michigan	25	12	9	7	1	2	0	3	3	–	24	27	pending			5 ft @ 7 s NW	14 kt SSE
3	**Marquette**	Michigan	21	17	17	11	1	1	0	2	5	–	21	19	pending			6 ft @ 7 s N	15 kt S
4	**Au Train**	Michigan	19	16	14	6	1	1	0	2	5	–	24	19	pending			5 ft @ 7 s NNW	15 kt S
5	**Ontonagon**	Michigan	19	9	12	11	1	0	0	1	3	–	23	18	pending			5 ft @ 7 s NW	13 kt WSW
6	**Agawa Bay**	Ontario	20	7	4	3	1	1	0	1	2	–	21	26	pending			5 ft @ 8 s W	11 kt SSE
7	**Terrace Bay**	Ontario	12	9	6	4	2	1	1	1	4	–	20	15	pending			5 ft @ 7 s SSW	13 kt NNE
8	**Little Girl's Point**	Michigan	6	3	8	7	1	0	0	0	0	–	7	7	pending			5 ft @ 7 s N	13 kt W
9	**Stoney Point**	Minnesota	2	3	4	5	1	0	0	0	0	–	1	2	pending			6 ft @ 7 s ENE	8 kt SSW
10	**Park Point**	Duluth, Minnesota	2	2	5	5	1	0	0	0	0	–	1	2	pending			6 ft @ 7 s ENE	8 kt SSW
11	**Grand Marais MN**	Minnesota	3	1	0	1	1	0	0	0	0	–	2	1	pending			5 ft @ 7 s S	12 kt ENE

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical good morning = medians over good mornings: total wave height @ dominant period and the direction waves come from. Months marked – are still downloading.


Coasts are ranked by year-round good %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Mediterranean: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with surfable windswell



Mediterranean: coast by coast

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.


#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Ajaccio**	Corsica, France	54	45	35	31	23	18	23	19	24	–	53	37	pending			5 ft @ 7 s W	11 kt WNW
2	**Capo Mannu**	Sardinia, Italy	47	40	35	29	26	13	22	22	20	–	53	37	pending			5 ft @ 7 s WNW	11 kt ESE
3	**Susa**	Cyrenaica, Libya	50	45	37	32	19	13	29	15	24	–	30	38	pending			4 ft @ 7 s NNW	8 kt W
4	**Tabarka**	Tunisia	51	49	30	33	23	9	19	14	17	–	49	37	pending			5 ft @ 7 s NW	12 kt W
5	**Alexandria**	Egypt	35	45	39	30	26	19	33	21	21	–	21	33	pending			4 ft @ 7 s NW	9 kt NW
6	**Paphos**	Cyprus	37	43	36	32	28	15	19	25	27	–	21	34	pending			4 ft @ 6 s W	9 kt W
7	**Falasarna**	Crete, Greece	45	37	32	31	14	9	19	11	14	–	37	43	pending			5 ft @ 6 s W	14 kt NE
8	**Benghazi**	Libya	43	32	28	30	19	11	16	8	12	–	33	48	pending			5 ft @ 6 s NW	10 kt NW
9	**Algiers**	Algeria	34	40	41	29	16	7	11	7	12	–	36	32	pending			4 ft @ 7 s N	10 kt WSW
10	**Gaza**	Palestine	35	39	33	21	23	7	13	10	10	–	19	35	pending			4 ft @ 7 s WNW	7 kt WSW
11	**Bizerte**	Tunisia	39	35	23	27	15	5	17	11	11	–	30	31	pending			5 ft @ 7 s NNW	13 kt WNW
12	**Tripoli**	Libya	43	32	26	30	14	8	10	5	13	–	21	37	pending			4 ft @ 6 s N	10 kt ESE
13	**Beirut**	Lebanon	37	36	33	21	25	5	7	13	11	–	15	32	pending			4 ft @ 7 s W	10 kt SW
14	**Livorno**	Tuscany, Italy	33	26	21	19	14	9	13	12	23	–	37	27	pending			5 ft @ 7 s WSW	10 kt ENE
15	**Kalo Nero**	Messinia, Greece	40	35	21	20	8	3	5	3	9	–	39	39	pending			4 ft @ 7 s WSW	9 kt NNW
16	**Santa Marinella**	Lazio, Italy	26	23	25	20	12	7	8	10	15	–	27	27	pending			4 ft @ 6 s SW	11 kt NNE
17	**Marseille**	France	25	26	17	16	9	5	5	5	10	–	26	15	pending			5 ft @ 6 s WSW	16 kt NNW
18	**Barcelona**	Spain	23	19	37	17	2	3	0	5	8	–	17	12	pending			4 ft @ 7 s ESE	10 kt W
19	**Varazze**	Liguria, Italy	22	20	21	12	7	5	3	5	7	–	21	14	pending			4 ft @ 7 s SSW	10 kt N
20	**Kelibia**	Cap Bon, Tunisia	11	9	29	18	14	2	1	1	6	–	9	14	pending			4 ft @ 6 s ESE	13 kt NW
21	**Mondello**	Sicily, Italy	14	11	9	11	2	1	3	1	9	–	10	15	pending			4 ft @ 6 s NNW	9 kt WSW
22	**Hammamet**	Tunisia	6	7	25	15	8	2	1	1	1	–	1	7	pending			4 ft @ 7 s ESE	11 kt NW
23	**Kastro**	Kyllini, Greece	14	11	3	5	3	1	0	0	1	–	16	14	pending			4 ft @ 6 s SW	8 kt NNW
24	**Antalya**	Turkey	14	7	3	2	1	0	0	0	0	–	4	12	pending			4 ft @ 6 s SSE	4 kt NNW

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from. Months marked – are still downloading.
Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Greece: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.






#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Zakynthos west**	Ionian Islands	48	38	28	33	17	8	17	12	14	–	46	45	pending			5 ft @ 6 s W	11 kt NNW
2	**Falasarna**	West Crete	45	37	32	32	14	9	18	14	15	–	37	43	pending			5 ft @ 6 s W	14 kt NE
3	**Komos**	South Crete	36	33	28	27	15	5	8	7	7	–	28	40	pending			5 ft @ 6 s W	9 kt N
4	**Kalo Nero**	Messinia, W Peloponnese	40	35	25	22	10	5	7	3	9	–	39	39	pending			4 ft @ 7 s WSW	9 kt NNW
5	**Kokkini Hani**	North Crete	31	30	28	15	14	7	13	10	19	–	23	23	pending			4 ft @ 6 s N	11 kt W
6	**Agios Gordios**	Corfu, west coast	34	28	21	23	12	3	8	4	12	–	33	35	pending			4 ft @ 6 s WSW	12 kt NNW
7	**Chania**	North Crete	22	27	26	16	17	11	14	15	20	–	23	16	pending			5 ft @ 6 s N	12 kt N
8	**Kymi**	East Evia	22	24	23	11	8	1	3	9	15	–	16	17	pending			5 ft @ 6 s NE	13 kt NNW
9	**Tinos north**	Cyclades	19	23	20	10	9	5	6	8	17	–	16	11	pending			5 ft @ 5 s N	15 kt NNW
10	**Kastro**	Kyllini, NW Peloponnese	14	13	3	3	1	0	0	0	1	–	16	14	pending			4 ft @ 6 s SSW	8 kt NNW
11	**Ialysos**	Rhodes, northwest	6	5	4	1	2	0	0	0	0	–	4	2	pending			4 ft @ 6 s W	9 kt NNW

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from. Months marked – are still downloading.
Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Caspian Sea: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.






#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Nabran**	Azerbaijan	37	26	19	15	9	5	6	8	17	–	29	36	pending			5 ft @ 6 s NE	12 kt NNW
2	**Ramsar**	Iran	17	18	15	8	8	7	11	5	15	–	13	14	pending			4 ft @ 6 s N	5 kt NNW
3	**Babolsar**	Iran	15	17	12	8	6	8	14	6	13	–	16	14	pending			4 ft @ 7 s NNW	7 kt E
4	**Novkhani**	Absheron, Azerbaijan	17	19	14	9	6	5	8	8	10	–	18	14	pending			5 ft @ 6 s N	12 kt NNW
5	**Derbent**	Dagestan, Russia	22	9	8	9	3	1	2	6	11	–	15	27	pending			4 ft @ 6 s E	12 kt NNW
6	**Awaza**	Turkmenistan	12	14	12	8	8	5	8	5	7	–	13	12	pending			4 ft @ 6 s NW	10 kt NNW
7	**Makhachkala**	Dagestan, Russia	22	8	7	5	1	0	1	1	7	–	15	27	pending			4 ft @ 6 s ESE	12 kt SE
8	**Aktau**	Kazakhstan	15	10	8	3	6	0	1	0	1	–	20	14	pending			5 ft @ 6 s WSW	13 kt ESE
9	**Lankaran**	Azerbaijan	8	8	4	1	1	0	1	1	2	–	4	3	pending			4 ft @ 6 s ENE	9 kt S
10	**Shikhov**	Baku, Azerbaijan	1	0	0	0	0	0	0	0	0	–	0	1	pending			4 ft @ 5 s SE	12 kt N


Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from. Months marked – are still downloading.
Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Black Sea: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with surfable windswell



#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Kilyos**	Istanbul, Turkey	33	36	26	18	18	9	12	34	38	–	24	35	pending			5 ft @ 6 s NNE	11 kt NNE
2	**Şile**	Istanbul, Turkey	32	35	23	15	14	7	10	27	33	–	19	33	pending			5 ft @ 6 s NNE	10 kt NE
3	**Vama Veche**	Romania	22	32	21	9	13	4	3	15	27	–	16	27	pending			5 ft @ 6 s ENE	13 kt NNW
4	**Sozopol**	Bulgaria	23	30	18	9	11	3	2	19	27	–	16	24	pending			4 ft @ 6 s ENE	12 kt N
5	**Batumi**	Georgia	25	27	15	15	5	7	6	3	13	–	19	19	pending			4 ft @ 7 s WNW	8 kt E
6	**Sochi**	Russia	31	28	9	13	4	4	3	1	6	–	24	25	pending			5 ft @ 6 s WSW	8 kt SE
7	**Varna**	Bulgaria	19	25	16	8	10	1	1	14	19	–	9	21	pending			4 ft @ 6 s ENE	13 kt N
8	**Anapa**	Russia	23	17	9	15	3	3	1	0	4	–	16	23	pending			5 ft @ 6 s WSW	13 kt ENE
9	**Odesa**	Ukraine	12	6	4	3	2	1	1	0	1	–	11	10	pending			4 ft @ 6 s SSE	12 kt NNE

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from. Months marked – are still downloading.
Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.