
Year-round: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 1,803 mornings, Nov 2021 to Oct 2026



Best named breaks year-round

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt S	**78%**
2	**Iquique** Chile · 7 kt S	**78%**
3	**Piha** New Zealand · 14 kt SW	**63%**
4	**Punta Hermosa** Peru · 7 kt SSE	**60%**
5	**Skeleton Bay** Namibia · 10 kt S	**60%**
6	**Elands Bay** South Africa · 9 kt SSE	**57%**
7	**Punta de Lobos** Chile · 12 kt S	**56%**
8	**Short Sands** Oregon · 14 kt N	**56%**
9	**Krui** Lampung, Sumatra · 10 kt SE	**52%**
10	**Mentawais** Sumatra, Indonesia · 6 kt SE	**49%**
11	**Cactus Beach** South Australia · 12 kt SE	**47%**
12	**Montara** California · 12 kt NW	**47%**
13	**Chicama** Peru · 10 kt SSE	**46%**
14	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**46%**
15	**Nias** Sumatra, Indonesia · 6 kt SE	**46%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 9 kt N	**78%**
◆2	**Antofagasta coast** Chile · 8 kt S	**78%**
◆3	**La Ligua coast** Chile · 12 kt S	**77%**
◆4	**Camaná coast** Peru · 10 kt SE	**74%**
◆5	**Toltén coast** Chile · 13 kt S	**72%**
◆6	**La Serena coast** Chile · 7 kt SSW	**64%**
◆7	**Chañaral coast** Chile · 6 kt SSW	**62%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea good = 4 ft @ 6 s



Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

Popular breaks, month by month

The 40 well-known breaks with the strongest three-month season (best one per 100 miles of coast), listed by year-round consistency. Breaks in the enclosed seas are on the map insets.

#	Break	Region	J	F	M	A	M	J	J	A	S	O	N	D	Year	Worst	Best 3 months	Typical good morning	Morning wind
1	**Arica**	Chile	70	74	83	85	90	77	82	79	81	79	75	66	**78%**	66%	Mar–May 86%	6 ft @ 13 s SSW, surf ~9 ft	4 kt S
2	**Iquique**	Chile	76	83	81	84	88	74	78	76	76	77	77	69	**78%**	69%	Mar–May 84%	6 ft @ 13 s SSW, surf ~9 ft	7 kt S
3	**Piha**	New Zealand	56	68	74	70	70	53	54	59	55	60	73	65	**63%**	53%	Mar–May 72%	7 ft @ 14 s SW, surf ~10 ft	14 kt SW
4	**Punta Hermosa**	Peru	48	45	70	73	79	61	60	65	59	55	58	42	**60%**	42%	Mar–May 74%	5 ft @ 14 s SSW, surf ~8 ft	7 kt SSE
5	**Skeleton Bay**	Namibia	63	71	48	56	68	61	61	66	57	52	59	53	**60%**	48%	May–Jul 63%	7 ft @ 12 s SSW, surf ~9 ft	10 kt S
6	**Elands Bay**	South Africa	55	57	44	49	71	55	63	66	58	50	55	56	**57%**	44%	May–Jul 63%	6 ft @ 13 s SW, surf ~8 ft	9 kt SSE
7	**Punta de Lobos**	Chile	45	45	61	55	68	59	64	57	56	52	53	52	**56%**	45%	May–Jul 64%	8 ft @ 13 s SW, surf ~11 ft	12 kt S
8	**Short Sands**	Oregon	79	79	63	70	61	40	16	19	53	56	69	66	**56%**	16%	Dec–Feb 75%	7 ft @ 12 s WNW, surf ~9 ft	14 kt N
9	**Krui**	Lampung, Sumatra	29	28	30	51	61	73	79	77	71	63	27	29	**52%**	27%	Jun–Aug 76%	5 ft @ 15 s SSW, surf ~8 ft	10 kt SE
10	**Mentawais**	Sumatra, Indonesia	33	29	31	49	57	73	77	76	59	53	15	29	**49%**	15%	Jun–Aug 75%	5 ft @ 14 s SSW, surf ~8 ft	6 kt SE
11	**Cactus Beach**	South Australia	23	41	42	67	57	46	50	58	54	49	36	37	**47%**	23%	Apr–Jun 57%	7 ft @ 14 s SW, surf ~10 ft	12 kt SE
12	**Montara**	California	68	64	50	45	46	37	19	17	39	53	67	58	**47%**	17%	Nov–Jan 64%	7 ft @ 12 s WNW, surf ~10 ft	12 kt NW
13	**Chicama**	Peru	24	31	53	61	61	45	50	49	53	45	49	31	**46%**	24%	Mar–May 58%	5 ft @ 14 s SSW, surf ~8 ft	10 kt SSE
14	**Cimaja**	Pelabuhan Ratu, West Java	31	25	41	53	59	66	59	57	47	50	28	32	**46%**	25%	May–Jul 61%	5 ft @ 14 s SSW, surf ~8 ft	4 kt E
15	**Nias**	Sumatra, Indonesia	31	26	23	50	54	71	74	72	53	47	9	32	**46%**	9%	Jun–Aug 72%	5 ft @ 14 s SSW, surf ~8 ft	6 kt SE
16	**Imsouane**	Morocco	69	62	67	51	27	16	15	16	39	44	59	72	**45%**	15%	Dec–Feb 68%	6 ft @ 13 s NW, surf ~9 ft	11 kt N
17	**Jeffreys Bay**	South Africa	43	43	38	35	49	44	43	45	40	40	44	46	**43%**	35%	May–Jul 45%	8 ft @ 12 s SW, surf ~11 ft	13 kt W
18	**G-Land**	Java, Indonesia	43	39	47	55	29	18	17	33	47	66	53	54	**42%**	17%	Oct–Dec 58%	6 ft @ 14 s SSW, surf ~9 ft	10 kt ESE
19	**Hossegor**	France	48	74	62	42	21	23	19	26	41	45	46	54	**41%**	19%	Jan–Mar 61%	6 ft @ 13 s NW, surf ~9 ft	9 kt W
20	**Shipstern Bluff**	Tasmania, Australia	48	46	46	46	39	34	33	35	31	27	47	37	**39%**	27%	Jan–Mar 47%	8 ft @ 13 s SW, surf ~12 ft	16 kt SW
21	**Ericeira**	Portugal	39	52	54	45	29	25	14	21	37	54	44	52	**39%**	14%	Oct–Dec 50%	7 ft @ 12 s NW, surf ~10 ft	13 kt N
22	**Desert Point**	Lombok, Indonesia	39	39	44	45	30	21	16	32	41	55	48	50	**38%**	16%	Oct–Dec 51%	5 ft @ 14 s SSW, surf ~8 ft	10 kt ESE
23	**Lakey Peak**	Sumbawa, Indonesia	44	39	39	31	23	17	25	41	45	56	31	43	**36%**	17%	Aug–Oct 47%	5 ft @ 14 s SSW, surf ~8 ft	8 kt E
24	**Tofino**	British Columbia	38	48	49	47	41	29	10	13	44	40	35	35	**36%**	10%	Feb–Apr 48%	7 ft @ 11 s W, surf ~10 ft	11 kt ESE
25	**Bundoran**	Ireland	55	44	41	36	32	28	18	19	37	37	41	38	**35%**	18%	Jan–Mar 46%	7 ft @ 12 s WNW, surf ~10 ft	13 kt S
26	**Safi**	Morocco	48	45	48	33	22	19	13	17	32	41	47	53	**35%**	13%	Nov–Jan 49%	7 ft @ 12 s NW, surf ~10 ft	12 kt NNE
27	**Margaret River**	Western Australia	31	26	26	39	37	33	24	29	36	51	33	35	**33%**	24%	Sep–Nov 40%	9 ft @ 13 s SW, surf ~13 ft	13 kt ESE
28	**Rapa Nui**	Easter Island	28	30	23	35	37	28	34	37	35	41	35	20	**32%**	20%	Aug–Oct 38%	7 ft @ 14 s SW, surf ~10 ft	11 kt E
29	**Santa Catalina**	Panama	9	13	31	47	53	34	41	33	35	37	27	11	**31%**	9%	Apr–Jun 45%	4 ft @ 15 s SSW, surf ~7 ft	7 kt NE
30	**Punta Roca**	El Salvador	7	9	31	47	54	38	39	36	40	33	29	6	**31%**	6%	Apr–Jun 46%	4 ft @ 15 s SSW, surf ~7 ft	7 kt ENE
31	**Todos Santos**	Baja California	63	60	51	35	27	17	5	2	5	18	36	49	**31%**	2%	Jan–Mar 58%	5 ft @ 14 s WNW, surf ~8 ft	8 kt NW
32	**Sayulita**	Nayarit, Mexico	28	18	24	35	43	42	35	36	33	34	21	15	**31%**	15%	Apr–Jun 40%	4 ft @ 15 s SW, surf ~7 ft	7 kt N
33	**Puerto Escondido**	Oaxaca, Mexico	5	18	22	36	42	41	44	40	37	43	29	9	**30%**	5%	May–Jul 42%	4 ft @ 15 s SSW, surf ~7 ft	6 kt WNW
34	**Pascuales**	Colima, Mexico	5	9	23	40	48	43	44	40	43	42	22	5	**30%**	5%	May–Jul 45%	4 ft @ 15 s SSW, surf ~7 ft	6 kt E
35	**Lhoknga**	Aceh, Sumatra	14	10	17	23	37	45	27	46	33	28	5	23	**26%**	5%	Jun–Aug 40%	4 ft @ 15 s SSW, surf ~7 ft	11 kt S
36	**Montañita**	Ecuador	16	17	23	33	46	30	26	26	29	29	24	11	**26%**	11%	Apr–Jun 37%	4 ft @ 15 s SW, surf ~7 ft	8 kt SSW
37	**Rincon**	California	50	42	32	23	21	13	3	5	5	12	27	37	**23%**	3%	Dec–Feb 43%	5 ft @ 14 s W, surf ~8 ft	8 kt NW
38	**La Santa**	Lanzarote, Canaries	38	32	34	22	9	7	4	4	15	25	22	39	**21%**	4%	Dec–Feb 36%	6 ft @ 13 s NW, surf ~9 ft	14 kt NNE
39	**Cloud 9**	Siargao, Philippines	30	34	33	39	10	1	1	3	3	11	24	48	**20%**	1%	Dec–Feb 37%	6 ft @ 10 s NE, surf ~8 ft	10 kt ENE
40	**Hanalei Bay**	Kauai, Hawaii	47	38	21	13	7	1	0	1	5	27	21	33	**18%**	0%	Dec–Feb 39%	7 ft @ 13 s NNW, surf ~11 ft	15 kt E

Month cells use the map colours (% of mornings good+). Typical good morning = medians over the good+ mornings: swell height @ peak period and the direction it comes from, and estimated breaking surf (Komar–Gaughan).
Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward.

Each break is read from its nearest seaward model cell (0.5° in East and Southeast Asia, 1° elsewhere) with swell inside its exposure window. The rule has no wind; local reefs, points and sheltering are not modelled.

Under-the-radar stretches, month by month

Coast where mornings are often good+ with swell aimed at the shore (≥40% year-round, or ≥65% over its best three months),
at least 300 km from the well-known breaks and within 150 km of a town of 5,000+. Listed by year-round consistency; numbers match the ◆ markers on the maps.


#	Stretch	Region	J	F	M	A	M	J	J	A	S	O	N	D	Year	Worst	Best 3 months	Typical good morning	Morning wind
◆1	**Ancud coast**	Chile	85	79	75	77	76	75	65	74	73	81	83	90	**78%**	65%	Nov–Jan 86%	7 ft @ 13 s WSW, surf ~10 ft	9 kt N
◆2	**Antofagasta coast**	Chile	79	79	82	84	84	77	79	73	73	77	75	70	**78%**	70%	Mar–May 83%	6 ft @ 13 s SW, surf ~9 ft	8 kt S
◆3	**La Ligua coast**	Chile	72	80	84	79	80	76	78	70	73	75	76	77	**77%**	70%	Feb–Apr 81%	7 ft @ 13 s SW, surf ~10 ft	12 kt S
◆4	**Camaná coast**	Peru	68	75	80	82	81	70	82	75	72	70	73	59	**74%**	59%	Mar–May 81%	6 ft @ 13 s SSW, surf ~9 ft	10 kt SE
◆5	**Toltén coast**	Chile	80	76	73	72	75	67	63	71	73	70	76	74	**72%**	63%	Dec–Feb 76%	8 ft @ 13 s SW, surf ~11 ft	13 kt S
◆6	**La Serena coast**	Chile	70	68	74	73	66	61	56	50	51	64	61	69	**64%**	50%	Feb–Apr 71%	7 ft @ 13 s SW, surf ~10 ft	7 kt SSW
◆7	**Chañaral coast**	Chile	62	59	62	61	69	65	69	63	65	65	53	48	**62%**	48%	May–Jul 68%	5 ft @ 14 s SW, surf ~8 ft	6 kt SSW
◆8	**Greymouth coast**	New Zealand	37	46	52	58	63	49	57	65	62	54	54	48	**54%**	37%	Jul–Sep 61%	6 ft @ 13 s W, surf ~9 ft	9 kt E
◆9	**Isabela coast**	Galápagos, Ecuador	50	44	56	63	69	51	47	50	42	55	46	40	**51%**	40%	Mar–May 62%	4 ft @ 15 s SW, surf ~8 ft	10 kt SSE
◆10	**Marcona coast**	Peru	48	57	65	71	59	39	45	43	38	43	54	41	**50%**	38%	Mar–May 65%	6 ft @ 13 s SSW, surf ~9 ft	13 kt SE
◆11	**Port Lincoln coast**	Australia	41	43	47	51	51	45	48	54	55	52	34	52	**48%**	34%	Aug–Oct 54%	7 ft @ 14 s SW, surf ~10 ft	12 kt SE
◆12	**Nzeto coast**	Angola	28	38	29	53	64	60	71	76	58	37	28	24	**47%**	24%	Jun–Aug 69%	5 ft @ 13 s SSW, surf ~8 ft	7 kt SSE
◆13	**Meningie coast**	Australia	35	57	52	62	47	39	42	49	46	44	44	48	**47%**	35%	Feb–Apr 57%	7 ft @ 14 s SW, surf ~11 ft	10 kt S
◆14	**Oranjemund coast**	Namibia	52	41	37	34	61	55	50	56	41	40	40	41	**46%**	34%	May–Jul 55%	7 ft @ 13 s SW, surf ~10 ft	9 kt SE
◆15	**Quellón coast**	Chile	63	55	38	42	42	36	33	39	32	51	52	58	**45%**	32%	Dec–Feb 58%	8 ft @ 13 s WSW, surf ~12 ft	13 kt W
◆16	**Pointe-Noire coast**	Congo (Brazzaville)	25	28	22	44	64	57	65	77	56	31	31	21	**44%**	21%	Jun–Aug 66%	5 ft @ 13 s SSW, surf ~7 ft	6 kt S
◆17	**Blitar coast**	Indonesia	45	37	44	56	35	25	17	35	57	63	47	49	**42%**	17%	Sep–Nov 56%	6 ft @ 14 s SSW, surf ~9 ft	12 kt ESE
◆18	**Port Denison coast**	Australia	32	46	37	48	45	49	37	37	45	53	46	36	**42%**	32%	Sep–Nov 48%	9 ft @ 14 s SW, surf ~12 ft	12 kt SSE
◆19	**Sumbe coast**	Angola	22	27	18	37	59	55	67	77	50	37	32	17	**42%**	17%	Jun–Aug 66%	5 ft @ 13 s SW, surf ~8 ft	5 kt S
◆20	**Nelson coast**	New Zealand	45	55	52	41	42	30	30	44	31	32	53	43	**41%**	30%	Jan–Mar 50%	8 ft @ 13 s WSW, surf ~11 ft	8 kt SSW
◆21	**Kaitaia coast**	New Zealand	37	49	52	36	33	28	33	41	42	38	53	51	**41%**	28%	Oct–Dec 47%	7 ft @ 14 s SW, surf ~11 ft	13 kt S
◆22	**Bredasdorp coast**	South Africa	46	44	39	44	48	35	37	32	38	32	47	46	**41%**	32%	Nov–Jan 46%	8 ft @ 12 s SW, surf ~12 ft	10 kt W
◆23	**Huarmey coast**	Peru	45	34	55	53	50	33	35	34	43	31	38	32	**40%**	31%	Mar–May 53%	5 ft @ 14 s SSW, surf ~9 ft	10 kt SSE
◆24	**Gwadar coast**	Pakistan	0	0	0	1	6	67	95	86	34	3	0	0	**25%**	0%	Jun–Aug 82%	8 ft @ 11 s SSW, surf ~10 ft	8 kt WSW
◆25	**Salalah coast**	Oman	0	0	0	0	7	55	95	66	21	0	0	0	**21%**	0%	Jun–Aug 72%	7 ft @ 10 s S, surf ~10 ft	6 kt SW
◆26	**Karachi coast**	Pakistan	0	0	0	0	2	43	86	69	28	2	0	0	**20%**	0%	Jun–Aug 66%	8 ft @ 11 s SW, surf ~11 ft	10 kt W
◆27	**Chabahar coast**	Iran	0	0	0	0	2	46	92	70	15	3	0	0	**19%**	0%	Jun–Aug 69%	7 ft @ 11 s S, surf ~9 ft	8 kt ESE

Month cells use the map colours (% of mornings good+). Typical good morning = medians over the good+ mornings: swell height @ peak period and the direction it comes from, and estimated breaking surf (Komar–Gaughan).
Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward.

Each stretch is read from one WW3 cell 50–120 km offshore. The rule has no wind, which flatters cold, steady coasts (Chile, Peru, Namibia) and storm coasts (southern Chile, New Zealand's West Coast); check local wind and access before booking a trip.

October–March: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 889 mornings, Nov 2021 to Oct 2026, these months only



Best named breaks, October–March

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Iquique** Chile · 7 kt S	**77%**
2	**Arica** Chile · 5 kt S	**74%**
3	**Short Sands** Oregon · 14 kt S	**69%**
4	**Piha** New Zealand · 13 kt SSW	**66%**
5	**Imsouane** Morocco · 8 kt NNE	**63%**
6	**Montara** California · 11 kt NNW	**60%**
7	**Skeleton Bay** Namibia · 8 kt S	**58%**
8	**Hossegor** France · 10 kt SSW	**55%**
9	**Punta Hermosa** Peru · 7 kt SSE	**53%**
10	**Elands Bay** South Africa · 9 kt S	**53%**
11	**Punta de Lobos** Chile · 12 kt SSW	**51%**
12	**G-Land** Java, Indonesia · 9 kt W	**50%**
13	**Ericeira** Portugal · 14 kt NNW	**49%**
14	**Safi** Morocco · 11 kt ENE	**47%**
15	**Todos Santos** Baja California · 6 kt NW	**47%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 8 kt SE	**82%**
◆3	**La Ligua coast** Chile · 11 kt S	**77%**
◆2	**Antofagasta coast** Chile · 7 kt SSW	**77%**
◆5	**Toltén coast** Chile · 13 kt S	**75%**
◆4	**Camaná coast** Peru · 9 kt SE	**71%**
◆6	**La Serena coast** Chile · 5 kt SSW	**68%**
◆7	**Chañaral coast** Chile · 5 kt SSW	**58%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

April–September: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 914 mornings, Apr 2022 to Sep 2026, these months only



Best named breaks, April–September

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt SSE	**82%**
2	**Iquique** Chile · 7 kt S	**79%**
3	**Krui** Lampung, Sumatra · 9 kt SE	**69%**
4	**Punta Hermosa** Peru · 8 kt S	**66%**
5	**Mentawais** Sumatra, Indonesia · 5 kt SE	**65%**
6	**Nias** Sumatra, Indonesia · 5 kt SE	**62%**
7	**Skeleton Bay** Namibia · 10 kt S	**62%**
8	**Elands Bay** South Africa · 8 kt SSE	**61%**
9	**Piha** New Zealand · 15 kt WSW	**60%**
10	**Punta de Lobos** Chile · 12 kt S	**60%**
11	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**57%**
12	**Cactus Beach** South Australia · 12 kt ENE	**55%**
13	**Chicama** Peru · 10 kt SSE	**53%**
14	**Short Sands** Oregon · 13 kt N	**43%**
15	**Pascuales** Colima, Mexico · 6 kt E	**43%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Antofagasta coast** Chile · 10 kt S	**78%**
◆4	**Camaná coast** Peru · 11 kt SE	**77%**
◆3	**La Ligua coast** Chile · 13 kt S	**76%**
◆1	**Ancud coast** Chile · 11 kt N	**73%**
◆5	**Toltén coast** Chile · 12 kt SSE	**70%**
◆7	**Chañaral coast** Chile · 8 kt SSW	**65%**
◆12	**Nzeto coast** Angola · 7 kt SSE	**64%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

November: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (135 mornings; 15 days missing from the archive)



Best named breaks in November

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Iquique** Chile · 7 kt S	**77%**
2	**Arica** Chile · 5 kt S	**75%**
3	**Piha** New Zealand · 11 kt SW	**73%**
4	**Short Sands** Oregon · 11 kt SSE	**69%**
5	**Montara** California · 11 kt NNW	**67%**
6	**Skeleton Bay** Namibia · 7 kt S	**59%**
7	**Imsouane** Morocco · 5 kt NNE	**59%**
8	**Punta Hermosa** Peru · 6 kt SSE	**58%**
9	**Elands Bay** South Africa · 9 kt S	**55%**
10	**G-Land** Java, Indonesia · 6 kt E	**53%**
11	**Punta de Lobos** Chile · 11 kt SSW	**53%**
12	**Chicama** Peru · 9 kt SSE	**49%**
13	**Desert Point** Lombok, Indonesia · 7 kt SE	**48%**
14	**Safi** Morocco · 11 kt NE	**47%**
15	**Shipstern Bluff** Tasmania, Australia · 15 kt WNW	**47%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 10 kt NNW	**83%**
◆3	**La Ligua coast** Chile · 12 kt S	**76%**
◆5	**Toltén coast** Chile · 11 kt S	**76%**
◆2	**Antofagasta coast** Chile · 9 kt SSW	**75%**
◆4	**Camaná coast** Peru · 10 kt SE	**73%**
◆6	**La Serena coast** Chile · 8 kt SSW	**61%**
◆8	**Greymouth coast** New Zealand · 5 kt ESE	**54%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

December: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (155 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in December

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Imsouane** Morocco · 7 kt NNE	**72%**
2	**Iquique** Chile · 7 kt S	**69%**
3	**Short Sands** Oregon · 14 kt SSE	**66%**
4	**Arica** Chile · 5 kt S	**66%**
5	**Piha** New Zealand · 11 kt N	**65%**
6	**Montara** California · 11 kt NNW	**58%**
7	**Elands Bay** South Africa · 9 kt S	**56%**
8	**Uluwatu** Bali, Indonesia · 9 kt W	**55%**
9	**Hossegor** France · 9 kt SE	**54%**
10	**Safi** Morocco · 12 kt ENE	**53%**
11	**Skeleton Bay** Namibia · 9 kt S	**53%**
12	**Ericeira** Portugal · 14 kt ENE	**52%**
13	**Punta de Lobos** Chile · 12 kt SSW	**52%**
14	**Todos Santos** Baja California · 5 kt NNW	**49%**
15	**Cloud 9** Siargao, Philippines · 10 kt ENE	**48%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 8 kt SSW	**90%**
◆3	**La Ligua coast** Chile · 12 kt S	**77%**
◆5	**Toltén coast** Chile · 14 kt S	**74%**
◆2	**Antofagasta coast** Chile · 7 kt S	**70%**
◆6	**La Serena coast** Chile · 6 kt SSW	**69%**
◆4	**Camaná coast** Peru · 9 kt SE	**59%**
◆15	**Quellón coast** Chile · 13 kt SW	**58%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

January: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Best named breaks in January

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Short Sands** Oregon · 14 kt S	**79%**
2	**Iquique** Chile · 6 kt S	**76%**
3	**Arica** Chile · 5 kt S	**70%**
4	**Imsouane** Morocco · 9 kt NNE	**69%**
5	**Montara** California · 13 kt SSE	**68%**
6	**Todos Santos** Baja California · 5 kt SE	**63%**
7	**Skeleton Bay** Namibia · 7 kt S	**63%**
8	**Piha** New Zealand · 13 kt ENE	**56%**
9	**Elands Bay** South Africa · 8 kt SSE	**55%**
10	**Bundoran** Ireland · 13 kt SSE	**55%**
11	**Rincon** California · 7 kt NW	**50%**
12	**Safi** Morocco · 11 kt ENE	**48%**
13	**Shipstern Bluff** Tasmania, Australia · 11 kt SW	**48%**
14	**Honolua Bay** Maui, Hawaii · 14 kt E	**48%**
15	**Punta Hermosa** Peru · 7 kt SSE	**48%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 8 kt SE	**85%**
◆5	**Toltén coast** Chile · 14 kt S	**80%**
◆2	**Antofagasta coast** Chile · 5 kt SSW	**79%**
◆3	**La Ligua coast** Chile · 10 kt S	**72%**
◆6	**La Serena coast** Chile · 4 kt SSW	**70%**
◆4	**Camaná coast** Peru · 9 kt ESE	**68%**
◆15	**Quellón coast** Chile · 11 kt SSW	**63%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

February: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (141 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in February

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Iquique** Chile · 7 kt S	**83%**
2	**Short Sands** Oregon · 13 kt SSE	**79%**
3	**Hossegor** France · 8 kt SSW	**74%**
4	**Arica** Chile · 5 kt S	**74%**
5	**Skeleton Bay** Namibia · 9 kt S	**71%**
6	**Piha** New Zealand · 11 kt SSW	**68%**
7	**Montara** California · 11 kt NNW	**64%**
8	**Imsouane** Morocco · 6 kt NNE	**62%**
9	**Todos Santos** Baja California · 6 kt NW	**60%**
10	**Elands Bay** South Africa · 9 kt S	**57%**
11	**Nazaré** Portugal · 12 kt WNW	**52%**
12	**Tofino** British Columbia · 10 kt ESE	**48%**
13	**Shipstern Bluff** Tasmania, Australia · 16 kt SW	**46%**
14	**Punta Hermosa** Peru · 7 kt SSE	**45%**
15	**Punta de Lobos** Chile · 14 kt SSW	**45%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆3	**La Ligua coast** Chile · 12 kt S	**80%**
◆2	**Antofagasta coast** Chile · 6 kt SSW	**79%**
◆1	**Ancud coast** Chile · 7 kt SE	**79%**
◆5	**Toltén coast** Chile · 14 kt S	**76%**
◆4	**Camaná coast** Peru · 8 kt ESE	**75%**
◆6	**La Serena coast** Chile · 5 kt SSW	**68%**
◆7	**Chañaral coast** Chile · 5 kt SSW	**59%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

March: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in March

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 5 kt S	**83%**
2	**Iquique** Chile · 7 kt S	**81%**
3	**Piha** New Zealand · 15 kt SSW	**74%**
4	**Punta Hermosa** Peru · 7 kt SSE	**70%**
5	**Imsouane** Morocco · 11 kt N	**67%**
6	**Short Sands** Oregon · 16 kt S	**63%**
7	**Hossegor** France · 13 kt W	**62%**
8	**Punta de Lobos** Chile · 14 kt SSW	**61%**
9	**Ericeira** Portugal · 12 kt SSW	**54%**
10	**Chicama** Peru · 9 kt SE	**53%**
11	**Todos Santos** Baja California · 10 kt NNW	**51%**
12	**Montara** California · 13 kt NNW	**50%**
13	**Tofino** British Columbia · 12 kt ESE	**49%**
14	**Uluwatu** Bali, Indonesia · 10 kt ENE	**49%**
15	**Safi** Morocco · 13 kt NE	**48%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆3	**La Ligua coast** Chile · 11 kt S	**84%**
◆2	**Antofagasta coast** Chile · 7 kt S	**82%**
◆4	**Camaná coast** Peru · 9 kt ESE	**80%**
◆1	**Ancud coast** Chile · 9 kt ESE	**75%**
◆6	**La Serena coast** Chile · 6 kt S	**74%**
◆5	**Toltén coast** Chile · 15 kt S	**73%**
◆10	**Marcona coast** Peru · 12 kt SE	**65%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

April: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in April

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt SSE	**85%**
2	**Iquique** Chile · 8 kt S	**84%**
3	**Punta Hermosa** Peru · 8 kt SSE	**73%**
4	**Short Sands** Oregon · 15 kt N	**70%**
5	**Piha** New Zealand · 16 kt ENE	**70%**
6	**Cactus Beach** South Australia · 11 kt ENE	**67%**
7	**Chicama** Peru · 10 kt SE	**61%**
8	**Skeleton Bay** Namibia · 10 kt S	**56%**
9	**Punta de Lobos** Chile · 12 kt S	**55%**
10	**G-Land** Java, Indonesia · 9 kt ESE	**55%**
11	**Cimaja** Pelabuhan Ratu, West Java · 4 kt ENE	**53%**
12	**Imsouane** Morocco · 10 kt N	**51%**
13	**Krui** Lampung, Sumatra · 6 kt ESE	**51%**
14	**Nias** Sumatra, Indonesia · 5 kt SSE	**50%**
15	**Elands Bay** South Africa · 8 kt S	**49%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Antofagasta coast** Chile · 10 kt S	**84%**
◆4	**Camaná coast** Peru · 11 kt ESE	**82%**
◆3	**La Ligua coast** Chile · 12 kt S	**79%**
◆1	**Ancud coast** Chile · 9 kt N	**77%**
◆6	**La Serena coast** Chile · 7 kt S	**73%**
◆5	**Toltén coast** Chile · 13 kt SSE	**72%**
◆10	**Marcona coast** Peru · 14 kt SE	**71%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

May: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in May

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 3 kt SSE	**90%**
2	**Iquique** Chile · 7 kt S	**88%**
3	**Punta Hermosa** Peru · 7 kt S	**79%**
4	**Elands Bay** South Africa · 8 kt SSE	**71%**
5	**Piha** New Zealand · 15 kt S	**70%**
6	**Punta de Lobos** Chile · 11 kt S	**68%**
7	**Skeleton Bay** Namibia · 10 kt S	**68%**
8	**Short Sands** Oregon · 14 kt NNW	**61%**
9	**Chicama** Peru · 10 kt SE	**61%**
10	**Krui** Lampung, Sumatra · 9 kt ESE	**61%**
11	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**59%**
12	**Mentawais** Sumatra, Indonesia · 5 kt WSW	**57%**
13	**Cactus Beach** South Australia · 11 kt E	**57%**
14	**Nias** Sumatra, Indonesia · 5 kt W	**54%**
15	**Punta Roca** El Salvador · 5 kt ESE	**54%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Antofagasta coast** Chile · 10 kt S	**84%**
◆4	**Camaná coast** Peru · 11 kt SE	**81%**
◆3	**La Ligua coast** Chile · 12 kt S	**80%**
◆1	**Ancud coast** Chile · 8 kt N	**76%**
◆5	**Toltén coast** Chile · 14 kt SSE	**75%**
◆7	**Chañaral coast** Chile · 8 kt SSW	**69%**
◆9	**Isabela coast** Galápagos, Ecuador · 9 kt SSE	**69%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.27× the mean period.

June: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Best named breaks in June

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 3 kt SSE	**77%**
2	**Iquique** Chile · 7 kt S	**74%**
3	**Mentawais** Sumatra, Indonesia · 4 kt SE	**73%**
4	**Krui** Lampung, Sumatra · 9 kt ESE	**73%**
5	**Nias** Sumatra, Indonesia · 4 kt SE	**71%**
6	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**66%**
7	**Punta Hermosa** Peru · 8 kt S	**61%**
8	**Skeleton Bay** Namibia · 10 kt SSE	**61%**
9	**Punta de Lobos** Chile · 14 kt S	**59%**
10	**Elands Bay** South Africa · 8 kt SE	**55%**
11	**Piha** New Zealand · 15 kt NE	**53%**
12	**Cactus Beach** South Australia · 14 kt N	**46%**
13	**Lhoknga** Aceh, Sumatra · 12 kt S	**45%**
14	**Chicama** Peru · 11 kt SSE	**45%**
15	**Jeffreys Bay** South Africa · 12 kt W	**44%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Antofagasta coast** Chile · 10 kt S	**77%**
◆3	**La Ligua coast** Chile · 13 kt S	**76%**
◆1	**Ancud coast** Chile · 12 kt N	**75%**
◆4	**Camaná coast** Peru · 11 kt SE	**70%**
◆5	**Toltén coast** Chile · 13 kt N	**67%**
◆24	**Gwadar coast** Pakistan · 8 kt WSW	**67%**
◆7	**Chañaral coast** Chile · 10 kt SSW	**65%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.27× the mean period.

July: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (154 mornings; 1 days missing from the archive)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in July

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt SSE	**82%**
2	**Krui** Lampung, Sumatra · 11 kt SE	**79%**
3	**Iquique** Chile · 7 kt S	**78%**
4	**Mentawais** Sumatra, Indonesia · 4 kt SE	**77%**
5	**Nias** Sumatra, Indonesia · 5 kt SE	**74%**
6	**Punta de Lobos** Chile · 12 kt S	**64%**
7	**Elands Bay** South Africa · 10 kt SSE	**63%**
8	**Skeleton Bay** Namibia · 11 kt S	**61%**
9	**Punta Hermosa** Peru · 8 kt SSE	**60%**
10	**Cimaja** Pelabuhan Ratu, West Java · 5 kt E	**59%**
11	**Piha** New Zealand · 15 kt SW	**54%**
12	**Chicama** Peru · 11 kt SSE	**50%**
13	**Cactus Beach** South Australia · 13 kt WSW	**50%**
14	**Puerto Escondido** Oaxaca, Mexico · 6 kt NW	**44%**
15	**Pascuales** Colima, Mexico · 6 kt NE	**44%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆24	**Gwadar coast** Pakistan · 7 kt WSW	**95%**
◆25	**Salalah coast** Oman · 7 kt SW	**95%**
◆27	**Chabahar coast** Iran · 8 kt ESE	**92%**
◆26	**Karachi coast** Pakistan · 11 kt WSW	**86%**
◆4	**Camaná coast** Peru · 11 kt SE	**82%**
◆2	**Antofagasta coast** Chile · 9 kt S	**79%**
◆3	**La Ligua coast** Chile · 13 kt S	**78%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

August: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in August

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt S	**79%**
2	**Krui** Lampung, Sumatra · 14 kt SE	**77%**
3	**Mentawais** Sumatra, Indonesia · 9 kt SE	**76%**
4	**Iquique** Chile · 8 kt S	**76%**
5	**Nias** Sumatra, Indonesia · 7 kt SE	**72%**
6	**Elands Bay** South Africa · 9 kt N	**66%**
7	**Skeleton Bay** Namibia · 10 kt S	**66%**
8	**Punta Hermosa** Peru · 8 kt SSE	**65%**
9	**Piha** New Zealand · 14 kt ESE	**59%**
10	**Cactus Beach** South Australia · 12 kt E	**58%**
11	**Punta de Lobos** Chile · 11 kt S	**57%**
12	**Cimaja** Pelabuhan Ratu, West Java · 5 kt E	**57%**
13	**Chicama** Peru · 11 kt SSE	**49%**
14	**Lhoknga** Aceh, Sumatra · 13 kt S	**46%**
15	**Jeffreys Bay** South Africa · 13 kt W	**45%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆24	**Gwadar coast** Pakistan · 8 kt WSW	**86%**
◆16	**Pointe-Noire coast** Congo (Brazzaville) · 6 kt S	**77%**
◆19	**Sumbe coast** Angola · 4 kt S	**77%**
◆12	**Nzeto coast** Angola · 7 kt SSE	**76%**
◆4	**Camaná coast** Peru · 11 kt SE	**75%**
◆1	**Ancud coast** Chile · 12 kt N	**74%**
◆2	**Antofagasta coast** Chile · 11 kt S	**73%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

September: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)



Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

Best named breaks in September

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt S	**81%**
2	**Iquique** Chile · 7 kt S	**76%**
3	**Krui** Lampung, Sumatra · 12 kt SE	**71%**
4	**Punta Hermosa** Peru · 8 kt S	**59%**
5	**Mentawais** Sumatra, Indonesia · 9 kt SE	**59%**
6	**Elands Bay** South Africa · 8 kt SE	**58%**
7	**Skeleton Bay** Namibia · 10 kt S	**57%**
8	**Punta de Lobos** Chile · 13 kt S	**56%**
9	**Piha** New Zealand · 15 kt WSW	**55%**
10	**Cactus Beach** South Australia · 11 kt ENE	**54%**
11	**Chicama** Peru · 9 kt SSE	**53%**
12	**Nias** Sumatra, Indonesia · 8 kt SSE	**53%**
13	**Short Sands** Oregon · 13 kt N	**53%**
14	**Cimaja** Pelabuhan Ratu, West Java · 4 kt E	**47%**
15	**G-Land** Java, Indonesia · 11 kt ESE	**47%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆2	**Antofagasta coast** Chile · 10 kt S	**73%**
◆3	**La Ligua coast** Chile · 14 kt S	**73%**
◆1	**Ancud coast** Chile · 11 kt WNW	**73%**
◆5	**Toltén coast** Chile · 11 kt N	**73%**
◆4	**Camaná coast** Peru · 11 kt SE	**72%**
◆7	**Chañaral coast** Chile · 7 kt SSW	**65%**
◆8	**Greymouth coast** New Zealand · 14 kt E	**62%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

October: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (148 mornings; 7 days missing from the archive)



Best named breaks in October

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


1	**Arica** Chile · 4 kt S	**79%**
2	**Iquique** Chile · 7 kt S	**77%**
3	**G-Land** Java, Indonesia · 7 kt ESE	**66%**
4	**Krui** Lampung, Sumatra · 9 kt SE	**63%**
5	**Piha** New Zealand · 14 kt NW	**60%**
6	**Short Sands** Oregon · 16 kt S	**56%**
7	**Lakey Peak** Sumbawa, Indonesia · 6 kt E	**56%**
8	**Punta Hermosa** Peru · 7 kt SSE	**55%**
9	**Desert Point** Lombok, Indonesia · 8 kt ESE	**55%**
10	**Ericeira** Portugal · 13 kt NNE	**54%**
11	**Mentawais** Sumatra, Indonesia · 4 kt SE	**53%**
12	**Montara** California · 11 kt NNW	**53%**
13	**Skeleton Bay** Namibia · 9 kt S	**52%**
14	**Punta de Lobos** Chile · 14 kt SSW	**52%**
15	**Margaret River** Western Australia · 9 kt SSW	**51%**

Under-the-radar stretches


◆ numbered on the map; month by month on page 3

◆1	**Ancud coast** Chile · 7 kt SSW	**81%**
◆2	**Antofagasta coast** Chile · 7 kt S	**77%**
◆3	**La Ligua coast** Chile · 14 kt S	**75%**
◆4	**Camaná coast** Peru · 11 kt SE	**70%**
◆5	**Toltén coast** Chile · 12 kt S	**70%**
◆7	**Chañaral coast** Chile · 7 kt SSW	**65%**
◆6	**La Serena coast** Chile · 7 kt SSW	**64%**


Great Lakes · good = 4 ft @ 6 s



Mediterranean & Black Sea · good = 4 ft @ 6 s



Caspian Sea
good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.


Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

Lake Michigan: when and where it's good

Good lake surf = total wave height 4–10 ft (1.22–3 m) with dominant period ≥ 6 s (taken as 1.29× the mean period), arriving within 75° of the shore's facing, on the ~08:00 sample. Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with good lake surf




#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical good morning	Morning wind
1	**Holland**	Michigan	30	22	17	11	3	2	1	1	5	11	28	26	**13%**	Nov–Jan 28%	0%	5 ft @ 7 s W	13 kt S
2	**Grand Haven**	Michigan	28	22	15	10	2	1	0	1	4	11	27	26	**12%**	Nov–Jan 27%	0%	5 ft @ 7 s W	13 kt S
3	**Ludington**	Michigan	23	21	12	9	3	1	1	1	3	11	28	27	**11%**	Nov–Jan 26%	0%	5 ft @ 7 s W	14 kt S
4	**Frankfort**	Michigan	23	20	14	9	3	1	1	1	4	10	26	25	**11%**	Nov–Jan 25%	0%	5 ft @ 7 s W	14 kt S
5	**St. Joseph**	Michigan	24	16	14	6	2	2	2	1	5	11	24	14	**10%**	Nov–Jan 21%	0%	5 ft @ 7 s NW	13 kt SSW
6	**New Buffalo**	Michigan	19	14	12	5	2	2	1	1	5	11	16	10	**8%**	Nov–Jan 15%	0%	5 ft @ 7 s NNW	13 kt SSW
7	**Portage**	Indiana	10	7	7	3	1	1	1	1	4	8	6	5	**5%**	Jan–Mar 8%	0%	5 ft @ 7 s N	13 kt SSW
8	**Milwaukee**	Wisconsin	6	7	8	6	1	1	1	2	5	5	5	5	**4%**	Jan–Mar 7%	0%	5 ft @ 7 s ENE	13 kt NW
9	**Sheboygan**	Wisconsin	6	7	9	7	1	1	0	1	4	6	4	5	**4%**	Feb–Apr 8%	0%	5 ft @ 7 s E	14 kt S
10	**Gary**	Indiana	8	5	6	3	1	1	1	1	3	6	5	4	**4%**	Jan–Mar 6%	0%	5 ft @ 7 s N	12 kt NW
11	**Racine**	Wisconsin	6	3	3	3	1	1	1	1	3	5	2	2	**2%**	Jan–Mar 4%	0%	5 ft @ 7 s NE	13 kt WNW
12	**Whiting**	Indiana	5	3	6	1	0	1	0	1	3	5	1	2	**2%**	Jan–Mar 4%	0%	5 ft @ 7 s N	12 kt NW
13	**Chicago**	Illinois	3	2	4	2	0	1	0	1	2	3	0	0	**1%**	Jan–Mar 3%	0%	5 ft @ 8 s NNE	13 kt SSW

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical good morning = medians over good mornings: total wave height @ dominant period and the direction waves come from.


Coasts are ranked by year-round good %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Lake Superior: when and where it's good

Good lake surf = total wave height 4–10 ft (1.22–3 m) with dominant period ≥ 6 s (taken as 1.29× the mean period), arriving within 75° of the shore's facing, on the ~08:00 sample. Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with good lake surf



#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical good morning	Morning wind
1	**Grand Marais MI**	Michigan	28	18	13	8	1	3	0	3	6	14	27	23	**12%**	Nov–Jan 26%	0%	5 ft @ 7 s NW	14 kt W
2	**Whitefish Point**	Michigan	25	12	9	7	1	2	0	3	3	13	24	27	**10%**	Nov–Jan 25%	0%	5 ft @ 7 s NW	14 kt SSE
3	**Marquette**	Michigan	21	17	17	11	1	1	0	2	5	11	21	19	**10%**	Nov–Jan 20%	0%	6 ft @ 7 s N	15 kt S
4	**Au Train**	Michigan	19	16	14	6	1	1	0	2	5	9	24	19	**9%**	Nov–Jan 21%	0%	5 ft @ 7 s NNW	15 kt S
5	**Ontonagon**	Michigan	19	9	12	11	1	0	0	1	3	10	23	18	**9%**	Nov–Jan 20%	0%	5 ft @ 7 s NW	13 kt WSW
6	**Agawa Bay**	Ontario	20	7	4	3	1	1	0	1	2	8	21	26	**8%**	Nov–Jan 22%	0%	5 ft @ 8 s W	11 kt SSE
7	**Terrace Bay**	Ontario	12	9	6	4	2	1	1	1	4	13	20	15	**7%**	Oct–Dec 16%	0%	5 ft @ 7 s SSW	13 kt NNE
8	**Little Girl's Point**	Michigan	6	3	8	7	1	0	0	0	0	5	7	7	**4%**	Nov–Jan 7%	0%	5 ft @ 7 s N	13 kt W
9	**Stoney Point**	Minnesota	2	3	4	5	1	0	0	0	0	3	1	2	**2%**	Feb–Apr 4%	0%	6 ft @ 7 s ENE	8 kt SSW
10	**Park Point**	Duluth, Minnesota	2	2	5	5	1	0	0	0	0	3	1	2	**2%**	Feb–Apr 4%	0%	6 ft @ 7 s ENE	8 kt SSW
11	**Grand Marais MN**	Minnesota	3	1	0	1	1	0	0	0	0	3	2	1	**1%**	Nov–Jan 2%	0%	5 ft @ 7 s SSW	12 kt ENE

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical good morning = medians over good mornings: total wave height @ dominant period and the direction waves come from.


Coasts are ranked by year-round good %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Mediterranean: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with surfable windswell



Mediterranean: coast by coast

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample. Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.

#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Ajaccio**	Corsica, France	54	45	35	31	23	18	23	19	24	24	53	37	**32%**	Nov–Jan 48%	6%	5 ft @ 7 s W	11 kt WNW
2	**Susa**	Cyrenaica, Libya	50	45	37	32	19	13	29	15	24	38	30	38	**31%**	Dec–Feb 44%	3%	4 ft @ 7 s NNW	8 kt W
3	**Capo Mannu**	Sardinia, Italy	47	40	35	29	26	13	22	22	20	26	53	37	**31%**	Nov–Jan 46%	4%	5 ft @ 7 s WNW	11 kt ESE
4	**Tabarka**	Tunisia	51	49	30	33	23	9	19	14	17	22	49	37	**29%**	Dec–Feb 46%	6%	5 ft @ 7 s NW	12 kt W
5	**Alexandria**	Egypt	35	45	39	30	26	19	33	21	21	24	21	33	**29%**	Jan–Mar 40%	2%	4 ft @ 7 s NW	9 kt NW
6	**Paphos**	Cyprus	37	43	36	32	28	15	19	25	27	18	21	34	**28%**	Jan–Mar 39%	1%	4 ft @ 6 s W	9 kt W
7	**Falasarna**	Crete, Greece	45	37	32	31	14	9	19	11	14	16	37	43	**25%**	Nov–Jan 41%	2%	5 ft @ 6 s W	14 kt NE
8	**Benghazi**	Libya	43	32	28	30	19	11	16	8	12	20	33	48	**25%**	Nov–Jan 41%	2%	5 ft @ 6 s NW	10 kt NW
9	**Algiers**	Algeria	34	40	41	29	16	7	11	7	12	16	36	32	**23%**	Jan–Mar 38%	3%	4 ft @ 7 s N	10 kt WSW
10	**Gaza**	Palestine	35	39	33	21	23	7	13	10	10	18	19	35	**22%**	Dec–Feb 36%	2%	4 ft @ 7 s WNW	7 kt WSW
11	**Bizerte**	Tunisia	39	35	23	27	15	5	17	11	11	14	30	31	**21%**	Dec–Feb 35%	1%	5 ft @ 7 s NNW	13 kt WNW
12	**Tripoli**	Libya	43	32	26	30	14	8	10	5	13	16	21	37	**21%**	Dec–Feb 37%	1%	4 ft @ 6 s N	10 kt ESE
13	**Beirut**	Lebanon	37	36	33	21	25	5	7	13	11	15	15	32	**21%**	Jan–Mar 35%	2%	4 ft @ 7 s W	10 kt SW
14	**Livorno**	Tuscany, Italy	33	26	21	19	14	9	13	12	23	18	37	27	**21%**	Nov–Jan 32%	2%	5 ft @ 7 s WSW	10 kt ENE
15	**Kalo Nero**	Messinia, Greece	40	35	21	20	8	3	5	3	9	14	39	39	**19%**	Nov–Jan 39%	1%	4 ft @ 7 s WSW	9 kt NNW
16	**Santa Marinella**	Lazio, Italy	26	23	25	20	12	7	8	10	15	19	27	27	**18%**	Nov–Jan 27%	0%	4 ft @ 6 s SW	11 kt NNE
17	**Marseille**	France	25	26	17	16	9	5	5	5	10	13	26	15	**14%**	Jan–Mar 23%	0%	5 ft @ 6 s WSW	16 kt NNW
18	**Barcelona**	Spain	23	19	37	17	2	3	0	5	8	16	17	12	**13%**	Jan–Mar 26%	1%	4 ft @ 6 s ESE	10 kt W
19	**Varazze**	Liguria, Italy	22	20	21	12	7	5	3	5	7	17	21	14	**13%**	Jan–Mar 21%	1%	4 ft @ 7 s SSW	10 kt N
20	**Kelibia**	Cap Bon, Tunisia	11	9	29	18	14	2	1	1	6	11	9	14	**10%**	Mar–May 20%	0%	4 ft @ 6 s ESE	13 kt NW
21	**Mondello**	Sicily, Italy	14	11	9	11	2	1	3	1	9	7	10	15	**8%**	Dec–Feb 13%	0%	4 ft @ 6 s NNW	9 kt WSW
22	**Hammamet**	Tunisia	6	7	25	15	8	2	1	1	1	7	1	7	**7%**	Mar–May 16%	0%	4 ft @ 6 s ESE	11 kt NW
23	**Kastro**	Kyllini, Greece	14	11	3	5	3	1	0	0	1	1	16	14	**6%**	Nov–Jan 15%	0%	4 ft @ 6 s SW	8 kt NNW
24	**Antalya**	Turkey	14	7	3	2	1	0	0	0	0	0	4	12	**4%**	Dec–Feb 11%	0%	4 ft @ 6 s SSE	4 kt NNW


Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.

Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Greece: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.






#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Zakynthos west**	Ionian Islands	48	38	28	33	17	8	17	12	14	27	46	45	**27%**	Nov–Jan 46%	1%	5 ft @ 6 s W	11 kt NNW
2	**Falasarna**	West Crete	45	37	32	32	14	9	18	14	15	17	37	43	**26%**	Nov–Jan 41%	2%	5 ft @ 6 s W	14 kt NE
3	**Komos**	South Crete	36	33	28	27	15	5	8	7	7	12	28	40	**21%**	Dec–Feb 36%	2%	4 ft @ 6 s W	9 kt N
4	**Kalo Nero**	Messinia, W Peloponnese	40	35	25	22	10	5	7	3	9	14	39	39	**20%**	Nov–Jan 39%	2%	4 ft @ 7 s WSW	9 kt NNW
5	**Kokkini Hani**	North Crete	31	30	28	15	14	7	13	10	19	25	23	23	**20%**	Jan–Mar 30%	0%	4 ft @ 6 s N	11 kt W
6	**Agios Gordios**	Corfu, west coast	34	28	21	23	12	3	8	4	12	19	33	35	**19%**	Nov–Jan 34%	0%	4 ft @ 6 s WSW	12 kt NNW
7	**Chania**	North Crete	22	27	26	16	17	11	14	15	20	25	23	16	**19%**	Jan–Mar 25%	0%	5 ft @ 6 s N	12 kt N
8	**Kymi**	East Evia	22	24	23	11	8	1	3	9	15	18	16	17	**14%**	Jan–Mar 23%	0%	5 ft @ 6 s NE	13 kt NNW
9	**Tinos north**	Cyclades	19	23	20	10	9	5	6	8	17	22	16	11	**14%**	Jan–Mar 21%	0%	5 ft @ 5 s N	15 kt NNW
10	**Kastro**	Kyllini, NW Peloponnese	14	13	3	3	1	0	0	0	1	2	16	14	**5%**	Nov–Jan 15%	0%	4 ft @ 6 s SSW	8 kt NNW
11	**Ialysos**	Rhodes, northwest	6	5	4	1	2	0	0	0	0	0	4	2	**2%**	Jan–Mar 5%	0%	4 ft @ 6 s W	9 kt NNW


Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.
Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Caspian Sea: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with surfable windswell




#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Nabran**	Azerbaijan	37	26	19	15	9	5	6	8	17	26	29	36	**19%**	Nov–Jan 34%	0%	5 ft @ 6 s NE	12 kt NNW
2	**Ramsar**	Iran	17	18	15	8	8	7	11	5	15	16	13	14	**12%**	Jan–Mar 17%	1%	4 ft @ 6 s N	5 kt NNW
3	**Babolsar**	Iran	15	17	12	8	6	8	14	6	13	11	16	14	**12%**	Dec–Feb 16%	0%	4 ft @ 7 s NNW	7 kt E
4	**Novkhani**	Absheron, Azerbaijan	17	19	14	9	6	5	8	8	10	14	18	14	**12%**	Dec–Feb 17%	0%	5 ft @ 6 s N	12 kt NNW
5	**Derbent**	Dagestan, Russia	22	9	8	9	3	1	2	6	11	17	15	27	**11%**	Nov–Jan 21%	0%	4 ft @ 6 s E	12 kt NNW
6	**Awaza**	Turkmenistan	12	14	12	8	8	5	8	5	7	8	13	12	**9%**	Jan–Mar 13%	0%	4 ft @ 6 s NW	10 kt NNW
7	**Makhachkala**	Dagestan, Russia	22	8	7	5	1	0	1	1	7	15	15	27	**9%**	Nov–Jan 21%	0%	4 ft @ 6 s ESE	12 kt SE
8	**Aktau**	Kazakhstan	15	10	8	3	6	0	1	0	1	7	20	14	**7%**	Nov–Jan 16%	0%	5 ft @ 6 s W	13 kt ESE
9	**Lankaran**	Azerbaijan	8	8	4	1	1	0	1	1	2	4	4	3	**3%**	Jan–Mar 6%	0%	4 ft @ 6 s ENE	9 kt S
10	**Shikhov**	Baku, Azerbaijan	1	0	0	0	0	0	0	0	0	0	0	1	**0%**	Nov–Jan 1%	0%	4 ft @ 5 s SE	12 kt N


Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.
Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Black Sea: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.



Average of these coasts: % of mornings with surfable windswell




#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical surfable morning	Morning wind
1	**Kilyos**	Istanbul, Turkey	33	36	26	18	18	9	12	34	38	29	24	35	**26%**	Dec–Feb 35%	0%	5 ft @ 6 s NNE	11 kt NNE
2	**Şile**	Istanbul, Turkey	32	35	23	15	14	7	10	27	33	28	19	33	**23%**	Dec–Feb 33%	0%	5 ft @ 6 s NNE	10 kt NE
3	**Vama Veche**	Romania	22	32	21	9	13	4	3	15	27	18	16	27	**17%**	Dec–Feb 27%	0%	5 ft @ 6 s ENE	13 kt NNW
4	**Sozopol**	Bulgaria	23	30	18	9	11	3	2	19	27	18	16	24	**17%**	Dec–Feb 26%	0%	4 ft @ 6 s ENE	12 kt N
5	**Batumi**	Georgia	25	27	15	15	5	7	6	3	13	15	19	19	**14%**	Dec–Feb 23%	1%	4 ft @ 7 s WNW	8 kt E
6	**Sochi**	Russia	31	28	9	13	4	4	3	1	6	9	24	25	**13%**	Dec–Feb 28%	1%	5 ft @ 6 s WSW	8 kt SE
7	**Varna**	Bulgaria	19	25	16	8	10	1	1	14	19	11	9	21	**13%**	Dec–Feb 21%	0%	4 ft @ 6 s ENE	13 kt N
8	**Anapa**	Russia	23	17	9	15	3	3	1	0	4	11	16	23	**10%**	Dec–Feb 21%	0%	5 ft @ 6 s WSW	13 kt ENE
9	**Odesa**	Ukraine	12	6	4	3	2	1	1	0	1	1	11	10	**4%**	Nov–Jan 11%	0%	4 ft @ 6 s SSE	12 kt NNE


Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.

Coasts are ranked by year-round surfable %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Madagascar: when and where it's good+

Good+ = swell ≥ 4 ft over 9 s, or ≥ 3 ft over 14 s; breaking under 16 ft; wind sea ≤ 0.35× swell; on the ~08:00 sample.
Open-ocean coasts: the world maps' good+ standard, read from Météo-France MFWAM at each coast's ~8 km cell, with swell aimed at the shore.







#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical good+ morning	Morning wind
1	**Lavanono**	Deep south	35	26	34	51	65	64	68	58	54	45	44	41	**49%**	May–Jul 66%	49%	6 ft @ 13 s SSW	9 kt E
2	**Anakao**	Toliara, southwest	19	24	23	43	50	49	54	52	41	26	34	31	**37%**	Jun–Aug 52%	37%	5 ft @ 13 s SSW	9 kt SE
3	**Manakara**	Southeast	34	40	37	49	37	46	51	44	36	30	19	19	**37%**	Jun–Aug 47%	37%	5 ft @ 10 s SE	9 kt NNE
4	**Ifaty**	Toliara, southwest	14	22	22	39	43	45	47	47	36	26	25	23	**33%**	Jun–Aug 46%	33%	5 ft @ 14 s SSW	10 kt SE
5	**Andavadoaka**	Morombe, west	14	13	21	35	37	45	42	37	29	20	16	21	**28%**	Jun–Aug 41%	28%	5 ft @ 13 s SSW	10 kt SSE
6	**Fort Dauphin**	Tolagnaro, southeast	17	6	15	24	23	29	25	17	10	13	7	12	**17%**	May–Jul 26%	17%	6 ft @ 11 s SSE	15 kt NNE
7	**Mahambo**	East	8	8	3	11	15	17	22	33	17	4	1	2	**12%**	Jun–Aug 24%	12%	5 ft @ 10 s ESE	10 kt SSE
8	**Morondava**	West	1	4	2	5	6	12	8	12	9	6	1	2	**6%**	Jun–Aug 11%	6%	5 ft @ 11 s SW	7 kt ENE
9	**Maintirano**	West	1	2	2	3	1	4	5	6	5	5	2	1	**3%**	Jul–Sep 5%	3%	5 ft @ 10 s SW	9 kt S
10	**Sainte-Marie**	Nosy Boraha, east	1	3	0	5	6	4	8	3	3	2	0	0	**3%**	May–Jul 6%	3%	5 ft @ 10 s ESE	10 kt SSE
11	**Ramena**	Antsiranana, north	1	0	1	0	0	2	0	1	0	0	0	1	**0%**	Jun–Aug 1%	0%	4 ft @ 10 s ESE	19 kt SE
12	**Mahajanga**	Northwest	1	0	0	0	0	0	0	0	0	0	0	0	**0%**	Jan–Mar 0%	0%	5 ft @ 10 s N	10 kt ESE
13	**Nosy Be**	Northwest	0	0	0	0	0	0	0	0	0	0	0	0	**0%**	Jan–Mar 0%	0%	–	5 kt E

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical good+ morning = medians over good+ mornings: swell height @ peak period and the direction it comes from.
Coasts are ranked by year-round good+ %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

Peru: when and where it's good+

Good+ = swell ≥ 4 ft over 9 s, or ≥ 3 ft over 14 s; breaking under 16 ft; wind sea ≤ 0.35× swell; on the ~08:00 sample.
Open-ocean coasts: the world maps' good+ standard, read from Météo-France MFWAM at each coast's ~8 km cell, with swell aimed at the shore.





#	Coast	Area	J	F	M	A	M	J	J	A	S	O	N	D	Year	Best 3 months	Ocean good+	Typical good+ morning	Morning wind
1	**Mollendo**	Arequipa	68	74	80	85	87	83	83	84	81	82	82	64	**79%**	Apr–Jun 85%	79%	5 ft @ 13 s SSW	9 kt SE
2	**Ilo**	Moquegua	66	70	81	84	85	84	81	85	81	83	78	61	**78%**	Apr–Jun 84%	78%	5 ft @ 13 s SSW	7 kt SE
3	**Camaná**	Arequipa	58	68	74	78	81	79	75	78	73	76	69	55	**72%**	Apr–Jun 79%	72%	5 ft @ 13 s SSW	10 kt SE
4	**Miraflores**	Lima	55	61	72	75	75	65	67	61	57	66	70	46	**64%**	Mar–May 74%	64%	5 ft @ 14 s SSW	11 kt SSE
5	**Cerro Azul**	Lima	45	49	70	69	79	65	68	64	58	67	61	42	**61%**	Mar–May 73%	61%	5 ft @ 14 s SSW	7 kt SSE
6	**Huarmey**	Áncash	45	48	65	71	74	63	58	58	58	58	62	37	**58%**	Mar–May 70%	58%	5 ft @ 14 s SSW	10 kt SSE
7	**Huanchaco**	Trujillo	41	40	61	69	75	71	67	61	62	61	54	33	**58%**	Apr–Jun 72%	58%	5 ft @ 14 s SSW	9 kt SSE
8	**Punta Hermosa**	Lima	39	42	59	67	76	61	66	57	56	61	59	38	**57%**	Apr–Jun 68%	57%	5 ft @ 14 s SSW	7 kt SSE
9	**Chicama**	La Libertad	27	28	54	59	65	54	55	47	51	49	43	23	**46%**	Mar–May 59%	46%	5 ft @ 14 s SSW	10 kt SSE
10	**Pacasmayo**	La Libertad	13	10	33	41	49	39	42	32	41	32	29	10	**31%**	May–Jul 43%	31%	4 ft @ 13 s SSW	10 kt SSE
11	**Máncora**	Piura	6	4	10	7	18	24	12	10	14	18	4	6	**11%**	May–Jul 18%	11%	4 ft @ 16 s SW	15 kt S
12	**Lobitos**	Piura	6	3	6	2	8	13	3	3	4	3	1	3	**4%**	May–Jul 8%	4%	4 ft @ 16 s SW	13 kt SSE
13	**Cabo Blanco**	Piura	6	1	1	0	4	1	1	2	2	1	0	3	**2%**	Dec–Feb 3%	2%	4 ft @ 17 s W	15 kt S

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical good+ morning = medians over good+ mornings: swell height @ peak period and the direction it comes from.
Coasts are ranked by year-round good+ %. Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.