
# Year-round: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 1,803 mornings, Nov 2021 to Oct 2026




## Best named breaks year-round

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Arica**<br>Chile · 4 kt S | **78%** |
| 2 | **Iquique**<br>Chile · 7 kt S | **78%** |
| 3 | **Piha**<br>New Zealand · 14 kt SW | **63%** |
| 4 | **Punta Hermosa**<br>Peru · 7 kt SSE | **60%** |
| 5 | **Skeleton Bay**<br>Namibia · 10 kt S | **60%** |
| 6 | **Elands Bay**<br>South Africa · 9 kt SSE | **57%** |
| 7 | **Punta de Lobos**<br>Chile · 12 kt S | **56%** |
| 8 | **Short Sands**<br>Oregon · 14 kt N | **56%** |
| 9 | **Krui**<br>Lampung, Sumatra · 10 kt SE | **52%** |
| 10 | **Mentawais**<br>Sumatra, Indonesia · 6 kt SE | **49%** |
| 11 | **Cactus Beach**<br>South Australia · 12 kt SE | **47%** |
| 12 | **Montara**<br>California · 12 kt NW | **47%** |
| 13 | **Chicama**<br>Peru · 10 kt SSE | **46%** |
| 14 | **Cimaja**<br>Pelabuhan Ratu, West Java · 4 kt E | **46%** |
| 15 | **Nias**<br>Sumatra, Indonesia · 6 kt SE | **46%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆1 | **Ancud coast**<br>Chile · 9 kt N | **78%** |
| ◆2 | **Antofagasta coast**<br>Chile · 8 kt S | **78%** |
| ◆3 | **La Ligua coast**<br>Chile · 12 kt S | **77%** |
| ◆4 | **Camaná coast**<br>Peru · 10 kt SE | **74%** |
| ◆5 | **Toltén coast**<br>Chile · 13 kt S | **72%** |
| ◆6 | **La Serena coast**<br>Chile · 7 kt SSW | **64%** |
| ◆7 | **Chañaral coast**<br>Chile · 6 kt SSW | **62%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea good = 4 ft @ 6 s




Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

# Popular breaks, month by month

The 40 well-known breaks with the strongest three-month season (best one per 100 miles of coast), listed by year-round consistency. Breaks in the enclosed seas are on the map insets.

| # | Break | Region | J | F | M | A | M | J | J | A | S | O | N | D | Year | Worst | Best 3 months | Tide range | Typical good morning | Morning wind |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | **Arica** | Chile | 70 | 74 | 83 | 85 | 90 | 77 | 82 | 79 | 81 | 79 | 75 | 66 | **78%** | 66% | Mar–May 86% | 3–4 ft | 6 ft @ 13 s SSW, surf ~9 ft | 4 kt S |
| 2 | **Iquique** | Chile | 76 | 83 | 81 | 84 | 88 | 74 | 78 | 76 | 76 | 77 | 77 | 69 | **78%** | 69% | Mar–May 84% | 3–4 ft | 6 ft @ 13 s SSW, surf ~9 ft | 7 kt S |
| 3 | **Piha** | New Zealand | 56 | 68 | 74 | 70 | 70 | 53 | 54 | 59 | 55 | 60 | 73 | 65 | **63%** | 53% | Mar–May 72% | 8–10 ft | 7 ft @ 14 s SW, surf ~10 ft | 14 kt SW |
| 4 | **Punta Hermosa** | Peru | 48 | 45 | 70 | 73 | 79 | 61 | 60 | 65 | 59 | 55 | 58 | 42 | **60%** | 42% | Mar–May 74% | 2–3 ft | 5 ft @ 14 s SSW, surf ~8 ft | 7 kt SSE |
| 5 | **Skeleton Bay** | Namibia | 63 | 71 | 48 | 56 | 68 | 61 | 61 | 66 | 57 | 52 | 59 | 53 | **60%** | 48% | May–Jul 63% | 4–5 ft | 7 ft @ 12 s SSW, surf ~9 ft | 10 kt S |
| 6 | **Elands Bay** | South Africa | 55 | 57 | 44 | 49 | 71 | 55 | 63 | 66 | 58 | 50 | 55 | 56 | **57%** | 44% | May–Jul 63% | 4–6 ft | 6 ft @ 13 s SW, surf ~8 ft | 9 kt SSE |
| 7 | **Punta de Lobos** | Chile | 45 | 45 | 61 | 55 | 68 | 59 | 64 | 57 | 56 | 52 | 53 | 52 | **56%** | 45% | May–Jul 64% | 4–5 ft | 8 ft @ 13 s SW, surf ~11 ft | 12 kt S |
| 8 | **Short Sands** | Oregon | 79 | 79 | 63 | 70 | 61 | 40 | 16 | 19 | 53 | 56 | 69 | 66 | **56%** | 16% | Dec–Feb 75% | 8–10 ft | 7 ft @ 12 s WNW, surf ~9 ft | 14 kt N |
| 9 | **Krui** | Lampung, Sumatra | 29 | 28 | 30 | 51 | 61 | 73 | 79 | 77 | 71 | 63 | 27 | 29 | **52%** | 27% | Jun–Aug 76% | 3–4 ft | 5 ft @ 15 s SSW, surf ~8 ft | 10 kt SE |
| 10 | **Mentawais** | Sumatra, Indonesia | 33 | 29 | 31 | 49 | 57 | 73 | 77 | 76 | 59 | 53 | 15 | 29 | **49%** | 15% | Jun–Aug 75% | 3–3 ft | 5 ft @ 14 s SSW, surf ~8 ft | 6 kt SE |
| 11 | **Cactus Beach** | South Australia | 23 | 41 | 42 | 67 | 57 | 46 | 50 | 58 | 54 | 49 | 36 | 37 | **47%** | 23% | Apr–Jun 57% | 4–5 ft | 7 ft @ 14 s SW, surf ~10 ft | 12 kt SE |
| 12 | **Montara** | California | 68 | 64 | 50 | 45 | 46 | 37 | 19 | 17 | 39 | 53 | 67 | 58 | **47%** | 17% | Nov–Jan 64% | 6–7 ft | 7 ft @ 12 s WNW, surf ~10 ft | 12 kt NW |
| 13 | **Chicama** | Peru | 24 | 31 | 53 | 61 | 61 | 45 | 50 | 49 | 53 | 45 | 49 | 31 | **46%** | 24% | Mar–May 58% | 3–4 ft | 5 ft @ 14 s SSW, surf ~8 ft | 10 kt SSE |
| 14 | **Cimaja** | Pelabuhan Ratu, West Java | 31 | 25 | 41 | 53 | 59 | 66 | 59 | 57 | 47 | 50 | 28 | 32 | **46%** | 25% | May–Jul 61% | 3–4 ft | 5 ft @ 14 s SSW, surf ~8 ft | 4 kt E |
| 15 | **Nias** | Sumatra, Indonesia | 31 | 26 | 23 | 50 | 54 | 71 | 74 | 72 | 53 | 47 | 9 | 32 | **46%** | 9% | Jun–Aug 72% | 2–3 ft | 5 ft @ 14 s SSW, surf ~8 ft | 6 kt SE |
| 16 | **Imsouane** | Morocco | 69 | 62 | 67 | 51 | 27 | 16 | 15 | 16 | 39 | 44 | 59 | 72 | **45%** | 15% | Dec–Feb 68% | 7–9 ft | 6 ft @ 13 s NW, surf ~9 ft | 11 kt N |
| 17 | **Jeffreys Bay** | South Africa | 43 | 43 | 38 | 35 | 49 | 44 | 43 | 45 | 40 | 40 | 44 | 46 | **43%** | 35% | May–Jul 45% | 4–6 ft | 8 ft @ 12 s SW, surf ~11 ft | 13 kt W |
| 18 | **G-Land** | Java, Indonesia | 43 | 39 | 47 | 55 | 29 | 18 | 17 | 33 | 47 | 66 | 53 | 54 | **42%** | 17% | Oct–Dec 58% | 6–8 ft | 6 ft @ 14 s SSW, surf ~9 ft | 10 kt ESE |
| 19 | **Hossegor** | France | 48 | 74 | 62 | 42 | 21 | 23 | 19 | 26 | 41 | 45 | 46 | 54 | **41%** | 19% | Jan–Mar 61% | 10–13 ft | 6 ft @ 13 s NW, surf ~9 ft | 9 kt W |
| 20 | **Shipstern Bluff** | Tasmania, Australia | 48 | 46 | 46 | 46 | 39 | 34 | 33 | 35 | 31 | 27 | 47 | 37 | **39%** | 27% | Jan–Mar 47% | 3–3 ft | 8 ft @ 13 s SW, surf ~12 ft | 16 kt SW |
| 21 | **Ericeira** | Portugal | 39 | 52 | 54 | 45 | 29 | 25 | 14 | 21 | 37 | 54 | 44 | 52 | **39%** | 14% | Oct–Dec 50% | 7–10 ft | 7 ft @ 12 s NW, surf ~10 ft | 13 kt N |
| 22 | **Desert Point** | Lombok, Indonesia | 39 | 39 | 44 | 45 | 30 | 21 | 16 | 32 | 41 | 55 | 48 | 50 | **38%** | 16% | Oct–Dec 51% | 4–6 ft | 5 ft @ 14 s SSW, surf ~8 ft | 10 kt ESE |
| 23 | **Lakey Peak** | Sumbawa, Indonesia | 44 | 39 | 39 | 31 | 23 | 17 | 25 | 41 | 45 | 56 | 31 | 43 | **36%** | 17% | Aug–Oct 47% | 7–10 ft | 5 ft @ 14 s SSW, surf ~8 ft | 8 kt E |
| 24 | **Tofino** | British Columbia | 38 | 48 | 49 | 47 | 41 | 29 | 10 | 13 | 44 | 40 | 35 | 35 | **36%** | 10% | Feb–Apr 48% | 8–11 ft | 7 ft @ 11 s W, surf ~10 ft | 11 kt ESE |
| 25 | **Bundoran** | Ireland | 55 | 44 | 41 | 36 | 32 | 28 | 18 | 19 | 37 | 37 | 41 | 38 | **35%** | 18% | Jan–Mar 46% | 9–12 ft | 7 ft @ 12 s WNW, surf ~10 ft | 13 kt S |
| 26 | **Safi** | Morocco | 48 | 45 | 48 | 33 | 22 | 19 | 13 | 17 | 32 | 41 | 47 | 53 | **35%** | 13% | Nov–Jan 49% | 7–10 ft | 7 ft @ 12 s NW, surf ~10 ft | 12 kt NNE |
| 27 | **Margaret River** | Western Australia | 31 | 26 | 26 | 39 | 37 | 33 | 24 | 29 | 36 | 51 | 33 | 35 | **33%** | 24% | Sep–Nov 40% | 2–2 ft | 9 ft @ 13 s SW, surf ~13 ft | 13 kt ESE |
| 28 | **Rapa Nui** | Easter Island | 28 | 30 | 23 | 35 | 37 | 28 | 34 | 37 | 35 | 41 | 35 | 20 | **32%** | 20% | Aug–Oct 38% | 2–2 ft | 7 ft @ 14 s SW, surf ~10 ft | 11 kt E |
| 29 | **Santa Catalina** | Panama | 9 | 13 | 31 | 47 | 53 | 34 | 41 | 33 | 35 | 37 | 27 | 11 | **31%** | 9% | Apr–Jun 45% | 9–11 ft | 4 ft @ 15 s SSW, surf ~7 ft | 7 kt NE |
| 30 | **Punta Roca** | El Salvador | 7 | 9 | 31 | 47 | 54 | 38 | 39 | 36 | 40 | 33 | 29 | 6 | **31%** | 6% | Apr–Jun 46% | 6–7 ft | 4 ft @ 15 s SSW, surf ~7 ft | 7 kt ENE |
| 31 | **Todos Santos** | Baja California | 63 | 60 | 51 | 35 | 27 | 17 | 5 | 2 | 5 | 18 | 36 | 49 | **31%** | 2% | Jan–Mar 58% | 5–7 ft | 5 ft @ 14 s WNW, surf ~8 ft | 8 kt NW |
| 32 | **Sayulita** | Nayarit, Mexico | 28 | 18 | 24 | 35 | 43 | 42 | 35 | 36 | 33 | 34 | 21 | 15 | **31%** | 15% | Apr–Jun 40% | 3–5 ft | 4 ft @ 15 s SW, surf ~7 ft | 7 kt N |
| 33 | **Puerto Escondido** | Oaxaca, Mexico | 5 | 18 | 22 | 36 | 42 | 41 | 44 | 40 | 37 | 43 | 29 | 9 | **30%** | 5% | May–Jul 42% | 3–4 ft | 4 ft @ 15 s SSW, surf ~7 ft | 6 kt WNW |
| 34 | **Pascuales** | Colima, Mexico | 5 | 9 | 23 | 40 | 48 | 43 | 44 | 40 | 43 | 42 | 22 | 5 | **30%** | 5% | May–Jul 45% | 2–3 ft | 4 ft @ 15 s SSW, surf ~7 ft | 6 kt E |
| 35 | **Lhoknga** | Aceh, Sumatra | 14 | 10 | 17 | 23 | 37 | 45 | 27 | 46 | 33 | 28 | 5 | 23 | **26%** | 5% | Jun–Aug 40% | 2–3 ft | 4 ft @ 15 s SSW, surf ~7 ft | 11 kt S |
| 36 | **Montañita** | Ecuador | 16 | 17 | 23 | 33 | 46 | 30 | 26 | 26 | 29 | 29 | 24 | 11 | **26%** | 11% | Apr–Jun 37% | 6–8 ft | 4 ft @ 15 s SW, surf ~7 ft | 8 kt SSW |
| 37 | **Rincon** | California | 50 | 42 | 32 | 23 | 21 | 13 | 3 | 5 | 5 | 12 | 27 | 37 | **23%** | 3% | Dec–Feb 43% | 5–7 ft | 5 ft @ 14 s W, surf ~8 ft | 8 kt NW |
| 38 | **La Santa** | Lanzarote, Canaries | 38 | 32 | 34 | 22 | 9 | 7 | 4 | 4 | 15 | 25 | 22 | 39 | **21%** | 4% | Dec–Feb 36% | 6–8 ft | 6 ft @ 13 s NW, surf ~9 ft | 14 kt NNE |
| 39 | **Cloud 9** | Siargao, Philippines | 30 | 34 | 33 | 39 | 10 | 1 | 1 | 3 | 3 | 11 | 24 | 48 | **20%** | 1% | Dec–Feb 37% | 5–6 ft | 6 ft @ 10 s NE, surf ~8 ft | 10 kt ENE |
| 40 | **Hanalei Bay** | Kauai, Hawaii | 47 | 38 | 21 | 13 | 7 | 1 | 0 | 1 | 5 | 27 | 21 | 33 | **18%** | 0% | Dec–Feb 39% | 2–2 ft | 7 ft @ 13 s NNW, surf ~11 ft | 15 kt E |

Month cells use the map colours (% of mornings good+). Typical good morning = medians over the good+ mornings: swell height @ peak period and the direction it comes from, and estimated breaking surf (Komar–Gaughan). Tide range = typical to big-tide daily range (EOT20, DGFI-TUM). Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward.

Each break is read from its nearest seaward model cell (0.5° in East and Southeast Asia, 1° elsewhere) with swell inside its exposure window. The rule has no wind; local reefs, points and sheltering are not modelled.

# Under-the-radar stretches, month by month

Coast where mornings are often good+ with swell aimed at the shore (≥40% year-round, or ≥65% over its best three months), at least 300 km from the well-known breaks and within 150 km of a town of 5,000+. Listed by year-round consistency; numbers match the ◆ markers on the maps.


| # | Stretch | Region | J | F | M | A | M | J | J | A | S | O | N | D | Year | Worst | Best 3 months | Tide range | Typical good morning | Morning wind |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| ◆1 | **Ancud coast** | Chile | 85 | 79 | 75 | 77 | 76 | 75 | 65 | 74 | 73 | 81 | 83 | 90 | **78%** | 65% | Nov–Jan 86% | 4–5 ft | 7 ft @ 13 s WSW, surf ~10 ft | 9 kt N |
| ◆2 | **Antofagasta coast** | Chile | 79 | 79 | 82 | 84 | 84 | 77 | 79 | 73 | 73 | 77 | 75 | 70 | **78%** | 70% | Mar–May 83% | 3–4 ft | 6 ft @ 13 s SW, surf ~9 ft | 8 kt S |
| ◆3 | **La Ligua coast** | Chile | 72 | 80 | 84 | 79 | 80 | 76 | 78 | 70 | 73 | 75 | 76 | 77 | **77%** | 70% | Feb–Apr 81% | 4–5 ft | 7 ft @ 13 s SW, surf ~10 ft | 12 kt S |
| ◆4 | **Camaná coast** | Peru | 68 | 75 | 80 | 82 | 81 | 70 | 82 | 75 | 72 | 70 | 73 | 59 | **74%** | 59% | Mar–May 81% | 3–3 ft | 6 ft @ 13 s SSW, surf ~9 ft | 10 kt SE |
| ◆5 | **Toltén coast** | Chile | 80 | 76 | 73 | 72 | 75 | 67 | 63 | 71 | 73 | 70 | 76 | 74 | **72%** | 63% | Dec–Feb 76% | 4–5 ft | 8 ft @ 13 s SW, surf ~11 ft | 13 kt S |
| ◆6 | **La Serena coast** | Chile | 70 | 68 | 74 | 73 | 66 | 61 | 56 | 50 | 51 | 64 | 61 | 69 | **64%** | 50% | Feb–Apr 71% | 4–5 ft | 7 ft @ 13 s SW, surf ~10 ft | 7 kt SSW |
| ◆7 | **Chañaral coast** | Chile | 62 | 59 | 62 | 61 | 69 | 65 | 69 | 63 | 65 | 65 | 53 | 48 | **62%** | 48% | May–Jul 68% | 3–4 ft | 5 ft @ 14 s SW, surf ~8 ft | 6 kt SSW |
| ◆8 | **Greymouth coast** | New Zealand | 37 | 46 | 52 | 58 | 63 | 49 | 57 | 65 | 62 | 54 | 54 | 48 | **54%** | 37% | Jul–Sep 61% | 7–9 ft | 6 ft @ 13 s W, surf ~9 ft | 9 kt E |
| ◆9 | **Isabela coast** | Galápagos, Ecuador | 50 | 44 | 56 | 63 | 69 | 51 | 47 | 50 | 42 | 55 | 46 | 40 | **51%** | 40% | Mar–May 62% | 4–5 ft | 4 ft @ 15 s SW, surf ~8 ft | 10 kt SSE |
| ◆10 | **Marcona coast** | Peru | 48 | 57 | 65 | 71 | 59 | 39 | 45 | 43 | 38 | 43 | 54 | 41 | **50%** | 38% | Mar–May 65% | 3–3 ft | 6 ft @ 13 s SSW, surf ~9 ft | 13 kt SE |
| ◆11 | **Port Lincoln coast** | Australia | 41 | 43 | 47 | 51 | 51 | 45 | 48 | 54 | 55 | 52 | 34 | 52 | **48%** | 34% | Aug–Oct 54% | 3–4 ft | 7 ft @ 14 s SW, surf ~10 ft | 12 kt SE |
| ◆12 | **Nzeto coast** | Angola | 28 | 38 | 29 | 53 | 64 | 60 | 71 | 76 | 58 | 37 | 28 | 24 | **47%** | 24% | Jun–Aug 69% | 4–5 ft | 5 ft @ 13 s SSW, surf ~8 ft | 7 kt SSE |
| ◆13 | **Meningie coast** | Australia | 35 | 57 | 52 | 62 | 47 | 39 | 42 | 49 | 46 | 44 | 44 | 48 | **47%** | 35% | Feb–Apr 57% | 2–3 ft | 7 ft @ 14 s SW, surf ~11 ft | 10 kt S |
| ◆14 | **Oranjemund coast** | Namibia | 52 | 41 | 37 | 34 | 61 | 55 | 50 | 56 | 41 | 40 | 40 | 41 | **46%** | 34% | May–Jul 55% | 4–6 ft | 7 ft @ 13 s SW, surf ~10 ft | 9 kt SE |
| ◆15 | **Quellón coast** | Chile | 63 | 55 | 38 | 42 | 42 | 36 | 33 | 39 | 32 | 51 | 52 | 58 | **45%** | 32% | Dec–Feb 58% | 4–5 ft | 8 ft @ 13 s WSW, surf ~12 ft | 13 kt W |
| ◆16 | **Pointe-Noire coast** | Congo (Brazzaville) | 25 | 28 | 22 | 44 | 64 | 57 | 65 | 77 | 56 | 31 | 31 | 21 | **44%** | 21% | Jun–Aug 66% | 4–5 ft | 5 ft @ 13 s SSW, surf ~7 ft | 6 kt S |
| ◆17 | **Blitar coast** | Indonesia | 45 | 37 | 44 | 56 | 35 | 25 | 17 | 35 | 57 | 63 | 47 | 49 | **42%** | 17% | Sep–Nov 56% | 5–7 ft | 6 ft @ 14 s SSW, surf ~9 ft | 12 kt ESE |
| ◆18 | **Port Denison coast** | Australia | 32 | 46 | 37 | 48 | 45 | 49 | 37 | 37 | 45 | 53 | 46 | 36 | **42%** | 32% | Sep–Nov 48% | 2–2 ft | 9 ft @ 14 s SW, surf ~12 ft | 12 kt SSE |
| ◆19 | **Sumbe coast** | Angola | 22 | 27 | 18 | 37 | 59 | 55 | 67 | 77 | 50 | 37 | 32 | 17 | **42%** | 17% | Jun–Aug 66% | 4–5 ft | 5 ft @ 13 s SW, surf ~8 ft | 5 kt S |
| ◆20 | **Nelson coast** | New Zealand | 45 | 55 | 52 | 41 | 42 | 30 | 30 | 44 | 31 | 32 | 53 | 43 | **41%** | 30% | Jan–Mar 50% | 8–11 ft | 8 ft @ 13 s WSW, surf ~11 ft | 8 kt SSW |
| ◆21 | **Kaitaia coast** | New Zealand | 37 | 49 | 52 | 36 | 33 | 28 | 33 | 41 | 42 | 38 | 53 | 51 | **41%** | 28% | Oct–Dec 47% | 7–9 ft | 7 ft @ 14 s SW, surf ~11 ft | 13 kt S |
| ◆22 | **Bredasdorp coast** | South Africa | 46 | 44 | 39 | 44 | 48 | 35 | 37 | 32 | 38 | 32 | 47 | 46 | **41%** | 32% | Nov–Jan 46% | 5–6 ft | 8 ft @ 12 s SW, surf ~12 ft | 10 kt W |
| ◆23 | **Huarmey coast** | Peru | 45 | 34 | 55 | 53 | 50 | 33 | 35 | 34 | 43 | 31 | 38 | 32 | **40%** | 31% | Mar–May 53% | 3–3 ft | 5 ft @ 14 s SSW, surf ~9 ft | 10 kt SSE |
| ◆24 | **Gwadar coast** | Pakistan | 0 | 0 | 0 | 1 | 6 | 67 | 95 | 86 | 34 | 3 | 0 | 0 | **25%** | 0% | Jun–Aug 82% | 6–8 ft | 8 ft @ 11 s SSW, surf ~10 ft | 8 kt WSW |
| ◆25 | **Salalah coast** | Oman | 0 | 0 | 0 | 0 | 7 | 55 | 95 | 66 | 21 | 0 | 0 | 0 | **21%** | 0% | Jun–Aug 72% | 4–5 ft | 7 ft @ 10 s S, surf ~10 ft | 6 kt SW |
| ◆26 | **Karachi coast** | Pakistan | 0 | 0 | 0 | 0 | 2 | 43 | 86 | 69 | 28 | 2 | 0 | 0 | **20%** | 0% | Jun–Aug 66% | 7–9 ft | 8 ft @ 11 s SW, surf ~11 ft | 10 kt W |
| ◆27 | **Chabahar coast** | Iran | 0 | 0 | 0 | 0 | 2 | 46 | 92 | 70 | 15 | 3 | 0 | 0 | **19%** | 0% | Jun–Aug 69% | 6–8 ft | 7 ft @ 11 s S, surf ~9 ft | 8 kt ESE |

Month cells use the map colours (% of mornings good+). Typical good morning = medians over the good+ mornings: swell height @ peak period and the direction it comes from, and estimated breaking surf (Komar–Gaughan). Tide range = typical to big-tide daily range (EOT20, DGFI-TUM). Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward.

Each stretch is read from one WW3 cell 50–120 km offshore. The rule has no wind, which flatters cold, steady coasts (Chile, Peru, Namibia) and storm coasts (southern Chile, New Zealand's West Coast); check local wind and access before booking a trip.

# October–March: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 889 mornings, Nov 2021 to Oct 2026, these months only




## Best named breaks, October–March

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Iquique**<br>Chile · 7 kt S | **77%** |
| 2 | **Arica**<br>Chile · 5 kt S | **74%** |
| 3 | **Short Sands**<br>Oregon · 14 kt S | **69%** |
| 4 | **Piha**<br>New Zealand · 13 kt SSW | **66%** |
| 5 | **Imsouane**<br>Morocco · 8 kt NNE | **63%** |
| 6 | **Montara**<br>California · 11 kt NNW | **60%** |
| 7 | **Skeleton Bay**<br>Namibia · 8 kt S | **58%** |
| 8 | **Hossegor**<br>France · 10 kt SSW | **55%** |
| 9 | **Punta Hermosa**<br>Peru · 7 kt SSE | **53%** |
| 10 | **Elands Bay**<br>South Africa · 9 kt S | **53%** |
| 11 | **Punta de Lobos**<br>Chile · 12 kt SSW | **51%** |
| 12 | **G-Land**<br>Java, Indonesia · 9 kt W | **50%** |
| 13 | **Ericeira**<br>Portugal · 14 kt NNW | **49%** |
| 14 | **Safi**<br>Morocco · 11 kt ENE | **47%** |
| 15 | **Todos Santos**<br>Baja California · 6 kt NW | **47%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆1 | **Ancud coast**<br>Chile · 8 kt SE | **82%** |
| ◆3 | **La Ligua coast**<br>Chile · 11 kt S | **77%** |
| ◆2 | **Antofagasta coast**<br>Chile · 7 kt SSW | **77%** |
| ◆5 | **Toltén coast**<br>Chile · 13 kt S | **75%** |
| ◆4 | **Camaná coast**<br>Peru · 9 kt SE | **71%** |
| ◆6 | **La Serena coast**<br>Chile · 5 kt SSW | **68%** |
| ◆7 | **Chañaral coast**<br>Chile · 5 kt SSW | **58%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea
### good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

# April–September: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 914 mornings, Apr 2022 to Sep 2026, these months only




## Best named breaks, April–September

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Arica**<br>Chile · 4 kt SSE | **82%** |
| 2 | **Iquique**<br>Chile · 7 kt S | **79%** |
| 3 | **Krui**<br>Lampung, Sumatra · 9 kt SE | **69%** |
| 4 | **Punta Hermosa**<br>Peru · 8 kt S | **66%** |
| 5 | **Mentawais**<br>Sumatra, Indonesia · 5 kt SE | **65%** |
| 6 | **Nias**<br>Sumatra, Indonesia · 5 kt SE | **62%** |
| 7 | **Skeleton Bay**<br>Namibia · 10 kt S | **62%** |
| 8 | **Elands Bay**<br>South Africa · 8 kt SSE | **61%** |
| 9 | **Piha**<br>New Zealand · 15 kt WSW | **60%** |
| 10 | **Punta de Lobos**<br>Chile · 12 kt S | **60%** |
| 11 | **Cimaja**<br>Pelabuhan Ratu, West Java · 4 kt E | **57%** |
| 12 | **Cactus Beach**<br>South Australia · 12 kt ENE | **55%** |
| 13 | **Chicama**<br>Peru · 10 kt SSE | **53%** |
| 14 | **Short Sands**<br>Oregon · 13 kt N | **43%** |
| 15 | **Pascuales**<br>Colima, Mexico · 6 kt E | **43%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆2 | **Antofagasta coast**<br>Chile · 10 kt S | **78%** |
| ◆4 | **Camaná coast**<br>Peru · 11 kt SE | **77%** |
| ◆3 | **La Ligua coast**<br>Chile · 13 kt S | **76%** |
| ◆1 | **Ancud coast**<br>Chile · 11 kt N | **73%** |
| ◆5 | **Toltén coast**<br>Chile · 12 kt SSE | **70%** |
| ◆7 | **Chañaral coast**<br>Chile · 8 kt SSW | **65%** |
| ◆12 | **Nzeto coast**<br>Angola · 7 kt SSE | **64%** |


**Great Lakes · good = 4 ft @ 6 s**




**Mediterranean & Black Sea · good = 4 ft @ 6 s**




**Caspian Sea**
**good = 4 ft @ 6 s**





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

# November: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (135 mornings; 15 days missing from the archive)




## Best named breaks in November

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Iquique**<br>Chile · 7 kt S | **77%** |
| 2 | **Arica**<br>Chile · 5 kt S | **75%** |
| 3 | **Piha**<br>New Zealand · 11 kt SW | **73%** |
| 4 | **Short Sands**<br>Oregon · 11 kt SSE | **69%** |
| 5 | **Montara**<br>California · 11 kt NNW | **67%** |
| 6 | **Skeleton Bay**<br>Namibia · 7 kt S | **59%** |
| 7 | **Imsouane**<br>Morocco · 5 kt NNE | **59%** |
| 8 | **Punta Hermosa**<br>Peru · 6 kt SSE | **58%** |
| 9 | **Elands Bay**<br>South Africa · 9 kt S | **55%** |
| 10 | **G-Land**<br>Java, Indonesia · 6 kt E | **53%** |
| 11 | **Punta de Lobos**<br>Chile · 11 kt SSW | **53%** |
| 12 | **Chicama**<br>Peru · 9 kt SSE | **49%** |
| 13 | **Desert Point**<br>Lombok, Indonesia · 7 kt SE | **48%** |
| 14 | **Safi**<br>Morocco · 11 kt NE | **47%** |
| 15 | **Shipstern Bluff**<br>Tasmania, Australia · 15 kt WNW | **47%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆1 | **Ancud coast**<br>Chile · 10 kt NNW | **83%** |
| ◆3 | **La Ligua coast**<br>Chile · 12 kt S | **76%** |
| ◆5 | **Toltén coast**<br>Chile · 11 kt S | **76%** |
| ◆2 | **Antofagasta coast**<br>Chile · 9 kt SSW | **75%** |
| ◆4 | **Camaná coast**<br>Peru · 10 kt SE | **73%** |
| ◆6 | **La Serena coast**<br>Chile · 8 kt SSW | **61%** |
| ◆8 | **Greymouth coast**<br>New Zealand · 5 kt ESE | **54%** |


**Great Lakes · good = 4 ft @ 6 s**




**Mediterranean & Black Sea · good = 4 ft @ 6 s**




**Caspian Sea**
**good = 4 ft @ 6 s**





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

# December: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (155 mornings)




Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

## Best named breaks in December

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Imsouane**<br>Morocco · 7 kt NNE | **72%** |
| 2 | **Iquique**<br>Chile · 7 kt S | **69%** |
| 3 | **Short Sands**<br>Oregon · 14 kt SSE | **66%** |
| 4 | **Arica**<br>Chile · 5 kt S | **66%** |
| 5 | **Piha**<br>New Zealand · 11 kt N | **65%** |
| 6 | **Montara**<br>California · 11 kt NNW | **58%** |
| 7 | **Elands Bay**<br>South Africa · 9 kt S | **56%** |
| 8 | **Uluwatu**<br>Bali, Indonesia · 9 kt W | **55%** |
| 9 | **Hossegor**<br>France · 9 kt SE | **54%** |
| 10 | **Safi**<br>Morocco · 12 kt ENE | **53%** |
| 11 | **Skeleton Bay**<br>Namibia · 9 kt S | **53%** |
| 12 | **Ericeira**<br>Portugal · 14 kt ENE | **52%** |
| 13 | **Punta de Lobos**<br>Chile · 12 kt SSW | **52%** |
| 14 | **Todos Santos**<br>Baja California · 5 kt NNW | **49%** |
| 15 | **Cloud 9**<br>Siargao, Philippines · 10 kt ENE | **48%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆1 | **Ancud coast**<br>Chile · 8 kt SSW | **90%** |
| ◆3 | **La Ligua coast**<br>Chile · 12 kt S | **77%** |
| ◆5 | **Toltén coast**<br>Chile · 14 kt S | **74%** |
| ◆2 | **Antofagasta coast**<br>Chile · 7 kt S | **70%** |
| ◆6 | **La Serena coast**<br>Chile · 6 kt SSW | **69%** |
| ◆4 | **Camaná coast**<br>Peru · 9 kt SE | **59%** |
| ◆15 | **Quellón coast**<br>Chile · 13 kt SW | **58%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

# January: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)




## Best named breaks in January

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Short Sands**<br>Oregon · 14 kt S | **79%** |
| 2 | **Iquique**<br>Chile · 6 kt S | **76%** |
| 3 | **Arica**<br>Chile · 5 kt S | **70%** |
| 4 | **Imsouane**<br>Morocco · 9 kt NNE | **69%** |
| 5 | **Montara**<br>California · 13 kt SSE | **68%** |
| 6 | **Todos Santos**<br>Baja California · 5 kt SE | **63%** |
| 7 | **Skeleton Bay**<br>Namibia · 7 kt S | **63%** |
| 8 | **Piha**<br>New Zealand · 13 kt ENE | **56%** |
| 9 | **Elands Bay**<br>South Africa · 8 kt SSE | **55%** |
| 10 | **Bundoran**<br>Ireland · 13 kt SSE | **55%** |
| 11 | **Rincon**<br>California · 7 kt NW | **50%** |
| 12 | **Safi**<br>Morocco · 11 kt ENE | **48%** |
| 13 | **Shipstern Bluff**<br>Tasmania, Australia · 11 kt SW | **48%** |
| 14 | **Honolua Bay**<br>Maui, Hawaii · 14 kt E | **48%** |
| 15 | **Punta Hermosa**<br>Peru · 7 kt SSE | **48%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆1 | **Ancud coast**<br>Chile · 8 kt SE | **85%** |
| ◆5 | **Toltén coast**<br>Chile · 14 kt S | **80%** |
| ◆2 | **Antofagasta coast**<br>Chile · 5 kt SSW | **79%** |
| ◆3 | **La Ligua coast**<br>Chile · 10 kt S | **72%** |
| ◆6 | **La Serena coast**<br>Chile · 4 kt SSW | **70%** |
| ◆4 | **Camaná coast**<br>Peru · 9 kt ESE | **68%** |
| ◆15 | **Quellón coast**<br>Chile · 11 kt SSW | **63%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea
good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

# February: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (141 mornings)




Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

## Best named breaks in February

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Iquique**<br>Chile · 7 kt S | **83%** |
| 2 | **Short Sands**<br>Oregon · 13 kt SSE | **79%** |
| 3 | **Hossegor**<br>France · 8 kt SSW | **74%** |
| 4 | **Arica**<br>Chile · 5 kt S | **74%** |
| 5 | **Skeleton Bay**<br>Namibia · 9 kt S | **71%** |
| 6 | **Piha**<br>New Zealand · 11 kt SSW | **68%** |
| 7 | **Montara**<br>California · 11 kt NNW | **64%** |
| 8 | **Imsouane**<br>Morocco · 6 kt NNE | **62%** |
| 9 | **Todos Santos**<br>Baja California · 6 kt NW | **60%** |
| 10 | **Elands Bay**<br>South Africa · 9 kt S | **57%** |
| 11 | **Nazaré**<br>Portugal · 12 kt WNW | **52%** |
| 12 | **Tofino**<br>British Columbia · 10 kt ESE | **48%** |
| 13 | **Shipstern Bluff**<br>Tasmania, Australia · 16 kt SW | **46%** |
| 14 | **Punta Hermosa**<br>Peru · 7 kt SSE | **45%** |
| 15 | **Punta de Lobos**<br>Chile · 14 kt SSW | **45%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆3 | **La Ligua coast**<br>Chile · 12 kt S | **80%** |
| ◆2 | **Antofagasta coast**<br>Chile · 6 kt SSW | **79%** |
| ◆1 | **Ancud coast**<br>Chile · 7 kt SE | **79%** |
| ◆5 | **Toltén coast**<br>Chile · 14 kt S | **76%** |
| ◆4 | **Camaná coast**<br>Peru · 8 kt ESE | **75%** |
| ◆6 | **La Serena coast**<br>Chile · 5 kt SSW | **68%** |
| ◆7 | **Chañaral coast**<br>Chile · 5 kt SSW | **59%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea
### good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

# March: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)




Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

## Best named breaks in March

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Arica**<br>Chile · 5 kt S | **83%** |
| 2 | **Iquique**<br>Chile · 7 kt S | **81%** |
| 3 | **Piha**<br>New Zealand · 15 kt SSW | **74%** |
| 4 | **Punta Hermosa**<br>Peru · 7 kt SSE | **70%** |
| 5 | **Imsouane**<br>Morocco · 11 kt N | **67%** |
| 6 | **Short Sands**<br>Oregon · 16 kt S | **63%** |
| 7 | **Hossegor**<br>France · 13 kt W | **62%** |
| 8 | **Punta de Lobos**<br>Chile · 14 kt SSW | **61%** |
| 9 | **Ericeira**<br>Portugal · 12 kt SSW | **54%** |
| 10 | **Chicama**<br>Peru · 9 kt SE | **53%** |
| 11 | **Todos Santos**<br>Baja California · 10 kt NNW | **51%** |
| 12 | **Montara**<br>California · 13 kt NNW | **50%** |
| 13 | **Tofino**<br>British Columbia · 12 kt ESE | **49%** |
| 14 | **Uluwatu**<br>Bali, Indonesia · 10 kt ENE | **49%** |
| 15 | **Safi**<br>Morocco · 13 kt NE | **48%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆3 | **La Ligua coast**<br>Chile · 11 kt S | **84%** |
| ◆2 | **Antofagasta coast**<br>Chile · 7 kt S | **82%** |
| ◆4 | **Camaná coast**<br>Peru · 9 kt ESE | **80%** |
| ◆1 | **Ancud coast**<br>Chile · 9 kt ESE | **75%** |
| ◆6 | **La Serena coast**<br>Chile · 6 kt S | **74%** |
| ◆5 | **Toltén coast**<br>Chile · 15 kt S | **73%** |
| ◆10 | **Marcona coast**<br>Peru · 12 kt SE | **65%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

# April: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)




Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

## Best named breaks in April

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Arica**<br>Chile · 4 kt SSE | **85%** |
| 2 | **Iquique**<br>Chile · 8 kt S | **84%** |
| 3 | **Punta Hermosa**<br>Peru · 8 kt SSE | **73%** |
| 4 | **Short Sands**<br>Oregon · 15 kt N | **70%** |
| 5 | **Piha**<br>New Zealand · 16 kt ENE | **70%** |
| 6 | **Cactus Beach**<br>South Australia · 11 kt ENE | **67%** |
| 7 | **Chicama**<br>Peru · 10 kt SE | **61%** |
| 8 | **Skeleton Bay**<br>Namibia · 10 kt S | **56%** |
| 9 | **Punta de Lobos**<br>Chile · 12 kt S | **55%** |
| 10 | **G-Land**<br>Java, Indonesia · 9 kt ESE | **55%** |
| 11 | **Cimaja**<br>Pelabuhan Ratu, West Java · 4 kt ENE | **53%** |
| 12 | **Imsouane**<br>Morocco · 10 kt N | **51%** |
| 13 | **Krui**<br>Lampung, Sumatra · 6 kt ESE | **51%** |
| 14 | **Nias**<br>Sumatra, Indonesia · 5 kt SSE | **50%** |
| 15 | **Elands Bay**<br>South Africa · 8 kt S | **49%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆2 | **Antofagasta coast**<br>Chile · 10 kt S | **84%** |
| ◆4 | **Camaná coast**<br>Peru · 11 kt ESE | **82%** |
| ◆3 | **La Ligua coast**<br>Chile · 12 kt S | **79%** |
| ◆1 | **Ancud coast**<br>Chile · 9 kt N | **77%** |
| ◆6 | **La Serena coast**<br>Chile · 7 kt S | **73%** |
| ◆5 | **Toltén coast**<br>Chile · 13 kt SSE | **72%** |
| ◆10 | **Marcona coast**<br>Peru · 14 kt SE | **71%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea
### good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

# May: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)




Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

## Best named breaks in May

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Arica**<br>Chile · 3 kt SSE | **90%** |
| 2 | **Iquique**<br>Chile · 7 kt S | **88%** |
| 3 | **Punta Hermosa**<br>Peru · 7 kt S | **79%** |
| 4 | **Elands Bay**<br>South Africa · 8 kt SSE | **71%** |
| 5 | **Piha**<br>New Zealand · 15 kt S | **70%** |
| 6 | **Punta de Lobos**<br>Chile · 11 kt S | **68%** |
| 7 | **Skeleton Bay**<br>Namibia · 10 kt S | **68%** |
| 8 | **Short Sands**<br>Oregon · 14 kt NNW | **61%** |
| 9 | **Chicama**<br>Peru · 10 kt SE | **61%** |
| 10 | **Krui**<br>Lampung, Sumatra · 9 kt ESE | **61%** |
| 11 | **Cimaja**<br>Pelabuhan Ratu, West Java · 4 kt E | **59%** |
| 12 | **Mentawais**<br>Sumatra, Indonesia · 5 kt WSW | **57%** |
| 13 | **Cactus Beach**<br>South Australia · 11 kt E | **57%** |
| 14 | **Nias**<br>Sumatra, Indonesia · 5 kt W | **54%** |
| 15 | **Punta Roca**<br>El Salvador · 5 kt ESE | **54%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆2 | **Antofagasta coast**<br>Chile · 10 kt S | **84%** |
| ◆4 | **Camaná coast**<br>Peru · 11 kt SE | **81%** |
| ◆3 | **La Ligua coast**<br>Chile · 12 kt S | **80%** |
| ◆1 | **Ancud coast**<br>Chile · 8 kt N | **76%** |
| ◆5 | **Toltén coast**<br>Chile · 14 kt SSE | **75%** |
| ◆7 | **Chañaral coast**<br>Chile · 8 kt SSW | **69%** |
| ◆9 | **Isabela coast**<br>Galápagos, Ecuador · 9 kt SSE | **69%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea
### good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.27× the mean period.

# June: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)




## Best named breaks in June

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Arica**<br>Chile · 3 kt SSE | **77%** |
| 2 | **Iquique**<br>Chile · 7 kt S | **74%** |
| 3 | **Mentawais**<br>Sumatra, Indonesia · 4 kt SE | **73%** |
| 4 | **Krui**<br>Lampung, Sumatra · 9 kt ESE | **73%** |
| 5 | **Nias**<br>Sumatra, Indonesia · 4 kt SE | **71%** |
| 6 | **Cimaja**<br>Pelabuhan Ratu, West Java · 4 kt E | **66%** |
| 7 | **Punta Hermosa**<br>Peru · 8 kt S | **61%** |
| 8 | **Skeleton Bay**<br>Namibia · 10 kt SSE | **61%** |
| 9 | **Punta de Lobos**<br>Chile · 14 kt S | **59%** |
| 10 | **Elands Bay**<br>South Africa · 8 kt SE | **55%** |
| 11 | **Piha**<br>New Zealand · 15 kt NE | **53%** |
| 12 | **Cactus Beach**<br>South Australia · 14 kt N | **46%** |
| 13 | **Lhoknga**<br>Aceh, Sumatra · 12 kt S | **45%** |
| 14 | **Chicama**<br>Peru · 11 kt SSE | **45%** |
| 15 | **Jeffreys Bay**<br>South Africa · 12 kt W | **44%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆2 | **Antofagasta coast**<br>Chile · 10 kt S | **77%** |
| ◆3 | **La Ligua coast**<br>Chile · 13 kt S | **76%** |
| ◆1 | **Ancud coast**<br>Chile · 12 kt N | **75%** |
| ◆4 | **Camaná coast**<br>Peru · 11 kt SE | **70%** |
| ◆5 | **Toltén coast**<br>Chile · 13 kt N | **67%** |
| ◆24 | **Gwadar coast**<br>Pakistan · 8 kt WSW | **67%** |
| ◆7 | **Chañaral coast**<br>Chile · 10 kt SSW | **65%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea
### good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.27× the mean period.

# July: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (154 mornings; 1 days missing from the archive)




Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

## Best named breaks in July

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Arica**<br>Chile · 4 kt SSE | **82%** |
| 2 | **Krui**<br>Lampung, Sumatra · 11 kt SE | **79%** |
| 3 | **Iquique**<br>Chile · 7 kt S | **78%** |
| 4 | **Mentawais**<br>Sumatra, Indonesia · 4 kt SE | **77%** |
| 5 | **Nias**<br>Sumatra, Indonesia · 5 kt SE | **74%** |
| 6 | **Punta de Lobos**<br>Chile · 12 kt S | **64%** |
| 7 | **Elands Bay**<br>South Africa · 10 kt SSE | **63%** |
| 8 | **Skeleton Bay**<br>Namibia · 11 kt S | **61%** |
| 9 | **Punta Hermosa**<br>Peru · 8 kt SSE | **60%** |
| 10 | **Cimaja**<br>Pelabuhan Ratu, West Java · 5 kt E | **59%** |
| 11 | **Piha**<br>New Zealand · 15 kt SW | **54%** |
| 12 | **Chicama**<br>Peru · 11 kt SSE | **50%** |
| 13 | **Cactus Beach**<br>South Australia · 13 kt WSW | **50%** |
| 14 | **Puerto Escondido**<br>Oaxaca, Mexico · 6 kt NW | **44%** |
| 15 | **Pascuales**<br>Colima, Mexico · 6 kt NE | **44%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆24 | **Gwadar coast**<br>Pakistan · 7 kt WSW | **95%** |
| ◆25 | **Salalah coast**<br>Oman · 7 kt SW | **95%** |
| ◆27 | **Chabahar coast**<br>Iran · 8 kt ESE | **92%** |
| ◆26 | **Karachi coast**<br>Pakistan · 11 kt WSW | **86%** |
| ◆4 | **Camaná coast**<br>Peru · 11 kt SE | **82%** |
| ◆2 | **Antofagasta coast**<br>Chile · 9 kt S | **79%** |
| ◆3 | **La Ligua coast**<br>Chile · 13 kt S | **78%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea
### good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.28× the mean period.

# August: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (155 mornings)




Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

## Best named breaks in August

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Arica**<br>Chile · 4 kt S | **79%** |
| 2 | **Krui**<br>Lampung, Sumatra · 14 kt SE | **77%** |
| 3 | **Mentawais**<br>Sumatra, Indonesia · 9 kt SE | **76%** |
| 4 | **Iquique**<br>Chile · 8 kt S | **76%** |
| 5 | **Nias**<br>Sumatra, Indonesia · 7 kt SE | **72%** |
| 6 | **Elands Bay**<br>South Africa · 9 kt N | **66%** |
| 7 | **Skeleton Bay**<br>Namibia · 10 kt S | **66%** |
| 8 | **Punta Hermosa**<br>Peru · 8 kt SSE | **65%** |
| 9 | **Piha**<br>New Zealand · 14 kt ESE | **59%** |
| 10 | **Cactus Beach**<br>South Australia · 12 kt E | **58%** |
| 11 | **Punta de Lobos**<br>Chile · 11 kt S | **57%** |
| 12 | **Cimaja**<br>Pelabuhan Ratu, West Java · 5 kt E | **57%** |
| 13 | **Chicama**<br>Peru · 11 kt SSE | **49%** |
| 14 | **Lhoknga**<br>Aceh, Sumatra · 13 kt S | **46%** |
| 15 | **Jeffreys Bay**<br>South Africa · 13 kt W | **45%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆24 | **Gwadar coast**<br>Pakistan · 8 kt WSW | **86%** |
| ◆16 | **Pointe-Noire coast**<br>Congo (Brazzaville) · 6 kt S | **77%** |
| ◆19 | **Sumbe coast**<br>Angola · 4 kt S | **77%** |
| ◆12 | **Nzeto coast**<br>Angola · 7 kt SSE | **76%** |
| ◆4 | **Camaná coast**<br>Peru · 11 kt SE | **75%** |
| ◆1 | **Ancud coast**<br>Chile · 12 kt N | **74%** |
| ◆2 | **Antofagasta coast**<br>Chile · 11 kt S | **73%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea
### good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

# September: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (150 mornings)




Share of mornings with good-or-better surf

0% 2% 5% 10% 15% 20% 30% 40% 50% 60% 80% 100%

No model data (sea ice, or a sea neither source covers)

→ Predominant swell direction (power-weighted; omitted where swell comes from mixed directions)

## Best named breaks in September

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Arica**<br>Chile · 4 kt S | **81%** |
| 2 | **Iquique**<br>Chile · 7 kt S | **76%** |
| 3 | **Krui**<br>Lampung, Sumatra · 12 kt SE | **71%** |
| 4 | **Punta Hermosa**<br>Peru · 8 kt S | **59%** |
| 5 | **Mentawais**<br>Sumatra, Indonesia · 9 kt SE | **59%** |
| 6 | **Elands Bay**<br>South Africa · 8 kt SE | **58%** |
| 7 | **Skeleton Bay**<br>Namibia · 10 kt S | **57%** |
| 8 | **Punta de Lobos**<br>Chile · 13 kt S | **56%** |
| 9 | **Piha**<br>New Zealand · 15 kt WSW | **55%** |
| 10 | **Cactus Beach**<br>South Australia · 11 kt ENE | **54%** |
| 11 | **Chicama**<br>Peru · 9 kt SSE | **53%** |
| 12 | **Nias**<br>Sumatra, Indonesia · 8 kt SSE | **53%** |
| 13 | **Short Sands**<br>Oregon · 13 kt N | **53%** |
| 14 | **Cimaja**<br>Pelabuhan Ratu, West Java · 4 kt E | **47%** |
| 15 | **G-Land**<br>Java, Indonesia · 11 kt ESE | **47%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆2 | **Antofagasta coast**<br>Chile · 10 kt S | **73%** |
| ◆3 | **La Ligua coast**<br>Chile · 14 kt S | **73%** |
| ◆1 | **Ancud coast**<br>Chile · 11 kt WNW | **73%** |
| ◆5 | **Toltén coast**<br>Chile · 11 kt N | **73%** |
| ◆4 | **Camaná coast**<br>Peru · 11 kt SE | **72%** |
| ◆7 | **Chañaral coast**<br>Chile · 7 kt SSW | **65%** |
| ◆8 | **Greymouth coast**<br>New Zealand · 14 kt E | **62%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea
### good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

# October: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2022–2026 (148 mornings; 7 days missing from the archive)




## Best named breaks in October

% of mornings good+ with swell inside the break's exposure window; best per 100 miles; then typical morning wind (GFS)


| | | |
|---|---|---|
| 1 | **Arica**<br>Chile · 4 kt S | **79%** |
| 2 | **Iquique**<br>Chile · 7 kt S | **77%** |
| 3 | **G-Land**<br>Java, Indonesia · 7 kt ESE | **66%** |
| 4 | **Krui**<br>Lampung, Sumatra · 9 kt SE | **63%** |
| 5 | **Piha**<br>New Zealand · 14 kt NW | **60%** |
| 6 | **Short Sands**<br>Oregon · 16 kt S | **56%** |
| 7 | **Lakey Peak**<br>Sumbawa, Indonesia · 6 kt E | **56%** |
| 8 | **Punta Hermosa**<br>Peru · 7 kt SSE | **55%** |
| 9 | **Desert Point**<br>Lombok, Indonesia · 8 kt ESE | **55%** |
| 10 | **Ericeira**<br>Portugal · 13 kt NNE | **54%** |
| 11 | **Mentawais**<br>Sumatra, Indonesia · 4 kt SE | **53%** |
| 12 | **Montara**<br>California · 11 kt NNW | **53%** |
| 13 | **Skeleton Bay**<br>Namibia · 9 kt S | **52%** |
| 14 | **Punta de Lobos**<br>Chile · 14 kt SSW | **52%** |
| 15 | **Margaret River**<br>Western Australia · 9 kt SSW | **51%** |

## Under-the-radar stretches


◆ numbered on the map; month by month on page 3

| | | |
|---|---|---|
| ◆1 | **Ancud coast**<br>Chile · 7 kt SSW | **81%** |
| ◆2 | **Antofagasta coast**<br>Chile · 7 kt S | **77%** |
| ◆3 | **La Ligua coast**<br>Chile · 14 kt S | **75%** |
| ◆4 | **Camaná coast**<br>Peru · 11 kt SE | **70%** |
| ◆5 | **Toltén coast**<br>Chile · 12 kt S | **70%** |
| ◆7 | **Chañaral coast**<br>Chile · 7 kt SSW | **65%** |
| ◆6 | **La Serena coast**<br>Chile · 7 kt SSW | **64%** |


### Great Lakes · good = 4 ft @ 6 s




### Mediterranean & Black Sea · good = 4 ft @ 6 s




### Caspian Sea
### good = 4 ft @ 6 s





Good+ morning: swell of at least 4 ft (1.22 m) with peak period over 9 s, or 3 ft (0.91 m) over 14 s; breaking under 16 ft (Komar–Gaughan from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Enclosed seas (MFWAM via Open-Meteo and Copernicus Marine; 0.75° lakes, 1.5–2° seas) use a relaxed standard: total waves 4–10 ft at a dominant period of at least 6 s.


Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Missing peak periods (Great Lakes, Copernicus years) are taken as 1.29× the mean period.

# Lake Michigan: when and where it's good

Good lake surf = total wave height 4–10 ft (1.22–3 m) with dominant period ≥ 6 s (taken as 1.29× the mean period), arriving within 75° of the shore's facing, on the ~08:00 sample. Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.




**Average of these coasts: % of mornings with good lake surf**





| # | Coast | Area | J | F | M | A | M | J | J | A | S | O | N | D | Year | Best 3 months | Ocean good+ | Tide range | Typical good morning | Morning wind |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | **Holland** | Michigan | 30 | 22 | 17 | 11 | 3 | 2 | 1 | 1 | 5 | 11 | 28 | 26 | **13%** | Nov–Jan 28% | 0% | none | 5 ft @ 7 s W | 13 kt S |
| 2 | **Grand Haven** | Michigan | 28 | 22 | 15 | 10 | 2 | 1 | 0 | 1 | 4 | 11 | 27 | 26 | **12%** | Nov–Jan 27% | 0% | none | 5 ft @ 7 s W | 13 kt S |
| 3 | **Ludington** | Michigan | 23 | 21 | 12 | 9 | 3 | 1 | 1 | 1 | 3 | 11 | 28 | 27 | **11%** | Nov–Jan 26% | 0% | none | 5 ft @ 7 s W | 14 kt S |
| 4 | **Frankfort** | Michigan | 23 | 20 | 14 | 9 | 3 | 1 | 1 | 1 | 4 | 10 | 26 | 25 | **11%** | Nov–Jan 25% | 0% | none | 5 ft @ 7 s W | 14 kt S |
| 5 | **St. Joseph** | Michigan | 24 | 16 | 14 | 6 | 2 | 2 | 2 | 1 | 5 | 11 | 24 | 14 | **10%** | Nov–Jan 21% | 0% | none | 5 ft @ 7 s NW | 13 kt SSW |
| 6 | **New Buffalo** | Michigan | 19 | 14 | 12 | 5 | 2 | 2 | 1 | 1 | 5 | 11 | 16 | 10 | **8%** | Nov–Jan 15% | 0% | none | 5 ft @ 7 s NNW | 13 kt SSW |
| 7 | **Portage** | Indiana | 10 | 7 | 7 | 3 | 1 | 1 | 1 | 1 | 4 | 8 | 6 | 5 | **5%** | Jan–Mar 8% | 0% | none | 5 ft @ 7 s N | 13 kt SSW |
| 8 | **Milwaukee** | Wisconsin | 6 | 7 | 8 | 6 | 1 | 1 | 1 | 2 | 5 | 5 | 5 | 5 | **4%** | Jan–Mar 7% | 0% | none | 5 ft @ 7 s ENE | 13 kt NW |
| 9 | **Sheboygan** | Wisconsin | 6 | 7 | 9 | 7 | 1 | 1 | 0 | 1 | 4 | 6 | 4 | 5 | **4%** | Feb–Apr 8% | 0% | none | 5 ft @ 7 s E | 14 kt S |
| 10 | **Gary** | Indiana | 8 | 5 | 6 | 3 | 1 | 1 | 1 | 1 | 3 | 6 | 5 | 4 | **4%** | Jan–Mar 6% | 0% | none | 5 ft @ 7 s N | 12 kt NW |
| 11 | **Racine** | Wisconsin | 6 | 3 | 3 | 3 | 1 | 1 | 1 | 1 | 3 | 5 | 2 | 2 | **2%** | Jan–Mar 4% | 0% | none | 5 ft @ 7 s NE | 13 kt WNW |
| 12 | **Whiting** | Indiana | 5 | 3 | 6 | 1 | 0 | 1 | 0 | 1 | 3 | 5 | 1 | 2 | **2%** | Jan–Mar 4% | 0% | none | 5 ft @ 7 s N | 12 kt NW |
| 13 | **Chicago** | Illinois | 3 | 2 | 4 | 2 | 0 | 1 | 0 | 1 | 2 | 3 | 0 | 0 | **1%** | Jan–Mar 3% | 0% | none | 5 ft @ 8 s NNE | 13 kt SSW |

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical good morning = medians over good mornings: total wave height @ dominant period and the direction waves come from.


Coasts are ranked by year-round good %. Tide range = typical to big-tide daily range (EOT20, DGFI-TUM). Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

# Lake Superior: when and where it's good

Good lake surf = total wave height 4–10 ft (1.22–3 m) with dominant period ≥ 6 s (taken as 1.29× the mean period), arriving within 75° of the shore's facing, on the ~08:00 sample. Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.




**Average of these coasts: % of mornings with good lake surf**




| # | Coast | Area | J | F | M | A | M | J | J | A | S | O | N | D | Year | Best 3 months | Ocean good+ | Tide range | Typical good morning | Morning wind |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | **Grand Marais MI** | Michigan | 28 | 18 | 13 | 8 | 1 | 3 | 0 | 3 | 6 | 14 | 27 | 23 | **12%** | Nov–Jan 26% | 0% | none | 5 ft @ 7 s NW | 14 kt W |
| 2 | **Whitefish Point** | Michigan | 25 | 12 | 9 | 7 | 1 | 2 | 0 | 3 | 3 | 13 | 24 | 27 | **10%** | Nov–Jan 25% | 0% | none | 5 ft @ 7 s NW | 14 kt SSE |
| 3 | **Marquette** | Michigan | 21 | 17 | 17 | 11 | 1 | 1 | 0 | 2 | 5 | 11 | 21 | 19 | **10%** | Nov–Jan 20% | 0% | none | 6 ft @ 7 s N | 15 kt S |
| 4 | **Au Train** | Michigan | 19 | 16 | 14 | 6 | 1 | 1 | 0 | 2 | 5 | 9 | 24 | 19 | **9%** | Nov–Jan 21% | 0% | none | 5 ft @ 7 s NNW | 15 kt S |
| 5 | **Ontonagon** | Michigan | 19 | 9 | 12 | 11 | 1 | 0 | 0 | 1 | 3 | 10 | 23 | 18 | **9%** | Nov–Jan 20% | 0% | none | 5 ft @ 7 s NW | 13 kt WSW |
| 6 | **Agawa Bay** | Ontario | 20 | 7 | 4 | 3 | 1 | 1 | 0 | 1 | 2 | 8 | 21 | 26 | **8%** | Nov–Jan 22% | 0% | none | 5 ft @ 8 s W | 11 kt SSE |
| 7 | **Terrace Bay** | Ontario | 12 | 9 | 6 | 4 | 2 | 1 | 1 | 1 | 4 | 13 | 20 | 15 | **7%** | Oct–Dec 16% | 0% | none | 5 ft @ 7 s SSW | 13 kt NNE |
| 8 | **Little Girl's Point** | Michigan | 6 | 3 | 8 | 7 | 1 | 0 | 0 | 0 | 0 | 5 | 7 | 7 | **4%** | Nov–Jan 7% | 0% | none | 5 ft @ 7 s N | 13 kt W |
| 9 | **Stoney Point** | Minnesota | 2 | 3 | 4 | 5 | 1 | 0 | 0 | 0 | 0 | 3 | 1 | 2 | **2%** | Feb–Apr 4% | 0% | none | 6 ft @ 7 s ENE | 8 kt SSW |
| 10 | **Park Point** | Duluth, Minnesota | 2 | 2 | 5 | 5 | 1 | 0 | 0 | 0 | 0 | 3 | 1 | 2 | **2%** | Feb–Apr 4% | 0% | none | 6 ft @ 7 s ENE | 8 kt SSW |
| 11 | **Grand Marais MN** | Minnesota | 3 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 3 | 2 | 1 | **1%** | Nov–Jan 2% | 0% | none | 5 ft @ 7 s SSW | 12 kt ENE |

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical good morning = medians over good mornings: total wave height @ dominant period and the direction waves come from.


Coasts are ranked by year-round good %. Tide range = typical to big-tide daily range (EOT20, DGFI-TUM). Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

# Mediterranean: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.




**Average of these coasts: % of mornings with surfable windswell**



# Mediterranean: coast by coast

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.


| # | Coast | Area | J | F | M | A | M | J | J | A | S | O | N | D | Year | Best 3 months | Ocean good+ | Tide range | Typical surfable morning | Morning wind |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | **Ajaccio** | Corsica, France | 54 | 45 | 35 | 31 | 23 | 18 | 23 | 19 | 24 | 24 | 53 | 37 | **32%** | Nov–Jan 48% | 6% | 1–1 ft | 5 ft @ 7 s W | 11 kt WNW |
| 2 | **Susa** | Cyrenaica, Libya | 50 | 45 | 37 | 32 | 19 | 13 | 29 | 15 | 24 | 38 | 30 | 38 | **31%** | Dec–Feb 44% | 3% | 0–0 ft | 4 ft @ 7 s NNW | 8 kt W |
| 3 | **Capo Mannu** | Sardinia, Italy | 47 | 40 | 35 | 29 | 26 | 13 | 22 | 22 | 20 | 26 | 53 | 37 | **31%** | Nov–Jan 46% | 4% | 1–1 ft | 5 ft @ 7 s WNW | 11 kt ESE |
| 4 | **Tabarka** | Tunisia | 51 | 49 | 30 | 33 | 23 | 9 | 19 | 14 | 17 | 22 | 49 | 37 | **29%** | Dec–Feb 46% | 6% | 1–1 ft | 5 ft @ 7 s NW | 12 kt W |
| 5 | **Alexandria** | Egypt | 35 | 45 | 39 | 30 | 26 | 19 | 33 | 21 | 21 | 24 | 21 | 33 | **29%** | Jan–Mar 40% | 2% | 1–1 ft | 4 ft @ 7 s NW | 9 kt NW |
| 6 | **Paphos** | Cyprus | 37 | 43 | 36 | 32 | 28 | 15 | 19 | 25 | 27 | 18 | 21 | 34 | **28%** | Jan–Mar 39% | 1% | 1–1 ft | 4 ft @ 6 s W | 9 kt W |
| 7 | **Falasarna** | Crete, Greece | 45 | 37 | 32 | 31 | 14 | 9 | 19 | 11 | 14 | 16 | 37 | 43 | **25%** | Nov–Jan 41% | 2% | 0–0 ft | 5 ft @ 6 s W | 14 kt NE |
| 8 | **Benghazi** | Libya | 43 | 32 | 28 | 30 | 19 | 11 | 16 | 8 | 12 | 20 | 33 | 48 | **25%** | Nov–Jan 41% | 2% | 0–1 ft | 5 ft @ 6 s NW | 10 kt NW |
| 9 | **Algiers** | Algeria | 34 | 40 | 41 | 29 | 16 | 7 | 11 | 7 | 12 | 16 | 36 | 32 | **23%** | Jan–Mar 38% | 3% | 0–0 ft | 4 ft @ 7 s N | 10 kt WSW |
| 10 | **Gaza** | Palestine | 35 | 39 | 33 | 21 | 23 | 7 | 13 | 10 | 10 | 18 | 19 | 35 | **22%** | Dec–Feb 36% | 2% | 1–1 ft | 4 ft @ 7 s WNW | 7 kt WSW |
| 11 | **Bizerte** | Tunisia | 39 | 35 | 23 | 27 | 15 | 5 | 17 | 11 | 11 | 14 | 30 | 31 | **21%** | Dec–Feb 35% | 1% | 1–1 ft | 5 ft @ 7 s NNW | 13 kt WNW |
| 12 | **Tripoli** | Libya | 43 | 32 | 26 | 30 | 14 | 8 | 10 | 5 | 13 | 16 | 21 | 37 | **21%** | Dec–Feb 37% | 1% | 1–1 ft | 4 ft @ 6 s N | 10 kt ESE |
| 13 | **Beirut** | Lebanon | 37 | 36 | 33 | 21 | 25 | 5 | 7 | 13 | 11 | 15 | 15 | 32 | **21%** | Jan–Mar 35% | 2% | 1–1 ft | 4 ft @ 7 s W | 10 kt SW |
| 14 | **Livorno** | Tuscany, Italy | 33 | 26 | 21 | 19 | 14 | 9 | 13 | 12 | 23 | 18 | 37 | 27 | **21%** | Nov–Jan 32% | 2% | 1–1 ft | 5 ft @ 7 s WSW | 10 kt ENE |
| 15 | **Kalo Nero** | Messinia, Greece | 40 | 35 | 21 | 20 | 8 | 3 | 5 | 3 | 9 | 14 | 39 | 39 | **19%** | Nov–Jan 39% | 1% | 0–0 ft | 4 ft @ 7 s WSW | 9 kt NNW |
| 16 | **Santa Marinella** | Lazio, Italy | 26 | 23 | 25 | 20 | 12 | 7 | 8 | 10 | 15 | 19 | 27 | 27 | **18%** | Nov–Jan 27% | 0% | 1–1 ft | 4 ft @ 6 s SW | 11 kt NNE |
| 17 | **Marseille** | France | 25 | 26 | 17 | 16 | 9 | 5 | 5 | 5 | 10 | 13 | 26 | 15 | **14%** | Jan–Mar 23% | 0% | 1–1 ft | 5 ft @ 6 s WSW | 16 kt NNW |
| 18 | **Barcelona** | Spain | 23 | 19 | 37 | 17 | 2 | 3 | 0 | 5 | 8 | 16 | 17 | 12 | **13%** | Jan–Mar 26% | 1% | 0–1 ft | 4 ft @ 6 s ESE | 10 kt W |
| 19 | **Varazze** | Liguria, Italy | 22 | 20 | 21 | 12 | 7 | 5 | 3 | 5 | 7 | 17 | 21 | 14 | **13%** | Jan–Mar 21% | 1% | 1–1 ft | 4 ft @ 7 s SSW | 10 kt N |
| 20 | **Kelibia** | Cap Bon, Tunisia | 11 | 9 | 29 | 18 | 14 | 2 | 1 | 1 | 6 | 11 | 9 | 14 | **10%** | Mar–May 20% | 0% | 0–0 ft | 4 ft @ 6 s ESE | 13 kt NW |
| 21 | **Mondello** | Sicily, Italy | 14 | 11 | 9 | 11 | 2 | 1 | 3 | 1 | 9 | 7 | 10 | 15 | **8%** | Dec–Feb 13% | 0% | 1–1 ft | 4 ft @ 6 s NNW | 9 kt WSW |
| 22 | **Hammamet** | Tunisia | 6 | 7 | 25 | 15 | 8 | 2 | 1 | 1 | 1 | 7 | 1 | 7 | **7%** | Mar–May 16% | 0% | 0–1 ft | 4 ft @ 6 s ESE | 11 kt NW |
| 23 | **Kastro** | Kyllini, Greece | 14 | 11 | 3 | 5 | 3 | 1 | 0 | 0 | 1 | 1 | 16 | 14 | **6%** | Nov–Jan 15% | 0% | 0–1 ft | 4 ft @ 6 s SW | 8 kt NNW |
| 24 | **Antalya** | Turkey | 14 | 7 | 3 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 4 | 12 | **4%** | Dec–Feb 11% | 0% | 1–1 ft | 4 ft @ 6 s SSE | 4 kt NNW |

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.
Coasts are ranked by year-round surfable %. Tide range = typical to big-tide daily range (EOT20, DGFI-TUM). Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out

# Greece: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.









| # | Coast | Area | J | F | M | A | M | J | J | A | S | O | N | D | Year | Best 3 months | Ocean good+ | Tide range | Typical surfable morning | Morning wind |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | **Zakynthos west** | Ionian Islands | 48 | 38 | 28 | 33 | 17 | 8 | 17 | 12 | 14 | 27 | 46 | 45 | **27%** | Nov–Jan 46% | 1% | 0–1 ft | 5 ft @ 6 s W | 11 kt NNW |
| 2 | **Falasarna** | West Crete | 45 | 37 | 32 | 32 | 14 | 9 | 18 | 14 | 15 | 17 | 37 | 43 | **26%** | Nov–Jan 41% | 2% | 0–0 ft | 5 ft @ 6 s W | 14 kt NE |
| 3 | **Komos** | South Crete | 36 | 33 | 28 | 27 | 15 | 5 | 8 | 7 | 7 | 12 | 28 | 40 | **21%** | Dec–Feb 36% | 2% | 0–0 ft | 4 ft @ 6 s W | 9 kt N |
| 4 | **Kalo Nero** | Messinia, W Peloponnese | 40 | 35 | 25 | 22 | 10 | 5 | 7 | 3 | 9 | 14 | 39 | 39 | **20%** | Nov–Jan 39% | 2% | 0–0 ft | 4 ft @ 7 s WSW | 9 kt NNW |
| 5 | **Kokkini Hani** | North Crete | 31 | 30 | 28 | 15 | 14 | 7 | 13 | 10 | 19 | 25 | 23 | 23 | **20%** | Jan–Mar 30% | 0% | 0–0 ft | 4 ft @ 6 s N | 11 kt W |
| 6 | **Agios Gordios** | Corfu, west coast | 34 | 28 | 21 | 23 | 12 | 3 | 8 | 4 | 12 | 19 | 33 | 35 | **19%** | Nov–Jan 34% | 0% | 1–1 ft | 4 ft @ 6 s WSW | 12 kt NNW |
| 7 | **Chania** | North Crete | 22 | 27 | 26 | 16 | 17 | 11 | 14 | 15 | 20 | 25 | 23 | 16 | **19%** | Jan–Mar 25% | 0% | 0–0 ft | 5 ft @ 6 s N | 12 kt N |
| 8 | **Kymi** | East Evia | 22 | 24 | 23 | 11 | 8 | 1 | 3 | 9 | 15 | 18 | 16 | 17 | **14%** | Jan–Mar 23% | 0% | 1–1 ft | 5 ft @ 6 s NE | 13 kt NNW |
| 9 | **Tinos north** | Cyclades | 19 | 23 | 20 | 10 | 9 | 5 | 6 | 8 | 17 | 22 | 16 | 11 | **14%** | Jan–Mar 21% | 0% | 0–1 ft | 5 ft @ 5 s N | 15 kt NNW |
| 10 | **Kastro** | Kyllini, NW Peloponnese | 14 | 13 | 3 | 3 | 1 | 0 | 0 | 0 | 1 | 2 | 16 | 14 | **5%** | Nov–Jan 15% | 0% | 0–1 ft | 4 ft @ 6 s SSW | 8 kt NNW |
| 11 | **Ialysos** | Rhodes, northwest | 6 | 5 | 4 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 4 | 2 | **2%** | Jan–Mar 5% | 0% | 1–1 ft | 4 ft @ 6 s W | 9 kt NNW |

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.
Coasts are ranked by year-round surfable %. Tide range = typical to big-tide daily range (EOT20, DGFI-TUM). Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out

# Caspian Sea: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.








| # | Coast | Area | J | F | M | A | M | J | J | A | S | O | N | D | Year | Best 3 months | Ocean good+ | Tide range | Typical surfable morning | Morning wind |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | **Nabran** | Azerbaijan | 37 | 26 | 19 | 15 | 9 | 5 | 6 | 8 | 17 | 26 | 29 | 36 | **19%** | Nov–Jan 34% | 0% | none | 5 ft @ 6 s NE | 12 kt NNW |
| 2 | **Ramsar** | Iran | 17 | 18 | 15 | 8 | 8 | 7 | 11 | 5 | 15 | 16 | 13 | 14 | **12%** | Jan–Mar 17% | 1% | none | 4 ft @ 6 s N | 5 kt NNW |
| 3 | **Babolsar** | Iran | 15 | 17 | 12 | 8 | 6 | 8 | 14 | 6 | 13 | 11 | 16 | 14 | **12%** | Dec–Feb 16% | 0% | none | 4 ft @ 7 s NNW | 7 kt E |
| 4 | **Novkhani** | Absheron, Azerbaijan | 17 | 19 | 14 | 9 | 6 | 5 | 8 | 8 | 10 | 14 | 18 | 14 | **12%** | Dec–Feb 17% | 0% | none | 5 ft @ 6 s N | 12 kt NNW |
| 5 | **Derbent** | Dagestan, Russia | 22 | 9 | 8 | 9 | 3 | 1 | 2 | 6 | 11 | 17 | 15 | 27 | **11%** | Nov–Jan 21% | 0% | none | 4 ft @ 6 s E | 12 kt NNW |
| 6 | **Awaza** | Turkmenistan | 12 | 14 | 12 | 8 | 8 | 5 | 8 | 5 | 7 | 8 | 13 | 12 | **9%** | Jan–Mar 13% | 0% | none | 4 ft @ 6 s NW | 10 kt NNW |
| 7 | **Makhachkala** | Dagestan, Russia | 22 | 8 | 7 | 5 | 1 | 0 | 1 | 1 | 7 | 15 | 15 | 27 | **9%** | Nov–Jan 21% | 0% | none | 4 ft @ 6 s ESE | 12 kt SE |
| 8 | **Aktau** | Kazakhstan | 15 | 10 | 8 | 3 | 6 | 0 | 1 | 0 | 1 | 7 | 20 | 14 | **7%** | Nov–Jan 16% | 0% | none | 5 ft @ 6 s W | 13 kt ESE |
| 9 | **Lankaran** | Azerbaijan | 8 | 8 | 4 | 1 | 1 | 0 | 1 | 1 | 2 | 4 | 4 | 3 | **3%** | Jan–Mar 6% | 0% | none | 4 ft @ 6 s ENE | 9 kt S |
| 10 | **Shikhov** | Baku, Azerbaijan | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | **0%** | Nov–Jan 1% | 0% | none | 4 ft @ 5 s SE | 12 kt N |


Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.
Coasts are ranked by year-round surfable %. Tide range = typical to big-tide daily range (EOT20, DGFI-TUM). Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out

# Black Sea: when and where it's surfable

Surfable windswell = total wave height 3–10 ft (0.9–3 m) with mean period ≥ 5 s, arriving within 75° of the shore's facing, on the ~08:00 sample.
Enclosed seas rarely see groundswell, so this regional standard sits alongside ocean good+, the open-ocean standard used on the other pages.




**Average of these coasts: % of mornings with surfable windswell**






| # | Coast | Area | J | F | M | A | M | J | J | A | S | O | N | D | Year | Best 3 months | Ocean good+ | Tide range | Typical surfable morning | Morning wind |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | **Kilyos** | Istanbul, Turkey | 33 | 36 | 26 | 18 | 18 | 9 | 12 | 34 | 38 | 29 | 24 | 35 | **26%** | Dec–Feb 35% | 0% | 0–0 ft | 5 ft @ 6 s NNE | 11 kt NNE |
| 2 | **Şile** | Istanbul, Turkey | 32 | 35 | 23 | 15 | 14 | 7 | 10 | 27 | 33 | 28 | 19 | 33 | **23%** | Dec–Feb 33% | 0% | 0–0 ft | 5 ft @ 6 s NNE | 10 kt NE |
| 3 | **Vama Veche** | Romania | 22 | 32 | 21 | 9 | 13 | 4 | 3 | 15 | 27 | 18 | 16 | 27 | **17%** | Dec–Feb 27% | 0% | 0–0 ft | 5 ft @ 6 s ENE | 13 kt NNW |
| 4 | **Sozopol** | Bulgaria | 23 | 30 | 18 | 9 | 11 | 3 | 2 | 19 | 27 | 18 | 16 | 24 | **17%** | Dec–Feb 26% | 0% | 0–0 ft | 4 ft @ 6 s ENE | 12 kt N |
| 5 | **Batumi** | Georgia | 25 | 27 | 15 | 15 | 5 | 7 | 6 | 3 | 13 | 15 | 19 | 19 | **14%** | Dec–Feb 23% | 1% | 0–0 ft | 4 ft @ 7 s WNW | 8 kt E |
| 6 | **Sochi** | Russia | 31 | 28 | 9 | 13 | 4 | 4 | 3 | 1 | 6 | 9 | 24 | 25 | **13%** | Dec–Feb 28% | 1% | 0–0 ft | 5 ft @ 6 s WSW | 8 kt SE |
| 7 | **Varna** | Bulgaria | 19 | 25 | 16 | 8 | 10 | 1 | 1 | 14 | 19 | 11 | 9 | 21 | **13%** | Dec–Feb 21% | 0% | 0–0 ft | 4 ft @ 6 s ENE | 13 kt N |
| 8 | **Anapa** | Russia | 23 | 17 | 9 | 15 | 3 | 3 | 1 | 0 | 4 | 11 | 16 | 23 | **10%** | Dec–Feb 21% | 0% | 0–0 ft | 5 ft @ 6 s WSW | 13 kt ENE |
| 9 | **Odesa** | Ukraine | 12 | 6 | 4 | 3 | 2 | 1 | 1 | 0 | 1 | 1 | 11 | 10 | **4%** | Nov–Jan 11% | 0% | 0–0 ft | 4 ft @ 6 s SSE | 12 kt NNE |

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical surfable morning = medians over surfable mornings: total wave height @ mean period and the direction waves come from.
Coasts are ranked by year-round surfable %. Tide range = typical to big-tide daily range (EOT20, DGFI-TUM). Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out

# Madagascar: when and where it's good+

Good+ = swell ≥ 4 ft over 9 s, or ≥ 3 ft over 14 s; breaking under 16 ft; wind sea ≤ 0.35× swell; on the ~08:00 sample.
Open-ocean coasts: the world maps' good+ standard, read from Météo-France MFWAM at each coast's ~8 km cell, with swell aimed at the shore.









| # | Coast | Area | J | F | M | A | M | J | J | A | S | O | N | D | Year | Best 3 months | Ocean good+ | Tide range | Typical good+ morning | Morning wind |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | **Lavanono** | Deep south | 35 | 26 | 34 | 51 | 65 | 64 | 68 | 58 | 54 | 45 | 44 | 41 | **49%** | May–Jul 66% | 49% | 5–6 ft | 6 ft @ 13 s SSW | 9 kt E |
| 2 | **Anakao** | Toliara, southwest | 19 | 24 | 23 | 43 | 50 | 49 | 54 | 52 | 41 | 26 | 34 | 31 | **37%** | Jun–Aug 52% | 37% | 6–9 ft | 5 ft @ 13 s SSW | 9 kt SE |
| 3 | **Manakara** | Southeast | 34 | 40 | 37 | 49 | 37 | 46 | 51 | 44 | 36 | 30 | 19 | 19 | **37%** | Jun–Aug 47% | 37% | 1–1 ft | 5 ft @ 10 s SE | 9 kt NNE |
| 4 | **Ifaty** | Toliara, southwest | 14 | 22 | 22 | 39 | 43 | 45 | 47 | 47 | 36 | 26 | 25 | 23 | **33%** | Jun–Aug 46% | 33% | 7–9 ft | 5 ft @ 14 s SSW | 10 kt SE |
| 5 | **Andavadoaka** | Morombe, west | 14 | 13 | 21 | 35 | 37 | 45 | 42 | 37 | 29 | 20 | 16 | 21 | **28%** | Jun–Aug 41% | 28% | 7–10 ft | 5 ft @ 13 s SSW | 10 kt SSE |
| 6 | **Fort Dauphin** | Tolagnaro, southeast | 17 | 6 | 15 | 24 | 23 | 29 | 25 | 17 | 10 | 13 | 7 | 12 | **17%** | May–Jul 26% | 17% | 1–1 ft | 6 ft @ 11 s SSE | 15 kt NNE |
| 7 | **Mahambo** | East | 8 | 8 | 3 | 11 | 15 | 17 | 22 | 33 | 17 | 4 | 1 | 2 | **12%** | Jun–Aug 24% | 12% | 2–2 ft | 5 ft @ 10 s ESE | 10 kt SSE |
| 8 | **Morondava** | West | 1 | 4 | 2 | 5 | 6 | 12 | 8 | 12 | 9 | 6 | 1 | 2 | **6%** | Jun–Aug 11% | 6% | 8–12 ft | 5 ft @ 11 s SW | 7 kt ENE |
| 9 | **Maintirano** | West | 1 | 2 | 2 | 3 | 1 | 4 | 5 | 6 | 5 | 5 | 2 | 1 | **3%** | Jul–Sep 5% | 3% | 10–13 ft | 5 ft @ 10 s SW | 9 kt S |
| 10 | **Sainte-Marie** | Nosy Boraha, east | 1 | 3 | 0 | 5 | 6 | 4 | 8 | 3 | 3 | 2 | 0 | 0 | **3%** | May–Jul 6% | 3% | 2–3 ft | 5 ft @ 10 s ESE | 10 kt SSE |
| 11 | **Ramena** | Antsiranana, north | 1 | 0 | 1 | 0 | 0 | 2 | 0 | 1 | 0 | 0 | 0 | 1 | **0%** | Jun–Aug 1% | 0% | 5–7 ft | 4 ft @ 10 s ESE | 19 kt SE |
| 12 | **Mahajanga** | Northwest | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | **0%** | Jan–Mar 0% | 0% | 9–12 ft | 5 ft @ 10 s N | 10 kt ESE |
| 13 | **Nosy Be** | Northwest | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | **0%** | Jan–Mar 0% | 0% | 8–11 ft | – | 5 kt E |

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical good+ morning = medians over good+ mornings: swell height @ peak period and the direction it comes from.
Coasts are ranked by year-round good+ %. Tide range = typical to big-tide daily range (EOT20, DGFI-TUM). Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.

# Peru: when and where it's good+

Good+ = swell ≥ 4 ft over 9 s, or ≥ 3 ft over 14 s; breaking under 16 ft; wind sea ≤ 0.35× swell; on the ~08:00 sample.
Open-ocean coasts: the world maps' good+ standard, read from Météo-France MFWAM at each coast's ~8 km cell, with swell aimed at the shore.




| # | Coast | Area | J | F | M | A | M | J | J | A | S | O | N | D | Year | Best 3 months | Ocean good+ | Tide range | Typical good+ morning | Morning wind |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | **Mollendo** | Arequipa | 68 | 74 | 80 | 85 | 87 | 83 | 83 | 84 | 81 | 82 | 82 | 64 | **79%** | Apr–Jun 85% | 79% | 3–4 ft | 5 ft @ 13 s SSW | 9 kt SE |
| 2 | **Ilo** | Moquegua | 66 | 70 | 81 | 84 | 85 | 84 | 81 | 85 | 81 | 83 | 78 | 61 | **78%** | Apr–Jun 84% | 78% | 3–4 ft | 5 ft @ 13 s SSW | 7 kt SE |
| 3 | **Camaná** | Arequipa | 58 | 68 | 74 | 78 | 81 | 79 | 75 | 78 | 73 | 76 | 69 | 55 | **72%** | Apr–Jun 79% | 72% | 3–3 ft | 5 ft @ 13 s SSW | 10 kt SE |
| 4 | **Miraflores** | Lima | 55 | 61 | 72 | 75 | 75 | 65 | 67 | 61 | 57 | 66 | 70 | 46 | **64%** | Mar–May 74% | 64% | 2–3 ft | 5 ft @ 14 s SSW | 11 kt SSE |
| 5 | **Cerro Azul** | Lima | 45 | 49 | 70 | 69 | 79 | 65 | 68 | 64 | 58 | 67 | 61 | 42 | **61%** | Mar–May 73% | 61% | 2–3 ft | 5 ft @ 14 s SSW | 7 kt SSE |
| 6 | **Huarmey** | Áncash | 45 | 48 | 65 | 71 | 74 | 63 | 58 | 58 | 58 | 58 | 62 | 37 | **58%** | Mar–May 70% | 58% | 3–3 ft | 5 ft @ 14 s SSW | 10 kt SSE |
| 7 | **Huanchaco** | Trujillo | 41 | 40 | 61 | 69 | 75 | 71 | 67 | 61 | 62 | 61 | 54 | 33 | **58%** | Apr–Jun 72% | 58% | 3–4 ft | 5 ft @ 14 s SSW | 9 kt SSE |
| 8 | **Punta Hermosa** | Lima | 39 | 42 | 59 | 67 | 76 | 61 | 66 | 57 | 56 | 61 | 59 | 38 | **57%** | Apr–Jun 68% | 57% | 2–3 ft | 5 ft @ 14 s SSW | 7 kt SSE |
| 9 | **Chicama** | La Libertad | 27 | 28 | 54 | 59 | 65 | 54 | 55 | 47 | 51 | 49 | 43 | 23 | **46%** | Mar–May 59% | 46% | 3–4 ft | 5 ft @ 14 s SSW | 10 kt SSE |
| 10 | **Pacasmayo** | La Libertad | 13 | 10 | 33 | 41 | 49 | 39 | 42 | 32 | 41 | 32 | 29 | 10 | **31%** | May–Jul 43% | 31% | 3–4 ft | 4 ft @ 13 s SSW | 10 kt SSE |
| 11 | **Máncora** | Piura | 6 | 4 | 10 | 7 | 18 | 24 | 12 | 10 | 14 | 18 | 4 | 6 | **11%** | May–Jul 18% | 11% | 5–6 ft | 4 ft @ 16 s SW | 15 kt S |
| 12 | **Lobitos** | Piura | 6 | 3 | 6 | 2 | 8 | 13 | 3 | 3 | 4 | 3 | 1 | 3 | **4%** | May–Jul 8% | 4% | 4–6 ft | 4 ft @ 16 s SW | 13 kt SSE |
| 13 | **Cabo Blanco** | Piura | 6 | 1 | 1 | 0 | 4 | 1 | 1 | 2 | 2 | 1 | 0 | 3 | **2%** | Dec–Feb 3% | 2% | 5–6 ft | 4 ft @ 17 s W | 15 kt S |

Météo-France MFWAM (via Open-Meteo and Copernicus Marine) at each coast's model cell, a few km offshore. Typical good+ morning = medians over good+ mornings: swell height @ peak period and the direction it comes from.
Coasts are ranked by year-round good+ %. Tide range = typical to big-tide daily range (EOT20, DGFI-TUM). Morning wind = median speed and most common direction over all mornings of the period, NOAA GFS 10 m wind just offshore, December 2022 onward. Neither standard uses wind; check it before paddling out.