
November: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (135 mornings; 15 days missing from the archive)



Best named breaks in November


% of mornings that are good+ with swell inside the break's exposure window

1	**Arica** Chile	**86%**
2	**Punta Hermosa** Peru	**75%**
3	**Piha** New Zealand	**74%**
4	**Elands Bay** South Africa	**68%**
5	**Short Sands** Oregon	**68%**
6	**Mavericks** California	**68%**
7	**Ocean Beach SF** California	**68%**
8	**Lakey Peak** Sumbawa, Indonesia	**67%**
9	**Imsouane** Morocco	**63%**
10	**Chicama** Peru	**61%**
11	**Anchor Point** Taghazout, Morocco	**59%**
12	**Todos Santos** Baja California	**58%**
13	**Raglan** New Zealand	**57%**
14	**Uluwatu** Bali, Indonesia	**57%**
15	**Keramas** Bali, Indonesia	**57%**
16	**Desert Point** Lombok, Indonesia	**56%**
17	**G-Land** Java, Indonesia	**54%**
18	**Punta de Lobos** Chile	**50%**
19	**Puerto Escondido** Oaxaca, Mexico	**49%**
20	**Hossegor** France	**48%**
21	**Shipstern Bluff** Tasmania, Australia	**48%**
22	**Zurriola** San Sebastián, Spain	**47%**
23	**Bundoran** Ireland	**46%**
24	**Santa Catalina** Panama	**46%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea




Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Open-Meteo has no swell peak period on the Great Lakes; it is taken as 1.29× the mean swell period, the median ratio in the other seas.

December: where the surf is good

Share of mornings with good-or-better surf in the wave-model record, 5 seasons, 2021–2025 (155 mornings)



Best named breaks in December


% of mornings that are good+ with swell inside the break's exposure window

1	**Arica** Chile	**81%**
2	**Imsouane** Morocco	**74%**
3	**Anchor Point** Taghazout, Morocco	**72%**
4	**Elands Bay** South Africa	**71%**
5	**Punta Hermosa** Peru	**69%**
6	**Piha** New Zealand	**68%**
7	**Hossegor** France	**64%**
8	**Lakey Peak** Sumbawa, Indonesia	**63%**
9	**Short Sands** Oregon	**63%**
10	**Desert Point** Lombok, Indonesia	**62%**
11	**Zurriola** San Sebastián, Spain	**61%**
12	**Uluwatu** Bali, Indonesia	**61%**
13	**Keramas** Bali, Indonesia	**61%**
14	**Mavericks** California	**60%**
15	**Ocean Beach SF** California	**60%**
16	**Todos Santos** Baja California	**60%**
17	**G-Land** Java, Indonesia	**59%**
18	**Chicama** Peru	**58%**
19	**Raglan** New Zealand	**57%**
20	**Mundaka** Basque Country, Spain	**54%**
21	**Safi** Morocco	**54%**
22	**Black's Beach** California	**52%**
23	**Ericeira** Portugal	**52%**
24	**Nias** Sumatra, Indonesia	**52%**


Great Lakes



Mediterranean & Black Sea



Caspian Sea



Good+ morning: swell peak period ≥ 10 s, breaking height 1.5–5 m (Komar–Gaughan, from swell height and period), and wind-sea height ≤ 0.35× swell height. Judged on one sample near 08:00 local solar time.

Ocean: NOAA WAVEWATCH III global model (Univ. of Hawai'i run, PacIOOS ERDDAP), 1° sampling. Great Lakes, Mediterranean, Black Sea, Caspian: Open-Meteo Marine API (Météo-France MFWAM), 0.75° lakes / 1.5–2° seas. Both judged by the same open-ocean standard.

Offshore sea state only: bathymetry, sheltering, tide and wind direction at each break are not modelled. Break percentages use the nearest seaward model cell within 250 km. Open-Meteo has no swell peak period on the Great Lakes; it is taken as 1.29× the mean swell period, the median ratio in the other seas.