

Deworming as HIV Prevention for Young Women: Evidence from Zimbabwe

Jon Denton-Schneider (Clark)

November 2, 2024

HIV Pandemic through Economic Lens

Most of world's HIV: East and Southern Africa, where **hugely disproportionate share of new infections among women 15-24**

HIV Pandemic through Economic Lens

Most of world's HIV: East and Southern Africa, where **hugely disproportionate share of new infections among women 15-24**

HIV status



Marriage market matching (partner age gap)

HIV Pandemic through Economic Lens

Most of world's HIV: East and Southern Africa, where **hugely disproportionate share of new infections among women 15-24**

HIV status



Marriage market matching (partner age gap)



Adult human capital (health and education)

HIV Pandemic through Economic Lens

Most of world's HIV: East and Southern Africa, where **hugely disproportionate share of new infections among women 15-24**

HIV status



Marriage market matching (partner age gap)



Adult human capital (health and education)



Childhood health (very cheap and effective interventions!)

HIV Pandemic through Economic Lens

Most of world's HIV: East and Southern Africa, where **hugely disproportionate share of new infections among women 15-24**

HIV status



Marriage market matching (partner age gap)



Adult human capital (health and education)



Childhood health (very cheap and effective interventions!)


Q: Can cheap improvements in girls' health (e.g., deworming) reduce their chances of contracting HIV as young women?

Theory: Urogenital Schistosomiasis → HIV

Disease: Inflammatory response to worm eggs getting trapped in nearby tissues → “anaemia, growth stunting, impaired cognition, ... organ-specific effects such as ... urogenital inflammation and scarring” (Colley et al., 2014, p. 2253)


Theory: Urogenital Schistosomiasis → HIV

Disease: Inflammatory response to worm eggs getting trapped in nearby tissues → “anaemia, growth stunting, impaired cognition, ... organ-specific effects such as ... urogenital inflammation and scarring” (Colley et al., 2014, p. 2253)




Theory: Urogenital Schistosomiasis → HIV

Disease: Inflammatory response to worm eggs getting trapped in nearby tissues → “anaemia, growth stunting, impaired cognition, ... organ-specific effects such as ... urogenital inflammation and scarring” (Colley et al., 2014, p. 2253)




Theory: Urogenital Schistosomiasis → HIV

Disease: Inflammatory response to worm eggs getting trapped in nearby tissues → “anaemia, growth stunting, impaired cognition, ... organ-specific effects such as ... urogenital inflammation and scarring” (Colley et al., 2014, p. 2253)



Novel: Linking childhood health to HIV via learning / attendance

Evidence: Urogenital Schistosomiasis → HIV




Heavy *S. haematobium* Infection:


Students, 2010

Source: Midzi et al. (2014)

Evidence: Urogenital Schistosomiasis → HIV



Heavy *S. haematobium* Infection:
Students, 2010
Source: Midzi et al. (2014)



HIV Prevalence:
Ages 15-49, 2005 and 2010
Source: DHS data

Quasi-Experiment: School-Based Deworming Program



Source: WHO (2012)

Wedza, 17 Sept. 2012 – In line with the new global momentum towards the control, elimination and eradication of neglected tropical diseases (NTDs), Zimbabwe launched a mass drug administration against schistosomiasis (bilharzia) and soil transmitted helminthes (intestinal worms) at a function held at Wedza High School.

The mass drug administration is the final phase of a process which started with a national prevalence survey in 2010, and the development of the master plan that began in 2011 and completed in 2012.

The National Prevalence Survey of 2010 showed that Mashonaland East Province, under which Wedza district falls was one of the highly affected. The mass drug administration will therefore target people, mainly under the age of 15, and will be delivered through the country's network of schools and health facilities in the high burden districts. The mass drug administration was made possible by WHO which donated to the Ministry of Health and Child Welfare Praziquantel (PZQ) used in the treatment of bilharzia, and Albendazole (ALB) for intestinal worms. A total of 2 583 000 PZQ tablets (600mg), and 2 450 200 ALB tablets (400mg) were donated. These drugs are expected to cover 3 794 638 people mainly under the age of 15 in the high burden districts.

End of 2012 school year: Mass deworming begins, planned to treat **3.8 million students (est. 5.2 million under age 15)**

Quasi-Experiment: School-Based Deworming Program



Source: WHO (2012)

Wedza, 17 Sept. 2012 – In line with the new global momentum towards the control, elimination and eradication of neglected tropical diseases (NTDs), Zimbabwe launched a mass drug administration against schistosomiasis (bilharzia) and soil transmitted helminthes (intestinal worms) at a function held at Wedza High School.

The mass drug administration is the final phase of a process which started with a national prevalence survey in 2010, and the development of the master plan that began in 2011 and completed in 2012.

The National Prevalence Survey of 2010 showed that Mashonaland East Province, under which Wedza district falls was one of the highly affected. The mass drug administration will therefore target people, mainly under the age of 15, and will be delivered through the country's network of schools and health facilities in the high burden districts. The mass drug administration was made possible by WHO which donated to the Ministry of Health and Child Welfare Praziquantel (PZQ) used in the treatment of bilharzia, and Albendazole (ALB) for intestinal worms. A total of 2 583 000 PZQ tablets (600mg), and 2 450 200 ALB tablets (400mg) were donated. These drugs are expected to cover 3 794 638 people mainly under the age of 15 in the high burden districts.

End of 2012 school year: Mass deworming begins, planned to treat 3.8 million students (est. 5.2 million under age 15)

Effects: Morbidity ↓ 85% after 1 round, sustained across subsequent rounds (Mduluzza et al., 2020)

Defining Comparison Groups

Prevalence category	Districts (IUs)	Comments and intervention strategies
≥10%	Murehwa, Shamva, Mwenezi, Shurugwi, Chikomba, Mutoko, UMP, Hwedza, Mazowe, Mt. Darwin, Zvimba, Chivi, Insiza, Mberengwa (n = 14)	Morbidity is highest, highest transmitting districts. Highest priority requiring uninterrupted intensified PCT with annual geographic coverage of 100% per district. Complementary strategies urgently required. The goal is to control morbidity (reduce prevalence of heavy infection by any schistosome to <5%) in the first 5 years and prevent transmission.
≥5% but <10%	Buhera, Chimanimani, Makoni, Mutare, Mudzi, Seke, Guruve, Muzarabani, Chegutu, Kariba, Kadoma, Chiredzi, Gutu, Masvingo, Zaka, Gwanda, Chirumhanzu, Zvishavane (n = 18)	Morbidity is high. High transmitting districts requiring MDA regularly according to WHO strategies with geographic coverage of 75–100% per district. Complementary strategies are required. The goal is to control morbidity by reducing the prevalence of heavy infection by any schistosome species in the first 5 years to <5% and prevent transmission.
≥1% but <5%	Mutasa, Nyanga, Goromonzi, Marondera, Rushinga, Makonde, Karoyi, Bikita, Hwange, Lupane, Gokwe North, Glenview/Mufakose, Highfields/Glen Norah, Maribereign/Warren Park, Mabvuku/Tafara, Chitungwiza-Zengeza, Mbare/Hatfield, Khami (n = 17)	Morbidity is moderate though unjustifiable. Moderate transmitting districts. Regular MDA according to WHO guidelines based on prevalence. In addition, identification of transmission foci for intensified PCT is recommended. Complementary strategies are required. The goal is to eliminate schistosomiasis as a public health problem.
<1%	Chipinge, Binga, Beitbridge, Chitungwiza-Seke (n = 4)	Morbidity is low. Low transmitting districts. PCT to be implemented according to WHO guidelines. In addition, monitoring and surveillance of schistosomiasis transmitting foci for intensified PCT is recommended. Complementary strategies are required. The goal is to interrupt transmission.
0%	Bubi, Nkayi, Tsholotsho, Umguza, Bulilima, Matobo, Magwe, Umzingwane, Gokwe South, Reigate, Imbizo, Mzilikazi, Sizinda, North Central (n = 15)	Detailed surveillance should be done to identify any transmitting foci for intensified PCT. Complementary strategies are required. The goal is to interrupt schistosomiasis.

Source: Midzi et al. (2014)

All districts treated simultaneously: “High schisto” vs “low schisto”

→ Treatment guidelines based on heavy infection rates

Defining Comparison Groups

Prevalence category	Districts (IUs)	Comments and intervention strategies
≥10%	Murehwa, Shamva, Mwenezi, Shurugwi, Chikomba, Mutoko, UMP, Hwedza, Mazowe, Mt. Darwin, Zvimba, Chivi, Insiza, Mberengwa (n = 14)	Morbidity is highest, highest transmitting districts. Highest priority requiring uninterrupted intensified PCT with annual geographic coverage of 100% per district. Complementary strategies urgently required. The goal is to control morbidity (reduce prevalence of heavy infection by any schistosome to <5%) in the first 5 years and prevent transmission.
≥5% but <10%	Buhera, Chimanimani, Makoni, Mutare, Mudzi, Seke, Guruve, Muzarabani, Chegutu, Kariba, Kadoma, Chiredzi, Gutu, Masvingo, Zaka, Gwanda, Chirumhanzu, Zvishavane (n = 18)	Morbidity is high. High transmitting districts requiring MDA regularly according to WHO strategies with geographic coverage of 75–100% per district. Complementary strategies are required. The goal is to control morbidity by reducing the prevalence of heavy infection by any schistosome species in the first 5 years to <5% and prevent transmission.
≥1% but <5%	Mutasa, Nyanga, Goromonzi, Marondera, Rushinga, Makonde, Karoyi, Bikita, Hwange, Lupane, Gokwe North, Glenview/Mufakose, Highfields/Glen Norah, Maribereign/Warren Park, Mabvuku/Tafara, Chitungwiza-Zengeza, Mbare/Hatfield, Khami (n = 17)	Morbidity is moderate though unjustifiable. Moderate transmitting districts. Regular MDA according to WHO guidelines based on prevalence. In addition, identification of transmission foci for intensified PCT is recommended. Complementary strategies are required. The goal is to eliminate schistosomiasis as a public health problem.
<1%	Chipinge, Binga, Beitbridge, Chitungwiza-Seke (n = 4)	Morbidity is low. Low transmitting districts. PCT to be implemented according to WHO guidelines. In addition, monitoring and surveillance of schistosomiasis transmitting foci for intensified PCT is recommended. Complementary strategies are required. The goal is to interrupt transmission.
0%	Bubi, Nkayi, Tsholotsho, Umguza, Bulilima, Matobo, Magwe, Umzingwane, Gokwe South, Reigate, Imbizo, Mzilikazi, Sizinda, North Central (n = 15)	Detailed surveillance should be done to identify any transmitting foci for intensified PCT. Complementary strategies are required. The goal is to interrupt schistosomiasis.

Source: Midzi et al. (2014)

All districts treated simultaneously: “High schisto” vs “low schisto”

→ Treatment guidelines based on heavy infection rates

Defining Comparison Groups


Prevalence category	Districts (IUs)	Comments and intervention strategies
≥10%	Murehwa, Shamva, Mwenezi, Shurugwi, Chikomba, Mutoko, UMP, Hwedza, Mazowe, Mt. Darwin, Zvimba, Chivi, Insiza, Mberengwa (n = 14)	Morbidity is highest, highest transmitting districts. Highest priority requiring uninterrupted intensified PCT with annual geographic coverage of 100% per district. Complementary strategies urgently required. The goal is to control morbidity (reduce prevalence of heavy infection by any schistosome to <5%) in the first 5 years and prevent transmission.
≥5% but <10%	Buhera, Chimanimani, Makoni, Mutare, Mudzi, Seke, Guruve, Muzarabani, Chegutu, Kariba, Kadoma, Chiredzi, Gutu, Masvingo, Zaka, Gwanda, Chirumhanzu, Zvishavane (n = 18)	Morbidity is high. High transmitting districts requiring MDA regularly according to WHO strategies with geographic coverage of 75–100% per district. Complementary strategies are required. The goal is to control morbidity by reducing the prevalence of heavy infection by any schistosome species in the first 5 years to <5% and prevent transmission.
≥1% but <5%	Mutasa, Nyanga, Goromonzi, Marondera, Rushinga, Makonde, Karoyi, Bikita, Hwange, Lupane, Gokwe North, Glenview/Mufakose, Highfields/Glen Norah, Marlborough/Warren Park, Mabvuku/Tafara, Chitungwiza-Zengeza, Mbare/Hatfield, Khami (n = 17)	Morbidity is moderate though unjustifiable. Moderate transmitting districts. Regular MDA according to WHO guidelines based on prevalence. In addition, identification of transmission foci for intensified PCT is recommended. Complementary strategies are required. The goal is to eliminate schistosomiasis as a public health problem.
<1%	Chipinge, Binga, Beitbridge, Chitungwiza-Seke (n = 4)	Morbidity is low. Low transmitting districts. PCT to be implemented according to WHO guidelines. In addition, monitoring and surveillance of schistosomiasis transmitting foci for intensified PCT is recommended. Complementary strategies are required. The goal is to interrupt transmission.
0%	Bubi, Nkayi, Tsholotsho, Umguza, Bulilima, Matobo, Magwe, Umzingwane, Gokwe South, Reigate, Imbizo, Mzilikazi, Sizinda, North Central (n = 15)	Detailed surveillance should be done to identify any transmitting foci for intensified PCT. Complementary strategies are required. The goal is to interrupt schistosomiasis.

Source: Midzi et al. (2014)

All districts treated simultaneously: “High schisto” vs “low schisto”

→ “High” ≥ 5% heavy infection (N = 43), “low” < 5% (N = 28)


Empirical Strategy: Diff-in-Diff with DHS Data



Data: Demographic and Health Surveys (DHS)

- ▶ Include HIV blood tests from random subset of respondents

Empirical Strategy: Diff-in-Diff with DHS Data




Data: Demographic and Health Surveys (DHS)

- ▶ Include HIV blood tests from random subset of respondents

Comparison: {Pre, Post} \times {High-schisto, Low-schisto}

- ▶ Use de Chaisemartin and D'Haultfœuille (2020) estimator
- ▶ Controls: Age, age², survey cluster lat./lon. (quadratic) and altitude

Empirical Strategy: Diff-in-Diff with DHS Data



Data: Demographic and Health Surveys (DHS)

- ▶ Include HIV blood tests from random subset of respondents


Comparison: {Pre, Post} \times {High-schisto, Low-schisto}

- ▶ Use de Chaisemartin and D'Haultfœuille (2020) estimator
- ▶ Controls: Age, age², survey cluster lat./lon. (quadratic) and altitude

Age-specific focus: Ages 12-17 in 2012 \rightarrow **ages 15-20 in 2015**

- ▶ 17 is last age at which most boys and girls were in school
- ▶ Note: Results robust to restricting sample to ages 15-18 (migration concerns) and rural areas (more affected by schisto)


Main Result: Young Women's HIV



Notes: Estimates and bootstrapped standard errors are obtained using the procedure developed by de Chaisemartin and D'Haultfœuille (2020). The 4,309 observations are clustered by the 71 districts in the sample. Districts are in the treatment group if at least 5 percent of students had heavy schistosome infections in 2010. Regressions control for age, age squared, altitude, and a quadratic in latitude and longitude. Prior to deworming, 6.4 percent of females 15-20 in high-schistosomiasis districts had HIV.

Women 15-20: HIV ↓ 2.9 p.p. (45%)


Social Channel: Older Girls' School Attendance



Notes: Estimates and bootstrapped standard errors are obtained using the procedure developed by de Chaisemartin and D'Haultfœuille (2020). The 6,261 observations are clustered by the 71 districts in the sample. Districts are in the treatment group if at least 5 percent of students had heavy schistosome infections in 2010. Regressions control for age, age squared, altitude, and a quadratic in latitude and longitude. Prior to deworming, 66.9 percent of females 13-18 in high-schistosomiasis districts were attending school.

Girls 13-18: Attendance \uparrow 6.8 p.p. (10%)


Social Channel: Young Women's Partner Age Gap



Notes: Estimates and bootstrapped standard errors are obtained using the procedure developed by de Chaisemartin and D'Haultfœuille (2020). The 1,308 observations are clustered by the 70 districts in the sample. Districts are in the treatment group if at least 5 percent of students had heavy schistosome infections in 2010. Regressions control for age, age squared, altitude, and a quadratic in latitude and longitude. Prior to deworming, 26.5 percent of females 15-20 in relationships had partner age gaps of 9 or more years (fourth quartile), and 40.0 percent had partner age gaps between 5 and 8 years (third quartile).

Partnered women 15-20: ↓ 22.1 p.p. in 4th quartile of age gaps

Social Channel: Young Women's Partner Age Gap



Notes: Estimates and bootstrapped standard errors are obtained using the procedure developed by de Chaisemartin and D'Haultfœuille (2020). The 1,308 observations are clustered by the 70 districts in the sample. Districts are in the treatment group if at least 5 percent of students had heavy schistosome infections in 2010. Regressions control for age, age squared, altitude, and a quadratic in latitude and longitude. Prior to deworming, 26.5 percent of females 15-20 in relationships had partner age gaps of 9 or more years (fourth quartile), and 40.0 percent had partner age gaps between 5 and 8 years (third quartile).

Partnered women 15-20: ↑ 19.4 p.p. in 3rd quartile of age gaps

What About Clinical Channel? And Effect Size?



Novel: Linking childhood health to HIV via learning / attendance

But: Data don't allow me to test for “clinical channel”

What About Clinical Channel? And Effect Size?



Novel: Linking childhood health to HIV via learning / attendance

But: Data don't allow me to test for “clinical channel”

→ **Epidemiological model (in prog.):** Sex-by-age compartments

$$\blacktriangleright \text{Force of transmission: } \mathbb{P}[\text{Having sex}] \times \underbrace{\mathbb{P}[\text{Partner HIV}]}_{\text{Social channel}} \times \underbrace{\mathbb{P}[\text{Transmission}]}_{\text{Clinical channel}}$$

What About Clinical Channel? And Effect Size?



Novel: Linking childhood health to HIV via learning / attendance

But: Data don't allow me to test for “clinical channel”

→ **Epidemiological model (in prog.):** Sex-by-age compartments

► Force of transmission: $\mathbb{P}[\text{Having sex}] \times \underbrace{\mathbb{P}[\text{Partner HIV}]}_{\text{Social channel}} \times \underbrace{\mathbb{P}[\text{Transmission}]}_{\text{Clinical channel}}$

Prelim. results: HIV ↓ 1.5-2.2 p.p., up to 50% via social channel

Conclusion

- ① **Cost-benefit analysis:** Cheap to improve childhood health → cheap to avert (very expensive) HIV infections for very high-risk group

Conclusion

- ① **Cost-benefit analysis:** Cheap to improve childhood health → cheap to avert (very expensive) HIV infections for very high-risk group
- ▶ Miguel and Kremer (2004) assumptions: \$/DALY saved without HIV effects = \$106
 - ▶ 2.9 p.p. of young women only avoid HIV through 2015, 9-year lag between infection and ART: \$/DALY saved with HIV effects = \$55

Conclusion

- ① **Cost-benefit analysis:** Cheap to improve childhood health → cheap to avert (very expensive) HIV infections for very high-risk group
- ▶ Miguel and Kremer (2004) assumptions: \$/DALY saved without HIV effects = \$106
 - ▶ 2.9 p.p. of young women only avoid HIV through 2015, 9-year lag between infection and ART: \$/DALY saved with HIV effects = \$55
 - ▶ Public finances: **Breakeven** if those add'l 2.9% of young women in high-schisto districts remain HIV negative through 2023

Conclusion

- ➊ **Cost-benefit analysis:** Cheap to improve childhood health → cheap to avert (very expensive) HIV infections for very high-risk group
 - ▶ Miguel and Kremer (2004) assumptions: \$/DALY saved without HIV effects = \$106
 - ▶ 2.9 p.p. of young women only avoid HIV through 2015, 9-year lag between infection and ART: \$/DALY saved with HIV effects = \$55
 - ▶ Public finances: **Breakeven** if those add'l 2.9% of young women in high-schisto districts remain HIV negative through 2023
- ➋ **Childhood health:** Improving it for girls lowers their chances of contracting HIV as young women, due in large part to “social channel” (health → schooling → marriage → HIV)

Conclusion

- ➊ **Cost-benefit analysis:** Cheap to improve childhood health → cheap to avert (very expensive) HIV infections for very high-risk group
 - ▶ Miguel and Kremer (2004) assumptions: \$/DALY saved without HIV effects = \$106
 - ▶ 2.9 p.p. of young women only avoid HIV through 2015, 9-year lag between infection and ART: \$/DALY saved with HIV effects = \$55
 - ▶ Public finances: Breakeven if those add'l 2.9% of young women in high-schisto districts remain HIV negative through 2023
- ➋ **Childhood health:** Improving it for girls lowers their chances of contracting HIV as young women, due in large part to “social channel” (health → schooling → marriage → HIV)
- ➌ **Marriage markets:** Helps us understand role of childhood health (as part of human capital), especially in non-Western context